

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	59,474	394.7	391.5-398.0	28,807	462.8	457.2-468.5	30,667	360.3	356.2-364.4	32,504	217.1	214.7-219.5	16,784	276.3	271.9-280.7	15,720	181.5	178.7-184.5
1977	59,448	394.5	391.3-397.8	28,896	464.7	459.1-470.4	30,552	358.2	354.1-362.3	32,787	218.5	216.1-220.9	16,936	278.3	273.9-282.8	15,851	182.3	179.4-185.2
1978	59,842	397.3	394.1-400.6	29,061	469.0	463.3-474.6	30,781	360.5	356.4-364.7	32,417	215.2	212.8-217.6	16,721	275.7	271.3-280.1	15,696	179.3	176.5-182.2
1979	60,989	404.0	400.8-407.3	29,594	477.2	471.6-482.9	31,395	366.7	362.6-370.9	32,331	214.4	212.0-216.8	16,522	272.3	268.0-276.7	15,809	180.1	177.3-183.0
1980	62,855	415.0	411.7-418.3	31,044	498.5	492.7-504.3	31,811	370.8	366.6-375.0	33,166	218.7	216.3-221.1	17,047	280.7	276.3-285.1	16,119	182.0	179.2-184.9
1981	63,006	416.2	412.9-419.5	30,686	494.8	489.1-500.6	32,320	376.0	371.8-380.3	32,665	215.2	212.8-217.5	16,659	274.4	270.1-278.8	16,006	180.6	177.8-183.5
1982	63,974	421.2	417.9-424.5	31,105	500.5	494.7-506.3	32,869	381.2	376.9-385.4	32,608	214.4	212.0-216.8	16,614	274.3	270.0-278.7	15,994	179.9	177.0-182.7
1983	65,236	428.4	425.1-431.8	31,771	508.4	502.6-514.2	33,465	387.9	383.7-392.2	32,659	213.7	211.4-216.1	16,571	273.3	269.0-277.7	16,088	179.6	176.8-182.5
1984	65,751	431.3	428.0-434.7	31,786	507.3	501.5-513.1	33,965	393.0	388.7-397.3	32,972	215.3	212.9-217.6	16,618	272.7	268.4-277.1	16,354	182.4	179.5-185.3
1985	67,375	441.9	438.5-445.3	32,894	522.4	516.6-528.3	34,481	399.5	395.2-403.9	32,944	214.8	212.4-217.1	16,682	272.8	268.5-277.2	16,262	181.3	178.5-184.2
1986	67,375	441.8	438.5-445.3	32,735	521.1	515.3-527.0	34,640	401.3	397.0-405.7	33,001	214.8	212.4-217.1	16,603	271.2	266.9-275.5	16,398	182.3	179.4-185.1
1987	67,742	443.1	439.7-446.5	32,705	518.8	513.0-524.7	35,037	404.5	400.2-408.9	32,849	213.6	211.2-215.9	16,550	270.6	266.3-274.9	16,299	180.3	177.5-183.2
1988	67,225	439.8	436.5-443.2	32,239	512.2	506.4-518.0	34,986	404.5	400.1-408.8	33,152	215.2	212.9-217.6	16,728	273.3	269.0-277.6	16,424	181.9	179.1-184.8
1989	67,346	441.4	438.1-444.8	32,752	518.5	512.7-524.3	34,594	401.5	397.2-405.9	32,321	210.2	207.9-212.5	16,163	263.8	259.6-268.0	16,158	179.3	176.5-182.2
1990	70,612	461.6	458.2-465.1	34,500	541.9	536.1-547.8	36,112	419.3	414.9-423.8	32,205	208.7	206.4-211.0	16,046	261.7	257.5-265.9	16,159	178.3	175.5-181.1
1991	72,106	471.0	467.5-474.5	35,762	560.3	554.3-566.3	36,344	422.6	418.1-427.1	32,411	209.8	207.5-212.2	16,200	263.3	259.1-267.5	16,211	179.0	176.2-181.9
1992	74,237	482.4	478.9-486.0	37,801	587.7	581.6-593.8	36,436	422.5	418.1-427.0	32,425	208.8	206.5-211.1	16,061	260.5	256.3-264.6	16,364	179.8	177.0-182.6
1993	73,414	476.1	472.6-479.6	37,036	570.6	564.7-576.5	36,378	420.6	416.2-425.1	32,092	205.8	203.5-208.1	15,924	256.6	252.5-260.7	16,168	176.7	174.0-179.5
1994	75,618	490.2	486.7-493.8	37,667	578.8	572.9-584.8	37,951	437.8	433.3-442.3	33,353	213.5	211.2-215.8	16,718	268.1	264.0-272.3	16,635	181.5	178.7-184.4
1995	74,605	483.6	480.1-487.1	36,512	557.8	552.0-563.6	38,093	440.7	436.3-445.3	32,897	210.0	207.8-212.3	16,245	259.7	255.7-263.8	16,652	181.4	178.6-184.2
1996	76,296	494.7	491.2-498.3	37,679	574.6	568.7-580.5	38,617	447.6	443.0-452.1	32,464	207.0	204.8-209.3	15,956	253.2	249.2-257.2	16,508	180.0	177.2-182.8
1997	77,509	500.3	496.8-503.8	38,305	580.0	574.2-585.9	39,204	451.6	447.1-456.2	31,618	200.5	198.3-202.8	15,645	246.2	242.3-250.2	15,973	173.2	170.5-175.9
1998	79,218	508.1	504.6-511.7	39,032	583.6	577.7-589.4	40,186	462.2	457.7-466.8	31,110	195.8	193.6-198.0	15,346	236.9	233.2-240.8	15,764	169.8	167.1-172.5
1999	79,809	507.5	503.9-511.0	39,475	585.9	580.0-591.7	40,334	459.2	454.7-463.8	31,490	196.0	193.8-198.2	15,562	238.6	234.8-242.4	15,928	170.1	167.4-172.8
2000	81,918	516.5	512.9-520.0	41,007	603.7	597.8-609.6	40,911	461.9	457.4-466.5	31,361	192.9	190.8-195.1	15,390	234.0	230.3-237.8	15,971	168.1	165.4-170.8
2001	81,089	508.8	505.3-512.3	40,515	590.8	585.0-596.6	40,574	457.4	452.9-461.9	30,891	189.0	186.9-191.1	15,242	229.6	225.9-233.3	15,649	164.2	161.6-166.9
2002	81,809	509.4	505.9-512.9	41,582	599.2	593.5-605.0	40,227	450.6	446.1-455.1	30,614	185.7	183.6-187.8	15,067	224.2	220.6-227.8	15,547	162.4	159.8-165.0
2003	81,307	503.2	499.7-506.7	40,793	582.7	577.1-588.5	40,514	452.0	447.6-456.5	30,009	180.6	178.6-182.7	14,562	214.1	210.6-217.6	15,447	160.1	157.6-162.7
2004	81,737	503.6	500.1-507.1	40,948	579.4	573.8-585.1	40,789	455.0	450.5-459.5	29,720	177.9	175.9-180.0	14,581	211.9	208.5-215.4	15,139	156.6	154.1-159.2
2005	82,699	508.2	504.7-511.7	41,566	583.8	578.2-589.5	41,133	458.3	453.8-462.9	29,189	174.0	172.0-176.1	14,385	207.8	204.4-211.2	14,804	152.6	150.1-155.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	85,063	522.3	518.7-525.8	43,362	605.1	599.4-610.9	41,701	466.4	461.9-471.0	29,189	173.8	171.8-175.8	14,350	205.6	202.2-209.0	14,839	153.3	150.8-155.8
2007	85,901	522.9	519.4-526.5	43,962	604.5	598.8-610.2	41,939	468.2	463.7-472.8	28,977	170.7	168.8-172.7	14,431	203.3	200.0-206.7	14,546	149.6	147.1-152.1
2008	87,104	525.6	522.1-529.1	44,072	598.2	592.5-603.8	43,032	478.0	473.4-482.6	29,109	169.7	167.7-171.7	14,576	202.8	199.5-206.1	14,533	147.9	145.4-150.4
2009	87,159	520.7	517.2-524.2	44,002	588.1	582.5-593.7	43,157	475.5	470.9-480.2	28,555	165.1	163.1-167.0	14,353	197.3	194.1-200.6	14,202	144.0	141.5-146.4
2010	84,854	502.6	499.2-506.1	42,673	562.3	556.9-567.7	42,181	463.2	458.7-467.8	28,761	164.1	162.1-166.0	14,212	192.0	188.9-195.3	14,549	145.6	143.2-148.1
2011	87,016	509.7	506.2-513.1	43,892	569.6	564.2-575.0	43,124	469.6	465.0-474.2	28,375	159.8	157.9-161.7	14,097	187.4	184.3-190.5	14,278	141.0	138.6-143.4
2012	84,947	492.0	488.6-495.3	42,378	541.2	536.0-546.5	42,569	461.3	456.7-465.8	28,698	160.2	158.3-162.1	14,417	189.2	186.1-192.4	14,281	140.5	138.1-142.9
2013	87,229	500.4	497.0-503.8	43,090	542.6	537.4-547.9	44,139	475.6	471.0-480.3	28,209	154.9	153.1-156.8	14,208	182.3	179.3-185.4	14,001	136.0	133.7-138.4
2014	86,843	491.9	488.6-495.3	42,771	529.2	524.1-534.4	44,072	470.6	466.1-475.2	27,598	149.5	147.7-151.3	14,009	176.9	174.0-179.9	13,589	130.6	128.3-132.9
2015	87,662	492.6	489.2-495.9	43,222	527.6	522.5-532.7	44,440	473.4	468.9-478.1	27,530	147.5	145.8-149.3	13,828	172.4	169.5-175.4	13,702	130.3	128.1-132.6
2016	88,119	490.3	487.0-493.7	44,148	531.2	526.1-536.3	43,971	466.1	461.6-470.7	27,605	146.6	144.8-148.3	13,892	170.6	167.7-173.5	13,713	130.2	128.0-132.5
2017	89,488	493.6	490.3-497.0	44,774	531.6	526.6-536.7	44,714	471.4	466.8-476.0	27,177	142.1	140.4-143.8	13,713	165.5	162.7-168.4	13,464	125.9	123.7-128.1
2018	88,092	481.2	477.9-484.5	44,326	518.5	513.6-523.5	43,766	457.8	453.3-462.3	26,352	136.3	134.6-138.0	13,231	157.3	154.6-160.0	13,121	121.9	119.8-124.1
2019	90,399	489.9	486.6-493.2	45,709	527.6	522.7-532.6	44,690	466.2	461.6-470.7	26,313	134.7	133.0-136.4	13,275	155.7	153.0-158.4	13,038	119.6	117.5-121.8
2020	79,886	430.9	427.8-434.0	40,367	463.4	458.7-468.0	39,519	410.8	406.5-415.1	25,283	128.9	127.3-130.5	12,736	148.8	146.1-151.4	12,547	114.9	112.9-117.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,630	10.6	10.0-11.1	1,107	16.8	15.8-17.9	523	6.0	5.4-6.5	708	4.6	4.2-4.9	507	7.7	7.0-8.5	201	2.2	1.9-2.6
1977	1,626	10.5	10.0-11.0	1,075	16.3	15.3-17.4	551	6.2	5.7-6.8	698	4.5	4.2-4.9	478	7.4	6.8-8.2	220	2.4	2.1-2.8
1978	1,614	10.4	9.9-11.0	1,062	16.1	15.1-17.2	552	6.3	5.8-6.9	693	4.5	4.2-4.8	473	7.3	6.6-8.0	220	2.5	2.2-2.8
1979	1,657	10.6	10.1-11.2	1,084	16.3	15.3-17.3	573	6.5	6.0-7.1	646	4.2	3.8-4.5	420	6.3	5.7-7.0	226	2.6	2.2-2.9
1980	1,617	10.5	10.0-11.0	1,049	16.0	15.0-17.0	568	6.5	6.0-7.1	645	4.1	3.8-4.5	424	6.5	5.9-7.2	221	2.4	2.1-2.8
1981	1,690	10.9	10.4-11.5	1,164	17.6	16.5-18.7	526	6.0	5.5-6.6	635	4.1	3.7-4.4	427	6.5	5.9-7.2	208	2.3	2.0-2.6
1982	1,757	11.5	11.0-12.1	1,180	18.2	17.1-19.3	577	6.7	6.1-7.3	627	4.1	3.7-4.4	437	6.8	6.2-7.5	190	2.1	1.8-2.5
1983	1,670	10.8	10.3-11.4	1,115	16.9	15.9-17.9	555	6.3	5.8-6.9	603	3.8	3.5-4.2	415	6.3	5.7-6.9	188	2.1	1.8-2.4
1984	1,677	11.0	10.4-11.5	1,129	17.4	16.3-18.5	548	6.3	5.7-6.8	596	3.9	3.5-4.2	402	6.3	5.6-6.9	194	2.2	1.9-2.5
1985	1,706	11.2	10.6-11.7	1,116	17.2	16.1-18.2	590	6.7	6.2-7.3	575	3.7	3.4-4.0	379	5.9	5.3-6.5	196	2.2	1.9-2.5
1986	1,612	10.6	10.1-11.2	1,069	16.6	15.6-17.7	543	6.2	5.6-6.7	591	3.8	3.5-4.1	390	6.0	5.4-6.6	201	2.2	1.9-2.6
1987	1,600	10.6	10.1-11.1	1,028	15.9	14.9-16.9	572	6.6	6.0-7.1	546	3.6	3.3-3.9	353	5.6	5.0-6.2	193	2.1	1.8-2.4
1988	1,552	10.2	9.7-10.8	988	15.1	14.2-16.1	564	6.5	5.9-7.0	591	3.9	3.6-4.2	382	6.2	5.6-6.9	209	2.3	2.0-2.7
1989	1,523	10.1	9.6-10.6	1,001	15.4	14.4-16.4	522	6.1	5.6-6.6	490	3.2	2.9-3.5	304	4.8	4.3-5.4	186	2.1	1.8-2.4
1990	1,544	10.3	9.8-10.8	993	15.3	14.4-16.4	551	6.5	5.9-7.0	494	3.2	2.9-3.5	312	4.9	4.3-5.5	182	2.0	1.7-2.3
1991	1,574	10.6	10.1-11.1	1,004	15.6	14.6-16.6	570	6.6	6.1-7.2	474	3.1	2.8-3.4	300	4.8	4.2-5.4	174	1.9	1.6-2.2
1992	1,605	10.7	10.2-11.2	1,044	16.0	15.0-17.0	561	6.6	6.0-7.1	444	2.9	2.6-3.1	272	4.2	3.7-4.7	172	1.8	1.6-2.1
1993	1,560	10.4	9.9-10.9	1,029	15.9	15.0-17.0	531	6.1	5.6-6.7	436	2.8	2.6-3.1	285	4.4	3.9-4.9	151	1.7	1.4-2.0
1994	1,515	10.0	9.5-10.6	995	15.2	14.2-16.1	520	6.1	5.6-6.6	438	2.9	2.6-3.1	307	4.7	4.2-5.3	131	1.5	1.2-1.7
1995	1,384	9.2	8.7-9.7	866	13.1	12.3-14.0	518	5.9	5.4-6.5	445	2.9	2.6-3.2	294	4.6	4.1-5.2	151	1.7	1.4-2.0
1996	1,537	10.1	9.6-10.7	1,007	15.0	14.1-16.0	530	6.2	5.6-6.7	413	2.7	2.5-3.0	257	3.9	3.5-4.5	156	1.8	1.5-2.1
1997	1,542	10.1	9.6-10.6	992	14.8	13.9-15.7	550	6.3	5.8-6.9	406	2.6	2.4-2.9	274	4.1	3.7-4.7	132	1.4	1.2-1.7
1998	1,532	10.0	9.5-10.5	961	14.2	13.3-15.1	571	6.6	6.0-7.2	418	2.6	2.4-2.9	265	4.0	3.5-4.5	153	1.6	1.4-1.9
1999	1,551	10.0	9.5-10.5	1,061	15.4	14.5-16.4	490	5.5	5.1-6.1	375	2.4	2.1-2.6	251	3.7	3.3-4.2	124	1.3	1.1-1.6
2000	1,461	9.3	8.8-9.8	957	13.9	13.0-14.8	504	5.7	5.2-6.2	342	2.1	1.9-2.4	216	3.2	2.8-3.7	126	1.3	1.1-1.6
2001	1,542	9.8	9.3-10.3	989	14.0	13.1-14.9	553	6.1	5.6-6.7	383	2.4	2.1-2.6	245	3.6	3.1-4.0	138	1.4	1.1-1.6
2002	1,584	10.0	9.5-10.5	1,031	14.5	13.6-15.4	553	6.1	5.6-6.7	359	2.2	2.0-2.4	228	3.3	2.9-3.7	131	1.3	1.1-1.6
2003	1,577	9.8	9.3-10.3	1,038	14.4	13.6-15.4	539	6.0	5.5-6.5	341	2.1	1.9-2.3	214	3.0	2.6-3.5	127	1.3	1.1-1.6
2004	1,636	10.1	9.6-10.6	1,089	15.0	14.1-15.9	547	6.0	5.5-6.6	405	2.4	2.2-2.7	260	3.6	3.2-4.1	145	1.5	1.3-1.8
2005	1,568	9.6	9.2-10.1	1,029	13.9	13.1-14.8	539	6.1	5.6-6.6	344	2.1	1.8-2.3	225	3.1	2.7-3.5	119	1.2	1.0-1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,700	10.4	9.9-10.9	1,167	15.7	14.8-16.6	533	5.9	5.4-6.5	342	2.1	1.8-2.3	227	3.1	2.7-3.6	115	1.2	1.0-1.4
2007	1,804	10.9	10.4-11.4	1,232	16.4	15.5-17.3	572	6.4	5.8-6.9	342	2.0	1.8-2.2	229	3.1	2.7-3.5	113	1.2	1.0-1.4
2008	1,776	10.7	10.2-11.2	1,185	15.6	14.7-16.5	591	6.5	6.0-7.1	345	2.0	1.8-2.2	250	3.3	2.9-3.7	95	1.0	0.8-1.2
2009	1,858	11.0	10.5-11.5	1,287	16.6	15.7-17.5	571	6.2	5.7-6.7	364	2.1	1.9-2.3	247	3.3	2.9-3.7	117	1.1	0.9-1.4
2010	1,878	10.9	10.4-11.5	1,254	15.9	15.0-16.8	624	6.8	6.2-7.4	381	2.2	2.0-2.4	252	3.3	2.9-3.7	129	1.3	1.1-1.5
2011	1,887	10.9	10.4-11.4	1,313	16.4	15.5-17.3	574	6.1	5.6-6.7	351	2.0	1.8-2.2	232	3.0	2.6-3.4	119	1.2	1.0-1.4
2012	1,922	11.0	10.5-11.5	1,309	16.2	15.3-17.1	613	6.4	5.9-7.0	391	2.2	2.0-2.4	271	3.4	3.0-3.9	120	1.1	0.9-1.4
2013	1,982	11.2	10.7-11.8	1,393	17.1	16.2-18.1	589	6.1	5.6-6.6	371	2.0	1.8-2.2	257	3.1	2.7-3.5	114	1.1	0.9-1.3
2014	2,063	11.5	11.0-12.0	1,382	16.6	15.7-17.5	681	7.1	6.5-7.6	330	1.8	1.6-2.0	231	2.8	2.5-3.2	99	0.9	0.8-1.1
2015	2,149	11.9	11.4-12.5	1,527	18.2	17.3-19.2	622	6.5	6.0-7.1	397	2.1	1.9-2.4	273	3.3	2.9-3.7	124	1.2	1.0-1.4
2016	2,261	12.3	11.8-12.9	1,569	18.4	17.5-19.4	692	7.1	6.6-7.7	424	2.2	2.0-2.5	300	3.5	3.1-4.0	124	1.1	0.9-1.4
2017	2,201	11.9	11.4-12.5	1,518	17.8	16.9-18.7	683	6.9	6.4-7.4	419	2.2	2.0-2.4	290	3.4	3.0-3.8	129	1.2	1.0-1.5
2018	2,175	11.7	11.2-12.2	1,501	17.3	16.4-18.2	674	6.8	6.2-7.3	414	2.1	1.9-2.3	276	3.2	2.8-3.6	138	1.3	1.1-1.5
2019	2,323	12.5	12.0-13.0	1,598	18.2	17.3-19.2	725	7.4	6.8-8.0	421	2.1	1.9-2.4	296	3.3	3.0-3.8	125	1.1	0.9-1.4
2020	2,115	11.1	10.6-11.6	1,481	16.6	15.7-17.5	634	6.3	5.8-6.9	425	2.2	2.0-2.4	290	3.2	2.9-3.7	135	1.3	1.1-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	631	4.1	3.8-4.4	455	7.0	6.3-7.7	176	2.0	1.7-2.3	584	3.8	3.5-4.1	423	6.6	5.9-7.3	161	1.8	1.6-2.2
1977	622	4.1	3.8-4.4	422	6.6	6.0-7.3	200	2.3	2.0-2.6	568	3.7	3.4-4.0	383	6.1	5.4-6.7	185	2.1	1.8-2.4
1978	618	4.0	3.7-4.4	421	6.6	6.0-7.3	197	2.2	1.9-2.5	627	4.1	3.8-4.4	434	6.9	6.3-7.6	193	2.1	1.9-2.5
1979	636	4.1	3.8-4.4	442	6.8	6.2-7.5	194	2.1	1.8-2.5	560	3.6	3.3-4.0	379	5.9	5.3-6.5	181	2.0	1.7-2.4
1980	668	4.3	4.0-4.6	450	7.1	6.4-7.8	218	2.4	2.0-2.7	563	3.6	3.3-3.9	382	6.0	5.4-6.7	181	2.0	1.7-2.3
1981	627	4.0	3.7-4.4	419	6.5	5.9-7.2	208	2.3	2.0-2.6	602	3.9	3.6-4.2	429	6.8	6.1-7.5	173	1.9	1.6-2.2
1982	630	4.0	3.7-4.3	406	6.3	5.7-7.0	224	2.4	2.1-2.7	558	3.6	3.3-3.9	379	6.0	5.4-6.7	179	1.9	1.7-2.2
1983	639	4.1	3.8-4.4	447	6.9	6.3-7.6	192	2.1	1.8-2.4	575	3.7	3.4-4.1	416	6.7	6.1-7.4	159	1.7	1.5-2.0
1984	638	4.1	3.7-4.4	436	6.8	6.2-7.5	202	2.2	1.9-2.5	584	3.7	3.4-4.0	394	6.1	5.5-6.8	190	2.0	1.7-2.3
1985	612	4.0	3.7-4.3	440	6.9	6.2-7.6	172	1.9	1.6-2.2	551	3.5	3.2-3.8	405	6.3	5.6-6.9	146	1.6	1.3-1.8
1986	668	4.3	4.0-4.7	449	7.0	6.4-7.7	219	2.4	2.1-2.8	564	3.6	3.3-4.0	413	6.6	5.9-7.2	151	1.6	1.4-1.9
1987	617	4.0	3.7-4.3	408	6.3	5.7-6.9	209	2.3	2.0-2.6	573	3.7	3.4-4.0	396	6.2	5.6-6.9	177	1.9	1.6-2.2
1988	604	3.9	3.6-4.2	414	6.4	5.8-7.0	190	2.0	1.8-2.4	623	4.0	3.7-4.3	455	7.1	6.5-7.8	168	1.8	1.5-2.1
1989	656	4.2	3.9-4.6	457	7.0	6.4-7.7	199	2.1	1.9-2.5	594	3.8	3.5-4.2	422	6.6	5.9-7.2	172	1.8	1.6-2.1
1990	696	4.5	4.2-4.8	461	7.1	6.4-7.7	235	2.5	2.2-2.9	587	3.7	3.4-4.1	411	6.3	5.7-7.0	176	1.9	1.6-2.2
1991	671	4.3	4.0-4.7	454	6.9	6.3-7.6	217	2.4	2.0-2.7	603	3.8	3.5-4.2	424	6.5	5.9-7.2	179	1.9	1.6-2.2
1992	739	4.8	4.4-5.1	494	7.5	6.8-8.2	245	2.7	2.3-3.0	636	4.1	3.8-4.4	461	7.0	6.4-7.7	175	1.9	1.6-2.2
1993	699	4.5	4.2-4.8	496	7.5	6.9-8.2	203	2.2	1.9-2.5	618	4.0	3.6-4.3	455	7.0	6.3-7.7	163	1.7	1.5-2.0
1994	753	4.8	4.5-5.2	538	8.2	7.5-8.9	215	2.3	2.0-2.6	647	4.1	3.8-4.5	460	7.0	6.4-7.7	187	2.0	1.7-2.3
1995	715	4.6	4.3-5.0	484	7.4	6.8-8.1	231	2.5	2.2-2.8	683	4.4	4.1-4.8	505	7.8	7.1-8.5	178	1.9	1.6-2.2
1996	752	4.8	4.5-5.2	520	7.9	7.2-8.6	232	2.5	2.2-2.8	669	4.3	4.0-4.7	466	7.1	6.5-7.8	203	2.1	1.9-2.5
1997	701	4.5	4.2-4.8	481	7.3	6.7-8.0	220	2.4	2.1-2.7	651	4.2	3.8-4.5	488	7.4	6.8-8.1	163	1.7	1.4-2.0
1998	773	4.9	4.6-5.3	570	8.5	7.8-9.3	203	2.2	1.9-2.5	679	4.3	4.0-4.7	502	7.5	6.9-8.2	177	1.9	1.6-2.2
1999	744	4.7	4.4-5.0	547	8.1	7.4-8.8	197	2.0	1.8-2.4	681	4.3	4.0-4.6	506	7.5	6.9-8.2	175	1.8	1.6-2.1
2000	786	4.9	4.6-5.3	563	8.3	7.6-9.0	223	2.3	2.0-2.6	698	4.3	4.0-4.7	495	7.3	6.7-8.0	203	2.1	1.8-2.4
2001	746	4.6	4.3-5.0	547	7.9	7.3-8.6	199	2.1	1.8-2.4	701	4.3	4.0-4.7	538	7.8	7.2-8.5	163	1.7	1.4-1.9
2002	800	4.9	4.6-5.3	596	8.5	7.8-9.2	204	2.1	1.8-2.4	679	4.1	3.8-4.5	507	7.3	6.7-8.0	172	1.7	1.5-2.0
2003	834	5.1	4.8-5.5	614	8.7	8.0-9.4	220	2.2	1.9-2.6	736	4.5	4.1-4.8	535	7.7	7.0-8.3	201	2.0	1.7-2.3
2004	842	5.1	4.8-5.5	624	8.8	8.1-9.5	218	2.2	1.9-2.5	746	4.5	4.2-4.9	567	8.0	7.3-8.7	179	1.8	1.5-2.1
2005	899	5.5	5.1-5.8	669	9.4	8.7-10.1	230	2.4	2.1-2.7	806	4.8	4.5-5.2	623	8.8	8.1-9.5	183	1.8	1.6-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	934	5.7	5.3-6.0	725	10.1	9.3-10.8	209	2.2	1.9-2.5	731	4.4	4.1-4.8	546	7.6	7.0-8.3	185	1.9	1.7-2.2
2007	874	5.2	4.9-5.6	655	8.9	8.2-9.6	219	2.3	2.0-2.6	775	4.6	4.3-4.9	578	8.0	7.3-8.7	197	2.0	1.7-2.3
2008	947	5.6	5.3-6.0	718	9.7	9.0-10.4	229	2.3	2.0-2.7	749	4.4	4.1-4.7	586	7.9	7.2-8.5	163	1.7	1.4-1.9
2009	932	5.4	5.1-5.8	721	9.5	8.8-10.2	211	2.1	1.8-2.4	755	4.4	4.1-4.7	590	7.9	7.2-8.5	165	1.6	1.4-1.9
2010	861	4.9	4.6-5.3	644	8.4	7.8-9.1	217	2.1	1.9-2.4	746	4.3	4.0-4.6	585	7.7	7.1-8.4	161	1.6	1.3-1.8
2011	837	4.8	4.4-5.1	629	8.0	7.4-8.7	208	2.1	1.8-2.4	775	4.4	4.1-4.7	585	7.6	7.0-8.2	190	1.8	1.6-2.1
2012	805	4.5	4.2-4.8	632	7.9	7.3-8.6	173	1.7	1.4-2.0	711	4.0	3.7-4.3	550	7.1	6.5-7.7	161	1.5	1.3-1.8
2013	891	4.9	4.6-5.3	684	8.4	7.8-9.1	207	2.0	1.7-2.3	737	4.0	3.7-4.3	570	7.1	6.5-7.7	167	1.6	1.4-1.9
2014	896	4.8	4.5-5.2	680	8.1	7.5-8.8	216	2.1	1.8-2.4	752	4.1	3.9-4.5	603	7.5	6.9-8.2	149	1.4	1.2-1.7
2015	935	5.0	4.7-5.4	703	8.4	7.7-9.0	232	2.3	2.0-2.6	707	3.8	3.5-4.1	530	6.4	5.8-7.0	177	1.6	1.4-1.9
2016	927	4.9	4.6-5.3	710	8.4	7.7-9.0	217	2.1	1.8-2.4	747	3.9	3.7-4.2	588	6.9	6.4-7.5	159	1.5	1.3-1.8
2017	863	4.6	4.3-4.9	664	7.7	7.1-8.3	199	1.9	1.7-2.2	720	3.7	3.5-4.0	579	6.7	6.1-7.2	141	1.3	1.1-1.6
2018	896	4.7	4.4-5.0	703	8.1	7.5-8.7	193	1.8	1.5-2.0	685	3.5	3.3-3.8	529	6.1	5.6-6.6	156	1.4	1.2-1.7
2019	935	4.8	4.5-5.1	717	8.1	7.5-8.8	218	1.9	1.7-2.2	699	3.5	3.2-3.8	555	6.2	5.7-6.8	144	1.3	1.1-1.5
2020	831	4.2	3.9-4.5	643	7.3	6.7-7.9	188	1.7	1.5-2.0	676	3.5	3.2-3.7	539	6.2	5.7-6.8	137	1.2	1.0-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,814	12.3	11.8-12.9	1,038	17.2	16.2-18.4	776	9.0	8.3-9.6	1,431	9.8	9.3-10.3	793	13.5	12.5-14.5	638	7.4	6.8-8.0
1977	1,801	12.3	11.8-12.9	1,043	17.7	16.6-18.8	758	8.8	8.2-9.4	1,397	9.6	9.1-10.2	797	13.8	12.8-14.9	600	6.9	6.4-7.5
1978	1,765	12.0	11.4-12.5	984	16.7	15.6-17.8	781	8.9	8.3-9.5	1,300	8.9	8.5-9.5	721	12.5	11.5-13.5	579	6.6	6.1-7.2
1979	1,803	12.0	11.5-12.6	1,021	16.9	15.8-18.0	782	8.7	8.1-9.4	1,325	8.9	8.4-9.4	746	12.7	11.7-13.6	579	6.4	5.9-7.0
1980	1,770	11.9	11.3-12.4	1,045	17.4	16.3-18.6	725	8.2	7.6-8.8	1,322	8.9	8.4-9.4	751	12.8	11.8-13.8	571	6.4	5.9-7.0
1981	1,807	12.0	11.5-12.6	1,051	17.5	16.4-18.6	756	8.4	7.8-9.1	1,284	8.6	8.1-9.1	751	12.8	11.8-13.7	533	5.9	5.4-6.4
1982	1,729	11.4	10.8-11.9	995	16.5	15.4-17.6	734	8.0	7.4-8.6	1,197	8.0	7.5-8.4	704	12.0	11.1-12.9	493	5.4	4.9-5.9
1983	1,787	11.8	11.3-12.4	1,013	16.8	15.8-17.9	774	8.4	7.9-9.1	1,198	8.0	7.5-8.4	690	11.8	10.9-12.7	508	5.5	5.1-6.0
1984	1,702	11.1	10.6-11.7	985	15.9	14.9-17.0	717	7.8	7.2-8.4	1,217	8.0	7.5-8.4	677	11.0	10.2-11.9	540	5.8	5.3-6.3
1985	1,690	11.1	10.5-11.6	1,017	16.7	15.6-17.8	673	7.3	6.8-7.9	1,148	7.6	7.1-8.0	655	10.9	10.1-11.8	493	5.4	4.9-5.9
1986	1,627	10.6	10.1-11.2	964	15.8	14.8-16.9	663	7.2	6.6-7.7	1,106	7.2	6.8-7.7	645	10.7	9.9-11.6	461	4.9	4.5-5.4
1987	1,660	10.8	10.2-11.3	974	15.9	14.8-16.9	686	7.3	6.7-7.9	1,103	7.2	6.7-7.6	652	10.8	10.0-11.7	451	4.8	4.4-5.3
1988	1,580	10.3	9.8-10.8	892	14.5	13.5-15.5	688	7.4	6.8-8.0	1,064	6.9	6.5-7.4	578	9.5	8.7-10.3	486	5.2	4.7-5.7
1989	1,511	9.8	9.3-10.4	934	15.1	14.2-16.2	577	6.1	5.6-6.7	1,109	7.2	6.8-7.6	645	10.7	9.9-11.6	464	4.9	4.5-5.4
1990	1,705	11.1	10.6-11.7	1,029	16.6	15.6-17.7	676	7.3	6.7-7.9	1,001	6.6	6.2-7.0	613	10.1	9.3-11.0	388	4.1	3.7-4.5
1991	1,590	10.3	9.8-10.8	965	15.7	14.7-16.7	625	6.6	6.1-7.2	1,005	6.5	6.1-6.9	615	10.2	9.4-11.1	390	4.1	3.7-4.5
1992	1,567	10.1	9.6-10.6	942	15.3	14.3-16.3	625	6.6	6.1-7.2	1,025	6.6	6.2-7.0	614	10.2	9.4-11.1	411	4.3	3.9-4.7
1993	1,637	10.5	10.0-11.0	974	15.7	14.7-16.7	663	7.0	6.4-7.5	998	6.4	6.0-6.8	587	9.8	9.0-10.6	411	4.3	3.9-4.7
1994	1,622	10.4	9.9-10.9	941	14.9	14.0-15.9	681	7.2	6.7-7.8	1,044	6.7	6.3-7.1	610	10.0	9.2-10.8	434	4.5	4.1-5.0
1995	1,531	9.8	9.3-10.3	897	14.2	13.2-15.1	634	6.7	6.2-7.2	1,003	6.4	6.0-6.8	559	9.2	8.4-10.0	444	4.6	4.2-5.1
1996	1,546	9.9	9.4-10.4	900	14.2	13.3-15.2	646	6.7	6.2-7.3	939	6.0	5.6-6.4	515	8.3	7.6-9.1	424	4.4	4.0-4.8
1997	1,511	9.6	9.1-10.1	898	14.0	13.1-15.0	613	6.4	5.9-7.0	888	5.6	5.3-6.0	490	7.7	7.1-8.5	398	4.2	3.7-4.6
1998	1,577	9.9	9.5-10.4	959	14.6	13.7-15.6	618	6.5	6.0-7.1	884	5.5	5.2-5.9	495	7.7	7.1-8.5	389	4.0	3.6-4.4
1999	1,488	9.3	8.8-9.8	915	13.9	13.0-14.9	573	6.0	5.5-6.5	838	5.2	4.8-5.5	487	7.5	6.9-8.2	351	3.6	3.2-4.0
2000	1,470	9.0	8.6-9.5	861	13.0	12.1-13.9	609	6.2	5.7-6.7	811	4.9	4.6-5.3	439	6.8	6.2-7.4	372	3.7	3.4-4.2
2001	1,450	8.9	8.5-9.4	883	13.1	12.2-14.0	567	6.0	5.5-6.5	789	4.8	4.5-5.2	434	6.5	5.9-7.2	355	3.6	3.2-4.0
2002	1,385	8.5	8.0-8.9	821	12.0	11.2-12.9	564	5.9	5.4-6.4	726	4.4	4.1-4.7	413	6.2	5.6-6.8	313	3.1	2.7-3.5
2003	1,478	9.0	8.5-9.4	877	12.7	11.8-13.5	601	6.2	5.7-6.8	724	4.3	4.0-4.7	401	5.9	5.3-6.5	323	3.3	2.9-3.7
2004	1,424	8.6	8.2-9.1	835	12.1	11.3-12.9	589	6.1	5.6-6.7	777	4.6	4.3-5.0	428	6.3	5.7-6.9	349	3.5	3.2-3.9
2005	1,349	8.1	7.7-8.6	801	11.4	10.6-12.2	548	5.7	5.2-6.2	715	4.2	3.9-4.6	404	5.9	5.3-6.5	311	3.1	2.7-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,313	8.0	7.5-8.4	785	11.1	10.3-11.9	528	5.6	5.1-6.1	631	3.8	3.5-4.1	358	5.1	4.6-5.7	273	2.8	2.5-3.2
2007	1,386	8.3	7.9-8.8	846	11.7	11.0-12.6	540	5.7	5.2-6.2	606	3.6	3.3-3.9	347	4.8	4.3-5.4	259	2.7	2.3-3.0
2008	1,279	7.6	7.1-8.0	809	11.2	10.4-12.0	470	4.9	4.4-5.4	642	3.7	3.4-4.0	396	5.5	5.0-6.1	246	2.4	2.1-2.7
2009	1,237	7.3	6.9-7.7	754	10.3	9.6-11.1	483	4.9	4.5-5.4	593	3.5	3.2-3.8	353	4.9	4.4-5.5	240	2.4	2.1-2.7
2010	1,314	7.7	7.3-8.1	813	10.8	10.1-11.6	501	5.2	4.7-5.7	581	3.3	3.0-3.6	313	4.3	3.8-4.8	268	2.7	2.3-3.0
2011	1,315	7.6	7.2-8.0	797	10.6	9.9-11.4	518	5.3	4.9-5.8	555	3.1	2.9-3.4	311	4.2	3.7-4.7	244	2.4	2.1-2.7
2012	1,336	7.6	7.2-8.1	817	10.6	9.9-11.4	519	5.4	4.9-5.9	578	3.2	3.0-3.5	337	4.4	4.0-4.9	241	2.3	2.0-2.7
2013	1,429	8.1	7.7-8.5	874	11.2	10.4-11.9	555	5.6	5.2-6.1	581	3.2	2.9-3.4	353	4.5	4.0-5.0	228	2.2	1.9-2.5
2014	1,324	7.5	7.1-7.9	860	10.9	10.1-11.6	464	4.8	4.3-5.3	588	3.2	3.0-3.5	354	4.5	4.0-5.0	234	2.3	2.0-2.6
2015	1,318	7.3	6.9-7.7	794	9.8	9.1-10.5	524	5.3	4.8-5.8	516	2.8	2.6-3.1	301	3.8	3.4-4.2	215	2.1	1.8-2.4
2016	1,372	7.6	7.2-8.0	826	10.1	9.4-10.8	546	5.7	5.2-6.2	535	2.9	2.6-3.1	340	4.1	3.7-4.6	195	1.9	1.6-2.2
2017	1,417	7.8	7.4-8.2	836	10.1	9.4-10.8	581	5.9	5.4-6.4	538	2.9	2.6-3.2	304	3.7	3.3-4.1	234	2.3	2.0-2.6
2018	1,312	7.2	6.8-7.6	782	9.4	8.7-10.1	530	5.5	5.0-6.0	507	2.7	2.5-3.0	301	3.7	3.3-4.1	206	1.9	1.7-2.3
2019	1,326	7.1	6.7-7.5	826	9.7	9.0-10.4	500	5.0	4.5-5.5	513	2.8	2.5-3.0	312	3.8	3.4-4.3	201	1.9	1.7-2.3
2020	1,189	6.5	6.1-6.9	743	8.8	8.2-9.5	446	4.6	4.2-5.1	465	2.5	2.2-2.7	279	3.3	2.9-3.7	186	1.8	1.5-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	10,107	67.6	66.2-68.9	4,935	80.9	78.5-83.3	5,172	59.0	57.4-60.7	5,252	35.8	34.8-36.8	2,572	44.0	42.2-45.8	2,680	30.8	29.6-32.0
1977	10,169	67.6	66.3-69.0	4,976	81.7	79.3-84.1	5,193	58.7	57.1-60.4	5,119	34.9	33.9-35.8	2,475	42.6	40.8-44.4	2,644	30.1	28.9-31.3
1978	10,132	67.3	66.0-68.7	4,952	81.5	79.2-83.9	5,180	58.3	56.7-59.9	5,165	34.9	34.0-35.9	2,568	43.8	42.1-45.7	2,597	29.4	28.3-30.6
1979	10,284	68.0	66.7-69.3	4,962	81.4	79.1-83.8	5,322	59.4	57.7-61.0	5,131	34.4	33.4-35.4	2,452	42.0	40.2-43.7	2,679	29.8	28.7-31.0
1980	10,655	69.9	68.6-71.3	5,332	86.8	84.3-89.2	5,323	58.9	57.3-60.5	5,120	34.2	33.3-35.2	2,530	43.4	41.6-45.2	2,590	28.7	27.6-29.8
1981	10,695	70.3	69.0-71.7	5,302	87.6	85.1-90.1	5,393	59.4	57.8-61.0	4,995	33.3	32.3-34.2	2,405	41.0	39.3-42.7	2,590	28.4	27.3-29.6
1982	10,708	70.0	68.6-71.3	5,245	85.8	83.4-88.2	5,463	59.8	58.2-61.4	4,936	32.7	31.8-33.6	2,438	41.6	39.9-43.4	2,498	27.3	26.2-28.4
1983	10,747	69.9	68.6-71.2	5,321	86.7	84.2-89.1	5,426	59.1	57.5-60.7	4,892	32.3	31.3-33.2	2,373	40.4	38.7-42.1	2,519	27.2	26.2-28.3
1984	10,701	69.6	68.2-70.9	5,180	84.1	81.7-86.5	5,521	60.1	58.5-61.7	5,044	33.0	32.1-34.0	2,509	42.3	40.6-44.1	2,535	27.4	26.3-28.5
1985	10,943	70.7	69.4-72.1	5,499	88.7	86.3-91.2	5,444	59.0	57.4-60.6	4,832	31.6	30.7-32.5	2,357	39.4	37.7-41.1	2,475	26.5	25.5-27.6
1986	10,735	69.4	68.0-70.7	5,316	86.2	83.8-88.6	5,419	58.6	57.1-60.2	4,779	31.2	30.3-32.1	2,360	39.7	38.1-41.4	2,419	25.9	24.9-27.0
1987	10,274	66.2	64.9-67.5	5,094	82.4	80.1-84.8	5,180	55.8	54.3-57.4	4,739	30.7	29.9-31.6	2,344	39.3	37.7-41.0	2,395	25.4	24.4-26.5
1988	9,927	63.9	62.6-65.2	4,837	77.9	75.6-80.2	5,090	54.6	53.1-56.2	4,645	30.2	29.3-31.0	2,342	39.1	37.5-40.8	2,303	24.6	23.6-25.6
1989	9,653	62.3	61.1-63.6	4,799	77.3	75.1-79.6	4,854	52.4	50.9-53.9	4,319	27.9	27.1-28.8	2,059	34.4	32.9-36.0	2,260	23.9	22.9-25.0
1990	9,657	61.9	60.7-63.2	4,681	74.5	72.3-76.7	4,976	53.2	51.7-54.7	4,238	27.3	26.5-28.1	2,114	35.1	33.6-36.7	2,124	22.4	21.4-23.4
1991	9,598	61.6	60.4-62.9	4,787	76.9	74.7-79.2	4,811	51.7	50.2-53.2	4,171	26.8	26.0-27.7	2,030	33.8	32.3-35.3	2,141	22.5	21.6-23.5
1992	9,376	60.0	58.8-61.3	4,624	73.8	71.6-76.0	4,752	50.9	49.4-52.4	3,964	25.3	24.6-26.1	1,913	31.5	30.1-33.0	2,051	21.4	20.5-22.4
1993	9,450	60.2	59.0-61.4	4,570	71.9	69.8-74.1	4,880	52.1	50.6-53.6	4,001	25.4	24.6-26.2	1,954	32.0	30.5-33.5	2,047	21.2	20.3-22.2
1994	9,922	63.1	61.9-64.4	4,801	75.7	73.6-77.9	5,121	54.7	53.2-56.2	4,193	26.6	25.8-27.4	2,014	33.1	31.7-34.7	2,179	22.5	21.6-23.5
1995	9,572	60.8	59.6-62.0	4,663	73.3	71.2-75.5	4,909	52.2	50.7-53.7	4,142	26.1	25.3-26.9	1,977	32.3	30.9-33.8	2,165	22.3	21.4-23.3
1996	9,322	59.2	58.0-60.4	4,637	72.5	70.4-74.6	4,685	49.9	48.5-51.4	3,794	23.9	23.1-24.6	1,862	29.9	28.6-31.4	1,932	19.8	18.9-20.8
1997	9,785	61.8	60.6-63.1	4,768	73.6	71.5-75.7	5,017	53.3	51.8-54.8	3,758	23.5	22.8-24.3	1,858	29.6	28.2-31.0	1,900	19.4	18.6-20.3
1998	10,180	63.9	62.7-65.2	4,990	76.1	74.0-78.3	5,190	55.2	53.7-56.8	3,690	22.9	22.2-23.7	1,754	27.3	26.1-28.7	1,936	19.7	18.8-20.6
1999	10,085	62.7	61.5-64.0	4,964	75.0	72.9-77.1	5,121	54.0	52.5-55.5	3,570	21.9	21.2-22.6	1,734	26.9	25.6-28.2	1,836	18.5	17.7-19.4
2000	10,128	62.5	61.3-63.7	5,033	75.1	73.0-77.2	5,095	53.2	51.7-54.7	3,633	22.0	21.3-22.8	1,812	27.8	26.5-29.1	1,821	18.0	17.2-18.9
2001	9,669	59.3	58.1-60.5	4,772	70.7	68.7-72.7	4,897	51.1	49.7-52.6	3,453	20.8	20.1-21.5	1,651	25.1	23.9-26.4	1,802	17.9	17.0-18.7
2002	9,527	58.1	56.9-59.3	4,726	68.8	66.8-70.8	4,801	49.8	48.4-51.2	3,456	20.7	20.0-21.4	1,688	25.3	24.1-26.5	1,768	17.5	16.7-18.4
2003	9,156	55.4	54.2-56.5	4,483	64.7	62.8-66.7	4,673	48.4	47.0-49.8	3,263	19.4	18.7-20.0	1,578	23.3	22.2-24.5	1,685	16.6	15.8-17.4
2004	8,897	53.5	52.4-54.7	4,305	61.6	59.7-63.5	4,592	47.5	46.1-48.9	3,073	18.1	17.5-18.8	1,478	21.5	20.4-22.6	1,595	15.5	14.7-16.3
2005	8,479	50.7	49.7-51.8	4,226	60.0	58.2-61.9	4,253	43.9	42.6-45.3	2,843	16.6	16.0-17.2	1,368	20.0	18.9-21.1	1,475	14.2	13.5-15.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	8,122	48.7	47.6-49.8	4,029	56.9	55.2-58.7	4,093	42.4	41.1-43.7	2,839	16.6	16.0-17.3	1,378	20.0	19.0-21.1	1,461	14.3	13.5-15.0
2007	8,209	48.9	47.8-50.0	4,054	56.2	54.5-58.0	4,155	43.2	41.9-44.6	2,740	15.9	15.3-16.5	1,332	18.9	17.9-20.0	1,408	13.6	12.9-14.4
2008	7,931	46.8	45.8-47.9	3,931	53.9	52.3-55.7	4,000	41.3	40.0-42.7	2,753	15.8	15.3-16.5	1,343	18.9	17.9-19.9	1,410	13.6	12.8-14.3
2009	7,572	44.3	43.3-45.3	3,737	50.6	49.0-52.3	3,835	39.3	38.0-40.6	2,646	15.1	14.5-15.7	1,279	17.8	16.8-18.8	1,367	13.2	12.5-13.9
2010	7,521	43.7	42.7-44.7	3,710	49.5	47.9-51.2	3,811	39.0	37.7-40.3	2,652	14.9	14.3-15.5	1,324	18.0	17.0-19.0	1,328	12.6	11.9-13.3
2011	7,197	41.6	40.6-42.6	3,557	47.1	45.6-48.7	3,640	37.3	36.0-38.5	2,584	14.3	13.7-14.9	1,283	17.0	16.0-17.9	1,301	12.1	11.5-12.8
2012	6,926	39.6	38.7-40.6	3,566	46.2	44.7-47.8	3,360	34.3	33.1-35.5	2,473	13.6	13.0-14.1	1,180	15.5	14.6-16.4	1,293	12.1	11.4-12.8
2013	7,109	40.2	39.3-41.2	3,643	46.6	45.1-48.2	3,466	35.2	34.0-36.4	2,448	13.3	12.8-13.9	1,258	16.3	15.4-17.2	1,190	10.9	10.3-11.6
2014	6,896	38.6	37.7-39.6	3,554	44.7	43.2-46.2	3,342	33.7	32.5-34.9	2,378	12.7	12.2-13.3	1,201	15.2	14.4-16.1	1,177	10.8	10.2-11.5
2015	6,907	38.6	37.7-39.6	3,482	43.5	42.0-45.0	3,425	34.7	33.5-35.9	2,378	12.6	12.1-13.2	1,219	15.3	14.4-16.2	1,159	10.6	10.0-11.2
2016	6,932	38.6	37.7-39.6	3,599	44.4	42.9-45.9	3,333	33.8	32.6-35.1	2,275	12.0	11.5-12.5	1,137	14.0	13.2-14.9	1,138	10.4	9.8-11.1
2017	6,887	38.1	37.1-39.0	3,533	43.0	41.6-44.5	3,354	34.0	32.8-35.2	2,420	12.7	12.2-13.2	1,182	14.4	13.6-15.3	1,238	11.4	10.7-12.0
2018	6,609	36.4	35.5-37.3	3,448	41.8	40.4-43.3	3,161	31.7	30.5-32.9	2,106	11.1	10.6-11.6	1,064	12.9	12.2-13.8	1,042	9.7	9.1-10.3
2019	6,718	37.0	36.1-37.9	3,533	42.5	41.1-44.0	3,185	32.2	31.1-33.4	2,135	11.0	10.5-11.5	1,108	13.2	12.5-14.1	1,027	9.0	8.5-9.6
2020	5,958	32.7	31.8-33.6	3,133	37.5	36.2-38.9	2,825	28.5	27.4-29.7	2,096	10.9	10.4-11.4	1,107	13.2	12.4-14.0	989	9.0	8.4-9.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	7,072	47.6	46.4-48.7	3,288	54.6	52.7-56.6	3,784	43.2	41.8-44.6	4,137	28.3	27.4-29.2	1,938	33.3	31.8-34.9	2,199	25.3	24.2-26.4
1977	7,146	47.9	46.8-49.1	3,342	56.0	54.1-58.1	3,804	43.1	41.7-44.5	4,177	28.5	27.6-29.4	1,973	34.0	32.4-35.6	2,204	25.1	24.1-26.2
1978	7,177	48.0	46.8-49.1	3,365	56.2	54.2-58.2	3,812	42.9	41.6-44.3	4,194	28.4	27.5-29.3	2,018	34.6	33.0-36.2	2,176	24.6	23.5-25.6
1979	7,346	48.9	47.8-50.1	3,393	56.7	54.7-58.7	3,953	44.2	42.8-45.6	4,253	28.5	27.6-29.4	1,971	33.7	32.1-35.3	2,282	25.4	24.4-26.5
1980	7,562	49.9	48.8-51.1	3,659	60.6	58.5-62.6	3,903	43.2	41.8-44.6	4,256	28.5	27.6-29.3	2,069	35.6	34.0-37.2	2,187	24.2	23.2-25.2
1981	7,621	50.3	49.1-51.4	3,629	60.6	58.5-62.7	3,992	44.0	42.6-45.4	4,243	28.3	27.5-29.2	2,010	34.4	32.8-36.0	2,233	24.6	23.5-25.6
1982	7,554	49.6	48.5-50.8	3,509	58.4	56.4-60.5	4,045	44.2	42.8-45.6	4,152	27.5	26.7-28.4	2,030	34.7	33.2-36.3	2,122	23.2	22.2-24.2
1983	7,631	49.9	48.7-51.0	3,583	59.3	57.3-61.4	4,048	44.1	42.7-45.5	4,108	27.1	26.2-27.9	1,959	33.4	31.8-34.9	2,149	23.2	22.2-24.2
1984	7,707	50.2	49.1-51.4	3,549	58.5	56.5-60.5	4,158	45.1	43.7-46.5	4,269	27.9	27.1-28.8	2,087	35.2	33.7-36.8	2,182	23.5	22.6-24.6
1985	7,936	51.4	50.3-52.6	3,809	62.2	60.2-64.3	4,127	44.6	43.2-46.0	4,128	27.0	26.1-27.8	1,981	33.0	31.5-34.6	2,147	23.0	22.1-24.0
1986	7,823	50.7	49.6-51.8	3,686	60.4	58.4-62.5	4,137	44.7	43.3-46.1	4,086	26.6	25.8-27.5	1,983	33.4	31.9-35.0	2,103	22.5	21.6-23.5
1987	7,369	47.5	46.4-48.6	3,506	57.4	55.4-59.4	3,863	41.3	40.0-42.7	4,054	26.3	25.5-27.1	2,010	33.7	32.2-35.3	2,044	21.7	20.8-22.7
1988	7,152	46.0	45.0-47.1	3,301	53.7	51.8-55.6	3,851	41.1	39.8-42.4	4,014	26.0	25.2-26.9	2,008	33.5	32.0-35.1	2,006	21.4	20.4-22.4
1989	6,886	44.5	43.5-45.6	3,303	54.1	52.2-56.0	3,583	38.5	37.2-39.8	3,730	24.1	23.3-24.9	1,742	29.2	27.8-30.7	1,988	21.1	20.1-22.0
1990	6,951	44.6	43.5-45.7	3,226	52.1	50.2-53.9	3,725	39.7	38.4-41.0	3,704	23.9	23.1-24.6	1,842	30.7	29.3-32.2	1,862	19.6	18.7-20.5
1991	6,996	44.8	43.8-45.9	3,359	54.5	52.6-56.4	3,637	38.8	37.5-40.1	3,549	22.9	22.1-23.6	1,708	28.5	27.1-29.9	1,841	19.3	18.4-20.2
1992	6,762	43.3	42.2-44.3	3,183	51.4	49.5-53.2	3,579	38.1	36.9-39.4	3,389	21.6	20.9-22.4	1,597	26.4	25.0-27.7	1,792	18.7	17.8-19.6
1993	6,900	43.9	42.9-45.0	3,178	50.5	48.7-52.3	3,722	39.5	38.2-40.8	3,453	22.0	21.2-22.7	1,669	27.5	26.1-28.9	1,784	18.5	17.7-19.4
1994	7,261	46.1	45.1-47.2	3,405	54.2	52.4-56.1	3,856	40.9	39.6-42.2	3,643	23.0	22.3-23.8	1,724	28.4	27.1-29.8	1,919	19.8	18.9-20.7
1995	6,999	44.3	43.3-45.3	3,292	52.2	50.4-54.0	3,707	39.0	37.7-40.3	3,537	22.3	21.5-23.0	1,659	27.3	25.9-28.6	1,878	19.3	18.4-20.2
1996	6,728	42.5	41.5-43.6	3,193	50.4	48.6-52.2	3,535	37.3	36.0-38.5	3,252	20.4	19.7-21.1	1,587	25.6	24.4-26.9	1,665	17.0	16.2-17.9
1997	7,254	45.7	44.6-46.8	3,382	52.6	50.8-54.4	3,872	40.8	39.5-42.1	3,267	20.4	19.8-21.2	1,580	25.1	23.9-26.4	1,687	17.2	16.4-18.1
1998	7,417	46.4	45.4-47.5	3,512	54.1	52.3-55.9	3,905	41.2	39.9-42.6	3,179	19.7	19.0-20.4	1,491	23.3	22.2-24.6	1,688	17.1	16.3-17.9
1999	7,313	45.3	44.2-46.3	3,456	52.6	50.8-54.4	3,857	40.2	38.9-41.5	3,022	18.5	17.9-19.2	1,448	22.5	21.4-23.7	1,574	15.8	15.0-16.6
2000	7,387	45.3	44.3-46.4	3,489	52.5	50.7-54.3	3,898	40.3	39.0-41.6	3,106	18.8	18.1-19.5	1,529	23.6	22.4-24.8	1,577	15.5	14.8-16.3
2001	7,053	43.1	42.1-44.1	3,359	50.0	48.4-51.8	3,694	38.0	36.8-39.3	2,931	17.6	17.0-18.3	1,373	21.0	19.9-22.1	1,558	15.4	14.6-16.2
2002	6,938	42.1	41.1-43.1	3,301	48.4	46.8-50.1	3,637	37.4	36.2-38.7	2,894	17.3	16.7-18.0	1,401	21.1	20.0-22.2	1,493	14.7	14.0-15.5
2003	6,652	40.0	39.1-41.0	3,123	45.4	43.8-47.1	3,529	36.1	34.9-37.3	2,791	16.5	15.9-17.2	1,324	19.7	18.6-20.8	1,467	14.4	13.6-15.1
2004	6,413	38.4	37.5-39.4	2,973	42.9	41.3-44.5	3,440	35.1	33.9-36.3	2,586	15.2	14.6-15.8	1,215	17.8	16.8-18.8	1,371	13.2	12.5-14.0
2005	6,219	37.1	36.2-38.0	2,969	42.4	40.9-44.0	3,250	33.2	32.0-34.4	2,376	13.9	13.3-14.4	1,119	16.4	15.5-17.4	1,257	12.1	11.4-12.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	5,919	35.3	34.4-36.2	2,824	40.3	38.8-41.8	3,095	31.6	30.4-32.7	2,401	14.0	13.5-14.6	1,144	16.7	15.7-17.7	1,257	12.2	11.5-12.9
2007	5,963	35.4	34.5-36.3	2,831	39.6	38.1-41.1	3,132	32.1	31.0-33.3	2,327	13.4	12.9-14.0	1,110	15.8	14.9-16.8	1,217	11.7	11.1-12.4
2008	5,738	33.7	32.8-34.6	2,716	37.5	36.1-38.9	3,022	31.0	29.8-32.1	2,262	13.0	12.5-13.6	1,088	15.3	14.4-16.3	1,174	11.3	10.6-12.0
2009	5,423	31.5	30.7-32.4	2,521	34.4	33.0-35.8	2,902	29.3	28.2-30.4	2,200	12.6	12.0-13.1	1,044	14.6	13.7-15.5	1,156	11.1	10.5-11.8
2010	5,384	31.1	30.3-32.0	2,534	34.1	32.8-35.5	2,850	28.6	27.5-29.7	2,202	12.3	11.8-12.8	1,065	14.5	13.6-15.4	1,137	10.6	10.0-11.3
2011	5,145	29.6	28.8-30.4	2,433	32.5	31.2-33.8	2,712	27.3	26.3-28.4	2,132	11.7	11.2-12.3	1,030	13.6	12.8-14.5	1,102	10.2	9.6-10.9
2012	4,993	28.4	27.6-29.2	2,427	31.7	30.4-33.0	2,566	25.7	24.7-26.8	2,044	11.2	10.7-11.7	942	12.4	11.6-13.3	1,102	10.3	9.6-10.9
2013	5,023	28.2	27.4-29.0	2,457	31.6	30.4-32.9	2,566	25.6	24.6-26.7	2,009	10.9	10.4-11.4	1,004	13.1	12.3-13.9	1,005	9.2	8.6-9.8
2014	4,897	27.2	26.5-28.0	2,403	30.5	29.3-31.8	2,494	24.7	23.7-25.7	1,924	10.3	9.8-10.7	948	12.1	11.3-12.9	976	8.9	8.3-9.5
2015	4,883	27.2	26.5-28.0	2,333	29.4	28.2-30.7	2,550	25.5	24.5-26.5	1,936	10.2	9.8-10.7	964	12.1	11.3-12.9	972	8.8	8.2-9.4
2016	4,966	27.5	26.7-28.3	2,454	30.4	29.2-31.7	2,512	25.1	24.1-26.2	1,807	9.5	9.1-10.0	875	10.8	10.1-11.6	932	8.5	7.9-9.1
2017	4,912	26.9	26.2-27.7	2,391	29.2	28.0-30.5	2,521	25.1	24.1-26.2	1,930	10.1	9.6-10.6	920	11.2	10.5-12.0	1,010	9.2	8.6-9.8
2018	4,672	25.5	24.8-26.3	2,343	28.6	27.4-29.8	2,329	22.9	21.9-23.9	1,656	8.7	8.2-9.1	804	9.8	9.1-10.5	852	7.8	7.3-8.4
2019	4,761	25.9	25.1-26.6	2,383	28.6	27.4-29.8	2,378	23.5	22.6-24.6	1,664	8.5	8.1-9.0	832	10.0	9.3-10.7	832	7.3	6.8-7.8
2020	4,197	22.8	22.1-23.5	2,099	25.2	24.1-26.3	2,098	20.8	19.8-21.7	1,598	8.2	7.8-8.7	822	9.8	9.1-10.5	776	7.0	6.5-7.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3,035	20.0	19.3-20.8	1,647	26.2	24.9-27.6	1,388	15.9	15.0-16.7	1,115	7.5	7.1-8.0	634	10.7	9.8-11.6	481	5.5	5.0-6.0
1977	3,023	19.7	19.0-20.5	1,634	25.7	24.4-27.0	1,389	15.6	14.8-16.5	942	6.4	6.0-6.8	502	8.6	7.8-9.4	440	5.0	4.5-5.5
1978	2,955	19.4	18.7-20.1	1,587	25.4	24.1-26.7	1,368	15.3	14.5-16.2	971	6.6	6.2-7.0	550	9.2	8.4-10.1	421	4.8	4.4-5.3
1979	2,938	19.1	18.4-19.8	1,569	24.7	23.5-26.0	1,369	15.2	14.4-16.0	878	5.9	5.5-6.3	481	8.3	7.5-9.1	397	4.4	4.0-4.9
1980	3,093	20.0	19.3-20.7	1,673	26.2	24.9-27.5	1,420	15.7	14.9-16.6	864	5.8	5.4-6.2	461	7.8	7.0-8.5	403	4.5	4.0-4.9
1981	3,074	20.0	19.3-20.8	1,673	27.0	25.7-28.4	1,401	15.4	14.6-16.3	752	5.0	4.6-5.3	395	6.6	6.0-7.3	357	3.9	3.5-4.3
1982	3,154	20.4	19.6-21.1	1,736	27.4	26.1-28.8	1,418	15.6	14.8-16.5	784	5.2	4.8-5.6	408	6.9	6.2-7.6	376	4.1	3.7-4.5
1983	3,116	20.0	19.3-20.7	1,738	27.3	26.0-28.7	1,378	15.0	14.2-15.9	784	5.2	4.8-5.6	414	7.0	6.3-7.8	370	4.0	3.6-4.5
1984	2,994	19.3	18.6-20.1	1,631	25.6	24.3-26.9	1,363	15.0	14.2-15.9	775	5.1	4.7-5.5	422	7.1	6.4-7.9	353	3.8	3.4-4.2
1985	3,007	19.3	18.6-20.0	1,690	26.5	25.2-27.8	1,317	14.4	13.6-15.2	704	4.6	4.3-5.0	376	6.3	5.7-7.0	328	3.5	3.1-3.9
1986	2,912	18.7	18.0-19.4	1,630	25.8	24.5-27.1	1,282	14.0	13.2-14.8	693	4.5	4.2-4.9	377	6.3	5.7-7.0	316	3.4	3.0-3.8
1987	2,905	18.8	18.1-19.5	1,588	25.0	23.8-26.3	1,317	14.5	13.7-15.3	685	4.5	4.1-4.8	334	5.6	5.0-6.3	351	3.7	3.3-4.1
1988	2,775	17.9	17.2-18.6	1,536	24.2	23.0-25.5	1,239	13.6	12.8-14.4	631	4.1	3.8-4.4	334	5.6	5.0-6.2	297	3.2	2.8-3.6
1989	2,767	17.8	17.1-18.5	1,496	23.2	22.0-24.5	1,271	13.9	13.2-14.7	589	3.8	3.5-4.2	317	5.2	4.6-5.8	272	2.9	2.5-3.3
1990	2,706	17.3	16.7-18.0	1,455	22.4	21.3-23.6	1,251	13.5	12.7-14.3	534	3.4	3.2-3.8	272	4.4	3.9-5.0	262	2.8	2.5-3.2
1991	2,602	16.8	16.1-17.5	1,428	22.4	21.3-23.7	1,174	12.9	12.1-13.7	622	4.0	3.7-4.3	322	5.3	4.7-5.9	300	3.2	2.9-3.6
1992	2,614	16.8	16.1-17.4	1,441	22.4	21.2-23.6	1,173	12.7	12.0-13.5	575	3.7	3.4-4.0	316	5.2	4.6-5.8	259	2.7	2.4-3.1
1993	2,550	16.3	15.7-17.0	1,392	21.5	20.4-22.7	1,158	12.6	11.9-13.4	548	3.5	3.2-3.8	285	4.5	4.0-5.1	263	2.7	2.4-3.1
1994	2,661	17.0	16.4-17.7	1,396	21.5	20.4-22.7	1,265	13.8	13.1-14.6	550	3.5	3.2-3.8	290	4.7	4.2-5.3	260	2.7	2.4-3.1
1995	2,573	16.5	15.9-17.2	1,371	21.2	20.0-22.3	1,202	13.2	12.5-14.0	605	3.8	3.5-4.2	318	5.1	4.5-5.7	287	3.0	2.7-3.4
1996	2,594	16.7	16.0-17.3	1,444	22.1	21.0-23.3	1,150	12.7	11.9-13.4	542	3.4	3.1-3.7	275	4.3	3.8-4.9	267	2.8	2.5-3.2
1997	2,531	16.1	15.5-16.8	1,386	21.0	19.9-22.1	1,145	12.5	11.8-13.2	491	3.1	2.8-3.4	278	4.4	3.9-5.0	213	2.2	1.9-2.5
1998	2,763	17.5	16.9-18.2	1,478	22.1	20.9-23.2	1,285	14.0	13.2-14.8	511	3.2	2.9-3.5	263	4.0	3.5-4.5	248	2.6	2.3-2.9
1999	2,772	17.5	16.8-18.1	1,508	22.4	21.3-23.6	1,264	13.8	13.0-14.6	548	3.4	3.1-3.7	286	4.4	3.9-4.9	262	2.7	2.4-3.0
2000	2,741	17.1	16.5-17.8	1,544	22.6	21.5-23.8	1,197	12.9	12.2-13.7	527	3.2	3.0-3.5	283	4.2	3.8-4.8	244	2.5	2.2-2.8
2001	2,616	16.3	15.6-16.9	1,413	20.6	19.5-21.7	1,203	13.1	12.4-13.9	522	3.2	2.9-3.5	278	4.2	3.7-4.7	244	2.5	2.2-2.8
2002	2,589	16.0	15.4-16.6	1,425	20.4	19.3-21.5	1,164	12.4	11.7-13.1	562	3.4	3.1-3.7	287	4.2	3.8-4.8	275	2.8	2.5-3.2
2003	2,504	15.3	14.8-16.0	1,360	19.3	18.3-20.3	1,144	12.3	11.5-13.0	472	2.8	2.6-3.1	254	3.7	3.2-4.2	218	2.2	1.9-2.6
2004	2,484	15.1	14.5-15.7	1,332	18.7	17.7-19.7	1,152	12.4	11.7-13.1	487	2.9	2.7-3.2	263	3.8	3.3-4.3	224	2.2	1.9-2.5
2005	2,260	13.7	13.1-14.3	1,257	17.6	16.6-18.6	1,003	10.7	10.0-11.4	467	2.8	2.5-3.0	249	3.6	3.1-4.1	218	2.1	1.9-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	2,203	13.4	12.9-14.0	1,205	16.7	15.7-17.7	998	10.8	10.1-11.5	438	2.6	2.4-2.9	234	3.4	2.9-3.8	204	2.1	1.8-2.4
2007	2,246	13.5	13.0-14.1	1,223	16.7	15.7-17.6	1,023	11.1	10.4-11.8	413	2.4	2.2-2.7	222	3.1	2.7-3.5	191	1.9	1.6-2.2
2008	2,193	13.1	12.5-13.7	1,215	16.5	15.5-17.4	978	10.4	9.7-11.1	491	2.8	2.6-3.1	255	3.5	3.1-4.0	236	2.3	2.0-2.6
2009	2,149	12.8	12.3-13.4	1,216	16.2	15.3-17.2	933	10.0	9.4-10.7	446	2.6	2.3-2.8	235	3.2	2.8-3.6	211	2.1	1.8-2.4
2010	2,137	12.6	12.0-13.1	1,176	15.4	14.5-16.3	961	10.3	9.7-11.0	450	2.6	2.4-2.9	259	3.5	3.1-3.9	191	1.9	1.7-2.2
2011	2,052	12.0	11.5-12.6	1,124	14.6	13.8-15.5	928	9.9	9.3-10.6	452	2.5	2.3-2.8	253	3.3	2.9-3.8	199	1.9	1.6-2.2
2012	1,933	11.2	10.7-11.8	1,139	14.5	13.7-15.4	794	8.6	8.0-9.2	429	2.4	2.2-2.6	238	3.1	2.7-3.5	191	1.9	1.6-2.2
2013	2,086	12.0	11.5-12.6	1,186	15.0	14.1-15.9	900	9.6	8.9-10.2	439	2.4	2.2-2.7	254	3.2	2.8-3.6	185	1.8	1.5-2.1
2014	1,999	11.4	10.9-11.9	1,151	14.2	13.4-15.1	848	9.0	8.4-9.7	454	2.5	2.2-2.7	253	3.1	2.7-3.5	201	1.9	1.7-2.3
2015	2,024	11.4	10.9-11.9	1,149	14.0	13.2-14.9	875	9.2	8.6-9.9	442	2.4	2.2-2.7	255	3.2	2.8-3.6	187	1.8	1.5-2.1
2016	1,966	11.1	10.6-11.7	1,145	14.0	13.2-14.9	821	8.7	8.1-9.4	468	2.5	2.3-2.8	262	3.2	2.8-3.7	206	2.0	1.7-2.3
2017	1,975	11.1	10.6-11.7	1,142	13.8	13.0-14.6	833	8.9	8.2-9.5	490	2.6	2.4-2.9	262	3.2	2.8-3.6	228	2.2	1.9-2.5
2018	1,937	10.9	10.4-11.4	1,105	13.3	12.5-14.1	832	8.8	8.2-9.5	450	2.4	2.2-2.7	260	3.1	2.8-3.6	190	1.9	1.6-2.2
2019	1,957	11.1	10.6-11.6	1,150	14.0	13.1-14.8	807	8.7	8.1-9.3	471	2.5	2.3-2.7	276	3.2	2.9-3.7	195	1.8	1.5-2.1
2020	1,761	9.9	9.4-10.4	1,034	12.4	11.6-13.2	727	7.8	7.2-8.4	498	2.6	2.4-2.9	285	3.4	3.0-3.9	213	1.9	1.7-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	457	3.0	2.8-3.3	259	4.1	3.6-4.7	198	2.2	1.9-2.6	408	2.7	2.4-3.0	229	3.6	3.1-4.1	179	2.1	1.8-2.4
1977	494	3.2	2.9-3.5	317	4.9	4.3-5.4	177	2.0	1.7-2.3	493	3.2	2.9-3.5	275	4.2	3.7-4.8	218	2.5	2.1-2.8
1978	455	3.0	2.7-3.2	261	4.1	3.6-4.6	194	2.2	1.9-2.5	467	3.0	2.8-3.3	274	4.3	3.8-4.9	193	2.2	1.9-2.5
1979	443	2.9	2.7-3.2	248	3.9	3.4-4.4	195	2.3	2.0-2.6	450	3.0	2.7-3.3	241	3.9	3.4-4.5	209	2.4	2.1-2.7
1980	453	3.0	2.7-3.3	262	4.3	3.8-4.9	191	2.2	1.9-2.5	432	2.8	2.6-3.1	222	3.6	3.2-4.2	210	2.3	2.0-2.7
1981	460	3.0	2.8-3.3	258	4.2	3.7-4.8	202	2.2	1.9-2.6	497	3.3	3.0-3.6	273	4.5	4.0-5.1	224	2.5	2.2-2.8
1982	435	2.8	2.6-3.1	281	4.5	4.0-5.1	154	1.7	1.4-2.0	461	3.0	2.7-3.3	280	4.6	4.0-5.2	181	2.0	1.7-2.3
1983	472	3.1	2.8-3.4	300	4.7	4.2-5.3	172	2.0	1.7-2.3	482	3.1	2.9-3.4	285	4.5	4.0-5.1	197	2.2	1.9-2.5
1984	461	3.0	2.7-3.3	269	4.2	3.7-4.8	192	2.1	1.8-2.5	477	3.1	2.8-3.4	266	4.3	3.7-4.8	211	2.3	2.0-2.6
1985	467	3.1	2.8-3.4	288	4.7	4.2-5.3	179	2.0	1.7-2.3	502	3.3	3.0-3.6	302	5.0	4.4-5.6	200	2.2	1.9-2.5
1986	497	3.3	3.0-3.6	312	5.0	4.5-5.6	185	2.1	1.8-2.4	519	3.4	3.1-3.7	293	4.7	4.2-5.3	226	2.5	2.2-2.9
1987	474	3.1	2.8-3.4	294	4.6	4.1-5.2	180	2.0	1.7-2.3	539	3.5	3.2-3.9	312	5.2	4.6-5.8	227	2.5	2.2-2.8
1988	557	3.6	3.3-4.0	351	5.6	5.0-6.2	206	2.3	2.0-2.6	555	3.6	3.3-3.9	329	5.3	4.7-5.9	226	2.5	2.1-2.8
1989	502	3.2	3.0-3.5	314	4.9	4.3-5.4	188	2.0	1.8-2.4	484	3.1	2.8-3.4	286	4.5	3.9-5.0	198	2.1	1.8-2.4
1990	590	3.9	3.5-4.2	370	5.9	5.3-6.6	220	2.4	2.1-2.7	584	3.8	3.5-4.1	346	5.7	5.1-6.4	238	2.5	2.2-2.9
1991	599	3.9	3.6-4.2	369	5.6	5.0-6.2	230	2.5	2.2-2.9	585	3.7	3.4-4.0	347	5.4	4.8-6.0	238	2.5	2.2-2.9
1992	655	4.3	3.9-4.6	386	6.1	5.5-6.8	269	3.0	2.6-3.4	620	4.0	3.7-4.3	370	5.9	5.3-6.6	250	2.7	2.4-3.1
1993	654	4.2	3.9-4.5	402	6.3	5.6-6.9	252	2.7	2.4-3.1	666	4.2	3.9-4.6	379	5.9	5.3-6.6	287	3.0	2.7-3.4
1994	662	4.3	3.9-4.6	426	6.6	6.0-7.3	236	2.5	2.2-2.9	604	3.9	3.6-4.2	362	5.8	5.2-6.4	242	2.6	2.3-3.0
1995	738	4.7	4.4-5.1	472	7.2	6.6-7.9	266	2.9	2.5-3.2	667	4.2	3.9-4.6	406	6.3	5.7-7.0	261	2.7	2.4-3.1
1996	723	4.7	4.3-5.0	467	7.2	6.5-7.9	256	2.8	2.4-3.1	645	4.1	3.8-4.4	391	6.0	5.4-6.7	254	2.6	2.3-3.0
1997	805	5.2	4.9-5.6	520	7.9	7.2-8.6	285	3.1	2.7-3.4	633	4.0	3.7-4.4	388	5.9	5.4-6.6	245	2.5	2.2-2.9
1998	900	5.7	5.4-6.1	590	8.8	8.1-9.6	310	3.3	3.0-3.7	660	4.2	3.9-4.5	402	6.0	5.5-6.7	258	2.7	2.4-3.1
1999	863	5.5	5.1-5.9	567	8.4	7.7-9.1	296	3.1	2.8-3.5	689	4.3	4.0-4.7	420	6.3	5.7-6.9	269	2.8	2.5-3.2
2000	857	5.4	5.0-5.8	566	8.2	7.6-8.9	291	3.1	2.8-3.5	684	4.2	3.9-4.6	422	6.2	5.6-6.8	262	2.7	2.4-3.1
2001	910	5.7	5.3-6.1	628	9.1	8.4-9.8	282	3.0	2.7-3.4	710	4.4	4.1-4.7	455	6.7	6.1-7.3	255	2.6	2.3-3.0
2002	902	5.6	5.2-6.0	631	9.0	8.3-9.7	271	2.9	2.6-3.3	772	4.7	4.4-5.1	498	7.2	6.6-7.9	274	2.8	2.5-3.1
2003	973	6.0	5.7-6.4	636	8.9	8.2-9.6	337	3.6	3.3-4.1	713	4.3	4.0-4.7	447	6.4	5.8-7.0	266	2.7	2.4-3.1
2004	1,014	6.2	5.8-6.6	726	10.0	9.3-10.8	288	3.1	2.7-3.5	779	4.7	4.4-5.0	481	6.7	6.1-7.3	298	3.0	2.7-3.4
2005	1,077	6.6	6.2-7.0	761	10.4	9.6-11.1	316	3.4	3.0-3.8	798	4.8	4.4-5.1	517	7.1	6.5-7.8	281	2.8	2.5-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,124	6.8	6.4-7.2	790	10.6	9.9-11.4	334	3.6	3.2-4.0	842	5.0	4.7-5.4	520	7.2	6.6-7.9	322	3.3	3.0-3.7
2007	1,181	7.1	6.7-7.5	853	11.3	10.5-12.1	328	3.5	3.1-3.9	825	4.9	4.6-5.2	550	7.5	6.9-8.1	275	2.8	2.5-3.2
2008	1,224	7.2	6.8-7.7	872	11.3	10.6-12.1	352	3.7	3.4-4.2	884	5.2	4.8-5.5	579	7.7	7.1-8.3	305	3.1	2.7-3.4
2009	1,227	7.2	6.8-7.6	860	11.1	10.3-11.8	367	3.8	3.4-4.2	909	5.3	4.9-5.6	594	7.9	7.2-8.5	315	3.2	2.8-3.6
2010	1,216	7.0	6.6-7.4	907	11.5	10.7-12.3	309	3.2	2.9-3.6	916	5.2	4.9-5.6	598	7.7	7.1-8.4	318	3.2	2.8-3.5
2011	1,272	7.2	6.8-7.6	895	10.9	10.2-11.7	377	3.9	3.5-4.4	935	5.2	4.9-5.6	590	7.4	6.8-8.0	345	3.4	3.0-3.7
2012	1,382	7.7	7.3-8.1	1,001	12.0	11.3-12.8	381	3.9	3.5-4.4	997	5.5	5.2-5.9	640	8.0	7.4-8.6	357	3.5	3.1-3.9
2013	1,447	7.9	7.5-8.3	1,009	12.0	11.3-12.8	438	4.4	4.0-4.8	1,024	5.6	5.2-5.9	689	8.4	7.8-9.1	335	3.2	2.9-3.6
2014	1,499	8.1	7.7-8.5	1,070	12.5	11.7-13.3	429	4.3	3.9-4.7	991	5.3	5.0-5.6	668	7.9	7.3-8.5	323	3.2	2.8-3.5
2015	1,497	8.0	7.5-8.4	1,062	12.3	11.5-13.1	435	4.3	3.9-4.7	1,037	5.5	5.1-5.8	679	8.1	7.5-8.7	358	3.3	3.0-3.7
2016	1,564	8.2	7.8-8.6	1,129	12.8	12.0-13.6	435	4.3	3.9-4.7	1,070	5.6	5.3-5.9	698	8.1	7.5-8.8	372	3.5	3.1-3.9
2017	1,547	8.0	7.6-8.5	1,099	12.4	11.7-13.2	448	4.3	3.9-4.7	962	5.0	4.7-5.4	662	7.7	7.1-8.3	300	2.9	2.6-3.2
2018	1,457	7.5	7.1-7.9	1,034	11.5	10.8-12.3	423	4.0	3.6-4.5	962	4.9	4.6-5.2	639	7.2	6.7-7.8	323	3.0	2.6-3.3
2019	1,502	7.7	7.3-8.1	1,040	11.5	10.8-12.3	462	4.4	4.0-4.9	983	5.0	4.7-5.3	651	7.3	6.8-7.9	332	3.0	2.7-3.4
2020	1,392	7.0	6.7-7.4	954	10.5	9.8-11.2	438	4.1	3.7-4.5	954	4.8	4.5-5.1	611	6.8	6.3-7.4	343	3.1	2.8-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,873	12.4	11.8-13.0	976	15.7	14.7-16.8	897	10.2	9.5-10.9	1,760	11.7	11.2-12.3	887	14.4	13.4-15.5	873	9.9	9.2-10.6
1977	1,926	12.9	12.4-13.6	976	16.3	15.2-17.4	950	10.7	10.1-11.5	1,850	12.4	11.8-13.0	928	15.3	14.3-16.4	922	10.4	9.8-11.1
1978	1,899	12.5	12.0-13.1	967	15.4	14.4-16.4	932	10.4	9.8-11.1	1,801	11.8	11.3-12.4	918	14.7	13.8-15.8	883	9.9	9.2-10.5
1979	1,973	13.0	12.4-13.6	1,009	16.3	15.3-17.4	964	10.7	10.0-11.4	1,892	12.5	11.9-13.0	966	15.8	14.8-16.9	926	10.2	9.5-10.9
1980	1,971	12.9	12.3-13.5	968	15.7	14.6-16.7	1,003	11.0	10.4-11.8	1,832	12.0	11.5-12.6	923	15.0	14.0-16.1	909	10.0	9.3-10.7
1981	1,887	12.4	11.9-13.0	949	15.4	14.4-16.5	938	10.4	9.8-11.1	1,803	11.9	11.3-12.4	883	14.4	13.4-15.4	920	10.1	9.5-10.8
1982	1,917	12.6	12.0-13.1	902	14.9	13.9-16.0	1,015	11.0	10.4-11.7	1,701	11.1	10.6-11.6	837	13.7	12.7-14.7	864	9.4	8.7-10.0
1983	1,821	11.8	11.3-12.4	882	14.1	13.1-15.1	939	10.2	9.6-10.9	1,783	11.5	11.0-12.1	868	13.7	12.8-14.7	915	9.9	9.3-10.6
1984	1,895	12.3	11.7-12.9	901	14.8	13.8-15.8	994	10.7	10.0-11.4	1,758	11.4	10.9-11.9	826	13.5	12.5-14.5	932	10.0	9.4-10.7
1985	1,942	12.5	11.9-13.1	901	14.2	13.3-15.2	1,041	11.2	10.5-11.9	1,718	11.1	10.6-11.6	811	13.2	12.2-14.1	907	9.8	9.1-10.4
1986	1,897	12.4	11.8-13.0	892	14.4	13.5-15.5	1,005	10.9	10.2-11.6	1,745	11.2	10.7-11.8	822	13.1	12.2-14.1	923	9.8	9.2-10.5
1987	1,893	12.2	11.6-12.7	870	14.1	13.2-15.1	1,023	10.8	10.2-11.5	1,700	11.0	10.4-11.5	817	13.2	12.3-14.1	883	9.4	8.7-10.0
1988	1,923	12.4	11.9-13.0	891	14.4	13.4-15.4	1,032	11.0	10.3-11.7	1,738	11.2	10.6-11.7	825	13.4	12.5-14.4	913	9.6	9.0-10.2
1989	1,838	11.8	11.3-12.4	875	14.0	13.0-15.0	963	10.4	9.7-11.0	1,680	10.8	10.3-11.4	813	13.1	12.2-14.1	867	9.3	8.6-9.9
1990	1,889	12.2	11.6-12.8	890	14.3	13.3-15.3	999	10.7	10.0-11.4	1,695	10.9	10.4-11.4	794	12.8	11.9-13.7	901	9.6	8.9-10.2
1991	1,948	12.5	12.0-13.1	909	14.4	13.5-15.4	1,039	11.1	10.4-11.8	1,684	10.8	10.3-11.4	789	12.7	11.8-13.6	895	9.5	8.9-10.1
1992	1,922	12.4	11.8-13.0	909	14.6	13.7-15.6	1,013	10.8	10.1-11.5	1,680	10.8	10.2-11.3	766	12.3	11.4-13.2	914	9.6	8.9-10.2
1993	1,922	12.3	11.8-12.9	902	14.2	13.2-15.1	1,020	10.8	10.2-11.5	1,798	11.4	10.9-12.0	860	13.6	12.7-14.5	938	9.9	9.2-10.5
1994	2,056	13.1	12.5-13.7	981	15.4	14.4-16.4	1,075	11.3	10.6-12.0	1,856	11.8	11.3-12.3	868	13.7	12.8-14.7	988	10.3	9.6-10.9
1995	1,961	12.4	11.9-13.0	873	13.6	12.7-14.5	1,088	11.5	10.8-12.2	1,790	11.3	10.8-11.8	821	12.7	11.9-13.7	969	10.0	9.4-10.7
1996	1,931	12.3	11.7-12.9	942	14.8	13.8-15.7	989	10.5	9.9-11.2	1,815	11.5	11.0-12.1	851	13.5	12.6-14.4	964	10.1	9.5-10.8
1997	1,988	12.6	12.0-13.1	943	14.5	13.6-15.5	1,045	11.0	10.4-11.7	1,811	11.4	10.9-11.9	871	13.7	12.8-14.6	940	9.8	9.1-10.4
1998	2,088	13.1	12.6-13.7	999	15.2	14.3-16.2	1,089	11.5	10.8-12.2	1,797	11.2	10.7-11.8	861	13.3	12.4-14.2	936	9.8	9.1-10.4
1999	2,054	12.8	12.2-13.3	982	14.8	13.9-15.8	1,072	11.2	10.5-11.9	1,867	11.6	11.1-12.1	927	14.0	13.1-14.9	940	9.8	9.1-10.4
2000	2,113	13.0	12.5-13.6	982	14.7	13.8-15.6	1,131	11.8	11.1-12.5	1,822	11.1	10.6-11.7	831	12.5	11.7-13.4	991	10.1	9.4-10.7
2001	2,073	12.7	12.1-13.2	937	13.8	12.9-14.7	1,136	11.7	11.0-12.4	1,859	11.3	10.8-11.8	849	12.7	11.9-13.6	1,010	10.2	9.6-10.9
2002	2,115	12.8	12.3-13.4	991	14.4	13.5-15.3	1,124	11.5	10.9-12.2	1,829	11.1	10.5-11.6	850	12.4	11.6-13.3	979	9.9	9.3-10.5
2003	2,200	13.2	12.7-13.8	1,046	15.1	14.2-16.1	1,154	11.8	11.1-12.5	1,829	10.9	10.5-11.5	855	12.4	11.6-13.3	974	9.8	9.1-10.4
2004	2,180	13.0	12.5-13.6	1,027	14.7	13.8-15.6	1,153	11.8	11.1-12.5	1,863	11.1	10.6-11.6	906	13.0	12.2-13.9	957	9.7	9.0-10.3
2005	2,182	13.0	12.5-13.6	1,072	15.1	14.2-16.0	1,110	11.3	10.7-12.0	1,899	11.2	10.7-11.8	900	12.8	12.0-13.7	999	10.0	9.4-10.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	2,238	13.3	12.8-13.9	1,062	14.9	14.0-15.9	1,176	12.0	11.3-12.7	1,900	11.2	10.7-11.7	914	12.9	12.1-13.8	986	9.8	9.2-10.5
2007	2,323	13.6	13.1-14.2	1,133	15.7	14.8-16.6	1,190	12.0	11.3-12.7	1,908	11.1	10.6-11.6	922	12.8	12.0-13.7	986	9.9	9.3-10.5
2008	2,453	14.3	13.8-14.9	1,195	16.4	15.5-17.4	1,258	12.6	11.9-13.4	1,987	11.5	11.0-12.0	938	12.9	12.1-13.8	1,049	10.2	9.6-10.9
2009	2,448	14.1	13.6-14.7	1,252	16.8	15.9-17.8	1,196	12.0	11.3-12.7	2,011	11.5	11.0-12.0	1,015	13.7	12.9-14.6	996	9.8	9.2-10.5
2010	2,470	14.2	13.6-14.8	1,196	15.9	15.0-16.8	1,274	12.7	12.0-13.5	2,039	11.5	11.0-12.0	987	13.0	12.2-13.9	1,052	10.1	9.5-10.8
2011	2,496	14.1	13.6-14.7	1,258	16.4	15.5-17.4	1,238	12.2	11.5-12.9	2,015	11.3	10.8-11.8	986	13.0	12.2-13.8	1,029	9.9	9.3-10.6
2012	2,521	14.1	13.6-14.7	1,275	16.5	15.6-17.5	1,246	12.3	11.6-13.0	2,072	11.5	11.0-12.0	1,066	13.8	13.0-14.7	1,006	9.7	9.1-10.3
2013	2,557	14.0	13.5-14.6	1,279	16.1	15.2-17.0	1,278	12.3	11.6-13.0	2,058	11.2	10.7-11.7	1,028	13.1	12.3-13.9	1,030	9.7	9.1-10.4
2014	2,666	14.5	14.0-15.1	1,365	16.8	15.9-17.8	1,301	12.6	11.9-13.3	2,094	11.2	10.8-11.7	1,066	13.3	12.5-14.1	1,028	9.5	9.0-10.2
2015	2,638	14.2	13.6-14.7	1,327	16.1	15.3-17.1	1,311	12.5	11.8-13.2	1,998	10.6	10.1-11.1	1,004	12.3	11.5-13.1	994	9.1	8.5-9.7
2016	2,718	14.6	14.0-15.2	1,378	16.6	15.7-17.6	1,340	13.0	12.3-13.7	2,140	11.3	10.8-11.8	1,056	12.7	12.0-13.5	1,084	10.1	9.5-10.8
2017	2,874	15.1	14.5-15.7	1,510	17.9	17.0-18.9	1,364	12.8	12.1-13.6	2,164	11.2	10.8-11.7	1,090	13.0	12.2-13.8	1,074	9.8	9.2-10.4
2018	2,827	14.8	14.2-15.3	1,421	16.6	15.7-17.5	1,406	13.2	12.5-13.9	2,122	10.8	10.4-11.3	1,080	12.6	11.9-13.4	1,042	9.4	8.8-10.0
2019	2,866	14.8	14.2-15.3	1,445	16.7	15.8-17.6	1,421	13.1	12.4-13.8	2,190	11.1	10.7-11.6	1,128	13.0	12.2-13.8	1,062	9.5	8.9-10.1
2020	2,796	14.3	13.7-14.8	1,463	16.6	15.7-17.5	1,333	12.3	11.6-13.0	2,148	10.8	10.4-11.3	1,079	12.4	11.6-13.1	1,069	9.5	9.0-10.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Larynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	857	5.3	5.0-5.7	720	10.5	9.7-11.3	137	1.5	1.3-1.8	312	2.0	1.8-2.2	253	3.9	3.4-4.4	59	0.7	0.5-0.9
1977	846	5.3	5.0-5.7	704	10.4	9.6-11.2	142	1.6	1.3-1.9	309	1.9	1.7-2.2	266	4.0	3.6-4.6	43	0.5	0.3-0.6
1978	865	5.5	5.1-5.9	710	10.6	9.8-11.5	155	1.7	1.5-2.0	312	2.0	1.8-2.2	247	3.8	3.3-4.4	65	0.7	0.6-0.9
1979	926	5.8	5.4-6.2	740	10.8	10.0-11.6	186	2.1	1.8-2.5	340	2.1	1.9-2.4	276	4.2	3.7-4.8	64	0.7	0.5-0.9
1980	946	6.0	5.7-6.5	776	11.6	10.8-12.5	170	2.0	1.7-2.3	306	1.9	1.7-2.2	252	3.9	3.4-4.5	54	0.6	0.5-0.8
1981	970	6.2	5.8-6.6	785	11.8	10.9-12.7	185	2.1	1.8-2.4	298	1.9	1.6-2.1	248	3.8	3.3-4.3	50	0.5	0.4-0.7
1982	896	5.7	5.4-6.1	717	10.8	10.0-11.6	179	2.1	1.8-2.4	312	1.9	1.7-2.2	249	3.8	3.3-4.3	63	0.7	0.5-0.9
1983	884	5.7	5.3-6.1	697	10.3	9.5-11.1	187	2.2	1.9-2.6	292	1.8	1.6-2.1	222	3.4	3.0-3.9	70	0.8	0.6-1.0
1984	932	6.0	5.6-6.4	732	10.9	10.1-11.7	200	2.3	2.0-2.7	268	1.7	1.5-1.9	211	3.2	2.8-3.7	57	0.6	0.5-0.8
1985	936	6.0	5.6-6.4	734	10.9	10.1-11.7	202	2.4	2.0-2.7	296	1.8	1.6-2.1	232	3.5	3.0-4.0	64	0.7	0.5-0.9
1986	884	5.7	5.3-6.1	683	10.3	9.5-11.1	201	2.4	2.0-2.7	248	1.6	1.4-1.8	197	3.1	2.7-3.6	51	0.5	0.4-0.7
1987	904	5.9	5.5-6.3	719	10.9	10.1-11.8	185	2.2	1.9-2.5	275	1.8	1.6-2.0	216	3.4	2.9-3.9	59	0.6	0.5-0.8
1988	859	5.6	5.2-5.9	663	9.9	9.1-10.7	196	2.3	2.0-2.7	267	1.7	1.5-2.0	211	3.3	2.9-3.8	56	0.6	0.5-0.8
1989	901	5.9	5.6-6.4	688	10.5	9.7-11.3	213	2.6	2.2-2.9	267	1.7	1.5-1.9	186	2.8	2.4-3.3	81	0.9	0.7-1.1
1990	923	6.0	5.6-6.4	713	10.7	9.9-11.5	210	2.5	2.1-2.8	237	1.6	1.4-1.8	196	3.1	2.7-3.6	41	0.4	0.3-0.6
1991	841	5.6	5.2-5.9	675	10.3	9.5-11.1	166	1.9	1.7-2.3	256	1.6	1.4-1.9	198	3.0	2.6-3.5	58	0.6	0.5-0.8
1992	856	5.6	5.3-6.0	651	9.8	9.1-10.6	205	2.5	2.1-2.8	314	2.0	1.8-2.3	239	3.8	3.3-4.3	75	0.8	0.7-1.1
1993	813	5.3	5.0-5.7	606	9.2	8.5-10.0	207	2.4	2.1-2.8	267	1.7	1.5-1.9	212	3.3	2.8-3.8	55	0.6	0.5-0.8
1994	788	5.2	4.8-5.5	609	9.3	8.5-10.1	179	2.1	1.8-2.4	264	1.7	1.5-1.9	203	3.2	2.7-3.6	61	0.7	0.5-0.9
1995	760	5.0	4.6-5.4	583	8.7	8.0-9.4	177	2.1	1.8-2.4	255	1.7	1.5-1.9	195	3.0	2.6-3.4	60	0.7	0.5-0.8
1996	765	5.0	4.7-5.4	595	8.9	8.2-9.7	170	2.0	1.7-2.4	264	1.7	1.5-1.9	205	3.1	2.7-3.6	59	0.7	0.5-0.8
1997	741	4.9	4.5-5.2	587	8.8	8.1-9.5	154	1.8	1.6-2.2	216	1.4	1.2-1.6	161	2.5	2.1-2.9	55	0.6	0.5-0.8
1998	708	4.6	4.2-4.9	522	7.7	7.0-8.4	186	2.2	1.9-2.5	234	1.5	1.3-1.7	178	2.7	2.3-3.1	56	0.6	0.5-0.8
1999	736	4.7	4.4-5.1	590	8.6	7.9-9.3	146	1.7	1.4-2.0	232	1.5	1.3-1.7	176	2.6	2.2-3.0	56	0.6	0.5-0.8
2000	735	4.7	4.3-5.0	572	8.3	7.6-9.0	163	1.9	1.6-2.2	225	1.4	1.2-1.6	181	2.7	2.3-3.1	44	0.5	0.4-0.7
2001	699	4.4	4.1-4.8	538	7.7	7.1-8.4	161	1.8	1.6-2.2	199	1.2	1.1-1.4	163	2.4	2.0-2.8	36	0.4	0.3-0.5
2002	674	4.2	3.9-4.5	535	7.6	7.0-8.3	139	1.6	1.3-1.9	223	1.4	1.2-1.5	176	2.6	2.2-3.0	47	0.5	0.4-0.7
2003	677	4.2	3.9-4.5	526	7.4	6.7-8.0	151	1.7	1.4-2.0	207	1.3	1.1-1.4	156	2.2	1.9-2.6	51	0.5	0.4-0.7
2004	681	4.2	3.9-4.5	554	7.6	7.0-8.2	127	1.4	1.2-1.7	197	1.2	1.0-1.4	159	2.3	1.9-2.6	38	0.4	0.3-0.5
2005	719	4.4	4.1-4.7	560	7.6	7.0-8.3	159	1.8	1.5-2.1	209	1.3	1.1-1.5	155	2.2	1.9-2.6	54	0.6	0.4-0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Larynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	676	4.1	3.8-4.5	522	7.2	6.6-7.8	154	1.8	1.5-2.1	207	1.2	1.1-1.4	161	2.3	1.9-2.6	46	0.5	0.4-0.7
2007	639	3.8	3.5-4.1	504	6.8	6.2-7.4	135	1.4	1.2-1.7	189	1.1	1.0-1.3	154	2.1	1.8-2.4	35	0.4	0.2-0.5
2008	724	4.3	4.0-4.7	568	7.6	6.9-8.2	156	1.7	1.5-2.0	173	1.0	0.9-1.2	140	1.9	1.6-2.2	33	0.3	0.2-0.5
2009	656	3.9	3.6-4.2	505	6.6	6.0-7.2	151	1.7	1.4-1.9	196	1.1	1.0-1.3	145	1.9	1.6-2.3	51	0.5	0.4-0.7
2010	631	3.7	3.4-4.0	502	6.5	5.9-7.1	129	1.4	1.2-1.7	204	1.1	1.0-1.3	153	2.0	1.7-2.4	51	0.5	0.4-0.7
2011	596	3.4	3.1-3.6	484	6.0	5.4-6.5	112	1.2	1.0-1.5	192	1.1	0.9-1.3	144	1.9	1.6-2.2	48	0.5	0.3-0.6
2012	652	3.7	3.4-4.0	520	6.5	5.9-7.1	132	1.4	1.1-1.6	174	1.0	0.8-1.1	135	1.7	1.4-2.0	39	0.4	0.3-0.6
2013	650	3.6	3.3-3.9	525	6.4	5.8-7.0	125	1.3	1.1-1.6	209	1.1	1.0-1.3	170	2.1	1.8-2.4	39	0.4	0.3-0.6
2014	616	3.4	3.1-3.7	479	5.7	5.2-6.3	137	1.4	1.2-1.7	179	1.0	0.8-1.1	143	1.7	1.5-2.1	36	0.4	0.3-0.5
2015	629	3.4	3.1-3.7	498	5.9	5.4-6.5	131	1.3	1.1-1.6	165	0.9	0.7-1.0	126	1.5	1.3-1.8	39	0.4	0.3-0.5
2016	579	3.1	2.9-3.4	470	5.6	5.1-6.1	109	1.1	0.9-1.4	183	1.0	0.8-1.1	152	1.8	1.6-2.2	31	0.3	0.2-0.4
2017	599	3.2	2.9-3.5	467	5.4	4.9-5.9	132	1.4	1.1-1.6	148	0.7	0.6-0.9	117	1.3	1.1-1.6	31	0.3	0.2-0.4
2018	556	2.9	2.6-3.1	433	4.8	4.4-5.3	123	1.2	1.0-1.5	166	0.8	0.7-1.0	127	1.5	1.2-1.7	39	0.4	0.3-0.5
2019	561	2.9	2.7-3.2	448	5.0	4.5-5.5	113	1.1	0.9-1.4	155	0.8	0.7-0.9	126	1.4	1.2-1.7	29	0.3	0.2-0.4
2020	472	2.4	2.2-2.7	373	4.1	3.7-4.6	99	1.0	0.8-1.2	130	0.6	0.5-0.8	104	1.2	0.9-1.4	26	0.2	0.1-0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	8,493	54.1	53.0-55.3	6,065	92.7	90.3-95.2	2,428	27.5	26.4-28.6	6,853	43.9	42.8-44.9	4,988	77.5	75.2-79.8	1,865	21.1	20.1-22.1
1977	8,493	54.1	53.0-55.3	5,964	91.6	89.2-94.1	2,529	28.7	27.5-29.8	7,129	45.6	44.6-46.7	5,157	80.1	77.9-82.4	1,972	22.3	21.3-23.3
1978	8,644	55.2	54.0-56.4	6,150	94.8	92.3-97.2	2,494	28.3	27.2-29.5	7,125	45.7	44.6-46.8	5,103	79.7	77.4-82.0	2,022	22.8	21.8-23.8
1979	8,847	56.6	55.4-57.8	6,064	93.8	91.4-96.3	2,783	31.6	30.4-32.9	7,070	45.3	44.2-46.3	4,958	77.5	75.3-79.8	2,112	23.7	22.7-24.8
1980	9,144	58.3	57.1-59.5	6,291	96.7	94.3-99.2	2,853	32.1	30.9-33.3	7,557	48.2	47.1-49.4	5,268	82.0	79.7-84.4	2,289	25.6	24.5-26.7
1981	9,073	58.0	56.8-59.2	6,076	94.3	91.9-96.8	2,997	33.9	32.6-35.1	7,423	47.5	46.4-48.6	5,159	80.7	78.4-83.0	2,264	25.4	24.4-26.5
1982	9,486	60.6	59.3-61.8	6,240	96.8	94.3-99.3	3,246	36.6	35.3-37.9	7,622	48.8	47.7-49.9	5,096	80.1	77.9-82.4	2,526	28.3	27.2-29.5
1983	9,773	62.3	61.0-63.5	6,330	97.7	95.3-100.3	3,443	38.5	37.2-39.9	7,763	49.5	48.4-50.6	5,066	79.7	77.5-82.0	2,697	29.9	28.7-31.0
1984	9,711	62.1	60.9-63.4	6,275	96.8	94.3-99.3	3,436	39.0	37.6-40.3	7,751	49.5	48.4-50.6	5,095	79.6	77.3-81.9	2,656	29.6	28.4-30.8
1985	10,032	63.9	62.7-65.2	6,356	97.8	95.3-100.3	3,676	41.5	40.1-42.9	8,100	51.7	50.5-52.8	5,199	81.4	79.1-83.7	2,901	32.2	31.0-33.5
1986	9,929	63.6	62.4-64.9	6,201	96.0	93.5-98.5	3,728	42.1	40.7-43.5	8,051	51.3	50.2-52.4	5,081	79.4	77.2-81.7	2,970	33.2	32.0-34.5
1987	10,166	65.0	63.8-66.3	6,273	96.7	94.3-99.2	3,893	43.9	42.5-45.3	8,098	51.8	50.6-52.9	5,135	80.5	78.2-82.8	2,963	32.9	31.7-34.1
1988	10,269	65.7	64.5-67.1	6,206	96.4	94.0-98.9	4,063	45.7	44.3-47.2	8,246	52.7	51.6-53.9	5,146	80.6	78.3-82.9	3,100	34.6	33.4-35.9
1989	10,374	66.7	65.4-68.0	6,163	95.1	92.6-97.5	4,211	47.7	46.2-49.2	8,239	52.9	51.7-54.0	5,046	79.0	76.7-81.2	3,193	35.7	34.4-37.0
1990	10,427	66.9	65.6-68.2	6,208	96.1	93.7-98.6	4,219	47.5	46.0-49.0	8,367	53.5	52.4-54.7	5,022	78.4	76.2-80.7	3,345	37.1	35.8-38.4
1991	10,566	68.1	66.8-69.4	6,105	94.5	92.1-97.0	4,461	50.4	48.9-52.0	8,594	55.1	54.0-56.3	5,088	79.7	77.4-81.9	3,506	39.0	37.7-40.3
1992	10,731	68.9	67.6-70.2	6,174	95.1	92.6-97.5	4,557	51.4	49.9-52.9	8,567	54.8	53.6-56.0	4,988	77.9	75.7-80.2	3,579	39.6	38.3-40.9
1993	10,616	68.2	66.9-69.5	5,943	91.3	89.0-93.7	4,673	52.7	51.2-54.3	8,419	53.8	52.6-54.9	4,832	75.3	73.1-77.4	3,587	39.8	38.5-41.1
1994	10,962	70.3	69.0-71.6	6,073	93.0	90.6-95.4	4,889	55.3	53.7-56.9	8,662	55.2	54.1-56.4	5,075	78.8	76.6-81.0	3,587	39.6	38.3-41.0
1995	10,801	69.1	67.8-70.4	6,022	91.6	89.3-94.0	4,779	53.5	52.0-55.1	8,598	54.7	53.6-55.9	4,904	75.7	73.5-77.9	3,694	40.8	39.4-42.1
1996	10,892	69.8	68.5-71.2	6,046	92.2	89.9-94.6	4,846	54.6	53.1-56.2	8,436	53.8	52.7-55.0	4,782	73.6	71.5-75.8	3,654	40.4	39.1-41.7
1997	10,932	69.8	68.5-71.2	5,944	90.0	87.7-92.4	4,988	56.1	54.5-57.7	8,198	52.0	50.9-53.2	4,587	70.3	68.3-72.4	3,611	39.7	38.4-41.0
1998	11,026	69.9	68.6-71.3	5,908	88.2	85.9-90.5	5,118	57.2	55.6-58.8	8,248	52.0	50.9-53.1	4,582	69.0	67.0-71.1	3,666	40.1	38.8-41.4
1999	10,985	69.1	67.8-70.4	5,840	86.7	84.5-89.0	5,145	57.0	55.5-58.6	8,174	51.1	50.0-52.2	4,450	66.8	64.8-68.8	3,724	40.6	39.3-41.9
2000	11,266	70.2	68.9-71.6	5,938	87.4	85.2-89.6	5,328	58.5	56.9-60.1	8,293	51.3	50.2-52.4	4,513	67.1	65.1-69.1	3,780	40.4	39.1-41.7
2001	11,018	68.3	67.1-69.6	5,673	83.0	80.8-85.2	5,345	58.5	56.9-60.1	8,204	50.5	49.4-51.6	4,447	65.9	63.9-67.9	3,757	40.2	38.9-41.5
2002	11,225	69.0	67.8-70.3	5,895	85.2	83.1-87.5	5,330	58.0	56.4-59.6	8,280	50.6	49.5-51.7	4,411	64.3	62.4-66.3	3,869	41.2	39.9-42.5
2003	11,184	68.4	67.2-69.7	5,885	84.3	82.2-86.5	5,299	57.4	55.8-59.0	8,067	49.0	47.9-50.0	4,301	62.1	60.3-64.0	3,766	39.8	38.6-41.1
2004	11,181	67.9	66.7-69.2	5,720	81.3	79.1-83.4	5,461	58.8	57.2-60.4	7,967	48.1	47.0-49.1	4,160	59.6	57.8-61.5	3,807	40.1	38.8-41.4
2005	11,235	68.1	66.9-69.4	5,751	81.3	79.2-83.5	5,484	59.0	57.5-60.6	7,880	47.5	46.4-48.5	4,174	59.7	57.9-61.5	3,706	39.0	37.7-40.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	11,457	69.6	68.3-70.9	5,686	80.0	77.9-82.1	5,771	62.6	60.9-64.2	7,850	47.2	46.2-48.3	4,183	59.3	57.5-61.1	3,667	38.7	37.5-40.0
2007	11,504	69.1	67.8-70.4	5,821	81.0	78.9-83.1	5,683	61.0	59.4-62.6	7,930	47.2	46.1-48.2	4,102	57.3	55.6-59.1	3,828	40.1	38.8-41.4
2008	11,700	69.6	68.3-70.9	5,842	80.3	78.3-82.4	5,858	62.4	60.8-64.0	7,866	46.2	45.2-47.2	4,082	56.4	54.7-58.2	3,784	39.2	37.9-40.5
2009	11,579	68.1	66.8-69.3	5,866	79.5	77.4-81.6	5,713	60.1	58.5-61.7	7,565	44.2	43.2-45.2	3,924	53.6	51.9-55.3	3,641	37.6	36.3-38.8
2010	11,313	66.0	64.7-67.2	5,609	74.7	72.8-76.7	5,704	60.0	58.4-61.6	7,595	43.9	42.9-44.9	3,873	52.1	50.4-53.8	3,722	38.1	36.9-39.4
2011	11,552	66.3	65.1-67.5	5,724	75.1	73.1-77.1	5,828	60.2	58.6-61.8	7,472	42.5	41.5-43.5	3,829	50.7	49.1-52.4	3,643	36.6	35.4-37.8
2012	11,400	64.5	63.3-65.7	5,607	72.3	70.4-74.3	5,793	59.3	57.7-60.9	7,621	42.9	42.0-43.9	3,993	52.0	50.4-53.7	3,628	36.5	35.3-37.7
2013	11,543	64.5	63.3-65.7	5,636	71.3	69.4-73.2	5,907	59.7	58.2-61.3	7,407	40.9	40.0-41.9	3,783	48.1	46.6-49.7	3,624	35.7	34.5-36.9
2014	11,218	61.4	60.3-62.6	5,468	67.9	66.1-69.7	5,750	57.0	55.5-58.5	6,961	38.0	37.1-38.9	3,580	44.9	43.4-46.4	3,381	32.9	31.8-34.1
2015	11,645	63.2	62.0-64.3	5,631	69.1	67.3-71.0	6,014	59.2	57.7-60.8	6,970	37.4	36.5-38.3	3,559	43.9	42.5-45.4	3,411	32.6	31.5-33.8
2016	11,514	61.4	60.3-62.6	5,545	66.6	64.9-68.5	5,969	58.1	56.6-59.6	6,560	34.8	34.0-35.7	3,313	40.5	39.1-41.9	3,247	30.7	29.6-31.8
2017	11,927	62.8	61.7-64.0	5,694	67.4	65.6-69.2	6,233	59.8	58.3-61.4	6,320	32.9	32.0-33.7	3,186	37.9	36.6-39.2	3,134	29.1	28.1-30.2
2018	11,643	60.3	59.2-61.4	5,480	63.4	61.7-65.1	6,163	58.4	56.9-59.9	6,172	31.8	31.0-32.6	3,103	36.4	35.1-37.7	3,069	28.3	27.3-29.4
2019	11,548	58.7	57.6-59.8	5,404	61.3	59.7-63.0	6,144	57.2	55.8-58.7	6,023	30.3	29.6-31.1	3,000	34.5	33.2-35.8	3,023	27.3	26.3-28.3
2020	10,123	51.1	50.1-52.1	4,739	53.5	51.9-55.1	5,384	49.5	48.2-50.9	5,543	27.7	27.0-28.5	2,802	31.8	30.7-33.1	2,741	24.7	23.8-25.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	922	6.3	5.9-6.7	473	7.2	6.6-7.9	449	5.6	5.1-6.2	379	2.6	2.3-2.9	211	3.3	2.8-3.7	168	2.1	1.8-2.4
1977	1,077	7.3	6.9-7.8	546	8.3	7.6-9.0	531	6.6	6.0-7.2	388	2.6	2.4-2.9	231	3.6	3.2-4.1	157	1.9	1.6-2.3
1978	1,011	7.0	6.6-7.5	503	7.8	7.1-8.5	508	6.6	6.0-7.2	332	2.3	2.0-2.5	208	3.3	2.9-3.8	124	1.5	1.2-1.8
1979	1,067	7.4	7.0-7.9	555	8.5	7.8-9.3	512	6.6	6.0-7.2	376	2.6	2.3-2.9	208	3.3	2.8-3.8	168	2.1	1.8-2.4
1980	1,245	8.6	8.1-9.1	662	10.3	9.5-11.1	583	7.5	6.8-8.1	387	2.6	2.4-2.9	222	3.4	3.0-3.9	165	2.0	1.7-2.4
1981	1,181	8.2	7.7-8.6	608	9.4	8.6-10.2	573	7.3	6.7-7.9	379	2.6	2.3-2.9	219	3.4	3.0-3.9	160	2.0	1.7-2.3
1982	1,227	8.4	8.0-8.9	618	9.6	8.8-10.4	609	7.7	7.1-8.3	382	2.6	2.3-2.9	230	3.6	3.1-4.1	152	1.8	1.5-2.1
1983	1,349	9.3	8.8-9.8	723	11.2	10.4-12.1	626	8.1	7.4-8.7	377	2.6	2.3-2.8	213	3.4	2.9-3.9	164	2.1	1.8-2.4
1984	1,289	8.8	8.3-9.3	686	10.5	9.7-11.3	603	7.7	7.0-8.3	412	2.8	2.5-3.1	238	3.7	3.2-4.2	174	2.1	1.8-2.4
1985	1,338	9.2	8.7-9.7	732	11.3	10.4-12.1	606	7.7	7.1-8.4	417	2.8	2.6-3.1	239	3.8	3.3-4.3	178	2.1	1.8-2.5
1986	1,412	9.6	9.1-10.2	737	11.4	10.5-12.3	675	8.6	8.0-9.3	412	2.8	2.5-3.1	230	3.6	3.1-4.1	182	2.2	1.9-2.6
1987	1,340	9.2	8.7-9.7	729	11.3	10.5-12.2	611	7.7	7.0-8.3	438	3.0	2.7-3.3	262	4.1	3.6-4.6	176	2.2	1.9-2.6
1988	1,276	8.7	8.2-9.2	717	11.2	10.3-12.0	559	6.9	6.3-7.5	416	2.8	2.5-3.1	260	4.1	3.6-4.6	156	1.8	1.6-2.2
1989	1,328	9.1	8.6-9.6	728	11.3	10.4-12.1	600	7.4	6.8-8.1	417	2.8	2.6-3.1	266	4.3	3.8-4.8	151	1.8	1.5-2.1
1990	1,367	9.2	8.7-9.7	736	11.2	10.4-12.1	631	7.9	7.3-8.5	412	2.8	2.5-3.1	246	3.8	3.4-4.4	166	2.0	1.7-2.4
1991	1,313	8.8	8.3-9.3	749	11.4	10.6-12.2	564	6.9	6.3-7.5	410	2.7	2.5-3.0	245	3.9	3.4-4.4	165	1.9	1.6-2.3
1992	1,282	8.6	8.1-9.0	723	11.0	10.2-11.8	559	6.8	6.2-7.4	389	2.6	2.3-2.8	230	3.5	3.1-4.0	159	1.8	1.5-2.1
1993	1,295	8.5	8.1-9.0	739	11.1	10.3-12.0	556	6.7	6.1-7.3	374	2.5	2.2-2.7	232	3.7	3.2-4.2	142	1.6	1.4-2.0
1994	1,454	9.6	9.1-10.2	844	12.7	11.9-13.6	610	7.4	6.8-8.0	405	2.7	2.4-2.9	257	4.0	3.5-4.5	148	1.7	1.4-2.0
1995	1,546	10.3	9.8-10.8	918	13.7	12.8-14.6	628	7.7	7.1-8.3	426	2.8	2.5-3.1	248	3.8	3.4-4.4	178	2.0	1.7-2.4
1996	1,654	10.9	10.4-11.5	928	13.9	13.0-14.9	726	8.8	8.2-9.5	431	2.8	2.6-3.1	254	3.8	3.3-4.3	177	2.1	1.8-2.4
1997	1,774	11.7	11.1-12.2	1,022	15.3	14.4-16.3	752	9.1	8.5-9.8	412	2.7	2.4-2.9	264	4.0	3.6-4.6	148	1.7	1.4-2.0
1998	1,870	12.3	11.7-12.9	1,047	15.5	14.5-16.4	823	10.1	9.4-10.8	418	2.7	2.4-2.9	259	3.9	3.4-4.4	159	1.8	1.5-2.1
1999	1,905	12.4	11.8-13.0	1,087	15.9	15.0-16.9	818	9.8	9.2-10.5	390	2.5	2.3-2.8	231	3.4	3.0-3.9	159	1.8	1.5-2.1
2000	2,168	14.0	13.4-14.6	1,216	17.7	16.7-18.7	952	11.5	10.8-12.2	428	2.7	2.4-3.0	268	3.9	3.5-4.4	160	1.8	1.5-2.1
2001	2,344	15.1	14.5-15.7	1,305	18.8	17.8-19.8	1,039	12.5	11.7-13.3	468	2.9	2.7-3.2	289	4.3	3.8-4.8	179	2.0	1.7-2.3
2002	2,333	14.8	14.3-15.5	1,326	19.0	18.0-20.0	1,007	12.0	11.3-12.8	412	2.6	2.3-2.8	218	3.1	2.7-3.6	194	2.2	1.9-2.5
2003	2,322	14.7	14.1-15.3	1,321	18.7	17.7-19.8	1,001	11.8	11.1-12.6	419	2.6	2.3-2.8	241	3.5	3.0-3.9	178	1.9	1.7-2.3
2004	2,863	18.1	17.5-18.8	1,611	22.7	21.6-23.8	1,252	15.0	14.2-15.9	421	2.6	2.3-2.8	250	3.5	3.1-4.0	171	1.9	1.6-2.2
2005	3,143	19.9	19.2-20.7	1,794	25.2	24.1-26.4	1,349	16.2	15.3-17.1	431	2.6	2.4-2.9	258	3.7	3.2-4.2	173	1.9	1.6-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	3,490	22.1	21.3-22.8	1,987	27.7	26.5-29.0	1,503	18.1	17.1-19.0	450	2.7	2.5-3.0	288	4.1	3.6-4.6	162	1.8	1.5-2.1
2007	3,220	20.2	19.5-20.9	1,816	25.1	24.0-26.3	1,404	16.9	16.0-17.8	433	2.6	2.4-2.9	298	4.2	3.7-4.7	135	1.4	1.2-1.7
2008	3,474	21.6	20.9-22.3	1,981	27.3	26.1-28.5	1,493	17.8	16.8-18.7	445	2.7	2.4-2.9	285	3.9	3.5-4.4	160	1.7	1.4-2.0
2009	3,729	22.9	22.2-23.7	2,146	29.1	27.8-30.3	1,583	18.7	17.8-19.7	447	2.7	2.4-2.9	269	3.7	3.3-4.2	178	1.9	1.6-2.2
2010	3,490	21.4	20.7-22.1	1,974	26.5	25.4-27.8	1,516	17.7	16.8-18.7	448	2.6	2.4-2.9	271	3.6	3.2-4.1	177	1.9	1.6-2.2
2011	3,949	23.6	22.9-24.4	2,266	29.8	28.6-31.1	1,683	19.3	18.4-20.3	450	2.6	2.3-2.8	284	3.8	3.3-4.2	166	1.7	1.4-2.0
2012	3,743	22.3	21.6-23.0	2,108	27.3	26.2-28.6	1,635	18.8	17.9-19.8	518	2.9	2.7-3.2	319	4.2	3.8-4.7	199	2.0	1.7-2.3
2013	4,154	24.5	23.7-25.2	2,384	30.4	29.2-31.7	1,770	20.3	19.4-21.4	480	2.7	2.5-3.0	316	4.1	3.7-4.6	164	1.6	1.4-1.9
2014	4,101	24.0	23.3-24.8	2,358	30.0	28.7-31.2	1,743	19.8	18.8-20.8	434	2.4	2.2-2.6	284	3.6	3.2-4.0	150	1.4	1.2-1.7
2015	4,039	23.2	22.5-24.0	2,360	29.5	28.3-30.8	1,679	18.7	17.8-19.7	359	2.0	1.8-2.2	233	2.9	2.5-3.3	126	1.3	1.1-1.6
2016	4,094	23.4	22.6-24.1	2,398	29.6	28.4-30.8	1,696	19.0	18.1-20.0	399	2.1	1.9-2.4	248	3.0	2.7-3.5	151	1.4	1.2-1.7
2017	4,143	23.6	22.9-24.4	2,444	30.1	28.9-31.4	1,699	19.0	18.1-20.0	377	2.0	1.8-2.2	239	2.9	2.6-3.4	138	1.3	1.1-1.5
2018	4,141	23.1	22.3-23.8	2,386	28.6	27.4-29.8	1,755	19.1	18.1-20.0	375	1.9	1.7-2.1	230	2.7	2.4-3.1	145	1.3	1.1-1.6
2019	4,271	23.7	23.0-24.5	2,483	29.4	28.3-30.6	1,788	19.8	18.8-20.8	344	1.8	1.6-2.0	219	2.7	2.3-3.1	125	1.2	1.0-1.4
2020	3,603	19.9	19.2-20.5	2,036	24.1	23.0-25.2	1,567	16.9	16.0-17.8	365	1.9	1.7-2.2	236	2.9	2.5-3.3	129	1.2	1.0-1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	8,578	104.1	101.9-106.4	3,310	38.8	37.5-40.2
1977	8,419	102.4	100.1-104.6	3,351	39.5	38.2-40.9
1978	8,570	104.2	102.0-106.5	3,359	39.4	38.0-40.8
1979	8,449	102.8	100.6-105.1	3,196	37.8	36.4-39.1
1980	8,564	104.2	102.0-106.5	3,300	38.5	37.1-39.9
1981	8,752	106.7	104.4-109.1	3,267	38.2	36.8-39.5
1982	8,789	106.7	104.4-109.0	3,212	37.4	36.1-38.8
1983	9,149	111.1	108.8-113.5	3,178	37.0	35.7-38.4
1984	9,509	115.0	112.6-117.4	3,312	38.5	37.2-39.9
1985	9,827	119.5	117.0-121.9	3,315	38.6	37.2-40.0
1986	9,891	120.0	117.5-122.4	3,297	38.1	36.8-39.5
1987	10,213	123.4	120.9-125.9	3,265	37.5	36.2-38.9
1988	10,342	125.0	122.5-127.5	3,327	38.4	37.1-39.8
1989	10,155	123.4	121.0-125.9	3,255	37.9	36.6-39.2
1990	10,790	131.4	128.8-133.9	3,156	36.4	35.1-37.7
1991	10,730	130.2	127.7-132.8	3,040	35.1	33.9-36.4
1992	10,766	130.7	128.2-133.3	3,020	34.7	33.4-35.9
1993	10,571	127.9	125.5-130.4	3,013	34.2	33.0-35.5
1994	11,004	132.9	130.3-135.4	3,007	34.2	33.0-35.5
1995	11,254	136.3	133.7-138.9	2,963	33.8	32.5-35.0
1996	11,884	143.7	141.0-146.3	3,013	34.0	32.8-35.3
1997	11,869	142.5	139.9-145.1	2,750	31.0	29.8-32.2
1998	12,205	145.5	142.9-148.2	2,706	30.1	29.0-31.3
1999	12,113	143.1	140.5-145.7	2,558	28.1	27.0-29.2
2000	12,086	141.3	138.8-143.9	2,571	27.9	26.8-29.0
2001	11,788	137.3	134.8-139.8	2,536	27.5	26.4-28.6
2002	11,714	135.7	133.2-138.2	2,446	26.1	25.1-27.2
2003	11,580	133.5	131.1-136.0	2,420	25.8	24.8-26.9
2004	11,407	131.2	128.7-133.6	2,298	24.3	23.3-25.3
2005	11,464	131.2	128.7-133.6	2,226	23.5	22.5-24.5
2006	11,648	134.2	131.7-136.7	2,155	22.8	21.8-23.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	11,771	134.7	132.3-137.3	2,167	22.7	21.7-23.7
2008	12,113	138.0	135.5-140.5	2,094	21.8	20.8-22.8
2009	12,364	139.2	136.7-141.7	2,090	21.6	20.6-22.6
2010	11,645	131.0	128.6-133.5	2,134	21.7	20.7-22.7
2011	12,188	136.1	133.6-138.6	2,050	20.6	19.7-21.6
2012	11,965	133.0	130.6-135.5	2,001	20.0	19.1-21.0
2013	12,438	137.7	135.2-140.3	1,996	19.9	19.0-20.9
2014	12,700	139.6	137.1-142.1	1,813	17.8	17.0-18.7
2015	12,773	140.5	138.0-143.1	1,882	18.5	17.7-19.4
2016	12,398	135.5	133.0-138.0	1,941	19.0	18.1-19.9
2017	12,845	140.8	138.3-143.4	1,827	17.5	16.7-18.4
2018	12,949	140.8	138.2-143.3	1,911	18.3	17.4-19.2
2019	13,360	145.0	142.4-147.6	1,829	17.5	16.7-18.4
2020	11,637	126.6	124.2-129.1	1,786	17.1	16.3-18.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	983	12.5	11.7-13.3	363	4.3	3.9-4.8
1977	862	10.9	10.1-11.6	319	3.8	3.4-4.2
1978	862	11.1	10.4-11.9	349	4.2	3.8-4.7
1979	867	11.3	10.6-12.1	318	3.9	3.5-4.4
1980	871	11.2	10.5-12.0	314	3.9	3.4-4.3
1981	767	9.7	9.0-10.5	330	4.0	3.6-4.5
1982	776	10.2	9.5-11.0	300	3.6	3.2-4.0
1983	803	10.5	9.7-11.3	257	3.1	2.7-3.5
1984	822	10.8	10.0-11.6	241	3.0	2.6-3.4
1985	784	10.3	9.5-11.0	242	3.0	2.6-3.5
1986	705	9.2	8.5-10.0	248	3.0	2.7-3.4
1987	821	10.6	9.9-11.4	251	3.1	2.7-3.5
1988	808	10.6	9.8-11.4	247	3.1	2.7-3.5
1989	774	10.1	9.4-10.9	215	2.6	2.3-3.0
1990	837	11.0	10.2-11.8	257	3.1	2.7-3.5
1991	831	10.8	10.1-11.6	210	2.6	2.3-3.0
1992	787	10.1	9.4-10.9	225	2.8	2.4-3.2
1993	791	10.2	9.5-11.0	209	2.6	2.2-3.0
1994	814	10.5	9.8-11.3	252	3.1	2.7-3.5
1995	899	11.5	10.8-12.3	204	2.5	2.2-2.9
1996	879	11.3	10.6-12.1	243	3.0	2.6-3.4
1997	801	10.2	9.5-10.9	240	2.9	2.6-3.3
1998	735	9.4	8.7-10.1	197	2.4	2.1-2.8
1999	720	9.1	8.5-9.8	238	2.9	2.5-3.3
2000	666	8.5	7.8-9.1	202	2.4	2.1-2.8
2001	652	8.2	7.6-8.9	193	2.2	1.9-2.6
2002	618	7.8	7.2-8.5	203	2.4	2.1-2.8
2003	648	8.2	7.6-8.9	191	2.2	1.9-2.6
2004	573	7.3	6.7-7.9	191	2.2	1.9-2.6
2005	647	8.3	7.6-8.9	197	2.2	1.9-2.6
2006	601	7.7	7.1-8.3	189	2.2	1.9-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	609	7.8	7.2-8.5	184	2.2	1.9-2.5
2008	605	7.8	7.2-8.4	160	1.8	1.5-2.1
2009	627	8.0	7.4-8.7	150	1.7	1.5-2.0
2010	567	7.4	6.8-8.1	177	2.0	1.7-2.3
2011	503	6.4	5.8-7.0	178	2.1	1.8-2.4
2012	558	7.1	6.5-7.7	201	2.3	2.0-2.7
2013	570	7.4	6.7-8.0	174	1.9	1.7-2.3
2014	547	6.9	6.3-7.5	191	2.1	1.8-2.5
2015	556	7.2	6.6-7.9	162	1.8	1.5-2.1
2016	532	6.9	6.3-7.6	182	2.1	1.8-2.4
2017	571	7.5	6.8-8.1	171	1.9	1.6-2.2
2018	498	6.5	5.9-7.1	146	1.6	1.4-1.9
2019	482	6.5	5.9-7.2	136	1.6	1.3-1.9
2020	438	5.7	5.1-6.3	123	1.3	1.1-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Corpus uteris and uterus, not otherwise specified**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2,371	27.3	26.2-28.5	533	6.0	5.5-6.5
1977	2,141	24.5	23.4-25.6	484	5.4	4.9-5.9
1978	1,922	21.8	20.8-22.8	440	4.9	4.4-5.4
1979	2,115	24.3	23.2-25.4	459	5.0	4.6-5.5
1980	2,147	25.0	23.9-26.1	488	5.3	4.8-5.8
1981	2,200	25.4	24.3-26.5	454	5.0	4.5-5.5
1982	2,154	24.8	23.7-25.9	464	5.1	4.6-5.6
1983	2,105	24.3	23.2-25.4	445	4.7	4.3-5.2
1984	2,084	24.0	23.0-25.1	454	4.9	4.5-5.4
1985	2,031	23.3	22.3-24.4	411	4.5	4.0-4.9
1986	2,018	23.5	22.5-24.6	412	4.4	3.9-4.8
1987	2,105	24.5	23.4-25.6	424	4.6	4.1-5.0
1988	1,980	23.1	22.1-24.2	400	4.3	3.9-4.7
1989	1,966	22.9	21.9-23.9	406	4.4	3.9-4.8
1990	1,994	23.6	22.5-24.7	369	3.9	3.5-4.4
1991	2,100	25.3	24.2-26.5	415	4.5	4.1-5.0
1992	2,168	25.8	24.7-26.9	418	4.5	4.1-5.0
1993	2,075	24.6	23.6-25.7	439	4.7	4.2-5.1
1994	2,256	26.5	25.4-27.7	435	4.7	4.3-5.2
1995	2,255	26.9	25.8-28.0	453	4.8	4.4-5.3
1996	2,339	27.7	26.5-28.8	417	4.5	4.0-4.9
1997	2,338	27.8	26.6-28.9	421	4.5	4.1-5.0
1998	2,532	30.0	28.9-31.3	417	4.5	4.1-5.0
1999	2,462	28.8	27.6-29.9	401	4.2	3.8-4.7
2000	2,399	27.8	26.7-28.9	409	4.3	3.8-4.7
2001	2,615	30.4	29.2-31.5	394	4.1	3.7-4.5
2002	2,646	30.3	29.1-31.5	422	4.5	4.1-4.9
2003	2,629	29.9	28.8-31.1	444	4.6	4.2-5.1
2004	2,689	30.5	29.3-31.6	418	4.4	3.9-4.8
2005	2,723	30.6	29.5-31.8	435	4.6	4.2-5.1
2006	2,691	30.0	28.9-31.2	461	4.8	4.3-5.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Corpus uteris and uterus, not otherwise specified**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	2,722	30.4	29.2-31.6	450	4.6	4.2-5.1
2008	3,002	33.2	32.0-34.5	455	4.7	4.2-5.1
2009	2,814	30.8	29.7-32.0	429	4.4	4.0-4.8
2010	2,830	30.8	29.6-32.0	480	4.8	4.4-5.3
2011	2,972	31.8	30.6-33.0	500	5.1	4.6-5.6
2012	3,004	31.6	30.5-32.8	495	4.9	4.4-5.3
2013	3,229	33.8	32.7-35.1	534	5.2	4.8-5.7
2014	3,145	32.5	31.3-33.7	511	4.9	4.5-5.4
2015	3,146	32.4	31.3-33.6	508	4.9	4.5-5.3
2016	3,229	33.4	32.2-34.6	567	5.4	5.0-5.9
2017	3,180	32.5	31.4-33.7	545	5.1	4.7-5.6
2018	3,015	30.5	29.4-31.7	549	5.1	4.7-5.6
2019	3,052	30.5	29.4-31.7	585	5.5	5.0-6.0
2020	2,874	28.6	27.6-29.8	574	5.2	4.8-5.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,437	16.9	16.1-17.9	1,033	11.9	11.1-12.6
1977	1,359	16.1	15.3-17.0	949	10.9	10.2-11.6
1978	1,368	16.3	15.4-17.2	931	10.6	9.9-11.3
1979	1,406	16.9	16.0-17.8	911	10.4	9.7-11.1
1980	1,395	16.5	15.7-17.5	937	10.7	10.0-11.4
1981	1,494	17.9	16.9-18.8	960	10.8	10.2-11.6
1982	1,438	17.3	16.4-18.3	933	10.6	9.9-11.3
1983	1,492	17.8	16.9-18.7	932	10.5	9.8-11.2
1984	1,425	17.0	16.1-17.9	929	10.5	9.8-11.2
1985	1,485	18.0	17.1-19.0	884	10.1	9.5-10.8
1986	1,423	17.2	16.3-18.2	883	10.1	9.4-10.8
1987	1,449	17.5	16.5-18.4	915	10.4	9.8-11.2
1988	1,429	17.2	16.3-18.1	960	10.8	10.1-11.6
1989	1,426	17.1	16.2-18.0	892	10.1	9.4-10.8
1990	1,487	17.9	17.0-18.9	915	10.3	9.6-11.0
1991	1,530	18.5	17.5-19.4	967	11.0	10.3-11.7
1992	1,486	18.0	17.1-19.0	940	10.8	10.1-11.6
1993	1,460	17.5	16.6-18.4	889	9.9	9.3-10.6
1994	1,512	18.0	17.1-19.0	892	10.1	9.4-10.8
1995	1,571	18.8	17.9-19.8	917	10.3	9.7-11.0
1996	1,525	18.2	17.3-19.1	880	9.9	9.3-10.6
1997	1,484	17.7	16.8-18.7	939	10.5	9.8-11.2
1998	1,362	16.2	15.4-17.1	821	9.2	8.5-9.8
1999	1,400	16.4	15.6-17.3	839	9.3	8.6-9.9
2000	1,416	16.3	15.5-17.2	898	9.8	9.1-10.4
2001	1,450	16.7	15.8-17.6	875	9.5	8.8-10.1
2002	1,407	16.1	15.2-17.0	867	9.4	8.7-10.0
2003	1,369	15.6	14.8-16.5	896	9.6	8.9-10.2
2004	1,355	15.4	14.6-16.2	874	9.3	8.7-9.9
2005	1,268	14.4	13.6-15.2	897	9.6	8.9-10.2
2006	1,311	14.8	14.0-15.6	873	9.3	8.7-10.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1,306	14.7	13.9-15.5	804	8.5	7.9-9.1
2008	1,292	14.4	13.6-15.2	847	8.8	8.2-9.4
2009	1,235	13.8	13.0-14.6	790	8.1	7.5-8.7
2010	1,268	14.0	13.2-14.8	780	8.1	7.5-8.7
2011	1,278	13.9	13.1-14.7	802	8.0	7.5-8.6
2012	1,169	12.7	12.0-13.5	777	7.8	7.3-8.4
2013	1,291	14.1	13.3-14.9	816	8.1	7.6-8.7
2014	1,153	12.5	11.7-13.2	717	7.0	6.5-7.6
2015	1,189	12.9	12.2-13.7	719	7.1	6.6-7.6
2016	1,166	12.5	11.8-13.3	721	7.0	6.5-7.6
2017	1,131	11.8	11.1-12.6	698	6.8	6.3-7.3
2018	1,028	11.1	10.4-11.9	680	6.5	6.0-7.0
2019	1,064	11.5	10.8-12.3	678	6.3	5.9-6.9
2020	932	10.1	9.4-10.8	658	6.2	5.7-6.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	4,131	74.4	72.0-76.8	1,501	29.3	27.8-30.9
1977	4,197	74.3	71.9-76.7	1,500	28.8	27.3-30.4
1978	4,148	73.6	71.3-76.0	1,566	30.4	28.8-32.0
1979	4,368	77.8	75.4-80.3	1,527	29.7	28.2-31.3
1980	4,814	83.8	81.3-86.3	1,548	29.9	28.4-31.5
1981	4,776	83.3	80.8-85.8	1,508	29.2	27.7-30.8
1982	4,883	84.6	82.2-87.1	1,530	29.2	27.7-30.7
1983	5,096	86.9	84.5-89.5	1,680	31.7	30.1-33.3
1984	5,066	86.2	83.7-88.7	1,746	33.0	31.4-34.6
1985	5,326	89.3	86.8-91.8	1,623	30.1	28.6-31.7
1986	5,424	91.3	88.8-93.8	1,695	31.0	29.5-32.5
1987	5,761	95.7	93.2-98.3	1,687	30.9	29.4-32.5
1988	5,774	96.7	94.1-99.3	1,767	32.3	30.8-33.9
1989	5,996	99.2	96.6-101.8	1,766	32.4	30.8-34.0
1990	7,154	115.3	112.6-118.1	1,802	32.9	31.3-34.5
1991	8,483	134.2	131.3-137.2	1,888	34.0	32.4-35.6
1992	10,692	165.6	162.4-168.8	1,898	34.1	32.5-35.7
1993	10,219	156.5	153.4-159.6	1,902	34.0	32.4-35.6
1994	9,670	147.1	144.2-150.1	1,965	34.9	33.3-36.5
1995	9,318	140.4	137.5-143.3	1,886	33.3	31.8-34.9
1996	10,042	151.2	148.2-154.2	1,852	32.1	30.7-33.7
1997	10,589	158.5	155.4-161.6	1,798	30.9	29.5-32.4
1998	10,485	155.2	152.2-158.2	1,680	28.3	26.9-29.7
1999	10,998	161.8	158.8-164.9	1,597	26.5	25.2-27.8
2000	12,122	177.1	174.0-180.3	1,632	26.9	25.6-28.3
2001	11,945	172.6	169.5-175.7	1,612	26.2	24.9-27.5
2002	12,245	174.8	171.7-177.9	1,531	24.6	23.4-25.9
2003	11,311	160.2	157.2-163.2	1,535	24.1	22.9-25.4
2004	11,018	154.0	151.1-156.9	1,462	22.8	21.6-24.0
2005	11,189	154.7	151.8-157.6	1,364	20.9	19.8-22.1
2006	12,582	173.0	169.9-176.0	1,324	20.1	19.0-21.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	12,831	172.7	169.7-175.7	1,385	20.5	19.5-21.6
2008	12,366	163.2	160.3-166.1	1,318	19.2	18.2-20.3
2009	12,160	156.9	154.1-159.8	1,265	18.2	17.2-19.3
2010	11,440	144.9	142.2-147.6	1,371	19.3	18.3-20.3
2011	11,916	148.2	145.5-151.0	1,305	18.0	17.1-19.0
2012	10,199	123.6	121.2-126.1	1,222	16.7	15.8-17.7
2013	9,928	118.2	115.9-120.6	1,165	15.5	14.6-16.5
2014	9,655	112.2	109.9-114.5	1,266	16.6	15.6-17.5
2015	10,064	115.0	112.7-117.3	1,227	15.9	15.0-16.8
2016	10,418	117.3	115.0-119.6	1,308	16.5	15.6-17.4
2017	11,141	122.9	120.6-125.3	1,317	16.4	15.5-17.3
2018	11,665	127.4	125.0-129.7	1,241	15.0	14.2-15.9
2019	12,328	132.3	129.9-134.7	1,196	14.4	13.6-15.2
2020	10,543	112.1	109.9-114.3	1,174	14.0	13.2-14.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Testis**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	249	3.4	2.9-3.8	48	0.6	0.4-0.8
1977	242	3.3	2.9-3.8	44	0.6	0.4-0.8
1978	263	3.7	3.3-4.2	49	0.7	0.5-0.9
1979	303	4.3	3.8-4.8	47	0.7	0.5-0.9
1980	324	4.5	4.0-5.0	44	0.6	0.4-0.8
1981	317	4.3	3.8-4.8	32	0.4	0.3-0.6
1982	321	4.4	3.9-4.9	32	0.5	0.3-0.7
1983	339	4.6	4.1-5.1	30	0.4	0.3-0.6
1984	381	5.2	4.7-5.8	27	0.4	0.3-0.6
1985	353	4.8	4.3-5.3	34	0.5	0.3-0.7
1986	356	4.7	4.2-5.2	34	0.5	0.3-0.7
1987	382	5.1	4.6-5.6	27	0.4	0.2-0.5
1988	351	4.6	4.2-5.2	23	0.3	0.2-0.5
1989	363	4.8	4.3-5.4	31	0.4	0.3-0.6
1990	408	5.4	4.9-6.0	24	0.3	0.2-0.5
1991	389	5.2	4.7-5.7	22	0.3	0.2-0.4
1992	400	5.4	4.9-6.0	28	0.4	0.3-0.6
1993	421	5.7	5.1-6.3	32	0.5	0.3-0.7
1994	412	5.6	5.1-6.2	21	0.3	0.2-0.5
1995	415	5.7	5.2-6.3	22	0.3	0.2-0.5
1996	410	5.6	5.1-6.2	26	0.4	0.2-0.5
1997	427	5.9	5.4-6.5	12	0.2	0.1-0.3
1998	430	6.0	5.5-6.6	25	0.3	0.2-0.5
1999	470	6.6	6.0-7.2	17	0.2	0.1-0.4
2000	422	5.9	5.4-6.5	15	0.2	0.1-0.3
2001	437	6.2	5.6-6.8	19	0.3	0.2-0.4
2002	427	6.1	5.5-6.7	25	0.4	0.2-0.5
2003	478	6.9	6.3-7.5	19	0.3	0.2-0.4
2004	464	6.7	6.1-7.3	19	0.3	0.2-0.4
2005	489	7.1	6.5-7.8	14	0.2	0.1-0.3
2006	464	6.8	6.2-7.4	23	0.3	0.2-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Testis**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	500	7.3	6.7-8.0	7	0.1	0.0-0.2
2008	518	7.6	7.0-8.3	15	0.2	0.1-0.4
2009	506	7.4	6.8-8.1	15	0.2	0.1-0.4
2010	510	7.5	6.9-8.2	28	0.4	0.3-0.6
2011	509	7.4	6.8-8.1	16	0.2	0.1-0.3
2012	540	8.0	7.3-8.7	18	0.3	0.2-0.4
2013	478	7.1	6.4-7.8	13	0.2	0.1-0.3
2014	499	7.4	6.7-8.1	17	0.3	0.1-0.4
2015	499	7.3	6.7-8.0	18	0.2	0.1-0.4
2016	515	7.7	7.0-8.4	14	0.2	0.1-0.3
2017	503	7.6	7.0-8.3	14	0.2	0.1-0.3
2018	526	8.0	7.3-8.7	16	0.2	0.1-0.3
2019	487	7.4	6.8-8.2	27	0.4	0.2-0.5
2020	460	7.2	6.6-7.9	23	0.4	0.2-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3,084	20.5	19.7-21.2	2,249	37.1	35.5-38.8	835	9.4	8.8-10.1	950	6.7	6.3-7.2	673	12.4	11.4-13.4	277	3.3	2.9-3.7
1977	3,236	21.5	20.7-22.2	2,339	38.2	36.5-39.8	897	10.3	9.6-11.0	967	6.8	6.3-7.2	677	12.3	11.4-13.3	290	3.4	3.0-3.8
1978	3,388	22.5	21.8-23.3	2,455	40.6	39.0-42.3	933	10.7	10.0-11.4	948	6.6	6.1-7.0	653	11.9	10.9-12.9	295	3.4	3.0-3.8
1979	3,419	22.6	21.8-23.4	2,505	41.6	39.9-43.3	914	10.2	9.6-10.9	925	6.4	6.0-6.8	631	11.4	10.5-12.4	294	3.3	3.0-3.7
1980	3,594	23.5	22.8-24.3	2,662	43.6	41.9-45.4	932	10.4	9.7-11.1	913	6.1	5.7-6.6	641	11.4	10.5-12.4	272	3.0	2.7-3.4
1981	3,509	23.0	22.2-23.7	2,589	42.4	40.7-44.1	920	10.3	9.6-11.0	916	6.2	5.8-6.6	638	11.5	10.6-12.5	278	3.1	2.7-3.5
1982	3,577	23.3	22.5-24.1	2,588	42.0	40.3-43.7	989	10.9	10.3-11.7	881	5.9	5.5-6.3	623	11.1	10.2-12.1	258	2.8	2.5-3.2
1983	3,635	23.6	22.8-24.4	2,655	43.6	41.9-45.4	980	10.7	10.0-11.4	844	5.6	5.2-6.0	598	10.9	10.0-11.9	246	2.6	2.3-3.0
1984	3,521	22.7	21.9-23.5	2,546	41.0	39.3-42.7	975	10.7	10.0-11.4	852	5.7	5.3-6.1	575	10.4	9.6-11.4	277	3.0	2.6-3.3
1985	3,521	22.8	22.0-23.6	2,548	41.3	39.6-43.0	973	10.7	10.0-11.4	854	5.7	5.3-6.0	585	10.5	9.7-11.4	269	2.9	2.5-3.2
1986	3,682	23.7	23.0-24.5	2,676	43.1	41.4-44.8	1,006	11.1	10.4-11.8	834	5.5	5.1-5.9	567	10.4	9.5-11.3	267	2.8	2.5-3.2
1987	3,564	22.8	22.0-23.6	2,555	40.8	39.1-42.4	1,009	10.9	10.2-11.6	773	5.1	4.7-5.5	528	9.4	8.6-10.3	245	2.6	2.3-2.9
1988	3,530	22.7	21.9-23.4	2,534	40.5	38.9-42.2	996	10.8	10.1-11.5	829	5.3	5.0-5.7	543	9.4	8.6-10.2	286	2.9	2.6-3.3
1989	3,517	22.6	21.9-23.4	2,550	41.0	39.4-42.7	967	10.5	9.8-11.2	818	5.3	4.9-5.7	563	10.0	9.2-10.9	255	2.6	2.3-3.0
1990	3,645	23.5	22.7-24.3	2,588	41.5	39.9-43.2	1,057	11.5	10.8-12.2	799	5.2	4.8-5.6	520	9.2	8.4-10.1	279	2.9	2.5-3.2
1991	3,647	23.4	22.6-24.2	2,634	41.9	40.3-43.6	1,013	11.0	10.3-11.7	765	4.9	4.6-5.3	525	9.1	8.3-9.9	240	2.5	2.2-2.8
1992	3,648	23.4	22.6-24.1	2,608	41.6	40.0-43.3	1,040	11.4	10.7-12.1	799	5.1	4.7-5.4	490	8.6	7.8-9.4	309	3.1	2.8-3.5
1993	3,710	23.6	22.9-24.4	2,622	41.2	39.6-42.8	1,088	11.8	11.1-12.5	830	5.3	4.9-5.6	561	9.7	8.9-10.6	269	2.7	2.4-3.1
1994	3,696	23.6	22.8-24.4	2,631	41.4	39.8-43.0	1,065	11.6	10.9-12.3	830	5.2	4.9-5.6	558	9.5	8.7-10.4	272	2.7	2.4-3.1
1995	3,829	24.4	23.6-25.2	2,710	42.5	40.9-44.1	1,119	12.1	11.4-12.9	835	5.2	4.9-5.6	548	9.2	8.4-10.0	287	2.9	2.5-3.2
1996	3,901	24.8	24.0-25.6	2,850	44.5	42.9-46.2	1,051	11.4	10.7-12.1	840	5.2	4.9-5.6	547	9.2	8.5-10.1	293	2.9	2.6-3.3
1997	4,175	26.5	25.7-27.3	2,999	46.3	44.6-48.0	1,176	12.7	11.9-13.4	857	5.3	5.0-5.7	572	9.4	8.7-10.2	285	2.9	2.6-3.2
1998	4,318	27.2	26.4-28.0	3,115	47.5	45.8-49.2	1,203	13.1	12.3-13.8	832	5.1	4.8-5.5	569	9.3	8.5-10.1	263	2.6	2.3-2.9
1999	4,249	26.5	25.8-27.4	3,055	46.2	44.5-47.9	1,194	12.8	12.1-13.6	830	5.0	4.7-5.4	573	9.3	8.5-10.1	257	2.5	2.2-2.8
2000	4,339	26.8	26.0-27.6	3,110	46.6	45.0-48.3	1,229	13.0	12.2-13.7	829	5.0	4.7-5.3	569	8.9	8.2-9.7	260	2.5	2.2-2.9
2001	4,278	26.3	25.5-27.1	3,121	46.4	44.8-48.1	1,157	12.2	11.5-13.0	847	5.1	4.7-5.4	597	9.4	8.6-10.1	250	2.4	2.1-2.7
2002	4,401	26.9	26.1-27.7	3,261	47.9	46.3-49.6	1,140	12.0	11.3-12.7	817	4.8	4.5-5.2	543	8.4	7.7-9.2	274	2.7	2.4-3.0
2003	4,526	27.5	26.7-28.3	3,328	48.3	46.7-50.0	1,198	12.6	11.9-13.3	755	4.4	4.1-4.7	512	7.9	7.2-8.6	243	2.3	2.0-2.6
2004	4,662	28.1	27.3-29.0	3,419	49.3	47.6-51.0	1,243	13.0	12.3-13.7	830	4.8	4.5-5.2	576	8.6	7.9-9.3	254	2.4	2.1-2.7
2005	4,682	28.2	27.4-29.0	3,413	48.9	47.2-50.5	1,269	13.2	12.5-14.0	790	4.6	4.3-4.9	526	7.8	7.2-8.5	264	2.5	2.2-2.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	4,570	27.3	26.5-28.1	3,390	48.3	46.6-49.9	1,180	12.4	11.7-13.1	816	4.7	4.4-5.1	550	8.1	7.5-8.9	266	2.5	2.2-2.9
2007	4,618	27.5	26.7-28.3	3,397	47.8	46.2-49.4	1,221	12.8	12.1-13.6	882	5.0	4.7-5.4	619	8.9	8.3-9.7	263	2.5	2.2-2.8
2008	4,735	27.8	27.0-28.6	3,579	49.6	47.9-51.2	1,156	11.9	11.2-12.6	876	5.0	4.6-5.3	605	8.7	8.0-9.4	271	2.5	2.2-2.9
2009	4,752	27.6	26.8-28.4	3,536	48.4	46.8-50.0	1,216	12.3	11.6-13.1	856	4.8	4.4-5.1	628	8.9	8.2-9.6	228	2.0	1.8-2.3
2010	4,473	25.8	25.0-26.5	3,387	45.8	44.2-47.4	1,086	11.0	10.3-11.7	907	5.0	4.6-5.3	637	8.7	8.1-9.4	270	2.4	2.1-2.7
2011	4,754	27.2	26.4-28.0	3,538	47.1	45.5-48.7	1,216	12.4	11.7-13.1	838	4.5	4.2-4.9	587	8.0	7.4-8.7	251	2.3	2.0-2.6
2012	4,872	27.3	26.5-28.1	3,606	47.3	45.8-48.9	1,266	12.5	11.8-13.2	875	4.7	4.4-5.0	602	8.0	7.4-8.7	273	2.4	2.1-2.7
2013	4,883	27.2	26.4-28.0	3,642	47.2	45.7-48.8	1,241	12.2	11.5-12.9	864	4.6	4.3-4.9	621	8.2	7.6-8.9	243	2.1	1.8-2.4
2014	4,820	26.3	25.5-27.1	3,597	45.5	44.0-47.1	1,223	11.7	11.1-12.4	923	4.8	4.5-5.1	631	8.2	7.5-8.8	292	2.6	2.3-2.9
2015	4,806	26.0	25.3-26.8	3,621	45.1	43.6-46.6	1,185	11.6	10.9-12.3	863	4.5	4.2-4.8	592	7.5	6.9-8.1	271	2.4	2.1-2.7
2016	4,877	26.0	25.3-26.8	3,686	45.1	43.6-46.6	1,191	11.3	10.7-12.0	891	4.6	4.3-4.9	630	8.0	7.4-8.7	261	2.3	2.0-2.6
2017	4,821	25.4	24.6-26.1	3,608	43.6	42.1-45.0	1,213	11.4	10.8-12.1	861	4.3	4.0-4.6	604	7.5	6.9-8.1	257	2.1	1.9-2.4
2018	4,788	24.8	24.1-25.5	3,593	42.5	41.1-43.9	1,195	11.1	10.5-11.8	853	4.2	3.9-4.5	603	7.3	6.7-7.9	250	2.2	1.9-2.5
2019	4,797	24.6	23.9-25.4	3,571	41.5	40.2-43.0	1,226	11.4	10.7-12.0	806	3.9	3.7-4.2	584	7.0	6.4-7.6	222	1.9	1.6-2.2
2020	4,275	21.7	21.0-22.3	3,236	37.1	35.8-38.5	1,039	9.5	8.9-10.1	769	3.8	3.5-4.0	539	6.4	5.9-7.0	230	1.9	1.7-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,107	7.2	6.8-7.6	689	10.5	9.7-11.4	418	4.8	4.3-5.3	598	3.9	3.6-4.2	381	6.0	5.4-6.6	217	2.4	2.1-2.8
1977	1,160	7.6	7.1-8.0	728	11.1	10.3-12.0	432	5.0	4.5-5.5	633	4.2	3.8-4.5	393	6.2	5.5-6.8	240	2.7	2.4-3.1
1978	1,149	7.6	7.1-8.0	703	10.9	10.1-11.8	446	5.1	4.6-5.6	575	3.7	3.4-4.1	361	5.6	5.0-6.3	214	2.4	2.1-2.8
1979	1,149	7.5	7.1-8.0	705	10.9	10.1-11.8	444	4.9	4.5-5.4	636	4.2	3.9-4.6	398	6.3	5.6-6.9	238	2.7	2.4-3.1
1980	1,140	7.4	7.0-7.9	695	10.6	9.8-11.4	445	5.2	4.7-5.7	598	3.9	3.5-4.2	357	5.7	5.1-6.3	241	2.6	2.3-3.0
1981	1,216	8.1	7.6-8.6	714	11.3	10.5-12.2	502	5.8	5.3-6.4	593	3.9	3.6-4.2	358	5.6	5.0-6.3	235	2.6	2.3-3.0
1982	1,301	8.5	8.1-9.0	784	12.1	11.2-13.0	517	5.9	5.4-6.5	599	3.8	3.5-4.2	358	5.7	5.1-6.3	241	2.6	2.3-3.0
1983	1,356	8.9	8.4-9.4	834	12.8	11.9-13.8	522	6.0	5.5-6.5	588	3.8	3.5-4.1	350	5.5	4.9-6.1	238	2.7	2.3-3.0
1984	1,376	9.0	8.5-9.5	863	13.4	12.5-14.3	513	5.9	5.3-6.4	599	3.9	3.6-4.3	371	6.0	5.4-6.7	228	2.5	2.2-2.9
1985	1,456	9.5	9.0-10.0	885	13.7	12.8-14.6	571	6.5	5.9-7.0	614	4.0	3.7-4.3	350	5.6	5.0-6.3	264	3.0	2.6-3.3
1986	1,447	9.4	9.0-10.0	889	13.7	12.7-14.6	558	6.4	5.9-7.0	654	4.2	3.9-4.6	395	6.2	5.6-6.9	259	2.9	2.5-3.2
1987	1,573	10.2	9.7-10.8	933	14.3	13.4-15.3	640	7.3	6.7-7.9	652	4.2	3.9-4.6	404	6.5	5.9-7.2	248	2.7	2.4-3.1
1988	1,452	9.5	9.0-10.0	878	13.6	12.7-14.6	574	6.6	6.0-7.2	622	4.0	3.7-4.4	359	5.7	5.1-6.3	263	2.9	2.5-3.3
1989	1,465	9.7	9.2-10.2	911	14.2	13.3-15.2	554	6.3	5.8-6.9	626	4.1	3.8-4.4	376	6.1	5.4-6.7	250	2.8	2.4-3.1
1990	1,627	10.7	10.1-11.2	986	15.2	14.3-16.3	641	7.4	6.8-8.0	602	3.9	3.6-4.2	355	5.6	5.0-6.2	247	2.7	2.3-3.0
1991	1,654	10.9	10.3-11.4	981	15.1	14.1-16.1	673	7.8	7.2-8.4	626	4.0	3.7-4.4	370	5.8	5.2-6.5	256	2.8	2.4-3.1
1992	1,676	11.0	10.5-11.6	1,004	15.3	14.4-16.3	672	7.8	7.2-8.4	577	3.7	3.4-4.1	343	5.6	5.0-6.2	234	2.5	2.2-2.9
1993	1,726	11.3	10.7-11.8	1,056	16.0	15.1-17.0	670	7.7	7.1-8.3	579	3.7	3.4-4.1	366	5.6	5.1-6.3	213	2.3	2.0-2.7
1994	1,768	11.6	11.0-12.1	1,068	16.3	15.3-17.3	700	8.1	7.5-8.7	662	4.2	3.9-4.6	410	6.5	5.9-7.2	252	2.7	2.4-3.1
1995	1,748	11.4	10.9-12.0	1,042	15.7	14.7-16.7	706	8.1	7.5-8.8	628	4.0	3.7-4.4	363	5.7	5.2-6.4	265	2.9	2.5-3.3
1996	1,910	12.4	11.9-13.0	1,164	17.6	16.6-18.7	746	8.5	7.9-9.2	649	4.2	3.8-4.5	373	5.8	5.2-6.4	276	3.0	2.6-3.3
1997	1,977	12.8	12.3-13.4	1,218	18.4	17.4-19.4	759	8.7	8.1-9.4	633	4.0	3.7-4.4	392	6.1	5.5-6.7	241	2.6	2.3-3.0
1998	2,083	13.5	12.9-14.1	1,269	18.7	17.7-19.8	814	9.4	8.7-10.0	634	4.0	3.7-4.3	385	5.8	5.3-6.5	249	2.6	2.3-3.0
1999	2,177	14.0	13.4-14.6	1,347	19.7	18.7-20.8	830	9.4	8.8-10.1	548	3.4	3.1-3.7	343	5.2	4.6-5.8	205	2.1	1.8-2.4
2000	2,198	13.9	13.4-14.5	1,302	18.9	17.9-20.0	896	10.1	9.4-10.8	605	3.7	3.5-4.1	370	5.6	5.0-6.2	235	2.5	2.2-2.8
2001	2,285	14.4	13.8-15.0	1,419	20.3	19.3-21.4	866	9.7	9.1-10.4	638	3.9	3.6-4.2	374	5.5	5.0-6.1	264	2.8	2.5-3.2
2002	2,304	14.4	13.9-15.0	1,438	20.5	19.4-21.6	866	9.6	9.0-10.3	605	3.7	3.4-4.0	363	5.3	4.8-5.9	242	2.5	2.2-2.8
2003	2,425	15.1	14.5-15.7	1,529	21.4	20.3-22.5	896	9.9	9.3-10.6	604	3.6	3.4-4.0	361	5.2	4.7-5.7	243	2.4	2.1-2.7
2004	2,614	16.3	15.6-16.9	1,629	22.7	21.6-23.8	985	11.1	10.4-11.8	635	3.8	3.5-4.1	383	5.5	5.0-6.1	252	2.6	2.3-2.9
2005	2,607	16.2	15.5-16.8	1,600	22.3	21.2-23.5	1,007	11.3	10.6-12.0	602	3.6	3.3-3.9	379	5.4	4.9-6.0	223	2.3	2.0-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	2,815	17.3	16.7-18.0	1,787	24.6	23.4-25.7	1,028	11.4	10.7-12.2	632	3.7	3.5-4.0	381	5.5	4.9-6.0	251	2.6	2.2-2.9
2007	2,881	17.6	17.0-18.3	1,831	24.9	23.8-26.1	1,050	11.8	11.1-12.6	612	3.6	3.3-3.9	380	5.3	4.8-5.8	232	2.4	2.1-2.7
2008	2,865	17.3	16.6-17.9	1,806	24.2	23.1-25.3	1,059	11.6	10.9-12.4	606	3.5	3.2-3.8	391	5.4	4.9-5.9	215	2.1	1.8-2.4
2009	2,750	16.6	16.0-17.2	1,696	22.5	21.5-23.7	1,054	11.7	11.0-12.4	621	3.6	3.3-3.9	406	5.5	5.0-6.1	215	2.2	1.9-2.5
2010	2,830	16.8	16.2-17.5	1,775	23.2	22.1-24.3	1,055	11.6	10.9-12.3	593	3.4	3.1-3.6	378	5.0	4.5-5.5	215	2.1	1.8-2.4
2011	2,911	17.2	16.6-17.9	1,877	24.2	23.1-25.3	1,034	11.4	10.6-12.1	613	3.4	3.2-3.7	384	5.1	4.6-5.6	229	2.2	1.9-2.5
2012	3,022	17.6	17.0-18.3	1,929	24.5	23.4-25.7	1,093	11.8	11.1-12.6	631	3.5	3.2-3.8	417	5.4	4.9-6.0	214	2.1	1.8-2.4
2013	3,022	17.4	16.8-18.1	1,947	24.3	23.2-25.5	1,075	11.6	10.8-12.3	579	3.2	2.9-3.5	366	4.7	4.2-5.2	213	2.0	1.7-2.3
2014	3,093	17.6	17.0-18.3	2,006	24.7	23.6-25.8	1,087	11.6	10.9-12.3	580	3.1	2.8-3.4	368	4.5	4.1-5.0	212	2.0	1.7-2.3
2015	3,148	17.7	17.1-18.3	2,026	24.7	23.6-25.8	1,122	11.6	10.9-12.4	629	3.3	3.1-3.6	401	4.9	4.5-5.5	228	2.1	1.8-2.4
2016	3,121	17.6	17.0-18.2	2,058	24.9	23.8-26.0	1,063	11.3	10.6-12.0	530	2.8	2.6-3.0	347	4.2	3.7-4.6	183	1.7	1.5-2.0
2017	3,218	17.9	17.3-18.6	2,042	24.5	23.4-25.6	1,176	12.3	11.6-13.1	554	2.9	2.7-3.2	342	4.2	3.7-4.6	212	2.0	1.7-2.3
2018	3,270	18.2	17.6-18.9	2,122	25.2	24.1-26.3	1,148	12.2	11.5-13.0	575	3.0	2.7-3.2	373	4.4	4.0-4.9	202	1.8	1.6-2.1
2019	3,360	18.5	17.9-19.2	2,233	26.3	25.2-27.4	1,127	11.8	11.1-12.6	533	2.7	2.5-2.9	332	3.8	3.4-4.2	201	1.8	1.6-2.1
2020	2,895	15.9	15.3-16.5	1,918	22.6	21.5-23.6	977	10.1	9.4-10.7	521	2.7	2.4-2.9	344	4.1	3.6-4.5	177	1.5	1.3-1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	906	5.8	5.4-6.2	511	7.1	6.5-7.8	395	4.7	4.3-5.2	688	4.4	4.1-4.8	376	5.4	4.9-6.0	312	3.7	3.3-4.1
1977	917	6.0	5.6-6.4	515	7.4	6.7-8.0	402	4.8	4.3-5.3	692	4.4	4.1-4.8	376	5.4	4.8-5.9	316	3.7	3.3-4.1
1978	893	5.8	5.5-6.3	461	6.6	6.0-7.3	432	5.2	4.7-5.7	660	4.2	3.9-4.5	354	5.1	4.6-5.7	306	3.5	3.1-3.9
1979	954	6.2	5.8-6.6	532	7.8	7.1-8.5	422	5.0	4.6-5.6	646	4.1	3.8-4.5	363	5.3	4.8-5.9	283	3.3	2.9-3.7
1980	953	6.2	5.8-6.6	523	7.5	6.8-8.1	430	5.2	4.7-5.7	723	4.6	4.3-5.0	418	6.1	5.5-6.7	305	3.5	3.1-4.0
1981	961	6.4	6.0-6.8	536	8.0	7.3-8.8	425	5.1	4.6-5.6	671	4.4	4.0-4.7	368	5.5	4.9-6.1	303	3.5	3.1-3.9
1982	997	6.6	6.2-7.1	519	7.7	7.0-8.4	478	5.8	5.3-6.4	690	4.5	4.2-4.9	348	5.2	4.6-5.8	342	4.0	3.6-4.5
1983	949	6.4	6.0-6.8	482	7.1	6.5-7.8	467	5.7	5.2-6.3	657	4.3	3.9-4.6	352	5.3	4.7-5.9	305	3.4	3.1-3.9
1984	1,043	6.9	6.4-7.3	541	8.0	7.3-8.7	502	6.0	5.5-6.6	662	4.3	4.0-4.6	350	5.3	4.7-5.9	312	3.6	3.2-4.0
1985	1,016	6.8	6.4-7.3	548	8.2	7.5-8.9	468	5.7	5.2-6.2	652	4.3	4.0-4.7	325	5.0	4.4-5.6	327	3.8	3.4-4.3
1986	1,021	6.8	6.4-7.3	562	8.5	7.8-9.2	459	5.6	5.1-6.1	702	4.6	4.3-5.0	386	5.9	5.3-6.6	316	3.6	3.2-4.0
1987	1,102	7.4	7.0-7.9	573	8.5	7.8-9.3	529	6.4	5.8-6.9	688	4.5	4.2-4.9	351	5.3	4.8-5.9	337	3.9	3.5-4.3
1988	1,085	7.3	6.8-7.7	589	8.8	8.1-9.5	496	6.1	5.5-6.7	657	4.2	3.9-4.6	363	5.3	4.8-5.9	294	3.3	3.0-3.7
1989	1,085	7.3	6.9-7.8	611	9.1	8.4-9.9	474	5.9	5.4-6.5	644	4.2	3.9-4.6	352	5.3	4.8-5.9	292	3.4	3.0-3.8
1990	1,140	7.7	7.3-8.2	614	9.3	8.6-10.1	526	6.5	5.9-7.1	613	4.1	3.7-4.4	356	5.5	4.9-6.1	257	2.9	2.6-3.3
1991	1,079	7.3	6.9-7.8	574	8.7	8.0-9.5	505	6.3	5.7-6.8	655	4.4	4.1-4.8	367	5.7	5.1-6.3	288	3.5	3.1-3.9
1992	1,091	7.4	6.9-7.8	579	8.7	8.0-9.4	512	6.3	5.8-6.9	648	4.3	3.9-4.6	332	5.0	4.5-5.6	316	3.7	3.3-4.2
1993	1,105	7.4	7.0-7.9	591	8.8	8.1-9.5	514	6.3	5.8-6.9	627	4.1	3.8-4.5	343	5.2	4.6-5.8	284	3.3	2.9-3.8
1994	1,209	8.1	7.7-8.6	658	9.8	9.1-10.6	551	6.7	6.1-7.2	713	4.7	4.4-5.1	387	5.8	5.3-6.5	326	3.8	3.4-4.2
1995	1,274	8.5	8.1-9.0	649	9.6	8.9-10.4	625	7.6	7.0-8.3	666	4.4	4.1-4.8	359	5.4	4.9-6.0	307	3.6	3.2-4.0
1996	1,250	8.4	8.0-8.9	670	10.0	9.2-10.8	580	7.1	6.6-7.8	701	4.7	4.3-5.0	378	5.7	5.1-6.3	323	3.8	3.4-4.2
1997	1,166	7.8	7.4-8.3	624	9.2	8.5-10.0	542	6.7	6.1-7.3	699	4.6	4.3-5.0	362	5.4	4.8-6.0	337	3.9	3.5-4.4
1998	1,271	8.5	8.0-9.0	699	10.2	9.4-10.9	572	7.1	6.5-7.7	691	4.5	4.2-4.9	375	5.5	4.9-6.1	316	3.7	3.3-4.1
1999	1,134	7.5	7.1-8.0	595	8.7	8.0-9.4	539	6.6	6.0-7.1	696	4.5	4.2-4.9	380	5.5	5.0-6.1	316	3.6	3.3-4.1
2000	1,247	8.2	7.8-8.7	637	9.2	8.5-10.0	610	7.3	6.7-7.9	653	4.2	3.9-4.5	336	4.8	4.3-5.4	317	3.6	3.2-4.0
2001	1,155	7.6	7.1-8.0	614	8.8	8.1-9.5	541	6.5	6.0-7.1	634	4.0	3.7-4.4	337	4.8	4.3-5.4	297	3.4	3.0-3.8
2002	1,179	7.7	7.3-8.2	648	9.3	8.6-10.1	531	6.4	5.9-7.0	671	4.2	3.9-4.6	373	5.3	4.8-5.9	298	3.4	3.0-3.8
2003	1,179	7.7	7.3-8.2	610	8.7	8.0-9.4	569	6.9	6.4-7.6	619	3.9	3.6-4.3	351	4.9	4.4-5.5	268	3.1	2.7-3.5
2004	1,141	7.5	7.0-7.9	637	9.1	8.4-9.8	504	6.2	5.6-6.7	657	4.1	3.8-4.5	350	4.9	4.4-5.5	307	3.5	3.1-3.9
2005	1,197	7.8	7.4-8.3	682	9.7	9.0-10.5	515	6.3	5.7-6.8	698	4.4	4.1-4.7	385	5.4	4.9-6.0	313	3.6	3.2-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,197	7.9	7.4-8.3	676	9.5	8.8-10.3	521	6.4	5.8-7.0	661	4.1	3.8-4.4	391	5.4	4.9-6.0	270	3.0	2.6-3.4
2007	1,120	7.3	6.9-7.8	602	8.5	7.8-9.2	518	6.4	5.8-7.0	690	4.3	3.9-4.6	403	5.5	5.0-6.1	287	3.2	2.8-3.6
2008	1,168	7.6	7.2-8.1	634	8.9	8.2-9.6	534	6.5	5.9-7.1	638	3.9	3.6-4.2	369	5.0	4.5-5.6	269	3.0	2.6-3.4
2009	1,193	7.7	7.2-8.1	636	8.8	8.1-9.5	557	6.7	6.1-7.3	733	4.5	4.1-4.8	404	5.4	4.9-6.0	329	3.7	3.3-4.1
2010	1,129	7.3	6.9-7.8	619	8.6	7.9-9.3	510	6.2	5.7-6.8	700	4.2	3.9-4.6	376	5.0	4.5-5.6	324	3.6	3.2-4.0
2011	1,117	7.2	6.7-7.6	618	8.4	7.8-9.1	499	6.1	5.6-6.7	664	4.0	3.7-4.3	361	4.8	4.3-5.3	303	3.2	2.9-3.7
2012	1,178	7.4	6.9-7.8	637	8.5	7.8-9.2	541	6.5	5.9-7.1	753	4.4	4.1-4.8	403	5.2	4.7-5.8	350	3.8	3.4-4.2
2013	1,155	7.3	6.9-7.8	660	8.8	8.1-9.5	495	5.9	5.4-6.5	701	4.0	3.7-4.3	380	4.8	4.3-5.3	321	3.3	3.0-3.7
2014	1,277	8.0	7.5-8.4	709	9.3	8.6-10.0	568	6.8	6.2-7.4	762	4.4	4.1-4.7	419	5.2	4.7-5.7	343	3.7	3.3-4.1
2015	1,212	7.5	7.1-8.0	634	8.3	7.7-9.0	578	6.8	6.3-7.4	730	4.2	3.9-4.5	378	4.7	4.2-5.2	352	3.7	3.3-4.2
2016	1,191	7.3	6.9-7.8	686	8.9	8.2-9.6	505	5.9	5.4-6.5	754	4.3	3.9-4.6	437	5.4	4.9-5.9	317	3.3	2.9-3.7
2017	1,264	7.8	7.4-8.3	708	9.2	8.5-10.0	556	6.6	6.0-7.2	794	4.5	4.2-4.8	453	5.5	5.0-6.0	341	3.6	3.2-4.1
2018	1,122	6.9	6.5-7.3	630	8.0	7.4-8.7	492	5.9	5.3-6.5	762	4.3	4.0-4.6	430	5.2	4.7-5.8	332	3.5	3.1-3.9
2019	1,088	6.7	6.3-7.2	602	7.9	7.3-8.6	486	5.6	5.1-6.2	749	4.1	3.8-4.5	423	5.1	4.6-5.6	326	3.3	3.0-3.7
2020	1,092	6.8	6.3-7.2	603	7.7	7.1-8.4	489	5.9	5.3-6.5	752	4.2	3.9-4.5	434	5.3	4.8-5.8	318	3.2	2.9-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	493	3.4	3.1-3.7	155	2.4	2.0-2.8	338	4.3	3.9-4.8	102	0.7	0.6-0.9	41	0.7	0.5-0.9	61	0.7	0.5-0.9
1977	569	3.9	3.6-4.2	170	2.5	2.1-2.9	399	5.1	4.6-5.7	114	0.8	0.6-0.9	46	0.7	0.5-1.0	68	0.8	0.6-1.0
1978	560	3.8	3.5-4.2	158	2.4	2.0-2.8	402	5.1	4.6-5.7	95	0.7	0.5-0.8	31	0.5	0.4-0.8	64	0.7	0.6-0.9
1979	513	3.6	3.3-3.9	171	2.7	2.3-3.1	342	4.4	3.9-4.9	76	0.5	0.4-0.6	28	0.5	0.3-0.7	48	0.5	0.4-0.7
1980	539	3.7	3.4-4.0	162	2.5	2.1-2.9	377	4.8	4.3-5.3	109	0.7	0.6-0.9	48	0.8	0.6-1.1	61	0.7	0.5-0.9
1981	522	3.6	3.3-3.9	158	2.4	2.0-2.8	364	4.6	4.2-5.2	92	0.6	0.5-0.8	36	0.6	0.4-0.8	56	0.6	0.5-0.8
1982	512	3.6	3.3-3.9	160	2.4	2.1-2.9	352	4.6	4.1-5.1	97	0.6	0.5-0.8	41	0.6	0.5-0.9	56	0.6	0.5-0.8
1983	601	4.2	3.8-4.5	159	2.4	2.0-2.8	442	5.7	5.2-6.3	78	0.5	0.4-0.7	21	0.3	0.2-0.5	57	0.6	0.5-0.8
1984	650	4.5	4.1-4.8	186	2.9	2.5-3.4	464	6.0	5.4-6.5	101	0.7	0.5-0.8	37	0.6	0.4-0.9	64	0.7	0.5-0.9
1985	653	4.5	4.1-4.9	188	2.8	2.4-3.2	465	6.0	5.4-6.6	89	0.6	0.5-0.7	43	0.7	0.5-1.0	46	0.5	0.4-0.7
1986	630	4.3	4.0-4.6	176	2.6	2.2-3.0	454	5.8	5.3-6.4	81	0.5	0.4-0.7	28	0.5	0.3-0.7	53	0.6	0.4-0.8
1987	612	4.2	3.9-4.6	196	3.0	2.5-3.4	416	5.4	4.9-5.9	80	0.5	0.4-0.6	25	0.4	0.2-0.6	55	0.6	0.4-0.8
1988	608	4.2	3.9-4.5	174	2.7	2.3-3.1	434	5.7	5.1-6.2	78	0.5	0.4-0.6	29	0.5	0.3-0.7	49	0.5	0.4-0.7
1989	599	4.2	3.8-4.5	165	2.5	2.1-2.9	434	5.7	5.2-6.3	82	0.5	0.4-0.7	24	0.4	0.2-0.6	58	0.6	0.5-0.8
1990	641	4.4	4.1-4.8	185	2.8	2.4-3.2	456	5.9	5.3-6.5	77	0.5	0.4-0.6	29	0.4	0.3-0.6	48	0.5	0.4-0.7
1991	723	5.0	4.6-5.3	209	3.1	2.7-3.5	514	6.7	6.2-7.4	72	0.5	0.4-0.6	23	0.4	0.2-0.6	49	0.5	0.4-0.7
1992	744	5.1	4.7-5.4	199	3.0	2.6-3.5	545	7.0	6.4-7.6	81	0.5	0.4-0.7	26	0.4	0.3-0.6	55	0.6	0.4-0.7
1993	774	5.3	4.9-5.7	210	3.1	2.7-3.5	564	7.4	6.8-8.1	83	0.5	0.4-0.7	30	0.5	0.3-0.7	53	0.6	0.4-0.7
1994	839	5.7	5.3-6.1	224	3.3	2.9-3.8	615	8.1	7.4-8.7	81	0.5	0.4-0.7	35	0.6	0.4-0.8	46	0.5	0.3-0.6
1995	932	6.3	5.9-6.7	260	3.8	3.4-4.3	672	8.8	8.1-9.5	78	0.5	0.4-0.6	31	0.5	0.3-0.7	47	0.5	0.4-0.7
1996	987	6.8	6.3-7.2	252	3.7	3.2-4.2	735	9.7	9.0-10.5	112	0.7	0.6-0.9	38	0.6	0.4-0.8	74	0.8	0.6-1.0
1997	1,093	7.4	7.0-7.9	295	4.3	3.8-4.8	798	10.4	9.7-11.2	85	0.5	0.4-0.7	32	0.5	0.3-0.7	53	0.5	0.4-0.7
1998	1,121	7.6	7.2-8.1	316	4.6	4.1-5.1	805	10.5	9.8-11.2	78	0.5	0.4-0.6	25	0.4	0.2-0.6	53	0.5	0.4-0.7
1999	1,317	8.9	8.4-9.4	315	4.5	4.0-5.0	1,002	13.2	12.4-14.0	76	0.5	0.4-0.6	30	0.4	0.3-0.6	46	0.5	0.3-0.6
2000	1,442	9.7	9.2-10.2	370	5.2	4.7-5.8	1,072	14.1	13.2-14.9	80	0.5	0.4-0.6	25	0.4	0.2-0.6	55	0.6	0.4-0.7
2001	1,522	10.2	9.7-10.8	349	4.9	4.4-5.4	1,173	15.3	14.5-16.3	80	0.5	0.4-0.6	33	0.5	0.3-0.7	47	0.5	0.4-0.7
2002	1,572	10.5	10.0-11.0	422	5.9	5.3-6.5	1,150	15.0	14.1-15.9	80	0.5	0.4-0.6	29	0.4	0.3-0.6	51	0.5	0.4-0.7
2003	1,600	10.7	10.1-11.2	418	5.8	5.3-6.4	1,182	15.3	14.5-16.2	77	0.5	0.4-0.6	23	0.3	0.2-0.5	54	0.5	0.4-0.7
2004	1,785	11.9	11.3-12.5	438	6.0	5.5-6.6	1,347	17.5	16.6-18.5	68	0.4	0.3-0.5	30	0.4	0.3-0.6	38	0.4	0.2-0.5
2005	2,119	14.2	13.6-14.8	536	7.4	6.8-8.1	1,583	20.8	19.8-21.8	83	0.5	0.4-0.6	40	0.6	0.4-0.8	43	0.4	0.3-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	2,231	14.9	14.3-15.5	561	7.7	7.1-8.4	1,670	21.8	20.8-22.9	96	0.6	0.5-0.7	38	0.5	0.4-0.8	58	0.6	0.4-0.7
2007	2,448	16.3	15.6-16.9	618	8.5	7.8-9.2	1,830	23.8	22.7-24.9	93	0.6	0.4-0.7	37	0.5	0.4-0.7	56	0.6	0.4-0.8
2008	2,784	18.5	17.8-19.2	660	9.0	8.3-9.7	2,124	27.8	26.6-29.1	84	0.5	0.4-0.6	40	0.6	0.4-0.8	44	0.5	0.3-0.6
2009	2,992	19.9	19.1-20.6	770	10.5	9.7-11.3	2,222	29.0	27.8-30.2	96	0.5	0.4-0.7	43	0.6	0.4-0.8	53	0.5	0.4-0.7
2010	2,986	19.8	19.0-20.5	754	10.1	9.4-10.9	2,232	29.2	28.0-30.5	90	0.5	0.4-0.6	39	0.5	0.4-0.7	51	0.5	0.3-0.6
2011	3,102	20.4	19.7-21.2	782	10.5	9.7-11.3	2,320	30.0	28.8-31.3	101	0.6	0.5-0.7	49	0.7	0.5-0.9	52	0.5	0.4-0.6
2012	3,184	21.0	20.3-21.8	834	11.0	10.3-11.8	2,350	30.8	29.6-32.2	92	0.5	0.4-0.6	41	0.5	0.4-0.7	51	0.5	0.4-0.7
2013	3,335	22.0	21.2-22.8	918	12.1	11.3-13.0	2,417	31.8	30.5-33.1	102	0.6	0.4-0.7	46	0.6	0.4-0.8	56	0.5	0.4-0.7
2014	3,299	21.7	21.0-22.5	885	11.6	10.8-12.4	2,414	31.8	30.5-33.1	86	0.4	0.4-0.6	38	0.5	0.3-0.6	48	0.4	0.3-0.6
2015	3,246	21.6	20.8-22.4	861	11.3	10.5-12.1	2,385	31.8	30.5-33.2	97	0.5	0.4-0.6	45	0.5	0.4-0.7	52	0.5	0.4-0.6
2016	3,075	20.4	19.7-21.2	822	10.7	10.0-11.5	2,253	30.0	28.7-31.4	102	0.5	0.4-0.7	39	0.5	0.4-0.7	63	0.6	0.5-0.8
2017	2,942	19.7	18.9-20.4	783	10.4	9.7-11.2	2,159	28.8	27.5-30.1	94	0.5	0.4-0.6	41	0.5	0.4-0.7	53	0.5	0.4-0.7
2018	2,761	18.4	17.7-19.2	802	10.6	9.8-11.4	1,959	26.2	25.0-27.5	96	0.5	0.4-0.6	38	0.4	0.3-0.6	58	0.5	0.4-0.7
2019	2,969	19.9	19.2-20.7	867	11.3	10.5-12.1	2,102	28.6	27.3-29.9	102	0.5	0.4-0.6	50	0.6	0.4-0.8	52	0.4	0.3-0.6
2020	2,376	15.9	15.3-16.6	704	9.3	8.6-10.0	1,672	22.6	21.4-23.7	86	0.5	0.4-0.6	34	0.4	0.3-0.6	52	0.5	0.4-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	567	3.5	3.2-3.8	307	4.1	3.6-4.6	260	3.0	2.7-3.5	217	1.4	1.2-1.6	126	1.8	1.5-2.2	91	1.1	0.9-1.3
1977	537	3.4	3.1-3.7	302	4.1	3.6-4.6	235	2.8	2.5-3.2	225	1.5	1.3-1.7	128	1.9	1.6-2.3	97	1.2	1.0-1.5
1978	571	3.6	3.3-3.9	299	4.1	3.6-4.6	272	3.2	2.8-3.6	197	1.3	1.1-1.5	109	1.6	1.3-1.9	88	1.1	0.8-1.3
1979	575	3.8	3.4-4.1	313	4.5	4.0-5.0	262	3.2	2.8-3.6	170	1.1	1.0-1.3	97	1.5	1.2-1.8	73	0.9	0.7-1.1
1980	597	3.9	3.5-4.2	327	4.6	4.1-5.2	270	3.2	2.8-3.6	203	1.4	1.2-1.6	112	1.7	1.4-2.1	91	1.1	0.8-1.3
1981	564	3.6	3.3-4.0	300	4.1	3.7-4.6	264	3.2	2.8-3.6	176	1.2	1.0-1.4	98	1.5	1.2-1.8	78	0.9	0.7-1.2
1982	579	3.7	3.4-4.1	300	4.3	3.8-4.8	279	3.3	2.9-3.7	182	1.2	1.0-1.4	97	1.4	1.1-1.7	85	1.0	0.8-1.3
1983	546	3.6	3.3-3.9	312	4.4	3.9-5.0	234	2.8	2.5-3.2	152	1.0	0.8-1.2	88	1.3	1.0-1.6	64	0.7	0.5-0.9
1984	568	3.7	3.4-4.0	302	4.3	3.8-4.9	266	3.2	2.8-3.6	157	1.1	0.9-1.2	93	1.4	1.1-1.7	64	0.7	0.6-1.0
1985	606	4.0	3.6-4.3	330	4.6	4.1-5.2	276	3.4	3.0-3.8	155	1.0	0.8-1.2	84	1.2	0.9-1.5	71	0.8	0.6-1.1
1986	563	3.6	3.3-4.0	307	4.2	3.7-4.7	256	3.1	2.7-3.5	174	1.1	1.0-1.3	99	1.5	1.2-1.8	75	0.9	0.7-1.1
1987	521	3.4	3.1-3.8	268	3.8	3.3-4.3	253	3.2	2.8-3.6	140	0.9	0.8-1.1	86	1.2	1.0-1.5	54	0.7	0.5-0.9
1988	562	3.7	3.4-4.1	313	4.4	3.9-5.0	249	3.1	2.7-3.5	119	0.8	0.7-1.0	76	1.1	0.9-1.4	43	0.5	0.4-0.7
1989	557	3.7	3.4-4.0	311	4.4	3.9-4.9	246	3.1	2.7-3.5	149	1.0	0.9-1.2	95	1.4	1.2-1.8	54	0.7	0.5-0.9
1990	527	3.6	3.3-3.9	286	4.1	3.6-4.6	241	3.1	2.7-3.5	137	0.9	0.8-1.1	84	1.3	1.0-1.6	53	0.6	0.5-0.8
1991	546	3.7	3.4-4.0	291	4.2	3.7-4.7	255	3.3	2.9-3.7	110	0.7	0.6-0.9	61	0.9	0.7-1.2	49	0.6	0.4-0.8
1992	561	3.8	3.5-4.1	305	4.4	3.9-4.9	256	3.3	2.9-3.7	128	0.8	0.7-1.0	64	0.9	0.7-1.2	64	0.8	0.6-1.0
1993	505	3.5	3.2-3.8	251	3.6	3.2-4.1	254	3.3	2.9-3.7	102	0.7	0.5-0.8	55	0.8	0.6-1.1	47	0.6	0.4-0.8
1994	571	3.9	3.6-4.2	300	4.3	3.8-4.8	271	3.5	3.1-3.9	113	0.7	0.6-0.9	55	0.8	0.6-1.1	58	0.7	0.5-0.9
1995	514	3.6	3.3-3.9	277	4.0	3.5-4.5	237	3.2	2.8-3.6	100	0.7	0.5-0.8	50	0.7	0.5-1.0	50	0.6	0.4-0.8
1996	494	3.4	3.1-3.7	251	3.6	3.2-4.1	243	3.3	2.9-3.7	89	0.6	0.5-0.7	45	0.7	0.5-0.9	44	0.6	0.4-0.7
1997	482	3.4	3.1-3.7	268	3.8	3.4-4.3	214	2.9	2.5-3.4	96	0.6	0.5-0.8	53	0.8	0.6-1.0	43	0.5	0.4-0.7
1998	540	3.7	3.4-4.1	275	4.0	3.5-4.5	265	3.6	3.1-4.0	93	0.6	0.5-0.8	48	0.7	0.5-0.9	45	0.6	0.4-0.8
1999	503	3.5	3.2-3.8	269	3.8	3.4-4.3	234	3.1	2.7-3.6	83	0.5	0.4-0.7	39	0.6	0.4-0.8	44	0.5	0.4-0.7
2000	538	3.7	3.4-4.0	291	4.2	3.7-4.7	247	3.3	2.9-3.8	86	0.6	0.4-0.7	44	0.7	0.5-0.9	42	0.5	0.3-0.7
2001	538	3.7	3.4-4.0	278	4.0	3.5-4.5	260	3.5	3.1-4.0	85	0.6	0.4-0.7	45	0.6	0.5-0.9	40	0.5	0.3-0.6
2002	587	4.0	3.7-4.4	328	4.7	4.2-5.2	259	3.5	3.1-3.9	93	0.6	0.5-0.7	48	0.7	0.5-0.9	45	0.5	0.4-0.7
2003	516	3.6	3.3-3.9	282	4.0	3.6-4.5	234	3.1	2.7-3.6	70	0.5	0.4-0.6	32	0.5	0.3-0.6	38	0.5	0.3-0.6
2004	545	3.8	3.5-4.1	281	4.0	3.6-4.5	264	3.6	3.2-4.0	75	0.5	0.4-0.6	44	0.6	0.4-0.8	31	0.4	0.2-0.5
2005	554	3.9	3.5-4.2	305	4.4	3.9-4.9	249	3.4	3.0-3.9	70	0.4	0.3-0.6	34	0.5	0.3-0.7	36	0.4	0.3-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	526	3.6	3.3-4.0	293	4.2	3.7-4.7	233	3.1	2.7-3.6	71	0.5	0.4-0.6	41	0.6	0.4-0.8	30	0.4	0.2-0.5
2007	535	3.7	3.4-4.0	266	3.8	3.3-4.3	269	3.6	3.2-4.1	84	0.5	0.4-0.7	42	0.6	0.4-0.8	42	0.5	0.3-0.6
2008	520	3.6	3.3-3.9	274	3.9	3.4-4.3	246	3.3	2.9-3.8	84	0.5	0.4-0.6	50	0.7	0.5-0.9	34	0.4	0.2-0.5
2009	527	3.7	3.3-4.0	293	4.1	3.7-4.6	234	3.2	2.8-3.7	63	0.4	0.3-0.5	35	0.5	0.3-0.7	28	0.3	0.2-0.5
2010	512	3.5	3.2-3.8	285	4.0	3.5-4.5	227	3.0	2.6-3.4	66	0.4	0.3-0.5	38	0.5	0.4-0.7	28	0.3	0.2-0.5
2011	522	3.6	3.3-4.0	290	4.1	3.7-4.6	232	3.2	2.8-3.6	69	0.4	0.3-0.5	43	0.6	0.4-0.8	26	0.3	0.2-0.4
2012	520	3.6	3.3-4.0	301	4.2	3.8-4.8	219	3.1	2.7-3.5	67	0.4	0.3-0.5	31	0.4	0.3-0.6	36	0.4	0.3-0.5
2013	500	3.4	3.1-3.8	244	3.4	3.0-3.9	256	3.5	3.1-4.0	55	0.3	0.2-0.4	27	0.4	0.2-0.5	28	0.3	0.2-0.4
2014	548	3.7	3.4-4.1	292	4.1	3.6-4.6	256	3.4	3.0-3.8	59	0.3	0.3-0.5	25	0.3	0.2-0.5	34	0.4	0.2-0.5
2015	474	3.3	3.0-3.7	269	3.8	3.4-4.3	205	2.9	2.5-3.3	53	0.3	0.2-0.4	26	0.4	0.2-0.5	27	0.3	0.2-0.4
2016	533	3.7	3.3-4.0	270	3.8	3.3-4.3	263	3.6	3.1-4.0	62	0.4	0.3-0.5	36	0.5	0.3-0.6	26	0.3	0.2-0.4
2017	485	3.3	3.0-3.6	268	3.7	3.2-4.2	217	3.0	2.6-3.4	47	0.3	0.2-0.4	25	0.3	0.2-0.5	22	0.2	0.1-0.3
2018	433	2.9	2.6-3.2	251	3.4	3.0-3.9	182	2.4	2.1-2.8	56	0.3	0.2-0.4	38	0.5	0.3-0.7	18	0.2	0.1-0.3
2019	489	3.4	3.1-3.7	260	3.6	3.2-4.1	229	3.2	2.8-3.7	52	0.3	0.2-0.4	35	0.4	0.3-0.6	17	0.2	0.1-0.3
2020	425	3.0	2.8-3.4	237	3.4	3.0-3.9	188	2.7	2.3-3.1	41	0.2	0.2-0.3	24	0.3	0.2-0.4	17	0.2	0.1-0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,756	11.6	11.0-12.1	902	14.0	13.0-14.9	854	9.9	9.2-10.6	970	6.4	6.0-6.8	519	8.2	7.5-9.0	451	5.1	4.7-5.7
1977	1,787	11.7	11.2-12.3	909	13.9	13.0-14.9	878	10.1	9.4-10.8	972	6.4	6.0-6.8	475	7.4	6.7-8.1	497	5.6	5.1-6.1
1978	1,907	12.5	12.0-13.1	992	15.1	14.1-16.1	915	10.5	9.8-11.2	989	6.5	6.1-7.0	510	8.1	7.4-8.9	479	5.4	5.0-6.0
1979	1,863	12.4	11.8-13.0	913	14.4	13.4-15.4	950	10.9	10.2-11.6	994	6.6	6.2-7.0	521	8.3	7.6-9.1	473	5.3	4.9-5.9
1980	2,020	13.3	12.7-13.9	1,018	15.6	14.6-16.6	1,002	11.5	10.8-12.2	1,019	6.7	6.3-7.1	503	7.8	7.1-8.5	516	5.8	5.3-6.4
1981	2,032	13.4	12.8-14.0	988	15.4	14.5-16.5	1,044	11.9	11.1-12.6	1,014	6.6	6.2-7.1	505	8.0	7.3-8.8	509	5.6	5.1-6.1
1982	2,127	14.0	13.4-14.6	1,053	16.3	15.3-17.3	1,074	12.3	11.5-13.0	1,057	7.0	6.6-7.4	528	8.6	7.9-9.4	529	5.9	5.4-6.4
1983	2,230	14.7	14.0-15.3	1,118	17.4	16.4-18.5	1,112	12.5	11.7-13.3	1,119	7.4	7.0-7.8	573	9.1	8.3-9.9	546	6.1	5.6-6.7
1984	2,288	15.1	14.5-15.7	1,186	18.8	17.7-20.0	1,102	12.4	11.6-13.1	1,067	7.0	6.6-7.4	568	9.2	8.5-10.1	499	5.4	5.0-5.9
1985	2,453	16.2	15.5-16.9	1,302	20.3	19.2-21.5	1,151	12.9	12.2-13.7	1,181	7.7	7.3-8.2	625	10.0	9.2-10.8	556	6.1	5.6-6.6
1986	2,576	17.0	16.4-17.7	1,375	21.3	20.1-22.5	1,201	13.6	12.8-14.4	1,176	7.8	7.3-8.2	644	10.4	9.6-11.2	532	5.8	5.3-6.3
1987	2,555	16.9	16.3-17.6	1,366	21.3	20.2-22.5	1,189	13.4	12.6-14.2	1,192	7.8	7.4-8.3	598	9.5	8.7-10.3	594	6.5	6.0-7.0
1988	2,594	17.2	16.6-17.9	1,354	21.0	19.8-22.1	1,240	14.3	13.5-15.1	1,239	8.1	7.7-8.6	616	9.9	9.1-10.7	623	6.8	6.2-7.3
1989	2,693	18.0	17.3-18.7	1,472	22.7	21.6-24.0	1,221	14.1	13.3-14.9	1,243	8.2	7.7-8.6	677	10.8	10.0-11.7	566	6.2	5.7-6.7
1990	2,895	19.1	18.4-19.9	1,586	24.4	23.2-25.6	1,309	14.8	14.0-15.7	1,283	8.4	7.9-8.9	648	10.4	9.6-11.2	635	6.9	6.4-7.5
1991	3,002	19.9	19.2-20.6	1,629	25.4	24.2-26.7	1,373	15.7	14.9-16.6	1,280	8.4	7.9-8.8	674	10.8	10.0-11.7	606	6.5	6.0-7.1
1992	2,893	19.1	18.4-19.9	1,555	23.8	22.6-25.0	1,338	15.3	14.5-16.2	1,335	8.7	8.2-9.1	676	10.8	10.0-11.7	659	7.1	6.6-7.7
1993	2,981	19.7	19.0-20.5	1,633	24.8	23.6-26.1	1,348	15.4	14.6-16.3	1,243	8.0	7.6-8.5	631	10.0	9.2-10.8	612	6.6	6.0-7.1
1994	3,277	21.6	20.9-22.4	1,821	27.6	26.3-28.9	1,456	16.6	15.7-17.5	1,419	9.1	8.7-9.6	728	11.4	10.6-12.3	691	7.4	6.9-8.0
1995	3,220	21.2	20.5-21.9	1,721	26.1	24.9-27.4	1,499	17.1	16.3-18.0	1,409	9.1	8.6-9.5	731	11.6	10.8-12.5	678	7.3	6.7-7.8
1996	3,349	21.9	21.2-22.7	1,795	27.2	26.0-28.5	1,554	17.8	16.9-18.7	1,426	9.1	8.7-9.6	741	11.7	10.9-12.6	685	7.4	6.8-7.9
1997	3,354	21.8	21.1-22.6	1,736	26.1	24.9-27.4	1,618	18.3	17.4-19.2	1,478	9.4	8.9-9.9	747	11.5	10.7-12.4	731	7.8	7.3-8.4
1998	3,318	21.4	20.7-22.2	1,706	25.3	24.1-26.6	1,612	18.3	17.4-19.2	1,382	8.7	8.3-9.2	662	10.0	9.3-10.8	720	7.6	7.1-8.2
1999	3,368	21.6	20.8-22.3	1,699	25.1	24.0-26.4	1,669	18.8	17.9-19.7	1,312	8.2	7.7-8.6	637	9.8	9.0-10.6	675	7.1	6.6-7.7
2000	3,384	21.5	20.7-22.2	1,728	25.3	24.1-26.5	1,656	18.4	17.5-19.3	1,354	8.4	7.9-8.8	666	10.1	9.3-10.9	688	7.2	6.7-7.8
2001	3,353	21.1	20.4-21.8	1,731	25.3	24.1-26.5	1,622	17.9	17.1-18.9	1,261	7.7	7.3-8.2	688	10.2	9.5-11.0	573	5.8	5.3-6.3
2002	3,412	21.4	20.7-22.1	1,833	26.3	25.1-27.6	1,579	17.5	16.6-18.4	1,223	7.4	7.0-7.9	654	9.7	9.0-10.5	569	5.9	5.4-6.4
2003	3,609	22.5	21.7-23.2	1,847	26.3	25.1-27.6	1,762	19.3	18.4-20.3	1,134	6.8	6.4-7.2	572	8.5	7.8-9.2	562	5.6	5.1-6.1
2004	3,649	22.6	21.9-23.4	1,930	27.4	26.1-28.6	1,719	19.0	18.1-19.9	1,144	6.8	6.4-7.2	561	8.2	7.5-8.9	583	5.8	5.3-6.3
2005	3,743	23.1	22.4-23.9	1,963	27.6	26.4-28.9	1,780	19.5	18.6-20.5	1,111	6.6	6.2-7.0	582	8.4	7.8-9.2	529	5.3	4.8-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	3,867	23.9	23.1-24.6	2,035	28.5	27.3-29.8	1,832	20.0	19.1-20.9	1,207	7.2	6.8-7.6	617	8.8	8.1-9.6	590	5.8	5.3-6.3
2007	3,840	23.5	22.7-24.2	2,034	28.2	27.0-29.5	1,806	19.8	18.9-20.7	1,088	6.4	6.0-6.8	596	8.4	7.7-9.1	492	4.9	4.5-5.4
2008	3,911	23.7	23.0-24.5	2,080	28.5	27.3-29.8	1,831	20.0	19.1-21.0	1,130	6.6	6.2-7.0	601	8.4	7.7-9.1	529	5.3	4.8-5.8
2009	3,999	24.0	23.3-24.8	2,115	28.7	27.5-30.0	1,884	20.4	19.4-21.3	1,080	6.2	5.8-6.6	599	8.2	7.6-8.9	481	4.7	4.3-5.2
2010	3,967	23.6	22.8-24.4	2,098	28.2	27.0-29.5	1,869	19.8	18.9-20.8	1,082	6.1	5.8-6.5	564	7.7	7.1-8.4	518	5.0	4.6-5.5
2011	3,944	23.4	22.6-24.1	2,116	28.1	26.9-29.3	1,828	19.6	18.7-20.5	1,080	6.1	5.7-6.5	565	7.6	6.9-8.2	515	5.0	4.6-5.5
2012	3,979	23.2	22.5-24.0	2,165	28.4	27.2-29.6	1,814	19.1	18.2-20.1	1,042	5.8	5.5-6.2	575	7.7	7.1-8.4	467	4.4	4.0-4.8
2013	4,087	23.6	22.9-24.4	2,223	28.8	27.6-30.1	1,864	19.4	18.5-20.3	981	5.4	5.1-5.8	530	7.0	6.4-7.6	451	4.2	3.8-4.7
2014	4,152	23.7	22.9-24.4	2,268	29.1	27.8-30.3	1,884	19.4	18.5-20.3	1,051	5.7	5.3-6.0	579	7.5	6.9-8.1	472	4.3	3.9-4.7
2015	4,129	23.4	22.7-24.2	2,263	28.4	27.2-29.6	1,866	19.4	18.5-20.3	990	5.4	5.1-5.8	562	7.2	6.6-7.9	428	4.0	3.6-4.4
2016	4,192	23.5	22.8-24.3	2,225	27.6	26.5-28.8	1,967	20.2	19.3-21.2	1,047	5.5	5.2-5.9	563	7.0	6.4-7.6	484	4.4	4.0-4.8
2017	4,174	23.2	22.5-23.9	2,269	27.9	26.7-29.1	1,905	19.4	18.5-20.4	1,070	5.6	5.3-6.0	593	7.3	6.7-8.0	477	4.3	3.9-4.7
2018	4,051	22.5	21.8-23.2	2,132	25.9	24.8-27.1	1,919	19.7	18.8-20.7	914	4.7	4.4-5.0	523	6.4	5.8-7.0	391	3.5	3.1-3.9
2019	4,196	22.8	22.1-23.5	2,288	27.3	26.2-28.5	1,908	19.2	18.3-20.1	939	4.8	4.5-5.1	519	6.2	5.7-6.8	420	3.7	3.3-4.1
2020	3,692	20.1	19.4-20.8	1,992	23.7	22.7-24.8	1,700	17.2	16.3-18.0	888	4.5	4.2-4.8	474	5.6	5.1-6.2	414	3.6	3.3-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	638	4.2	3.8-4.5	322	5.2	4.6-5.8	316	3.5	3.1-3.9	409	2.8	2.5-3.0	211	3.6	3.1-4.1	198	2.3	2.0-2.6
1977	629	4.1	3.8-4.5	315	5.1	4.5-5.8	314	3.5	3.1-3.9	410	2.6	2.4-2.9	203	3.2	2.7-3.7	207	2.3	2.0-2.6
1978	645	4.3	3.9-4.6	304	4.9	4.4-5.5	341	3.8	3.4-4.3	400	2.6	2.4-2.9	194	3.1	2.7-3.6	206	2.3	2.0-2.6
1979	683	4.5	4.2-4.9	326	5.3	4.7-6.0	357	4.0	3.6-4.4	431	2.9	2.6-3.1	202	3.3	2.9-3.9	229	2.5	2.2-2.9
1980	650	4.3	3.9-4.6	325	5.3	4.7-6.0	325	3.6	3.2-4.0	478	3.1	2.8-3.4	228	3.8	3.3-4.3	250	2.7	2.4-3.1
1981	752	4.9	4.6-5.3	359	5.9	5.2-6.5	393	4.4	3.9-4.8	457	3.0	2.7-3.3	218	3.7	3.2-4.2	239	2.6	2.3-3.0
1982	713	4.6	4.3-4.9	360	5.8	5.2-6.4	353	3.8	3.4-4.3	445	2.9	2.7-3.2	212	3.5	3.0-4.0	233	2.5	2.2-2.9
1983	710	4.6	4.3-5.0	377	6.3	5.6-7.0	333	3.6	3.2-4.0	458	3.0	2.7-3.3	223	3.7	3.2-4.3	235	2.5	2.2-2.9
1984	716	4.7	4.3-5.0	367	6.0	5.4-6.6	349	3.8	3.4-4.2	487	3.1	2.9-3.4	239	3.8	3.4-4.4	248	2.7	2.3-3.0
1985	736	4.8	4.4-5.1	379	6.1	5.5-6.8	357	3.9	3.5-4.3	488	3.1	2.9-3.4	257	4.2	3.7-4.8	231	2.4	2.1-2.8
1986	700	4.5	4.2-4.9	374	6.0	5.4-6.6	326	3.5	3.1-3.9	487	3.1	2.8-3.4	237	3.9	3.4-4.5	250	2.6	2.3-3.0
1987	752	4.9	4.5-5.3	386	6.3	5.7-7.0	366	3.9	3.5-4.3	494	3.2	2.9-3.5	257	4.2	3.7-4.8	237	2.6	2.2-2.9
1988	722	4.7	4.4-5.1	379	6.2	5.6-6.9	343	3.7	3.3-4.1	502	3.2	2.9-3.5	273	4.6	4.0-5.1	229	2.4	2.1-2.7
1989	769	5.0	4.7-5.4	350	5.7	5.1-6.4	419	4.6	4.2-5.1	470	3.0	2.7-3.3	236	3.8	3.3-4.3	234	2.5	2.2-2.8
1990	810	5.2	4.8-5.6	414	6.5	5.9-7.2	396	4.3	3.9-4.8	461	2.9	2.7-3.2	232	3.8	3.4-4.4	229	2.4	2.1-2.7
1991	752	4.8	4.5-5.2	380	6.0	5.4-6.7	372	4.0	3.6-4.5	479	3.1	2.8-3.4	231	3.7	3.3-4.3	248	2.6	2.3-3.0
1992	830	5.3	4.9-5.7	428	6.8	6.2-7.5	402	4.3	3.9-4.8	502	3.2	2.9-3.5	259	4.3	3.8-4.8	243	2.6	2.2-2.9
1993	821	5.3	4.9-5.7	404	6.4	5.8-7.0	417	4.6	4.1-5.0	562	3.6	3.3-3.9	279	4.5	4.0-5.1	283	2.9	2.6-3.3
1994	802	5.2	4.8-5.5	409	6.4	5.8-7.1	393	4.3	3.9-4.7	565	3.6	3.3-3.9	256	4.1	3.6-4.6	309	3.3	3.0-3.7
1995	843	5.4	5.1-5.8	434	6.8	6.2-7.5	409	4.5	4.1-5.0	553	3.5	3.2-3.8	282	4.5	4.0-5.0	271	2.9	2.5-3.2
1996	864	5.5	5.1-5.9	439	6.9	6.3-7.6	425	4.6	4.2-5.1	536	3.4	3.1-3.7	261	4.2	3.7-4.8	275	2.9	2.6-3.3
1997	834	5.3	5.0-5.7	408	6.3	5.7-6.9	426	4.7	4.2-5.2	508	3.2	2.9-3.5	249	4.0	3.5-4.5	259	2.8	2.4-3.1
1998	955	6.0	5.7-6.4	482	7.3	6.7-8.0	473	5.1	4.7-5.6	526	3.3	3.0-3.6	261	4.0	3.6-4.6	265	2.8	2.4-3.1
1999	961	6.0	5.6-6.4	481	7.3	6.6-8.0	480	5.2	4.7-5.7	543	3.3	3.1-3.6	276	4.2	3.8-4.8	267	2.7	2.4-3.1
2000	958	5.9	5.6-6.3	494	7.3	6.7-8.0	464	5.0	4.5-5.5	578	3.5	3.2-3.8	290	4.4	3.9-5.0	288	3.0	2.6-3.3
2001	946	5.8	5.5-6.2	530	7.9	7.2-8.6	416	4.4	3.9-4.8	485	2.9	2.7-3.2	261	3.9	3.5-4.4	224	2.3	2.0-2.6
2002	1,024	6.3	5.9-6.7	565	8.2	7.6-9.0	459	4.9	4.4-5.3	537	3.2	3.0-3.5	289	4.4	3.9-4.9	248	2.5	2.2-2.9
2003	990	6.0	5.6-6.4	543	7.9	7.2-8.6	447	4.8	4.3-5.2	506	3.0	2.8-3.3	274	4.1	3.6-4.6	232	2.3	2.0-2.6
2004	1,025	6.2	5.8-6.6	553	7.8	7.2-8.5	472	5.0	4.5-5.5	485	2.9	2.6-3.1	256	3.8	3.3-4.2	229	2.3	2.0-2.6
2005	1,021	6.1	5.8-6.5	548	7.7	7.1-8.4	473	4.9	4.5-5.4	513	3.0	2.8-3.3	264	3.8	3.4-4.3	249	2.5	2.2-2.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,057	6.4	6.0-6.8	575	8.0	7.4-8.7	482	5.1	4.6-5.6	486	2.9	2.6-3.2	281	4.0	3.6-4.5	205	2.1	1.8-2.4
2007	1,107	6.6	6.2-7.0	604	8.4	7.7-9.1	503	5.3	4.8-5.8	497	2.9	2.7-3.2	261	3.7	3.2-4.1	236	2.3	2.0-2.7
2008	1,138	6.7	6.3-7.1	630	8.6	7.9-9.3	508	5.3	4.8-5.8	496	2.9	2.6-3.1	268	3.8	3.3-4.3	228	2.3	2.0-2.6
2009	1,218	7.1	6.7-7.5	655	8.9	8.2-9.6	563	5.7	5.3-6.2	497	2.8	2.6-3.1	269	3.7	3.3-4.2	228	2.2	1.9-2.5
2010	1,216	7.0	6.6-7.4	664	8.7	8.1-9.4	552	5.7	5.3-6.2	534	3.0	2.8-3.3	271	3.7	3.3-4.2	263	2.5	2.2-2.9
2011	1,285	7.4	7.0-7.8	733	9.6	8.9-10.3	552	5.6	5.2-6.1	481	2.7	2.4-2.9	250	3.3	2.9-3.7	231	2.2	1.9-2.5
2012	1,256	7.1	6.7-7.5	709	9.0	8.3-9.7	547	5.6	5.1-6.1	517	2.8	2.6-3.1	256	3.4	3.0-3.8	261	2.5	2.2-2.8
2013	1,342	7.5	7.1-7.9	747	9.4	8.7-10.1	595	6.0	5.5-6.5	491	2.7	2.4-2.9	266	3.4	3.0-3.9	225	2.1	1.9-2.5
2014	1,459	8.1	7.6-8.5	807	10.0	9.3-10.8	652	6.5	6.0-7.0	538	2.9	2.6-3.1	291	3.6	3.2-4.1	247	2.3	2.0-2.6
2015	1,351	7.4	7.0-7.8	742	9.2	8.5-9.9	609	6.0	5.6-6.6	519	2.8	2.5-3.0	298	3.8	3.3-4.2	221	2.1	1.8-2.4
2016	1,409	7.7	7.3-8.1	822	9.9	9.3-10.7	587	5.9	5.4-6.4	525	2.7	2.5-3.0	266	3.2	2.9-3.7	259	2.4	2.1-2.7
2017	1,398	7.5	7.1-7.9	798	9.5	8.8-10.2	600	5.8	5.4-6.3	518	2.7	2.4-2.9	277	3.4	3.0-3.8	241	2.1	1.9-2.4
2018	1,312	7.0	6.6-7.4	743	8.8	8.1-9.4	569	5.5	5.0-6.0	476	2.4	2.2-2.6	252	2.9	2.6-3.3	224	2.0	1.7-2.2
2019	1,403	7.4	7.0-7.8	778	9.0	8.4-9.7	625	6.1	5.6-6.6	482	2.4	2.2-2.6	275	3.2	2.8-3.6	207	1.8	1.5-2.0
2020	1,184	6.2	5.8-6.5	667	7.7	7.1-8.3	517	5.0	4.5-5.4	430	2.2	2.0-2.4	239	2.8	2.5-3.2	191	1.6	1.4-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,872	12.6	12.1-13.2	1,055	17.1	16.0-18.2	817	9.6	9.0-10.3	1,227	8.4	7.9-8.9	678	11.4	10.5-12.4	549	6.5	5.9-7.0
1977	1,756	11.9	11.4-12.5	967	15.5	14.5-16.6	789	9.4	8.7-10.1	1,243	8.5	8.0-9.0	674	11.0	10.2-12.0	569	6.8	6.2-7.3
1978	1,842	12.6	12.0-13.2	1,032	16.9	15.8-18.0	810	9.7	9.0-10.4	1,203	8.1	7.7-8.6	636	10.4	9.6-11.3	567	6.7	6.1-7.3
1979	1,914	12.8	12.3-13.4	1,050	16.6	15.6-17.7	864	10.1	9.5-10.9	1,246	8.5	8.0-9.0	659	10.9	10.0-11.8	587	7.0	6.4-7.6
1980	2,010	13.5	12.9-14.1	1,071	17.3	16.3-18.4	939	11.1	10.4-11.9	1,256	8.4	8.0-8.9	698	11.4	10.6-12.3	558	6.4	5.9-7.0
1981	1,826	12.3	11.7-12.9	1,022	16.5	15.5-17.6	804	9.4	8.8-10.1	1,248	8.5	8.0-9.0	689	11.4	10.5-12.3	559	6.6	6.0-7.2
1982	1,900	12.8	12.3-13.4	1,065	17.4	16.3-18.5	835	9.8	9.1-10.5	1,295	8.7	8.3-9.2	728	12.2	11.2-13.1	567	6.6	6.1-7.2
1983	1,835	12.3	11.8-12.9	1,006	16.1	15.1-17.2	829	9.8	9.1-10.5	1,219	8.2	7.8-8.7	638	10.7	9.9-11.6	581	6.6	6.1-7.2
1984	1,967	13.2	12.6-13.8	1,055	17.1	16.1-18.2	912	10.5	9.8-11.2	1,211	8.1	7.6-8.5	605	9.9	9.1-10.8	606	6.9	6.3-7.4
1985	1,989	13.3	12.7-13.9	1,115	17.9	16.9-19.1	874	10.0	9.4-10.7	1,206	8.0	7.5-8.5	673	11.1	10.2-12.0	533	6.0	5.4-6.5
1986	1,944	13.0	12.4-13.6	1,045	16.8	15.7-17.8	899	10.4	9.8-11.2	1,194	8.0	7.5-8.5	653	10.6	9.8-11.5	541	6.1	5.6-6.7
1987	1,785	11.8	11.3-12.4	936	14.9	13.9-15.9	849	9.8	9.1-10.5	1,144	7.6	7.1-8.0	606	9.9	9.1-10.8	538	6.1	5.5-6.6
1988	1,844	12.3	11.7-12.9	1,012	16.3	15.2-17.3	832	9.7	9.0-10.4	1,153	7.6	7.2-8.1	653	10.7	9.9-11.6	500	5.6	5.1-6.1
1989	1,902	12.6	12.0-13.2	1,048	17.0	15.9-18.0	854	9.7	9.0-10.4	1,163	7.7	7.2-8.1	625	10.2	9.4-11.1	538	6.0	5.5-6.6
1990	2,061	13.7	13.1-14.3	1,114	17.6	16.5-18.6	947	11.0	10.2-11.7	1,151	7.6	7.1-8.0	586	9.6	8.8-10.4	565	6.3	5.8-6.9
1991	1,950	13.0	12.4-13.6	1,070	17.0	16.0-18.1	880	10.2	9.5-10.9	1,141	7.5	7.1-7.9	603	9.9	9.1-10.8	538	5.9	5.4-6.5
1992	1,995	13.2	12.6-13.8	1,120	17.9	16.8-19.0	875	10.1	9.4-10.8	1,166	7.6	7.2-8.1	642	10.5	9.7-11.3	524	5.8	5.3-6.3
1993	1,956	12.9	12.4-13.5	1,047	16.3	15.3-17.3	909	10.6	9.9-11.4	1,083	7.1	6.6-7.5	595	9.6	8.8-10.4	488	5.4	5.0-6.0
1994	2,170	14.2	13.6-14.8	1,175	18.4	17.3-19.5	995	11.2	10.5-12.0	1,226	7.9	7.5-8.4	673	10.7	9.9-11.6	553	6.0	5.5-6.5
1995	2,066	13.5	12.9-14.1	1,123	17.4	16.4-18.4	943	10.8	10.1-11.5	1,213	7.8	7.4-8.3	671	10.9	10.0-11.7	542	5.9	5.4-6.4
1996	2,132	13.9	13.3-14.5	1,222	18.7	17.7-19.8	910	10.3	9.7-11.1	1,235	7.9	7.5-8.4	661	10.3	9.6-11.2	574	6.3	5.8-6.9
1997	2,184	14.2	13.6-14.8	1,180	18.1	17.1-19.2	1,004	11.3	10.6-12.1	1,204	7.7	7.2-8.1	669	10.5	9.7-11.3	535	5.7	5.2-6.2
1998	2,187	14.1	13.5-14.7	1,248	18.7	17.7-19.8	939	10.6	9.9-11.3	1,190	7.5	7.1-8.0	659	10.1	9.3-10.9	531	5.7	5.2-6.2
1999	2,400	15.3	14.7-16.0	1,313	19.8	18.8-20.9	1,087	12.1	11.4-12.9	1,260	7.9	7.4-8.3	685	10.5	9.7-11.3	575	6.1	5.6-6.6
2000	2,451	15.5	14.9-16.1	1,351	20.2	19.1-21.3	1,100	12.1	11.4-12.9	1,222	7.6	7.2-8.0	655	9.9	9.2-10.7	567	6.0	5.5-6.5
2001	2,448	15.5	14.8-16.1	1,330	19.7	18.6-20.8	1,118	12.4	11.7-13.1	1,214	7.5	7.0-7.9	648	9.8	9.1-10.6	566	5.9	5.4-6.4
2002	2,286	14.4	13.8-15.0	1,303	19.0	18.0-20.1	983	10.9	10.2-11.6	1,208	7.4	7.0-7.8	661	9.9	9.1-10.7	547	5.7	5.2-6.2
2003	2,520	15.8	15.1-16.4	1,394	20.3	19.2-21.4	1,126	12.5	11.7-13.2	1,155	7.0	6.6-7.4	648	9.7	8.9-10.4	507	5.2	4.8-5.7
2004	2,551	15.9	15.3-16.5	1,402	20.2	19.2-21.3	1,149	12.7	11.9-13.5	1,260	7.6	7.2-8.0	691	10.2	9.4-11.0	569	5.9	5.4-6.4
2005	2,608	16.2	15.6-16.8	1,449	20.6	19.6-21.7	1,159	12.6	11.9-13.4	1,218	7.3	6.9-7.7	693	10.1	9.4-10.9	525	5.3	4.9-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
White Individuals
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	2,573	16.0	15.4-16.6	1,467	21.0	19.9-22.1	1,106	12.2	11.4-12.9	1,230	7.4	7.0-7.8	671	9.7	9.0-10.5	559	5.7	5.2-6.2
2007	2,687	16.6	16.0-17.3	1,546	21.7	20.6-22.8	1,141	12.8	12.0-13.6	1,178	7.1	6.7-7.5	665	9.6	8.9-10.3	513	5.3	4.9-5.8
2008	2,622	16.1	15.5-16.8	1,493	20.8	19.7-21.9	1,129	12.6	11.8-13.4	1,219	7.2	6.8-7.6	683	9.7	9.0-10.4	536	5.4	5.0-6.0
2009	2,826	17.2	16.5-17.8	1,622	22.3	21.2-23.4	1,204	13.2	12.4-14.0	1,221	7.2	6.8-7.7	689	9.7	9.0-10.5	532	5.4	5.0-5.9
2010	2,959	17.8	17.2-18.5	1,706	23.3	22.2-24.4	1,253	13.6	12.8-14.4	1,154	6.7	6.3-7.1	651	9.0	8.3-9.8	503	5.0	4.6-5.5
2011	3,109	18.5	17.8-19.2	1,776	23.9	22.8-25.0	1,333	14.3	13.5-15.1	1,223	7.0	6.6-7.4	701	9.5	8.8-10.2	522	5.1	4.7-5.6
2012	2,974	17.5	16.9-18.2	1,735	22.9	21.8-24.0	1,239	13.4	12.6-14.2	1,196	6.8	6.4-7.2	695	9.4	8.7-10.2	501	4.8	4.4-5.3
2013	3,188	18.6	18.0-19.3	1,864	24.4	23.3-25.6	1,324	14.1	13.3-14.9	1,205	6.8	6.4-7.2	691	9.2	8.5-9.9	514	5.0	4.6-5.5
2014	3,233	18.7	18.0-19.3	1,880	24.2	23.1-25.4	1,353	14.3	13.5-15.1	1,149	6.3	5.9-6.7	631	8.3	7.6-8.9	518	4.9	4.5-5.4
2015	3,249	18.6	18.0-19.3	1,884	24.0	22.9-25.2	1,365	14.3	13.5-15.1	1,187	6.4	6.1-6.8	680	8.8	8.1-9.5	507	4.8	4.3-5.2
2016	3,261	18.3	17.6-18.9	1,939	24.1	23.0-25.2	1,322	13.5	12.8-14.3	1,149	6.2	5.9-6.6	641	8.2	7.6-8.9	508	4.8	4.4-5.3
2017	3,110	17.5	16.9-18.2	1,840	22.8	21.8-23.9	1,270	13.3	12.6-14.1	1,094	5.7	5.4-6.1	607	7.6	7.0-8.2	487	4.4	4.0-4.8
2018	2,968	16.8	16.2-17.4	1,747	21.7	20.6-22.8	1,221	12.8	12.0-13.5	1,111	5.8	5.5-6.2	626	7.7	7.1-8.4	485	4.5	4.1-4.9
2019	2,932	16.2	15.6-16.8	1,733	21.1	20.1-22.1	1,199	12.1	11.4-12.9	1,129	5.9	5.6-6.3	653	8.0	7.4-8.7	476	4.4	4.0-4.8
2020	2,714	15.2	14.6-15.8	1,619	19.8	18.8-20.8	1,095	11.3	10.6-12.1	1,039	5.4	5.0-5.7	618	7.4	6.9-8.1	421	3.8	3.4-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	23,699	388.9	383.8-394.1	11,220	448.2	439.5-457.0	12,479	359.6	353.1-366.3	13,251	217.6	213.8-221.4	6,744	275.0	268.1-282.0	6,507	183.0	178.4-187.7
1977	23,302	389.6	384.4-394.8	11,140	454.4	445.6-463.3	12,162	356.5	349.9-363.2	13,069	216.7	212.9-220.6	6,612	273.5	266.7-280.5	6,457	182.8	178.2-187.5
1978	23,175	393.0	387.8-398.2	11,086	460.2	451.3-469.2	12,089	358.9	352.2-365.6	12,490	210.0	206.2-213.8	6,360	267.3	260.5-274.3	6,130	175.8	171.3-180.4
1979	23,491	403.4	398.1-408.8	11,239	473.7	464.6-482.8	12,252	368.7	361.9-375.6	12,323	209.5	205.7-213.3	6,199	264.6	257.8-271.5	6,124	177.1	172.5-181.8
1980	23,953	416.6	411.2-422.1	11,600	496.6	487.3-506.0	12,353	376.2	369.2-383.2	12,604	215.8	211.9-219.7	6,332	275.9	268.9-283.0	6,272	181.0	176.4-185.7
1981	23,487	412.4	407.0-417.9	11,116	482.1	472.9-491.4	12,371	377.7	370.8-384.8	12,085	209.7	205.9-213.6	5,935	261.1	254.3-268.1	6,150	180.2	175.5-184.9
1982	23,670	420.8	415.3-426.4	11,185	492.7	483.3-502.2	12,485	385.4	378.4-392.6	11,758	205.5	201.7-209.4	5,831	259.5	252.7-266.5	5,927	175.1	170.5-179.8
1983	23,762	424.2	418.7-429.8	11,323	499.5	490.1-509.1	12,439	385.7	378.6-392.8	11,567	203.5	199.7-207.3	5,680	256.1	249.3-263.1	5,887	173.3	168.7-178.0
1984	23,785	428.3	422.7-433.9	11,335	503.9	494.4-513.5	12,450	390.1	383.0-397.4	11,662	206.3	202.5-210.2	5,792	261.6	254.7-268.6	5,870	174.5	169.9-179.2
1985	23,956	434.4	428.8-440.1	11,645	519.6	509.9-529.4	12,311	387.3	380.1-394.5	11,308	201.2	197.5-205.1	5,653	255.7	248.9-262.7	5,655	169.8	165.3-174.5
1986	23,885	437.7	432.0-443.4	11,542	521.4	511.7-531.2	12,343	392.1	384.9-399.5	11,001	196.7	193.0-200.5	5,321	242.7	236.1-249.5	5,680	170.6	166.1-175.3
1987	23,426	433.1	427.4-438.8	11,272	512.6	503.0-522.4	12,154	389.4	382.2-396.7	10,922	197.5	193.8-201.3	5,389	249.3	242.5-256.2	5,533	167.4	162.9-172.0
1988	22,774	427.7	422.1-433.4	10,750	498.2	488.7-507.9	12,024	390.2	383.0-397.6	10,932	200.8	197.0-204.7	5,374	253.5	246.6-260.5	5,558	170.9	166.3-175.6
1989	22,531	433.0	427.3-438.9	10,758	505.2	495.5-515.0	11,773	392.8	385.4-400.2	10,457	195.3	191.5-199.2	5,071	242.9	236.2-249.8	5,386	168.2	163.5-172.9
1990	23,510	457.4	451.4-463.4	11,246	533.8	523.8-544.0	12,264	416.7	409.1-424.5	10,336	196.0	192.1-199.8	4,991	242.9	236.1-249.9	5,345	169.7	165.1-174.5
1991	24,294	479.4	473.3-485.6	11,772	566.0	555.6-576.4	12,522	431.1	423.3-439.1	10,113	194.8	190.9-198.6	4,890	240.8	234.0-247.7	5,223	168.3	163.6-173.1
1992	24,467	485.7	479.5-491.9	12,121	584.4	573.9-595.1	12,346	430.1	422.2-438.0	9,914	191.5	187.7-195.4	4,859	240.6	233.8-247.6	5,055	163.3	158.7-168.0
1993	23,546	471.1	465.0-477.2	11,684	565.5	555.2-576.0	11,862	415.6	408.0-423.4	9,803	190.8	187.0-194.7	4,819	240.0	233.2-247.0	4,984	162.4	157.8-167.1
1994	25,297	509.6	503.3-516.0	12,302	597.3	586.7-608.1	12,995	456.0	448.0-464.1	10,303	202.7	198.7-206.7	5,061	253.8	246.8-261.0	5,242	172.8	168.0-177.7
1995	23,829	483.1	477.0-489.4	11,371	555.5	545.2-565.9	12,458	441.0	433.0-449.0	10,147	200.8	196.9-204.8	4,829	243.4	236.5-250.4	5,318	176.0	171.1-180.9
1996	23,774	486.0	479.8-492.2	11,292	555.9	545.6-566.3	12,482	446.9	438.9-455.0	9,864	196.6	192.7-200.5	4,605	232.6	225.8-239.4	5,259	176.0	171.2-181.0
1997	23,792	486.6	480.4-492.9	11,317	557.6	547.3-568.1	12,475	445.2	437.3-453.3	9,416	188.5	184.7-192.3	4,478	226.6	219.9-233.3	4,938	166.5	161.8-171.3
1998	24,169	493.5	487.2-499.7	11,574	564.7	554.4-575.1	12,595	451.3	443.3-459.4	9,359	186.8	183.0-190.6	4,494	224.8	218.2-231.5	4,865	163.4	158.7-168.1
1999	23,906	483.7	477.6-489.9	11,401	551.3	541.2-561.6	12,505	444.9	437.0-452.9	9,150	180.9	177.2-184.7	4,409	218.7	212.2-225.2	4,741	158.5	153.9-163.1
2000	24,447	492.0	485.8-498.3	11,977	575.5	565.2-585.9	12,470	442.1	434.2-450.1	9,085	177.8	174.2-181.5	4,319	212.3	206.0-218.8	4,766	156.7	152.2-161.3
2001	23,894	479.9	473.8-486.0	11,662	557.6	547.5-567.9	12,232	433.2	425.4-441.1	8,867	173.0	169.4-176.7	4,287	209.7	203.4-216.1	4,580	150.4	146.0-154.9
2002	23,609	473.2	467.1-479.3	11,615	550.8	540.8-560.9	11,994	423.7	416.0-431.5	8,602	167.4	163.9-171.0	4,150	201.5	195.4-207.8	4,452	146.2	141.9-150.7
2003	23,284	467.3	461.3-473.4	11,362	540.2	530.3-550.2	11,922	422.9	415.1-430.7	8,533	166.3	162.8-169.9	4,037	195.5	189.5-201.6	4,496	147.9	143.5-152.4
2004	23,355	469.3	463.2-475.4	11,356	537.0	527.1-547.0	11,999	427.9	420.1-435.8	8,285	161.6	158.1-165.2	3,993	193.2	187.3-199.4	4,292	142.1	137.8-146.5
2005	23,559	476.1	470.0-482.3	11,537	545.6	535.7-555.7	12,022	433.7	425.9-441.7	8,144	159.9	156.4-163.5	3,962	192.0	186.0-198.1	4,182	139.7	135.4-144.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	24,210	493.9	487.6-500.2	12,074	574.0	563.8-584.5	12,136	442.7	434.6-450.8	8,063	159.7	156.2-163.3	3,891	189.8	183.9-195.9	4,172	140.7	136.3-145.1
2007	24,334	496.8	490.5-503.1	12,103	572.9	562.7-583.3	12,231	449.6	441.5-457.8	8,050	159.0	155.5-162.6	3,920	189.8	183.9-195.9	4,130	139.5	135.1-143.9
2008	24,441	497.2	490.9-503.5	12,028	565.1	555.0-575.4	12,413	455.5	447.4-463.8	7,982	157.4	153.9-160.9	3,947	189.9	184.0-196.0	4,035	136.1	131.8-140.5
2009	24,685	500.4	494.1-506.8	12,007	559.6	549.5-569.8	12,678	464.9	456.6-473.3	7,842	154.1	150.7-157.6	3,823	183.7	177.9-189.7	4,019	135.1	130.8-139.4
2010	23,568	475.4	469.2-481.6	11,599	535.5	525.7-545.5	11,969	438.5	430.5-446.7	8,032	156.0	152.5-159.5	3,951	186.9	181.1-192.9	4,081	135.8	131.5-140.1
2011	23,975	477.4	471.2-483.5	11,717	532.1	522.4-541.9	12,258	444.9	436.9-453.1	7,726	148.3	145.0-151.7	3,795	176.8	171.2-182.6	3,931	129.1	125.0-133.4
2012	23,230	457.8	451.8-463.8	11,311	504.6	495.3-514.1	11,919	431.1	423.2-439.2	7,597	144.4	141.1-147.8	3,698	169.1	163.7-174.7	3,899	128.4	124.3-132.7
2013	24,010	467.7	461.7-473.8	11,505	507.1	497.8-516.6	12,505	447.6	439.5-455.7	7,502	140.5	137.3-143.8	3,671	165.6	160.3-171.2	3,831	124.1	120.0-128.2
2014	23,688	456.0	450.1-461.9	11,242	486.7	477.6-495.9	12,446	441.9	433.9-450.0	7,422	137.0	133.8-140.2	3,653	161.9	156.6-167.3	3,769	120.0	116.1-124.1
2015	23,674	451.8	446.0-457.7	11,371	485.6	476.6-494.8	12,303	436.6	428.6-444.6	7,364	134.8	131.6-137.9	3,620	157.8	152.6-163.1	3,744	119.4	115.5-123.5
2016	23,601	446.9	441.1-452.7	11,412	481.2	472.3-490.2	12,189	431.4	423.5-439.4	7,402	133.5	130.4-136.6	3,671	157.8	152.6-163.0	3,731	117.7	113.8-121.7
2017	24,138	453.9	448.1-459.8	11,602	482.4	473.5-491.4	12,536	441.3	433.3-449.4	7,204	128.8	125.8-131.9	3,526	149.5	144.5-154.6	3,678	115.5	111.7-119.4
2018	23,102	432.4	426.8-438.2	11,111	458.6	449.9-467.3	11,991	420.5	412.7-428.4	6,436	113.9	111.1-116.8	3,108	129.3	124.8-134.0	3,328	103.5	99.9-107.2
2019	23,742	442.2	436.4-447.9	11,534	470.8	462.1-479.6	12,208	427.7	419.9-435.7	6,853	120.5	117.7-123.5	3,376	139.4	134.6-144.2	3,477	107.2	103.5-111.0
2020	20,011	373.6	368.3-378.9	9,699	396.2	388.2-404.3	10,312	363.5	356.2-370.9	6,109	107.8	105.1-110.6	3,004	123.9	119.5-128.5	3,105	96.8	93.3-100.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	641	10.4	9.6-11.3	436	17.3	15.6-19.1	205	5.6	4.8-6.4	312	5.1	4.5-5.7	222	8.7	7.5-10.0	90	2.5	2.0-3.1
1977	645	10.6	9.8-11.5	417	16.9	15.3-18.7	228	6.2	5.4-7.1	302	5.0	4.5-5.6	208	8.5	7.4-9.8	94	2.6	2.1-3.2
1978	604	10.3	9.5-11.2	397	16.1	14.5-17.9	207	6.0	5.2-6.9	223	3.7	3.2-4.2	146	5.8	4.9-6.9	77	2.1	1.7-2.7
1979	602	10.4	9.6-11.3	410	16.9	15.2-18.7	192	5.7	4.9-6.6	233	4.0	3.4-4.5	163	6.6	5.6-7.8	70	2.0	1.6-2.6
1980	598	10.6	9.7-11.5	384	16.2	14.6-18.0	214	6.6	5.7-7.6	244	4.3	3.7-4.8	166	7.1	6.0-8.3	78	2.3	1.8-2.9
1981	630	11.1	10.2-12.1	439	18.5	16.7-20.3	191	5.8	4.9-6.7	240	4.2	3.7-4.8	156	6.7	5.6-7.8	84	2.4	1.9-3.0
1982	645	11.6	10.7-12.6	440	19.0	17.2-20.9	205	6.5	5.6-7.5	210	3.7	3.2-4.3	161	6.8	5.8-8.0	49	1.6	1.1-2.1
1983	621	11.2	10.3-12.1	425	18.5	16.7-20.4	196	5.7	4.9-6.6	205	3.5	3.1-4.1	154	6.4	5.4-7.6	51	1.4	1.0-1.9
1984	615	11.3	10.4-12.3	414	18.3	16.6-20.3	201	6.2	5.3-7.2	207	3.8	3.3-4.3	153	6.8	5.7-8.0	54	1.7	1.2-2.2
1985	647	12.0	11.1-13.0	417	18.6	16.8-20.5	230	7.3	6.3-8.4	198	3.5	3.0-4.1	139	6.1	5.1-7.3	59	1.7	1.3-2.2
1986	575	10.8	9.9-11.7	384	17.5	15.7-19.3	191	5.9	5.1-6.9	182	3.3	2.8-3.8	124	5.5	4.6-6.6	58	1.7	1.3-2.3
1987	563	10.8	9.9-11.8	375	17.0	15.3-18.9	188	6.2	5.3-7.2	164	3.1	2.6-3.6	112	5.1	4.2-6.2	52	1.5	1.1-2.1
1988	520	10.1	9.2-11.0	339	15.6	14.0-17.4	181	5.7	4.9-6.6	192	3.8	3.2-4.3	121	5.8	4.8-6.9	71	2.3	1.8-2.9
1989	524	10.5	9.6-11.4	339	16.2	14.5-18.0	185	6.3	5.4-7.3	158	3.1	2.6-3.6	103	4.9	4.0-6.0	55	1.7	1.3-2.2
1990	560	11.3	10.4-12.3	370	17.9	16.1-19.9	190	6.4	5.5-7.4	149	2.9	2.5-3.4	92	4.5	3.6-5.6	57	1.7	1.3-2.3
1991	561	11.6	10.6-12.6	359	17.6	15.8-19.6	202	6.9	6.0-8.0	131	2.5	2.1-3.0	91	4.5	3.6-5.6	40	1.2	0.9-1.7
1992	596	12.4	11.4-13.5	404	19.8	17.9-21.9	192	6.7	5.8-7.8	142	2.8	2.4-3.3	90	4.5	3.6-5.5	52	1.6	1.2-2.1
1993	554	11.6	10.7-12.6	366	18.2	16.4-20.2	188	6.7	5.8-7.8	140	2.9	2.5-3.5	110	5.5	4.5-6.6	30	0.9	0.6-1.4
1994	509	10.7	9.7-11.6	334	16.4	14.7-18.3	175	6.2	5.3-7.2	133	2.8	2.3-3.3	96	4.8	3.9-5.9	37	1.3	0.9-1.8
1995	447	9.4	8.6-10.3	296	14.6	13.0-16.4	151	5.3	4.5-6.3	123	2.5	2.1-3.0	80	4.0	3.2-5.0	43	1.4	1.0-1.9
1996	507	10.7	9.8-11.7	341	16.8	15.0-18.7	166	5.9	5.0-6.9	130	2.7	2.3-3.3	73	3.5	2.8-4.4	57	2.0	1.5-2.7
1997	456	9.6	8.7-10.5	289	14.2	12.6-15.9	167	6.0	5.1-7.0	134	2.8	2.3-3.3	97	4.8	3.9-5.8	37	1.3	0.9-1.8
1998	478	9.9	9.1-10.9	296	14.3	12.7-16.1	182	6.5	5.6-7.5	115	2.3	1.9-2.8	75	3.7	2.9-4.6	40	1.3	0.9-1.7
1999	523	10.8	9.9-11.8	366	17.4	15.7-19.3	157	5.5	4.7-6.5	114	2.3	1.9-2.8	84	4.1	3.2-5.0	30	1.0	0.7-1.4
2000	421	8.6	7.8-9.5	280	13.2	11.7-14.8	141	5.0	4.2-6.0	106	2.2	1.8-2.6	69	3.3	2.6-4.2	37	1.3	0.9-1.7
2001	463	9.4	8.5-10.3	301	14.2	12.6-15.9	162	5.5	4.7-6.5	109	2.2	1.8-2.6	73	3.5	2.7-4.4	36	1.1	0.7-1.5
2002	472	9.6	8.7-10.5	305	14.3	12.7-16.0	167	5.9	5.0-6.9	108	2.2	1.8-2.6	70	3.3	2.6-4.2	38	1.2	0.9-1.7
2003	461	9.3	8.5-10.2	302	14.1	12.5-15.7	159	5.6	4.7-6.6	101	2.0	1.6-2.5	65	3.1	2.4-3.9	36	1.2	0.8-1.7
2004	466	9.4	8.6-10.3	294	13.7	12.2-15.4	172	6.0	5.1-7.0	133	2.6	2.2-3.1	81	3.8	3.0-4.7	52	1.7	1.3-2.2
2005	447	9.1	8.2-10.0	311	14.3	12.7-16.0	136	5.0	4.1-5.9	103	2.0	1.7-2.5	64	3.0	2.3-3.9	39	1.3	0.9-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	486	9.9	9.0-10.8	317	14.7	13.1-16.4	169	5.9	5.0-6.9	93	1.9	1.5-2.3	66	3.2	2.5-4.1	27	0.9	0.6-1.3
2007	499	10.2	9.3-11.1	326	15.2	13.6-17.0	173	6.3	5.3-7.3	108	2.2	1.8-2.6	70	3.3	2.6-4.2	38	1.2	0.9-1.7
2008	495	10.2	9.3-11.2	315	14.7	13.1-16.4	180	6.6	5.7-7.7	95	1.9	1.6-2.4	68	3.2	2.5-4.0	27	0.9	0.6-1.4
2009	514	10.6	9.7-11.6	375	17.4	15.7-19.3	139	5.0	4.2-6.0	117	2.3	1.9-2.8	83	3.8	3.0-4.7	34	1.2	0.8-1.7
2010	486	9.8	8.9-10.7	311	14.0	12.5-15.7	175	6.3	5.4-7.4	119	2.4	1.9-2.8	82	3.8	3.0-4.7	37	1.3	0.9-1.8
2011	480	9.5	8.7-10.5	332	15.0	13.4-16.7	148	5.2	4.3-6.1	93	1.8	1.4-2.2	64	2.9	2.3-3.8	29	0.9	0.6-1.4
2012	494	9.6	8.8-10.5	326	14.2	12.6-15.8	168	5.9	5.0-6.9	112	2.2	1.8-2.6	74	3.3	2.6-4.1	38	1.3	0.9-1.8
2013	498	9.7	8.8-10.6	345	15.0	13.4-16.6	153	5.3	4.5-6.3	94	1.8	1.4-2.2	66	2.8	2.2-3.6	28	1.0	0.6-1.4
2014	544	10.4	9.5-11.3	334	14.2	12.7-15.8	210	7.3	6.3-8.4	104	2.0	1.6-2.4	72	3.2	2.5-4.0	32	1.0	0.7-1.5
2015	531	10.2	9.3-11.1	360	15.2	13.6-16.9	171	6.0	5.1-7.0	116	2.1	1.8-2.6	84	3.6	2.9-4.5	32	1.0	0.7-1.4
2016	555	10.4	9.6-11.3	381	16.0	14.4-17.7	174	5.9	5.0-6.9	115	2.1	1.7-2.6	82	3.5	2.8-4.3	33	1.1	0.7-1.5
2017	571	10.6	9.7-11.5	384	15.8	14.2-17.4	187	6.3	5.4-7.3	115	2.1	1.7-2.5	80	3.4	2.7-4.2	35	1.1	0.7-1.5
2018	514	9.5	8.7-10.4	346	14.1	12.6-15.7	168	5.6	4.7-6.5	102	1.8	1.5-2.2	65	2.7	2.1-3.4	37	1.1	0.8-1.6
2019	550	10.3	9.4-11.2	366	14.8	13.3-16.4	184	6.3	5.4-7.3	102	1.8	1.5-2.2	72	2.9	2.3-3.7	30	0.9	0.6-1.3
2020	462	8.4	7.6-9.2	308	12.2	10.9-13.7	154	5.2	4.4-6.1	115	2.1	1.7-2.5	79	3.2	2.5-4.0	36	1.2	0.8-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	269	4.3	3.8-4.8	179	6.8	5.8-8.0	90	2.5	2.0-3.0	254	4.1	3.6-4.7	171	6.6	5.6-7.7	83	2.4	1.9-3.0
1977	260	4.2	3.7-4.8	167	6.5	5.5-7.6	93	2.6	2.1-3.2	245	4.0	3.5-4.5	151	6.1	5.1-7.2	94	2.6	2.1-3.2
1978	265	4.4	3.9-5.0	170	6.8	5.8-7.9	95	2.6	2.1-3.2	256	4.2	3.7-4.8	163	6.7	5.7-7.8	93	2.6	2.1-3.2
1979	261	4.3	3.8-4.9	185	7.6	6.5-8.8	76	2.1	1.6-2.6	235	3.9	3.4-4.5	154	6.2	5.3-7.3	81	2.3	1.8-2.9
1980	293	4.9	4.3-5.5	190	8.0	6.9-9.3	103	2.8	2.3-3.4	224	3.8	3.3-4.3	154	6.6	5.5-7.7	70	1.8	1.4-2.3
1981	251	4.3	3.8-4.9	160	6.8	5.7-8.0	91	2.7	2.1-3.3	247	4.2	3.6-4.7	167	7.0	6.0-8.2	80	2.2	1.7-2.8
1982	254	4.4	3.9-5.0	155	6.8	5.7-8.0	99	2.7	2.2-3.3	204	3.5	3.0-4.0	141	6.2	5.2-7.3	63	1.8	1.3-2.3
1983	243	4.3	3.7-4.8	167	7.2	6.1-8.4	76	2.2	1.7-2.8	207	3.6	3.1-4.1	140	6.2	5.2-7.3	67	1.8	1.4-2.4
1984	236	4.1	3.5-4.6	161	7.0	5.9-8.2	75	2.1	1.6-2.6	213	3.7	3.2-4.3	138	5.8	4.9-6.9	75	2.1	1.7-2.7
1985	239	4.3	3.8-4.9	160	7.0	6.0-8.3	79	2.4	1.9-3.1	192	3.4	3.0-4.0	140	6.1	5.1-7.2	52	1.6	1.1-2.1
1986	244	4.4	3.9-5.0	150	6.9	5.8-8.1	94	2.8	2.2-3.4	194	3.5	3.0-4.0	149	6.9	5.8-8.2	45	1.3	0.9-1.8
1987	218	4.0	3.5-4.6	132	5.9	4.9-7.1	86	2.7	2.1-3.3	182	3.3	2.8-3.8	112	5.0	4.1-6.1	70	2.1	1.6-2.6
1988	187	3.4	2.9-3.9	115	5.2	4.3-6.2	72	2.1	1.7-2.7	199	3.7	3.2-4.3	138	6.4	5.4-7.6	61	1.9	1.4-2.4
1989	242	4.5	4.0-5.2	157	7.4	6.2-8.6	85	2.5	2.0-3.2	189	3.5	3.0-4.1	121	5.6	4.7-6.8	68	2.0	1.5-2.5
1990	230	4.4	3.9-5.1	141	6.7	5.6-7.9	89	2.8	2.2-3.5	192	3.6	3.1-4.2	124	6.0	5.0-7.2	68	2.0	1.5-2.5
1991	221	4.3	3.7-4.9	144	7.0	5.9-8.2	77	2.4	1.9-3.0	172	3.3	2.8-3.8	113	5.4	4.4-6.5	59	1.8	1.3-2.3
1992	225	4.5	3.9-5.1	143	6.8	5.7-8.1	82	2.7	2.1-3.4	176	3.4	2.9-4.0	127	6.1	5.1-7.3	49	1.5	1.1-2.0
1993	198	3.9	3.4-4.5	139	6.8	5.7-8.0	59	1.9	1.4-2.5	163	3.2	2.7-3.7	120	5.8	4.8-7.0	43	1.4	1.0-1.9
1994	239	4.8	4.2-5.5	170	8.4	7.1-9.7	69	2.2	1.7-2.8	187	3.7	3.2-4.3	135	6.6	5.5-7.9	52	1.6	1.2-2.1
1995	221	4.5	4.0-5.2	130	6.4	5.4-7.6	91	3.1	2.5-3.8	181	3.7	3.1-4.2	127	6.3	5.2-7.4	54	1.7	1.3-2.3
1996	220	4.4	3.9-5.1	147	7.3	6.1-8.6	73	2.3	1.8-2.9	195	3.9	3.4-4.5	129	6.4	5.3-7.6	66	2.1	1.6-2.7
1997	220	4.4	3.9-5.1	142	7.1	6.0-8.3	78	2.6	2.0-3.2	174	3.5	3.0-4.1	120	6.0	5.0-7.2	54	1.7	1.3-2.3
1998	228	4.6	4.0-5.2	159	7.9	6.7-9.2	69	2.3	1.8-2.9	194	3.9	3.4-4.5	136	6.8	5.7-8.0	58	1.9	1.4-2.5
1999	211	4.2	3.7-4.8	161	7.8	6.6-9.1	50	1.6	1.1-2.1	182	3.6	3.1-4.2	139	6.8	5.7-8.0	43	1.4	1.0-1.9
2000	203	4.0	3.5-4.6	137	6.6	5.5-7.8	66	2.1	1.6-2.6	194	3.8	3.3-4.4	129	6.3	5.2-7.4	65	2.0	1.6-2.6
2001	210	4.1	3.6-4.8	145	7.0	5.9-8.2	65	2.1	1.6-2.7	162	3.2	2.7-3.7	126	6.1	5.0-7.2	36	1.2	0.8-1.7
2002	215	4.3	3.7-4.9	155	7.3	6.2-8.6	60	1.9	1.5-2.5	172	3.3	2.9-3.9	123	5.9	4.9-7.1	49	1.5	1.1-2.0
2003	226	4.5	3.9-5.1	151	7.1	6.0-8.4	75	2.4	1.9-3.1	169	3.3	2.8-3.9	114	5.4	4.4-6.4	55	1.7	1.3-2.3
2004	215	4.2	3.7-4.9	145	6.8	5.7-8.0	70	2.3	1.8-2.9	160	3.1	2.7-3.7	115	5.5	4.5-6.6	45	1.4	1.0-1.9
2005	217	4.4	3.8-5.0	143	6.8	5.7-8.0	74	2.5	1.9-3.1	202	4.0	3.5-4.6	152	7.2	6.1-8.5	50	1.5	1.1-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	232	4.7	4.1-5.3	175	8.4	7.2-9.8	57	1.9	1.5-2.5	178	3.6	3.1-4.2	110	5.3	4.4-6.4	68	2.2	1.7-2.9
2007	211	4.2	3.6-4.8	143	6.7	5.6-7.9	68	2.2	1.7-2.8	190	3.8	3.3-4.4	144	6.9	5.8-8.1	46	1.6	1.1-2.1
2008	244	4.9	4.3-5.5	169	7.8	6.7-9.1	75	2.5	2.0-3.2	174	3.4	2.9-4.0	126	5.8	4.9-7.0	48	1.6	1.2-2.1
2009	196	3.9	3.4-4.5	145	6.7	5.6-7.9	51	1.6	1.2-2.2	188	3.8	3.2-4.3	135	6.4	5.4-7.6	53	1.7	1.3-2.2
2010	183	3.6	3.1-4.1	130	6.0	5.0-7.2	53	1.7	1.3-2.3	161	3.1	2.6-3.6	115	5.3	4.4-6.4	46	1.5	1.1-2.0
2011	186	3.7	3.1-4.2	143	6.6	5.5-7.8	43	1.5	1.1-2.0	152	3.0	2.5-3.5	108	5.0	4.1-6.1	44	1.4	1.0-1.9
2012	165	3.2	2.7-3.7	121	5.4	4.4-6.4	44	1.5	1.0-2.0	127	2.4	2.0-2.9	92	4.2	3.4-5.2	35	1.1	0.7-1.5
2013	208	4.0	3.5-4.6	149	6.6	5.6-7.8	59	1.9	1.4-2.5	148	2.8	2.4-3.3	113	4.9	4.0-5.9	35	1.1	0.8-1.6
2014	210	4.0	3.5-4.6	144	6.3	5.3-7.4	66	2.1	1.6-2.8	176	3.3	2.8-3.9	128	5.7	4.8-6.8	48	1.5	1.1-2.0
2015	191	3.6	3.1-4.1	133	5.6	4.7-6.7	58	1.9	1.4-2.5	148	2.7	2.3-3.2	94	4.1	3.3-5.0	54	1.6	1.2-2.1
2016	190	3.5	3.0-4.0	143	5.9	5.0-7.0	47	1.5	1.1-2.0	148	2.7	2.3-3.2	114	4.8	3.9-5.7	34	1.0	0.7-1.5
2017	185	3.4	2.9-3.9	139	5.8	4.9-6.9	46	1.5	1.1-2.0	131	2.4	2.0-2.8	100	4.1	3.4-5.0	31	1.0	0.6-1.4
2018	181	3.2	2.7-3.7	139	5.6	4.7-6.6	42	1.2	0.9-1.7	120	2.1	1.7-2.5	92	3.7	3.0-4.6	28	0.8	0.5-1.2
2019	202	3.6	3.1-4.2	143	5.8	4.9-6.9	59	1.8	1.3-2.3	140	2.5	2.1-2.9	109	4.4	3.6-5.4	31	0.9	0.6-1.3
2020	156	2.8	2.3-3.2	108	4.3	3.5-5.2	48	1.5	1.1-2.0	115	2.0	1.7-2.5	88	3.5	2.8-4.4	27	0.8	0.5-1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	852	14.1	13.2-15.1	484	20.0	18.2-22.0	368	10.2	9.2-11.3	669	11.2	10.3-12.1	375	15.8	14.2-17.6	294	8.2	7.3-9.2
1977	846	14.5	13.5-15.5	480	20.6	18.7-22.6	366	10.6	9.5-11.7	648	11.1	10.2-12.0	354	15.6	13.9-17.4	294	8.3	7.4-9.3
1978	822	14.0	13.0-15.0	454	19.7	17.9-21.7	368	10.3	9.2-11.4	602	10.4	9.5-11.2	331	14.7	13.1-16.4	271	7.6	6.7-8.6
1979	837	14.2	13.3-15.3	448	19.1	17.3-21.1	389	11.0	9.9-12.2	597	10.2	9.4-11.1	326	14.3	12.7-16.0	271	7.6	6.7-8.6
1980	850	14.7	13.7-15.7	488	21.4	19.5-23.5	362	10.4	9.3-11.6	591	10.2	9.4-11.1	332	14.7	13.1-16.5	259	7.4	6.5-8.4
1981	793	13.8	12.8-14.8	440	19.7	17.9-21.7	353	10.1	9.0-11.2	583	10.1	9.3-11.0	338	15.3	13.7-17.1	245	6.9	6.1-7.9
1982	740	12.8	11.9-13.8	403	17.9	16.2-19.8	337	9.5	8.5-10.6	495	8.7	7.9-9.5	271	12.4	11.0-14.1	224	6.4	5.6-7.3
1983	814	14.5	13.5-15.5	423	19.3	17.5-21.3	391	11.3	10.2-12.5	511	9.1	8.3-9.9	278	13.1	11.6-14.8	233	6.6	5.8-7.5
1984	741	13.0	12.1-14.0	399	17.7	16.0-19.6	342	9.9	8.8-11.0	492	8.6	7.9-9.5	251	11.4	10.0-13.0	241	6.8	6.0-7.8
1985	708	12.6	11.7-13.6	400	18.4	16.6-20.3	308	9.0	8.0-10.1	498	8.9	8.1-9.8	277	12.6	11.1-14.2	221	6.5	5.7-7.5
1986	665	11.9	11.0-12.9	359	16.6	14.9-18.4	306	8.9	7.9-10.0	406	7.2	6.5-8.0	220	10.4	9.0-11.9	186	5.3	4.6-6.2
1987	691	12.3	11.4-13.3	392	18.0	16.3-20.0	299	8.5	7.5-9.5	427	7.7	6.9-8.4	248	11.5	10.1-13.1	179	5.0	4.3-5.9
1988	645	11.8	10.9-12.7	358	16.8	15.1-18.7	287	8.3	7.4-9.4	418	7.6	6.9-8.3	214	10.0	8.7-11.4	204	5.9	5.1-6.8
1989	617	11.6	10.7-12.5	339	16.1	14.4-18.0	278	8.3	7.4-9.4	433	8.0	7.2-8.8	247	11.8	10.3-13.4	186	5.6	4.8-6.5
1990	709	13.5	12.5-14.5	397	19.4	17.5-21.4	312	9.6	8.6-10.8	382	7.3	6.5-8.0	223	11.0	9.6-12.6	159	4.8	4.1-5.7
1991	697	13.4	12.4-14.5	405	20.2	18.3-22.3	292	8.9	7.9-10.0	409	7.8	7.0-8.6	230	11.5	10.1-13.1	179	5.4	4.6-6.3
1992	643	12.5	11.6-13.5	358	17.9	16.1-19.9	285	9.1	8.0-10.2	409	7.9	7.1-8.7	226	11.5	10.0-13.1	183	5.6	4.8-6.6
1993	675	13.1	12.2-14.2	387	19.5	17.6-21.6	288	9.1	8.0-10.2	400	7.8	7.0-8.6	225	11.5	10.1-13.2	175	5.5	4.7-6.4
1994	700	13.8	12.8-14.8	375	18.6	16.7-20.6	325	10.5	9.3-11.7	430	8.4	7.6-9.2	247	12.6	11.0-14.2	183	5.8	5.0-6.8
1995	624	12.2	11.3-13.2	347	17.4	15.6-19.4	277	8.8	7.7-9.9	422	8.3	7.5-9.1	229	11.8	10.3-13.4	193	6.1	5.3-7.1
1996	649	12.8	11.9-13.9	348	17.6	15.8-19.6	301	9.6	8.6-10.8	388	7.6	6.9-8.5	195	10.0	8.7-11.5	193	6.1	5.3-7.1
1997	585	11.6	10.7-12.6	334	17.0	15.2-18.9	251	8.1	7.2-9.3	367	7.3	6.6-8.1	198	10.1	8.7-11.6	169	5.6	4.8-6.5
1998	646	12.9	11.9-13.9	364	18.2	16.4-20.2	282	9.4	8.3-10.6	379	7.5	6.7-8.2	208	10.5	9.1-12.1	171	5.4	4.6-6.4
1999	603	11.9	10.9-12.9	354	17.6	15.8-19.5	249	8.0	7.0-9.1	344	6.7	6.0-7.4	187	9.4	8.1-10.9	157	5.0	4.3-5.9
2000	565	11.0	10.2-12.0	298	14.6	13.0-16.3	267	8.6	7.6-9.8	313	6.0	5.4-6.8	156	7.7	6.5-9.0	157	5.0	4.2-5.9
2001	590	11.6	10.7-12.6	333	16.1	14.4-18.0	257	8.5	7.5-9.6	316	6.2	5.5-6.9	163	8.0	6.8-9.3	153	5.0	4.2-5.9
2002	528	10.4	9.5-11.3	297	14.2	12.6-15.9	231	7.6	6.7-8.7	285	5.5	4.9-6.2	158	7.6	6.5-8.9	127	4.0	3.3-4.8
2003	571	11.2	10.2-12.1	311	14.9	13.2-16.6	260	8.6	7.6-9.7	299	5.8	5.2-6.5	158	7.7	6.5-9.0	141	4.6	3.9-5.5
2004	562	11.1	10.2-12.1	310	14.9	13.3-16.7	252	8.6	7.5-9.7	307	6.0	5.3-6.7	151	7.3	6.2-8.6	156	5.3	4.5-6.2
2005	549	10.9	10.0-11.8	306	14.8	13.2-16.6	243	8.1	7.1-9.2	296	5.7	5.1-6.4	164	8.0	6.8-9.3	132	4.2	3.5-5.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	514	10.3	9.4-11.3	293	14.2	12.6-15.9	221	7.6	6.6-8.7	260	5.2	4.5-5.8	146	7.1	6.0-8.3	114	3.9	3.2-4.7
2007	549	11.0	10.1-12.0	302	14.6	13.0-16.4	247	8.4	7.4-9.6	239	4.7	4.1-5.3	129	6.3	5.2-7.4	110	3.6	2.9-4.4
2008	504	10.0	9.2-11.0	304	14.5	12.9-16.2	200	6.9	5.9-7.9	248	4.9	4.3-5.6	156	7.5	6.4-8.8	92	3.1	2.5-3.8
2009	488	9.8	8.9-10.7	271	12.9	11.4-14.6	217	7.5	6.5-8.6	244	4.8	4.2-5.5	137	6.7	5.6-7.9	107	3.5	2.8-4.3
2010	494	9.9	9.0-10.8	282	13.3	11.7-14.9	212	7.3	6.3-8.4	241	4.6	4.0-5.2	109	5.2	4.2-6.3	132	4.4	3.6-5.2
2011	431	8.5	7.7-9.4	243	11.4	10.0-13.0	188	6.5	5.5-7.5	235	4.6	4.0-5.2	117	5.5	4.6-6.6	118	3.9	3.2-4.7
2012	489	9.6	8.7-10.5	284	13.1	11.6-14.7	205	7.0	6.1-8.1	223	4.2	3.7-4.8	125	5.7	4.8-6.9	98	3.1	2.5-3.8
2013	505	9.7	8.8-10.5	285	12.8	11.3-14.3	220	7.2	6.3-8.3	227	4.2	3.7-4.8	139	6.2	5.2-7.4	88	2.7	2.2-3.4
2014	478	9.2	8.4-10.1	293	13.0	11.5-14.6	185	6.2	5.3-7.3	250	4.7	4.1-5.3	139	6.1	5.2-7.3	111	3.5	2.9-4.3
2015	471	8.9	8.1-9.8	253	11.0	9.7-12.5	218	7.3	6.4-8.4	213	3.9	3.4-4.5	114	5.1	4.2-6.1	99	3.1	2.5-3.8
2016	485	9.2	8.4-10.1	281	12.1	10.7-13.6	204	7.1	6.1-8.1	221	4.1	3.5-4.6	143	6.2	5.2-7.3	78	2.5	2.0-3.2
2017	498	9.3	8.5-10.2	266	11.2	9.9-12.6	232	7.8	6.8-9.0	234	4.3	3.8-4.9	134	5.7	4.8-6.8	100	3.3	2.6-4.0
2018	434	8.1	7.3-8.9	245	10.3	9.0-11.7	189	6.4	5.5-7.4	197	3.7	3.2-4.2	114	4.9	4.0-5.9	83	2.7	2.1-3.4
2019	434	8.2	7.4-9.0	255	10.7	9.4-12.1	179	6.2	5.3-7.2	205	3.8	3.3-4.4	124	5.1	4.2-6.1	81	2.8	2.2-3.5
2020	377	7.2	6.4-7.9	208	8.8	7.6-10.1	169	5.9	5.0-6.9	171	3.1	2.7-3.7	107	4.5	3.7-5.5	64	2.1	1.6-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	4,176	67.6	65.5-69.8	2,021	81.1	77.4-84.9	2,155	59.1	56.6-61.7	2,246	37.1	35.5-38.7	1,139	47.8	44.9-50.8	1,107	30.5	28.7-32.4
1977	4,102	67.4	65.3-69.5	2,032	84.1	80.3-88.0	2,070	56.9	54.4-59.5	2,104	35.1	33.6-36.7	1,037	44.4	41.6-47.4	1,067	29.5	27.7-31.4
1978	3,965	65.4	63.4-67.6	1,946	81.6	77.9-85.5	2,019	55.5	53.1-58.1	1,967	33.2	31.7-34.7	1,018	44.1	41.3-47.0	949	26.5	24.8-28.3
1979	4,055	67.8	65.7-70.0	1,984	84.3	80.5-88.3	2,071	57.6	55.1-60.2	1,979	33.5	32.0-35.0	942	41.1	38.4-43.9	1,037	28.9	27.1-30.7
1980	4,207	71.1	68.9-73.3	2,096	90.1	86.2-94.2	2,111	59.3	56.7-62.0	2,005	34.4	32.9-36.0	1,019	46.2	43.3-49.2	986	27.4	25.7-29.3
1981	4,092	70.0	67.9-72.3	1,989	87.7	83.7-91.7	2,103	59.2	56.6-61.8	1,831	31.5	30.0-33.0	862	39.1	36.4-41.9	969	27.0	25.3-28.8
1982	4,109	70.7	68.5-73.0	1,976	87.9	84.0-92.0	2,133	60.4	57.8-63.1	1,755	30.4	29.0-31.9	873	39.6	36.9-42.4	882	25.1	23.4-26.9
1983	3,996	68.7	66.5-70.9	1,947	86.1	82.2-90.1	2,049	58.0	55.5-60.7	1,767	30.9	29.4-32.4	847	38.9	36.2-41.7	920	26.0	24.3-27.8
1984	3,889	67.9	65.7-70.1	1,854	83.1	79.3-87.1	2,035	58.6	56.0-61.3	1,784	31.0	29.5-32.5	900	41.2	38.5-44.1	884	25.0	23.4-26.8
1985	3,889	67.5	65.3-69.7	1,950	87.3	83.3-91.3	1,939	55.5	53.0-58.1	1,658	29.0	27.6-30.4	816	36.9	34.4-39.6	842	23.9	22.3-25.7
1986	3,950	69.6	67.4-71.9	1,971	89.6	85.6-93.7	1,979	57.4	54.9-60.1	1,611	28.5	27.1-30.0	794	37.0	34.4-39.7	817	23.6	22.0-25.4
1987	3,534	62.7	60.6-64.8	1,751	80.1	76.3-84.0	1,783	51.6	49.2-54.2	1,585	28.0	26.6-29.4	766	35.9	33.4-38.6	819	23.3	21.7-25.0
1988	3,433	62.0	59.9-64.2	1,635	76.2	72.5-80.1	1,798	53.0	50.5-55.5	1,510	27.4	26.0-28.8	772	37.1	34.5-39.9	738	21.7	20.1-23.4
1989	3,215	59.4	57.3-61.5	1,567	74.2	70.5-78.0	1,648	49.8	47.4-52.4	1,411	25.8	24.4-27.2	653	31.9	29.5-34.5	758	22.2	20.7-23.9
1990	3,232	60.0	57.9-62.1	1,505	71.2	67.6-74.9	1,727	52.3	49.8-54.9	1,420	26.3	25.0-27.8	677	33.2	30.7-35.8	743	22.1	20.5-23.8
1991	3,218	60.7	58.6-62.8	1,612	79.0	75.1-83.0	1,606	49.4	47.0-52.0	1,309	24.6	23.3-26.0	647	32.6	30.1-35.2	662	20.0	18.5-21.6
1992	3,094	59.2	57.1-61.3	1,450	71.4	67.7-75.2	1,644	51.6	49.1-54.3	1,239	23.3	22.0-24.6	593	29.7	27.3-32.2	646	19.3	17.8-20.9
1993	3,076	59.2	57.1-61.4	1,476	72.3	68.6-76.1	1,600	50.6	48.1-53.2	1,213	23.1	21.8-24.4	598	30.0	27.6-32.6	615	18.8	17.3-20.4
1994	3,386	65.7	63.5-68.0	1,573	77.7	73.9-81.7	1,813	57.6	54.9-60.4	1,353	25.9	24.5-27.3	665	33.8	31.3-36.5	688	21.2	19.6-22.9
1995	3,135	61.5	59.4-63.7	1,466	73.4	69.6-77.2	1,669	53.6	51.0-56.3	1,353	26.1	24.8-27.6	645	33.0	30.5-35.6	708	22.1	20.4-23.8
1996	3,025	60.0	57.9-62.2	1,457	72.9	69.2-76.8	1,568	51.3	48.7-54.0	1,145	22.2	20.9-23.5	546	27.6	25.3-30.0	599	18.6	17.1-20.2
1997	3,152	62.5	60.4-64.8	1,463	73.2	69.5-77.1	1,689	55.0	52.3-57.7	1,242	24.3	23.0-25.7	610	31.0	28.6-33.6	632	19.9	18.3-21.5
1998	3,217	63.9	61.7-66.2	1,540	76.6	72.8-80.6	1,677	55.4	52.8-58.2	1,172	22.9	21.6-24.2	537	27.1	24.8-29.4	635	19.9	18.4-21.6
1999	3,063	60.5	58.4-62.7	1,473	72.3	68.6-76.1	1,590	52.5	49.9-55.2	1,092	21.2	19.9-22.5	518	25.8	23.6-28.1	574	18.2	16.7-19.8
2000	3,114	61.1	58.9-63.3	1,553	75.6	71.9-79.5	1,561	50.6	48.0-53.2	1,136	21.9	20.6-23.2	556	27.5	25.2-29.9	580	18.1	16.6-19.7
2001	2,988	58.4	56.3-60.6	1,472	71.6	68.0-75.4	1,516	49.8	47.2-52.4	1,042	20.0	18.8-21.2	513	25.3	23.1-27.5	529	16.4	15.0-17.9
2002	2,912	57.0	54.9-59.1	1,424	68.2	64.7-71.9	1,488	48.5	46.0-51.0	1,044	20.1	18.8-21.3	514	25.1	23.0-27.4	530	16.7	15.3-18.2
2003	2,840	55.6	53.6-57.7	1,414	68.0	64.4-71.6	1,426	46.9	44.4-49.4	1,065	20.4	19.2-21.7	495	24.1	22.0-26.3	570	17.8	16.3-19.3
2004	2,697	52.7	50.8-54.8	1,326	63.6	60.2-67.1	1,371	44.9	42.5-47.4	927	17.8	16.7-19.0	448	21.8	19.8-23.9	479	15.0	13.6-16.4
2005	2,609	51.4	49.4-53.4	1,278	61.4	58.0-64.8	1,331	44.1	41.7-46.6	869	16.7	15.6-17.8	406	19.9	18.0-22.0	463	14.4	13.1-15.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	2,528	50.5	48.5-52.5	1,259	61.0	57.7-64.5	1,269	42.9	40.5-45.4	909	17.8	16.6-19.0	457	22.5	20.5-24.7	452	14.7	13.4-16.2
2007	2,479	49.4	47.4-51.4	1,234	58.8	55.6-62.2	1,245	42.4	40.0-44.9	818	15.9	14.8-17.0	388	18.9	17.1-20.9	430	13.7	12.4-15.1
2008	2,371	47.4	45.5-49.3	1,194	56.7	53.5-60.0	1,177	40.3	38.0-42.8	863	16.8	15.7-18.0	432	21.0	19.1-23.1	431	13.9	12.6-15.3
2009	2,360	47.1	45.2-49.1	1,143	53.9	50.8-57.2	1,217	42.0	39.6-44.5	825	15.9	14.8-17.1	383	18.5	16.7-20.5	442	14.0	12.7-15.4
2010	2,219	44.0	42.1-45.9	1,117	52.2	49.2-55.4	1,102	37.6	35.3-40.0	852	16.4	15.3-17.5	446	21.3	19.3-23.3	406	13.0	11.7-14.4
2011	2,080	40.9	39.1-42.7	1,045	48.4	45.5-51.5	1,035	35.5	33.3-37.8	809	15.3	14.3-16.5	410	19.0	17.2-21.0	399	12.5	11.3-13.9
2012	2,016	39.1	37.3-40.8	1,043	47.1	44.3-50.1	973	32.9	30.8-35.1	738	13.9	12.9-14.9	341	15.7	14.1-17.5	397	12.5	11.3-13.9
2013	2,175	41.9	40.1-43.7	1,100	49.3	46.4-52.4	1,075	36.1	33.9-38.4	734	13.5	12.5-14.6	351	15.8	14.2-17.6	383	11.8	10.6-13.1
2014	2,021	38.7	37.0-40.4	1,054	46.1	43.3-49.0	967	32.7	30.6-35.0	711	13.0	12.0-14.0	347	15.4	13.8-17.1	364	11.2	10.0-12.5
2015	1,997	37.8	36.2-39.6	996	43.4	40.7-46.2	1,001	33.7	31.5-35.9	705	12.7	11.7-13.7	360	15.8	14.2-17.5	345	10.4	9.3-11.6
2016	1,960	37.1	35.4-38.8	1,052	45.2	42.5-48.1	908	31.0	28.9-33.1	706	12.5	11.6-13.5	365	15.7	14.1-17.4	341	10.2	9.1-11.4
2017	1,980	37.0	35.3-38.7	1,027	42.8	40.2-45.5	953	32.3	30.2-34.5	724	12.8	11.9-13.8	341	14.3	12.8-15.9	383	11.9	10.7-13.2
2018	1,905	35.5	33.9-37.2	1,001	41.9	39.3-44.6	904	30.3	28.3-32.4	566	10.2	9.3-11.1	287	12.0	10.7-13.5	279	8.6	7.6-9.8
2019	1,930	36.2	34.6-37.9	1,037	43.3	40.7-46.1	893	30.3	28.3-32.5	591	10.5	9.7-11.4	303	12.8	11.4-14.3	288	8.6	7.6-9.7
2020	1,609	30.0	28.5-31.5	832	34.4	32.0-36.8	777	26.1	24.2-28.1	586	10.4	9.6-11.3	324	13.5	12.0-15.1	262	8.0	7.0-9.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2,936	47.5	45.7-49.3	1,369	55.3	52.3-58.5	1,567	42.8	40.6-45.0	1,782	29.5	28.1-31.0	881	37.4	34.8-40.1	901	24.8	23.2-26.6
1977	2,909	48.0	46.2-49.8	1,398	58.5	55.4-61.8	1,511	41.6	39.4-43.8	1,735	28.9	27.5-30.3	841	36.0	33.5-38.6	894	24.7	23.1-26.5
1978	2,840	47.1	45.3-48.9	1,366	57.6	54.5-60.9	1,474	40.7	38.6-42.9	1,631	27.5	26.1-28.9	832	36.2	33.6-38.8	799	22.2	20.6-23.8
1979	2,926	49.1	47.2-50.9	1,385	59.5	56.3-62.8	1,541	42.8	40.7-45.1	1,656	28.0	26.6-29.4	760	33.1	30.7-35.6	896	24.9	23.3-26.6
1980	2,978	50.4	48.6-52.3	1,442	63.2	59.9-66.7	1,536	43.0	40.8-45.2	1,714	29.5	28.1-31.0	857	39.1	36.4-41.9	857	23.9	22.3-25.6
1981	2,891	49.3	47.5-51.2	1,354	60.1	56.8-63.5	1,537	42.9	40.7-45.2	1,580	27.2	25.8-28.6	731	33.1	30.7-35.7	849	23.7	22.1-25.4
1982	2,912	50.1	48.3-52.0	1,334	60.3	57.0-63.7	1,578	44.5	42.2-46.8	1,527	26.4	25.1-27.8	761	34.6	32.1-37.2	766	21.8	20.3-23.5
1983	2,872	49.4	47.5-51.2	1,341	59.7	56.4-63.1	1,531	43.1	41.0-45.4	1,490	25.9	24.6-27.3	708	32.3	29.9-34.8	782	22.0	20.5-23.7
1984	2,812	48.9	47.1-50.8	1,311	59.4	56.1-62.8	1,501	42.9	40.7-45.1	1,534	26.6	25.2-27.9	766	35.1	32.6-37.7	768	21.8	20.2-23.4
1985	2,901	50.4	48.6-52.3	1,386	62.7	59.3-66.2	1,515	43.3	41.1-45.6	1,463	25.6	24.2-26.9	712	32.2	29.8-34.7	751	21.4	19.9-23.0
1986	2,921	51.5	49.6-53.4	1,387	63.4	60.1-67.0	1,534	44.4	42.1-46.7	1,397	24.7	23.4-26.1	678	31.4	29.1-34.0	719	20.8	19.3-22.5
1987	2,586	45.5	43.7-47.3	1,231	56.5	53.4-59.9	1,355	38.7	36.6-40.9	1,393	24.6	23.3-26.0	688	32.3	29.8-34.8	705	20.1	18.6-21.6
1988	2,503	45.0	43.2-46.8	1,143	53.5	50.4-56.8	1,360	39.6	37.5-41.9	1,319	23.9	22.6-25.3	678	32.6	30.1-35.2	641	18.8	17.4-20.4
1989	2,341	43.1	41.3-44.9	1,104	52.6	49.5-55.9	1,237	37.2	35.1-39.4	1,262	23.0	21.8-24.4	580	28.5	26.1-30.9	682	20.0	18.5-21.6
1990	2,413	44.7	42.9-46.5	1,086	51.7	48.7-55.0	1,327	40.1	37.9-42.4	1,257	23.3	22.0-24.6	590	28.9	26.6-31.4	667	19.8	18.2-21.4
1991	2,400	45.0	43.2-46.9	1,161	57.0	53.7-60.4	1,239	37.9	35.7-40.1	1,133	21.2	20.0-22.5	557	28.1	25.8-30.6	576	17.3	15.9-18.8
1992	2,276	43.3	41.5-45.1	1,048	51.9	48.8-55.2	1,228	38.3	36.1-40.6	1,077	20.2	19.0-21.4	509	25.3	23.1-27.7	568	16.9	15.5-18.4
1993	2,293	43.9	42.1-45.8	1,061	52.0	48.9-55.3	1,232	38.7	36.5-41.0	1,063	20.2	19.0-21.5	514	25.9	23.7-28.3	549	16.8	15.3-18.3
1994	2,522	48.7	46.8-50.6	1,152	57.2	53.9-60.6	1,370	43.0	40.7-45.4	1,206	23.0	21.8-24.4	592	30.1	27.7-32.6	614	19.0	17.5-20.6
1995	2,333	45.5	43.7-47.4	1,062	53.5	50.3-56.9	1,271	40.2	38.0-42.5	1,182	22.8	21.5-24.1	563	28.9	26.5-31.4	619	19.2	17.7-20.8
1996	2,230	43.9	42.1-45.8	1,032	51.8	48.6-55.1	1,198	38.5	36.3-40.8	1,007	19.5	18.3-20.7	486	24.7	22.5-27.0	521	16.0	14.6-17.5
1997	2,370	46.8	44.9-48.7	1,075	54.2	50.9-57.5	1,295	41.7	39.4-44.1	1,092	21.3	20.1-22.6	529	26.9	24.6-29.3	563	17.7	16.2-19.2
1998	2,346	46.3	44.5-48.3	1,086	54.4	51.2-57.7	1,260	41.1	38.8-43.5	1,048	20.4	19.2-21.7	478	24.2	22.1-26.5	570	17.8	16.4-19.4
1999	2,251	44.2	42.4-46.1	1,062	52.5	49.4-55.7	1,189	38.9	36.7-41.2	949	18.4	17.2-19.6	451	22.5	20.5-24.7	498	15.8	14.4-17.3
2000	2,289	44.6	42.8-46.5	1,087	53.2	50.1-56.4	1,202	38.7	36.5-41.0	1,000	19.2	18.1-20.5	490	24.3	22.2-26.5	510	15.9	14.5-17.4
2001	2,208	42.9	41.1-44.7	1,064	52.0	48.9-55.2	1,144	36.8	34.6-39.0	891	17.0	15.9-18.2	428	21.1	19.1-23.2	463	14.2	13.0-15.6
2002	2,146	41.7	39.9-43.5	1,006	48.6	45.6-51.7	1,140	36.8	34.6-39.0	903	17.3	16.2-18.5	447	21.9	19.9-24.0	456	14.4	13.0-15.8
2003	2,072	40.3	38.6-42.1	999	48.3	45.3-51.4	1,073	34.8	32.7-37.0	931	17.8	16.6-19.0	430	21.0	19.1-23.1	501	15.5	14.1-16.9
2004	1,961	38.2	36.5-39.9	948	45.7	42.8-48.7	1,013	32.7	30.7-34.8	813	15.6	14.5-16.7	382	18.6	16.7-20.5	431	13.4	12.2-14.8
2005	1,936	37.8	36.2-39.6	906	43.6	40.8-46.5	1,030	33.7	31.7-35.9	753	14.4	13.4-15.5	348	17.1	15.4-19.0	405	12.5	11.3-13.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,802	35.8	34.2-37.5	859	42.0	39.2-44.9	943	31.5	29.5-33.6	799	15.6	14.5-16.7	394	19.4	17.6-21.5	405	13.1	11.8-14.5
2007	1,767	35.0	33.3-36.6	856	41.0	38.2-43.8	911	30.7	28.7-32.8	709	13.7	12.7-14.8	338	16.5	14.8-18.3	371	11.7	10.5-13.0
2008	1,702	33.8	32.2-35.4	834	39.8	37.2-42.7	868	29.3	27.3-31.4	722	14.1	13.0-15.1	360	17.6	15.8-19.5	362	11.7	10.5-13.0
2009	1,704	33.7	32.1-35.4	774	37.0	34.4-39.7	930	31.5	29.5-33.7	702	13.5	12.5-14.6	326	15.8	14.1-17.6	376	11.8	10.6-13.1
2010	1,576	31.0	29.5-32.6	767	35.9	33.4-38.6	809	27.0	25.1-29.0	736	14.1	13.0-15.1	374	17.9	16.1-19.8	362	11.4	10.3-12.7
2011	1,495	29.3	27.8-30.8	747	34.8	32.3-37.4	748	25.4	23.5-27.3	680	12.9	11.9-13.9	344	16.0	14.3-17.8	336	10.5	9.4-11.8
2012	1,456	28.0	26.5-29.5	719	32.7	30.3-35.2	737	24.4	22.6-26.3	618	11.5	10.6-12.5	282	13.1	11.6-14.7	336	10.4	9.3-11.7
2013	1,556	29.8	28.3-31.3	756	34.1	31.7-36.7	800	26.6	24.7-28.6	621	11.4	10.5-12.3	292	13.3	11.8-14.9	329	10.0	8.9-11.2
2014	1,449	27.5	26.0-28.9	741	32.7	30.3-35.1	708	23.4	21.7-25.3	590	10.7	9.8-11.6	281	12.5	11.0-14.0	309	9.4	8.3-10.6
2015	1,386	26.1	24.7-27.5	668	29.3	27.1-31.6	718	23.8	22.1-25.7	575	10.3	9.4-11.2	291	12.7	11.3-14.3	284	8.5	7.5-9.7
2016	1,396	26.1	24.7-27.6	738	31.8	29.5-34.3	658	22.0	20.3-23.8	584	10.3	9.5-11.2	298	12.8	11.4-14.4	286	8.5	7.5-9.6
2017	1,399	25.9	24.5-27.3	692	28.9	26.8-31.2	707	23.6	21.8-25.4	590	10.4	9.5-11.3	277	11.6	10.3-13.1	313	9.6	8.6-10.8
2018	1,364	25.1	23.8-26.6	715	29.9	27.7-32.2	649	21.3	19.6-23.1	463	8.2	7.5-9.0	231	9.7	8.5-11.1	232	7.0	6.1-8.1
2019	1,375	25.5	24.1-26.9	716	29.9	27.7-32.2	659	21.8	20.1-23.6	464	8.1	7.4-8.9	228	9.6	8.4-10.9	236	6.9	6.1-7.9
2020	1,133	20.9	19.6-22.1	554	22.8	20.9-24.8	579	19.1	17.5-20.8	452	8.0	7.2-8.7	246	10.2	8.9-11.5	206	6.3	5.5-7.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,240	20.1	19.0-21.3	652	25.8	23.8-28.0	588	16.3	15.0-17.8	464	7.5	6.8-8.3	258	10.4	9.1-11.9	206	5.7	4.9-6.5
1977	1,193	19.4	18.3-20.6	634	25.6	23.5-27.7	559	15.4	14.1-16.7	369	6.2	5.6-6.9	196	8.5	7.2-9.8	173	4.8	4.1-5.6
1978	1,125	18.4	17.3-19.5	580	24.0	22.0-26.1	545	14.8	13.5-16.1	336	5.7	5.1-6.4	186	7.9	6.8-9.2	150	4.3	3.6-5.1
1979	1,129	18.8	17.7-20.0	599	24.8	22.8-27.0	530	14.8	13.5-16.2	323	5.5	4.9-6.2	182	8.0	6.8-9.3	141	4.0	3.3-4.7
1980	1,229	20.6	19.5-21.9	654	27.0	24.9-29.2	575	16.4	15.0-17.8	291	4.9	4.4-5.5	162	7.1	6.0-8.3	129	3.5	3.0-4.2
1981	1,201	20.7	19.5-22.0	635	27.6	25.4-29.9	566	16.3	14.9-17.7	251	4.3	3.8-4.9	131	6.0	5.0-7.1	120	3.3	2.8-4.0
1982	1,197	20.6	19.4-21.8	642	27.6	25.5-29.9	555	15.9	14.6-17.4	228	4.0	3.5-4.5	112	5.0	4.1-6.1	116	3.3	2.7-4.0
1983	1,124	19.3	18.2-20.5	606	26.4	24.3-28.6	518	14.9	13.6-16.3	277	4.9	4.4-5.6	139	6.6	5.5-7.8	138	4.0	3.3-4.7
1984	1,077	19.0	17.8-20.2	543	23.8	21.7-25.9	534	15.7	14.4-17.2	250	4.4	3.9-5.0	134	6.1	5.1-7.3	116	3.3	2.7-4.0
1985	988	17.1	16.0-18.2	564	24.6	22.5-26.7	424	12.1	11.0-13.4	195	3.4	3.0-4.0	104	4.7	3.8-5.8	91	2.6	2.0-3.2
1986	1,029	18.1	17.0-19.3	584	26.1	24.0-28.4	445	13.1	11.8-14.4	214	3.8	3.3-4.3	116	5.6	4.6-6.7	98	2.8	2.3-3.4
1987	948	17.2	16.1-18.3	520	23.5	21.5-25.7	428	12.9	11.7-14.2	192	3.4	2.9-3.9	78	3.7	2.9-4.6	114	3.2	2.6-3.9
1988	930	17.1	16.0-18.2	492	22.7	20.7-24.8	438	13.3	12.1-14.7	191	3.5	3.0-4.0	94	4.5	3.6-5.6	97	2.9	2.3-3.5
1989	874	16.3	15.2-17.4	463	21.5	19.6-23.6	411	12.7	11.4-14.0	149	2.7	2.3-3.2	73	3.5	2.7-4.4	76	2.2	1.7-2.8
1990	819	15.3	14.2-16.4	419	19.5	17.6-21.4	400	12.2	11.0-13.5	163	3.0	2.6-3.6	87	4.2	3.4-5.2	76	2.3	1.8-3.0
1991	818	15.6	14.6-16.8	451	22.0	19.9-24.1	367	11.6	10.4-12.9	176	3.4	2.9-3.9	90	4.5	3.6-5.5	86	2.7	2.1-3.4
1992	818	15.9	14.8-17.0	402	19.5	17.6-21.5	416	13.3	12.0-14.7	162	3.1	2.7-3.7	84	4.3	3.4-5.4	78	2.4	1.9-3.0
1993	783	15.3	14.2-16.4	415	20.3	18.4-22.3	368	11.9	10.7-13.2	150	2.9	2.4-3.4	84	4.1	3.3-5.1	66	2.0	1.6-2.6
1994	864	17.0	15.9-18.2	421	20.5	18.6-22.6	443	14.7	13.3-16.1	147	2.8	2.4-3.4	73	3.8	2.9-4.7	74	2.2	1.8-2.8
1995	802	16.0	14.9-17.2	404	19.8	17.9-21.9	398	13.4	12.1-14.9	171	3.4	2.9-3.9	82	4.1	3.3-5.1	89	2.9	2.3-3.6
1996	795	16.1	15.0-17.3	425	21.1	19.2-23.3	370	12.8	11.5-14.2	138	2.7	2.3-3.2	60	2.9	2.2-3.8	78	2.6	2.0-3.3
1997	782	15.8	14.7-16.9	388	19.1	17.2-21.1	394	13.2	11.9-14.7	150	3.0	2.5-3.5	81	4.1	3.3-5.1	69	2.2	1.7-2.8
1998	871	17.6	16.4-18.8	454	22.2	20.2-24.4	417	14.3	12.9-15.8	124	2.5	2.0-2.9	59	2.8	2.2-3.7	65	2.1	1.6-2.7
1999	812	16.3	15.2-17.5	411	19.8	18.0-21.9	401	13.6	12.3-15.1	143	2.8	2.3-3.3	67	3.3	2.5-4.2	76	2.4	1.9-3.0
2000	825	16.4	15.3-17.6	466	22.4	20.4-24.6	359	11.9	10.7-13.2	136	2.7	2.2-3.1	66	3.2	2.5-4.1	70	2.2	1.7-2.8
2001	780	15.6	14.5-16.7	408	19.6	17.8-21.6	372	13.0	11.7-14.4	151	3.0	2.5-3.5	85	4.1	3.3-5.1	66	2.2	1.7-2.8
2002	766	15.3	14.2-16.4	418	19.6	17.8-21.6	348	11.7	10.5-13.0	141	2.7	2.3-3.2	67	3.2	2.5-4.1	74	2.3	1.8-3.0
2003	768	15.3	14.2-16.4	415	19.7	17.8-21.6	353	12.0	10.8-13.4	134	2.6	2.2-3.1	65	3.1	2.4-4.0	69	2.3	1.8-2.9
2004	736	14.6	13.5-15.7	378	17.9	16.1-19.8	358	12.2	11.0-13.6	114	2.2	1.8-2.7	66	3.2	2.5-4.1	48	1.5	1.1-2.1
2005	673	13.5	12.5-14.6	372	17.8	16.0-19.7	301	10.4	9.2-11.7	116	2.2	1.9-2.7	58	2.8	2.1-3.6	58	1.9	1.4-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	726	14.7	13.6-15.8	400	19.0	17.2-21.0	326	11.4	10.1-12.7	110	2.2	1.8-2.6	63	3.1	2.4-4.0	47	1.6	1.2-2.1
2007	712	14.4	13.4-15.5	378	17.9	16.1-19.8	334	11.7	10.5-13.1	109	2.2	1.8-2.6	50	2.4	1.8-3.2	59	1.9	1.4-2.5
2008	669	13.6	12.6-14.7	360	16.8	15.1-18.7	309	11.0	9.8-12.4	141	2.7	2.3-3.2	72	3.4	2.7-4.3	69	2.2	1.7-2.8
2009	656	13.4	12.3-14.4	369	17.0	15.3-18.8	287	10.5	9.3-11.8	123	2.4	2.0-2.9	57	2.7	2.0-3.5	66	2.2	1.7-2.9
2010	643	13.0	12.0-14.0	350	16.3	14.6-18.2	293	10.6	9.4-11.9	116	2.3	1.9-2.8	72	3.4	2.7-4.3	44	1.5	1.1-2.1
2011	585	11.6	10.7-12.6	298	13.6	12.1-15.3	287	10.1	9.0-11.4	129	2.5	2.0-2.9	66	3.1	2.4-3.9	63	2.0	1.5-2.6
2012	560	11.1	10.2-12.1	324	14.4	12.9-16.1	236	8.5	7.4-9.7	120	2.3	1.9-2.8	59	2.6	2.0-3.4	61	2.1	1.6-2.7
2013	619	12.1	11.1-13.1	344	15.2	13.6-16.9	275	9.5	8.4-10.7	113	2.2	1.8-2.6	59	2.5	1.9-3.3	54	1.8	1.3-2.4
2014	572	11.2	10.3-12.2	313	13.4	12.0-15.0	259	9.3	8.2-10.6	121	2.3	1.9-2.8	66	2.9	2.3-3.7	55	1.8	1.3-2.4
2015	611	11.7	10.8-12.7	328	14.0	12.5-15.7	283	9.8	8.7-11.1	130	2.4	2.0-2.9	69	3.1	2.4-3.9	61	1.9	1.4-2.4
2016	564	10.9	10.0-11.9	314	13.4	11.9-15.0	250	9.0	7.8-10.2	122	2.2	1.8-2.6	67	2.9	2.2-3.7	55	1.6	1.2-2.2
2017	581	11.1	10.2-12.1	335	13.8	12.4-15.4	246	8.8	7.7-10.0	134	2.4	2.0-2.9	64	2.7	2.1-3.4	70	2.3	1.7-2.9
2018	541	10.4	9.5-11.3	286	12.0	10.6-13.5	255	9.0	7.9-10.2	103	1.9	1.6-2.4	56	2.3	1.7-3.0	47	1.6	1.2-2.2
2019	555	10.7	9.8-11.7	321	13.4	12.0-15.0	234	8.5	7.4-9.7	127	2.4	2.0-2.9	75	3.2	2.5-4.0	52	1.6	1.2-2.2
2020	476	9.1	8.3-10.0	278	11.5	10.2-13.0	198	7.0	6.1-8.1	134	2.5	2.0-2.9	78	3.3	2.6-4.2	56	1.7	1.2-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	223	3.7	3.2-4.2	122	5.0	4.1-6.0	101	2.8	2.2-3.4	232	3.8	3.3-4.3	122	4.8	3.9-5.7	110	3.1	2.5-3.7
1977	216	3.6	3.1-4.1	131	5.3	4.4-6.3	85	2.3	1.9-2.9	256	4.2	3.7-4.7	136	5.4	4.5-6.5	120	3.3	2.7-4.0
1978	190	3.2	2.7-3.7	109	4.4	3.6-5.4	81	2.3	1.8-2.8	231	3.8	3.3-4.3	136	5.5	4.6-6.6	95	2.6	2.1-3.2
1979	191	3.3	2.8-3.8	116	4.9	4.0-5.9	75	2.2	1.7-2.8	251	4.3	3.8-4.9	133	5.7	4.7-6.8	118	3.4	2.8-4.1
1980	203	3.6	3.1-4.1	112	4.9	4.0-5.9	91	2.8	2.2-3.4	215	3.7	3.2-4.2	108	4.7	3.8-5.7	107	3.0	2.4-3.6
1981	206	3.6	3.1-4.2	110	4.9	4.0-5.9	96	2.7	2.2-3.3	261	4.5	4.0-5.1	142	6.2	5.2-7.4	119	3.4	2.8-4.1
1982	175	3.1	2.6-3.6	123	5.5	4.5-6.6	52	1.5	1.1-2.0	241	4.1	3.6-4.7	140	6.1	5.1-7.2	101	2.9	2.3-3.6
1983	196	3.5	3.0-4.0	114	5.0	4.1-6.0	82	2.5	2.0-3.1	251	4.5	3.9-5.1	132	5.8	4.8-6.9	119	3.6	2.9-4.3
1984	196	3.4	3.0-4.0	117	5.1	4.2-6.1	79	2.3	1.8-3.0	244	4.3	3.7-4.9	130	5.8	4.8-6.9	114	3.2	2.6-3.9
1985	211	3.9	3.4-4.5	126	5.8	4.8-7.0	85	2.5	2.0-3.1	263	4.7	4.1-5.3	153	7.0	6.0-8.3	110	3.1	2.6-3.8
1986	232	4.1	3.6-4.7	143	6.5	5.4-7.6	89	2.6	2.1-3.3	266	4.8	4.2-5.4	137	6.2	5.2-7.4	129	3.8	3.2-4.6
1987	193	3.5	3.1-4.1	118	5.5	4.5-6.6	75	2.2	1.7-2.8	258	4.7	4.1-5.3	144	6.8	5.7-8.0	114	3.4	2.8-4.1
1988	227	4.3	3.8-4.9	139	6.5	5.5-7.7	88	2.7	2.2-3.4	274	5.1	4.5-5.7	150	7.0	5.9-8.2	124	3.7	3.1-4.4
1989	190	3.6	3.1-4.1	118	5.5	4.5-6.6	72	2.2	1.7-2.8	227	4.2	3.6-4.7	130	6.0	5.0-7.1	97	2.9	2.3-3.6
1990	242	4.7	4.1-5.4	148	7.3	6.2-8.6	94	3.0	2.4-3.7	272	5.1	4.5-5.8	152	7.5	6.3-8.8	120	3.7	3.0-4.4
1991	257	5.0	4.4-5.7	160	7.6	6.4-8.9	97	3.1	2.5-3.9	275	5.2	4.6-5.8	159	7.6	6.4-8.9	116	3.5	2.9-4.2
1992	264	5.3	4.6-5.9	158	7.6	6.5-8.9	106	3.5	2.9-4.3	296	5.8	5.1-6.5	175	8.5	7.3-9.9	121	3.8	3.1-4.6
1993	242	4.8	4.2-5.5	144	7.0	5.9-8.2	98	3.2	2.6-3.9	265	5.1	4.5-5.8	139	6.7	5.6-7.9	126	4.0	3.3-4.7
1994	278	5.6	5.0-6.4	176	8.7	7.4-10.1	102	3.3	2.7-4.0	241	4.8	4.2-5.5	152	7.6	6.4-8.9	89	2.9	2.3-3.6
1995	293	6.0	5.3-6.7	185	9.1	7.8-10.5	108	3.6	2.9-4.4	269	5.4	4.7-6.1	155	7.7	6.5-9.0	114	3.6	2.9-4.3
1996	324	6.7	6.0-7.5	223	11.1	9.7-12.6	101	3.4	2.8-4.2	265	5.3	4.7-6.0	167	8.3	7.1-9.6	98	3.1	2.5-3.9
1997	342	7.0	6.3-7.8	220	10.9	9.5-12.4	122	4.0	3.3-4.8	247	4.9	4.3-5.6	145	7.2	6.1-8.5	102	3.2	2.6-3.9
1998	401	8.2	7.4-9.1	255	12.4	10.9-14.0	146	5.0	4.2-5.9	291	5.9	5.2-6.6	183	9.0	7.7-10.4	108	3.5	2.9-4.3
1999	374	7.7	6.9-8.5	247	11.9	10.4-13.5	127	4.2	3.5-5.0	276	5.5	4.9-6.2	170	8.3	7.1-9.6	106	3.5	2.8-4.2
2000	371	7.5	6.8-8.4	259	12.4	10.9-14.0	112	3.8	3.1-4.6	278	5.6	5.0-6.3	176	8.4	7.2-9.8	102	3.4	2.7-4.1
2001	402	8.2	7.4-9.0	269	12.7	11.3-14.3	133	4.6	3.8-5.4	296	5.9	5.3-6.6	186	8.9	7.6-10.2	110	3.6	2.9-4.3
2002	376	7.7	6.9-8.5	261	12.2	10.8-13.8	115	4.0	3.3-4.8	303	6.1	5.4-6.8	193	9.1	7.9-10.5	110	3.6	2.9-4.3
2003	442	9.0	8.1-9.8	296	13.9	12.4-15.6	146	5.0	4.2-5.9	302	5.9	5.3-6.7	186	8.9	7.6-10.2	116	3.7	3.1-4.5
2004	436	8.8	8.0-9.7	309	14.3	12.8-16.0	127	4.5	3.7-5.3	313	6.2	5.5-6.9	193	9.0	7.8-10.4	120	3.9	3.2-4.7
2005	465	9.5	8.6-10.4	335	15.6	14.0-17.4	130	4.5	3.7-5.3	317	6.3	5.6-7.1	204	9.5	8.2-10.9	113	3.7	3.0-4.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	500	10.2	9.3-11.1	334	15.4	13.8-17.2	166	5.8	5.0-6.8	337	6.8	6.1-7.6	213	10.1	8.8-11.5	124	4.2	3.5-5.1
2007	496	10.0	9.2-11.0	367	16.7	15.0-18.5	129	4.6	3.8-5.5	341	6.9	6.2-7.7	228	10.6	9.3-12.1	113	3.9	3.2-4.8
2008	486	9.8	9.0-10.8	353	16.0	14.4-17.8	133	4.8	4.0-5.7	303	6.0	5.3-6.7	206	9.4	8.1-10.7	97	3.3	2.7-4.0
2009	498	10.0	9.1-10.9	350	15.8	14.2-17.5	148	5.3	4.4-6.2	355	7.1	6.4-7.9	237	11.0	9.6-12.5	118	4.2	3.4-5.0
2010	524	10.4	9.5-11.4	382	17.1	15.4-18.9	142	5.0	4.2-5.9	362	7.1	6.4-7.9	250	11.4	10.0-12.9	112	3.6	3.0-4.4
2011	515	10.1	9.2-11.0	359	15.7	14.1-17.4	156	5.4	4.5-6.3	326	6.3	5.6-7.0	214	9.4	8.2-10.8	112	3.6	2.9-4.3
2012	554	10.7	9.8-11.6	391	16.7	15.1-18.5	163	5.7	4.8-6.7	347	6.6	5.9-7.3	217	9.6	8.3-11.0	130	4.3	3.5-5.1
2013	533	10.0	9.2-10.9	355	14.9	13.4-16.6	178	6.0	5.1-7.0	339	6.4	5.7-7.1	223	9.6	8.3-10.9	116	3.8	3.1-4.6
2014	550	10.2	9.4-11.1	377	15.4	13.9-17.1	173	5.8	4.9-6.7	356	6.5	5.8-7.2	230	9.5	8.3-10.8	126	4.0	3.3-4.8
2015	569	10.4	9.6-11.4	388	16.0	14.5-17.7	181	5.9	5.1-6.9	381	6.9	6.2-7.7	250	10.4	9.2-11.8	131	4.0	3.4-4.8
2016	559	10.1	9.3-11.0	401	16.1	14.5-17.8	158	5.2	4.4-6.1	380	6.9	6.2-7.7	243	10.1	8.9-11.5	137	4.3	3.6-5.1
2017	540	9.9	9.0-10.7	370	15.0	13.5-16.6	170	5.5	4.7-6.4	321	5.8	5.2-6.5	209	8.7	7.6-10.0	112	3.5	2.9-4.3
2018	530	9.5	8.7-10.4	374	15.0	13.5-16.6	156	4.9	4.2-5.8	287	5.1	4.5-5.7	184	7.4	6.3-8.6	103	3.2	2.6-3.9
2019	495	8.8	8.0-9.6	341	13.6	12.2-15.1	154	4.9	4.1-5.8	349	6.1	5.5-6.8	223	9.0	7.8-10.3	126	3.8	3.1-4.5
2020	427	7.6	6.9-8.3	294	11.6	10.3-13.0	133	4.2	3.5-5.0	285	5.0	4.4-5.6	181	7.2	6.2-8.4	104	3.1	2.5-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	821	13.3	12.3-14.2	410	16.5	14.8-18.2	411	11.1	10.1-12.3	737	12.0	11.2-13.0	366	14.8	13.3-16.5	371	10.2	9.2-11.3
1977	815	13.5	12.6-14.5	389	16.2	14.6-18.0	426	11.8	10.6-13.0	765	12.6	11.7-13.6	360	15.0	13.4-16.7	405	11.0	9.9-12.2
1978	784	12.7	11.8-13.7	392	15.7	14.1-17.4	392	10.7	9.6-11.9	715	11.6	10.8-12.6	353	14.3	12.8-15.9	362	9.9	8.9-11.1
1979	754	12.5	11.6-13.5	365	15.1	13.6-16.8	389	10.6	9.6-11.8	719	12.0	11.1-12.9	351	14.7	13.2-16.4	368	10.1	9.1-11.3
1980	801	13.5	12.5-14.4	391	16.8	15.1-18.6	410	11.3	10.2-12.5	732	12.2	11.3-13.2	369	15.6	14.0-17.3	363	10.0	8.9-11.1
1981	750	12.9	12.0-13.9	376	16.4	14.7-18.2	374	10.6	9.5-11.8	701	11.9	11.0-12.8	329	14.2	12.7-15.9	372	10.2	9.2-11.3
1982	787	13.7	12.7-14.7	348	15.8	14.2-17.7	439	12.3	11.1-13.5	664	11.4	10.5-12.3	304	13.6	12.0-15.2	360	10.0	9.0-11.1
1983	734	12.6	11.7-13.5	346	14.9	13.3-16.6	388	10.9	9.8-12.1	654	11.2	10.4-12.1	319	13.7	12.2-15.4	335	9.5	8.5-10.6
1984	765	13.3	12.3-14.3	342	15.4	13.8-17.2	423	11.8	10.7-13.0	669	11.6	10.7-12.5	302	13.4	11.9-15.0	367	10.3	9.2-11.4
1985	710	12.4	11.4-13.3	321	14.2	12.6-15.8	389	11.0	9.9-12.2	592	10.3	9.5-11.2	283	12.8	11.3-14.4	309	8.7	7.8-9.8
1986	730	13.0	12.1-14.0	318	14.7	13.1-16.5	412	11.9	10.8-13.2	612	10.6	9.8-11.5	262	11.9	10.5-13.5	350	9.7	8.7-10.8
1987	687	12.2	11.3-13.2	299	13.8	12.2-15.5	388	11.0	9.9-12.2	568	10.0	9.2-10.9	263	11.9	10.4-13.4	305	8.7	7.7-9.8
1988	677	12.2	11.3-13.2	284	13.3	11.7-14.9	393	11.3	10.2-12.5	558	10.0	9.2-10.9	257	12.1	10.6-13.7	301	8.6	7.6-9.6
1989	674	12.4	11.5-13.4	302	14.3	12.7-16.0	372	11.2	10.0-12.4	591	10.8	10.0-11.7	266	12.6	11.1-14.3	325	9.5	8.5-10.7
1990	657	12.2	11.3-13.2	298	14.2	12.6-15.9	359	10.9	9.7-12.1	551	10.2	9.3-11.1	250	12.0	10.6-13.7	301	8.9	7.9-10.0
1991	658	12.4	11.5-13.4	291	14.2	12.6-16.0	367	11.1	9.9-12.3	536	10.1	9.3-11.0	248	12.1	10.6-13.7	288	8.7	7.7-9.8
1992	681	13.0	12.0-14.0	318	15.7	14.0-17.6	363	11.1	9.9-12.3	527	10.0	9.1-10.9	230	11.3	9.9-12.9	297	8.9	7.9-10.0
1993	666	12.9	11.9-13.9	285	14.0	12.4-15.8	381	11.9	10.7-13.2	581	11.1	10.3-12.1	279	13.8	12.2-15.5	302	9.2	8.2-10.4
1994	711	14.0	12.9-15.0	309	15.5	13.8-17.3	402	12.6	11.4-13.9	600	11.6	10.7-12.6	244	12.2	10.7-13.9	356	11.0	9.9-12.3
1995	620	12.0	11.1-13.0	259	13.0	11.5-14.7	361	11.4	10.3-12.7	544	10.5	9.6-11.4	243	12.2	10.7-13.9	301	9.2	8.2-10.3
1996	655	12.9	12.0-14.0	295	14.9	13.2-16.7	360	11.7	10.5-13.0	623	12.2	11.3-13.3	275	13.9	12.3-15.7	348	11.2	10.0-12.5
1997	654	13.0	12.0-14.0	297	14.9	13.2-16.7	357	11.6	10.4-12.9	524	10.4	9.5-11.3	230	11.8	10.3-13.4	294	9.3	8.3-10.5
1998	676	13.4	12.4-14.5	322	16.1	14.4-18.0	354	11.5	10.4-12.8	548	10.8	9.9-11.7	247	12.4	10.9-14.1	301	9.7	8.7-10.9
1999	667	13.2	12.2-14.2	327	16.1	14.4-18.0	340	11.1	9.9-12.3	573	11.3	10.4-12.3	295	14.6	12.9-16.3	278	8.9	7.9-10.1
2000	721	14.1	13.1-15.2	327	16.0	14.3-17.8	394	12.7	11.5-14.1	545	10.5	9.7-11.5	237	11.6	10.1-13.1	308	9.6	8.6-10.8
2001	653	12.7	11.7-13.7	287	13.9	12.4-15.6	366	11.7	10.5-13.0	582	11.2	10.3-12.1	264	13.0	11.5-14.7	318	9.9	8.9-11.1
2002	624	12.1	11.2-13.1	290	13.9	12.4-15.6	334	10.5	9.4-11.7	510	9.8	9.0-10.7	217	10.5	9.2-12.0	293	9.2	8.1-10.3
2003	698	13.6	12.6-14.6	320	15.5	13.9-17.3	378	12.0	10.8-13.3	539	10.4	9.6-11.3	264	12.7	11.2-14.3	275	8.5	7.5-9.6
2004	699	13.6	12.6-14.6	309	14.9	13.2-16.6	390	12.6	11.4-14.0	554	10.6	9.8-11.6	258	12.5	11.0-14.1	296	9.2	8.2-10.4
2005	664	13.1	12.1-14.1	324	15.4	13.8-17.2	340	11.4	10.2-12.7	557	10.8	10.0-11.8	269	12.9	11.4-14.6	288	9.4	8.3-10.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	698	13.8	12.8-14.8	326	15.8	14.1-17.6	372	12.2	11.0-13.6	556	10.9	10.0-11.8	253	12.3	10.8-13.9	303	9.8	8.7-11.0
2007	716	13.9	12.9-15.0	348	16.6	14.9-18.4	368	11.9	10.7-13.3	564	10.9	10.0-11.9	262	12.6	11.1-14.2	302	9.8	8.7-11.0
2008	774	15.2	14.1-16.3	368	17.6	15.8-19.5	406	13.3	12.0-14.7	599	11.7	10.7-12.6	271	13.0	11.5-14.7	328	10.4	9.3-11.7
2009	769	15.1	14.0-16.2	381	18.0	16.2-19.9	388	12.9	11.6-14.3	616	12.0	11.1-13.0	317	15.1	13.5-16.9	299	9.8	8.7-11.0
2010	748	14.7	13.7-15.8	371	17.3	15.6-19.2	377	12.6	11.3-14.0	598	11.5	10.6-12.5	285	13.3	11.8-15.0	313	9.9	8.8-11.1
2011	741	14.3	13.3-15.4	364	16.8	15.1-18.6	377	12.4	11.2-13.8	604	11.6	10.6-12.5	298	13.8	12.3-15.5	306	9.9	8.8-11.1
2012	741	14.3	13.2-15.4	359	16.3	14.7-18.1	382	12.7	11.4-14.0	593	11.2	10.3-12.1	295	13.2	11.8-14.9	298	9.6	8.5-10.8
2013	727	13.6	12.6-14.7	359	15.8	14.2-17.6	368	11.8	10.6-13.1	546	10.1	9.3-11.0	271	12.2	10.7-13.7	275	8.6	7.6-9.7
2014	742	13.8	12.8-14.8	380	16.8	15.1-18.6	362	11.5	10.3-12.8	595	10.9	10.0-11.8	300	13.3	11.9-15.0	295	9.0	7.9-10.1
2015	754	13.8	12.9-14.9	386	16.6	14.9-18.3	368	11.7	10.5-13.0	556	10.0	9.1-10.8	270	11.6	10.2-13.1	286	8.8	7.8-9.9
2016	720	13.1	12.1-14.1	343	14.5	13.0-16.1	377	11.9	10.7-13.2	607	10.9	10.1-11.8	299	12.7	11.3-14.2	308	9.6	8.5-10.8
2017	822	14.8	13.8-15.9	430	18.1	16.4-19.9	392	12.3	11.0-13.6	609	10.8	9.9-11.7	299	12.7	11.3-14.2	310	9.4	8.3-10.5
2018	790	14.1	13.2-15.2	379	15.7	14.2-17.4	411	12.8	11.5-14.1	553	9.7	8.9-10.5	279	11.7	10.3-13.1	274	8.1	7.1-9.1
2019	742	13.2	12.2-14.2	354	14.3	12.9-15.9	388	12.0	10.8-13.3	594	10.4	9.6-11.3	291	11.9	10.5-13.3	303	9.1	8.1-10.2
2020	744	13.2	12.2-14.2	374	15.2	13.6-16.8	370	11.6	10.4-12.9	549	9.5	8.7-10.4	267	10.9	9.6-12.3	282	8.4	7.4-9.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Larynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	352	5.4	4.9-6.1	287	10.6	9.3-11.9	65	1.8	1.4-2.4	137	2.1	1.8-2.5	108	4.1	3.3-5.0	29	0.8	0.5-1.2
1977	336	5.5	4.9-6.1	287	11.1	9.8-12.5	49	1.4	1.0-1.9	140	2.2	1.8-2.6	125	4.8	4.0-5.8	15	0.4	0.2-0.7
1978	317	5.3	4.7-5.9	252	10.3	9.0-11.7	65	1.8	1.4-2.4	127	2.1	1.7-2.5	102	4.1	3.3-5.0	25	0.7	0.4-1.1
1979	359	6.0	5.4-6.7	295	11.6	10.3-13.0	64	2.0	1.5-2.6	131	2.1	1.8-2.5	106	4.3	3.5-5.3	25	0.7	0.4-1.1
1980	368	6.4	5.7-7.1	303	12.6	11.1-14.1	65	2.0	1.5-2.6	122	2.1	1.7-2.5	104	4.4	3.5-5.3	18	0.5	0.3-0.8
1981	341	6.0	5.3-6.7	271	11.4	10.0-12.9	70	2.1	1.6-2.7	114	1.9	1.6-2.3	93	3.8	3.1-4.7	21	0.6	0.4-0.9
1982	334	5.8	5.2-6.5	264	11.2	9.8-12.6	70	2.2	1.7-2.8	121	2.0	1.7-2.4	97	4.0	3.3-5.0	24	0.7	0.4-1.1
1983	314	5.6	5.0-6.3	240	10.0	8.8-11.4	74	2.3	1.8-3.0	109	1.9	1.5-2.3	83	3.5	2.8-4.4	26	0.7	0.5-1.1
1984	360	6.4	5.7-7.1	283	12.0	10.6-13.5	77	2.5	1.9-3.1	106	1.9	1.5-2.3	85	3.6	2.8-4.5	21	0.7	0.4-1.1
1985	347	6.4	5.7-7.1	268	11.6	10.2-13.1	79	2.6	2.0-3.3	118	2.1	1.7-2.5	95	4.2	3.3-5.1	23	0.7	0.4-1.1
1986	312	5.8	5.1-6.5	241	10.6	9.2-12.0	71	2.4	1.9-3.1	81	1.4	1.1-1.7	65	2.9	2.2-3.7	16	0.4	0.3-0.7
1987	295	5.7	5.1-6.4	232	10.5	9.2-12.0	63	2.3	1.7-3.0	99	1.8	1.5-2.2	77	3.5	2.7-4.4	22	0.7	0.4-1.0
1988	314	6.0	5.3-6.7	248	11.1	9.7-12.6	66	2.3	1.8-3.0	95	1.8	1.5-2.2	75	3.5	2.7-4.4	20	0.7	0.4-1.1
1989	294	6.0	5.3-6.7	217	10.2	8.9-11.7	77	2.9	2.3-3.6	96	1.8	1.5-2.2	68	3.1	2.4-3.9	28	0.9	0.6-1.4
1990	313	6.2	5.6-7.0	256	12.1	10.7-13.7	57	2.0	1.5-2.6	84	1.6	1.3-2.0	75	3.5	2.8-4.5	9	0.3	0.1-0.5
1991	289	5.9	5.3-6.7	234	11.3	9.9-12.9	55	2.0	1.5-2.7	97	1.9	1.6-2.4	75	3.6	2.8-4.5	22	0.7	0.5-1.1
1992	291	6.0	5.3-6.7	221	10.6	9.2-12.1	70	2.6	2.0-3.3	109	2.2	1.8-2.7	82	4.1	3.3-5.1	27	1.0	0.6-1.4
1993	270	5.6	4.9-6.3	212	10.3	8.9-11.7	58	2.1	1.6-2.8	99	1.9	1.6-2.4	81	3.9	3.1-4.8	18	0.6	0.3-0.9
1994	263	5.4	4.8-6.2	209	10.3	9.0-11.9	54	2.0	1.5-2.6	95	1.9	1.5-2.3	76	3.7	2.9-4.6	19	0.7	0.4-1.1
1995	216	4.5	3.9-5.1	170	8.1	7.0-9.5	46	1.6	1.2-2.2	90	1.9	1.5-2.3	68	3.4	2.6-4.3	22	0.7	0.5-1.1
1996	283	6.0	5.3-6.7	224	11.1	9.7-12.6	59	2.2	1.6-2.8	99	2.0	1.6-2.5	83	4.1	3.3-5.1	16	0.5	0.3-0.9
1997	248	5.2	4.6-5.9	205	10.1	8.8-11.6	43	1.6	1.1-2.2	75	1.6	1.2-2.0	57	2.9	2.2-3.7	18	0.7	0.4-1.1
1998	241	5.0	4.4-5.7	185	9.0	7.7-10.4	56	2.0	1.5-2.7	83	1.7	1.4-2.1	66	3.2	2.5-4.1	17	0.6	0.3-1.0
1999	214	4.4	3.8-5.1	175	8.4	7.2-9.8	39	1.4	1.0-2.0	77	1.6	1.2-2.0	60	2.9	2.2-3.7	17	0.6	0.3-1.0
2000	235	4.8	4.2-5.5	180	8.7	7.4-10.0	55	2.0	1.5-2.6	85	1.7	1.4-2.1	68	3.3	2.6-4.2	17	0.6	0.4-1.0
2001	228	4.6	4.0-5.3	171	8.2	7.0-9.5	57	2.1	1.6-2.7	61	1.2	0.9-1.5	48	2.3	1.7-3.1	13	0.4	0.2-0.7
2002	211	4.3	3.7-4.9	175	8.3	7.1-9.6	36	1.3	0.9-1.8	77	1.5	1.2-1.9	61	2.9	2.2-3.8	16	0.5	0.3-0.8
2003	227	4.6	4.0-5.2	176	8.3	7.1-9.7	51	1.8	1.3-2.4	78	1.5	1.2-1.9	60	2.9	2.2-3.7	18	0.6	0.3-0.9
2004	228	4.6	4.0-5.2	191	8.9	7.6-10.2	37	1.4	0.9-1.9	69	1.4	1.1-1.8	64	3.0	2.3-3.8	5	0.2	0.1-0.4
2005	209	4.2	3.7-4.9	163	7.5	6.4-8.8	46	1.7	1.2-2.2	65	1.3	1.0-1.6	52	2.5	1.8-3.2	13	0.4	0.2-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Larynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	190	3.9	3.4-4.5	157	7.4	6.3-8.7	33	1.3	0.9-1.8	68	1.4	1.1-1.7	58	2.8	2.1-3.6	10	0.3	0.1-0.6
2007	199	4.1	3.5-4.7	151	7.1	6.0-8.3	48	1.7	1.3-2.3	63	1.3	1.0-1.6	48	2.2	1.7-3.0	15	0.5	0.3-0.9
2008	229	4.7	4.1-5.4	177	8.3	7.1-9.7	52	1.9	1.4-2.6	61	1.3	1.0-1.6	53	2.5	1.9-3.3	8	0.3	0.1-0.6
2009	200	4.0	3.5-4.6	151	6.8	5.8-8.0	49	1.8	1.3-2.4	68	1.3	1.0-1.7	52	2.4	1.8-3.2	16	0.5	0.3-0.9
2010	194	3.9	3.3-4.5	160	7.3	6.2-8.6	34	1.3	0.9-1.8	62	1.2	0.9-1.5	47	2.2	1.6-2.9	15	0.5	0.3-0.8
2011	155	3.0	2.5-3.5	126	5.5	4.6-6.6	29	1.0	0.7-1.5	60	1.2	0.9-1.5	48	2.2	1.6-3.0	12	0.4	0.2-0.7
2012	197	3.8	3.3-4.4	164	7.1	6.1-8.3	33	1.2	0.8-1.7	57	1.1	0.8-1.4	44	1.9	1.4-2.6	13	0.5	0.3-0.8
2013	180	3.5	3.0-4.0	152	6.7	5.7-7.9	28	1.0	0.6-1.4	56	1.1	0.8-1.4	46	2.1	1.6-2.8	10	0.4	0.2-0.7
2014	173	3.3	2.8-3.8	132	5.7	4.8-6.8	41	1.4	1.0-1.9	42	0.8	0.6-1.1	35	1.5	1.0-2.1	7	0.2	0.1-0.5
2015	164	3.0	2.6-3.5	128	5.4	4.5-6.4	36	1.2	0.8-1.7	51	1.0	0.7-1.3	39	1.7	1.2-2.3	12	0.4	0.2-0.8
2016	161	3.0	2.5-3.5	134	5.6	4.7-6.7	27	0.9	0.6-1.4	58	1.0	0.8-1.3	52	2.2	1.6-2.8	6	0.2	0.1-0.4
2017	157	2.9	2.5-3.4	129	5.4	4.5-6.4	28	1.0	0.7-1.5	39	0.7	0.5-1.0	33	1.4	0.9-1.9	6	0.2	0.1-0.4
2018	128	2.3	1.9-2.8	97	3.9	3.1-4.7	31	1.1	0.7-1.6	41	0.7	0.5-1.0	34	1.4	1.0-2.0	7	0.2	0.1-0.5
2019	148	2.7	2.3-3.2	114	4.7	3.9-5.7	34	1.2	0.8-1.7	53	1.0	0.7-1.3	45	1.9	1.3-2.5	8	0.2	0.1-0.5
2020	99	1.8	1.5-2.2	77	3.1	2.4-3.9	22	0.8	0.5-1.2	30	0.5	0.3-0.7	24	0.9	0.6-1.4	6	0.2	0.1-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3,168	49.6	47.9-51.5	2,209	84.1	80.5-87.9	959	26.3	24.6-28.1	2,564	40.1	38.6-41.8	1,825	70.0	66.7-73.5	739	20.3	18.8-21.9
1977	3,154	50.3	48.5-52.1	2,154	84.6	80.9-88.4	1,000	27.5	25.8-29.4	2,596	41.2	39.6-42.8	1,853	72.5	69.1-76.0	743	20.4	18.9-22.0
1978	3,176	51.9	50.0-53.8	2,229	89.1	85.3-93.0	947	27.1	25.4-29.0	2,573	42.1	40.4-43.8	1,821	73.5	70.0-77.0	752	21.4	19.8-23.0
1979	3,220	53.0	51.1-54.9	2,161	87.8	84.0-91.7	1,059	30.3	28.4-32.2	2,554	42.0	40.4-43.7	1,767	72.2	68.8-75.8	787	22.4	20.8-24.1
1980	3,186	53.4	51.5-55.3	2,124	87.8	84.0-91.7	1,062	30.8	28.9-32.8	2,665	44.5	42.8-46.2	1,784	74.1	70.6-77.7	881	25.4	23.7-27.2
1981	3,058	51.9	50.0-53.8	1,975	83.5	79.8-87.4	1,083	31.2	29.3-33.3	2,484	42.2	40.5-44.0	1,676	71.1	67.6-74.7	808	23.4	21.8-25.2
1982	3,231	55.5	53.6-57.5	2,034	87.1	83.2-91.1	1,197	35.2	33.1-37.3	2,548	43.4	41.7-45.2	1,671	71.4	67.9-75.0	877	25.6	23.8-27.4
1983	3,231	55.8	53.9-57.8	2,044	88.0	84.1-92.0	1,187	35.0	33.0-37.2	2,510	43.2	41.4-44.9	1,535	67.1	63.7-70.7	975	28.1	26.3-30.0
1984	3,220	56.8	54.8-58.8	2,064	90.0	86.0-94.1	1,156	35.2	33.1-37.4	2,502	43.8	42.1-45.6	1,635	72.1	68.6-75.8	867	25.5	23.8-27.4
1985	3,220	56.5	54.6-58.6	2,012	87.7	83.8-91.7	1,208	36.3	34.2-38.5	2,508	43.9	42.2-45.7	1,550	68.4	65.0-72.0	958	28.5	26.6-30.4
1986	3,164	56.5	54.5-58.6	1,931	85.4	81.5-89.4	1,233	37.5	35.3-39.7	2,426	42.8	41.1-44.6	1,469	65.0	61.6-68.5	957	28.9	27.0-30.9
1987	3,212	57.7	55.7-59.8	1,951	86.9	83.0-91.0	1,261	38.7	36.5-41.0	2,495	44.7	42.9-46.5	1,549	69.7	66.2-73.4	946	28.7	26.8-30.6
1988	3,130	57.6	55.6-59.7	1,847	84.8	80.9-88.8	1,283	40.1	37.9-42.4	2,454	44.7	42.9-46.6	1,527	69.9	66.4-73.6	927	28.7	26.8-30.7
1989	3,116	58.6	56.5-60.8	1,802	83.5	79.6-87.5	1,314	42.3	40.0-44.7	2,388	44.4	42.6-46.3	1,433	67.2	63.7-70.8	955	29.9	28.0-32.0
1990	3,117	59.5	57.4-61.6	1,794	84.7	80.8-88.8	1,323	43.1	40.7-45.6	2,379	44.8	43.0-46.7	1,403	66.4	62.9-70.0	976	31.1	29.1-33.2
1991	3,152	61.5	59.4-63.7	1,764	84.7	80.8-88.8	1,388	46.1	43.6-48.7	2,414	46.5	44.6-48.4	1,368	66.1	62.6-69.7	1,046	33.8	31.8-36.0
1992	3,270	64.1	61.9-66.4	1,884	90.7	86.6-94.9	1,386	46.7	44.2-49.3	2,341	45.4	43.5-47.3	1,372	66.4	62.9-70.0	969	31.7	29.7-33.8
1993	3,150	62.5	60.3-64.7	1,759	85.7	81.7-89.8	1,391	47.4	44.9-50.0	2,381	46.6	44.7-48.5	1,344	66.0	62.5-69.7	1,037	34.2	32.1-36.4
1994	3,162	63.3	61.1-65.6	1,703	83.1	79.2-87.2	1,459	50.2	47.6-52.9	2,422	48.2	46.3-50.1	1,419	70.0	66.4-73.8	1,003	33.8	31.7-36.1
1995	3,047	60.9	58.7-63.1	1,683	82.4	78.4-86.4	1,364	46.1	43.7-48.7	2,352	46.7	44.8-48.6	1,306	64.3	60.8-67.9	1,046	34.8	32.7-37.1
1996	3,029	61.4	59.2-63.6	1,642	81.6	77.7-85.6	1,387	48.1	45.6-50.8	2,299	46.0	44.2-48.0	1,281	63.6	60.1-67.2	1,018	34.0	31.9-36.2
1997	3,024	61.5	59.3-63.7	1,640	81.1	77.2-85.2	1,384	48.0	45.5-50.7	2,161	43.5	41.7-45.4	1,229	61.2	57.8-64.7	932	31.6	29.6-33.8
1998	2,999	60.6	58.5-62.9	1,597	78.2	74.4-82.2	1,402	48.0	45.5-50.7	2,196	44.1	42.2-46.0	1,182	58.3	55.0-61.8	1,014	34.3	32.2-36.5
1999	2,879	57.7	55.6-59.8	1,481	72.1	68.5-75.9	1,398	47.8	45.3-50.5	2,155	43.0	41.2-44.9	1,155	56.7	53.4-60.0	1,000	34.0	31.9-36.2
2000	3,004	60.0	57.8-62.2	1,575	76.2	72.5-80.1	1,429	49.0	46.4-51.6	2,113	41.6	39.8-43.4	1,151	56.0	52.8-59.4	962	31.5	29.5-33.6
2001	2,947	58.6	56.5-60.7	1,517	72.9	69.3-76.7	1,430	48.5	46.0-51.2	2,127	41.8	40.1-43.7	1,146	55.5	52.4-58.8	981	32.4	30.3-34.5
2002	2,937	58.2	56.1-60.4	1,539	73.4	69.7-77.1	1,398	47.5	45.0-50.1	2,081	40.8	39.1-42.6	1,117	53.5	50.4-56.7	964	31.7	29.7-33.8
2003	2,816	55.9	53.8-58.0	1,506	72.2	68.6-75.9	1,310	44.2	41.9-46.8	2,054	40.4	38.7-42.2	1,095	52.5	49.5-55.7	959	31.8	29.8-33.9
2004	2,790	55.4	53.3-57.5	1,488	71.1	67.5-74.8	1,302	44.5	42.1-47.0	1,938	38.1	36.4-39.8	1,071	51.5	48.4-54.6	867	28.8	26.9-30.9
2005	2,819	56.2	54.2-58.3	1,486	71.2	67.6-74.9	1,333	46.1	43.7-48.7	1,895	37.6	35.9-39.3	1,056	50.9	47.9-54.1	839	28.6	26.6-30.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	2,800	56.4	54.4-58.6	1,445	69.8	66.2-73.5	1,355	47.2	44.7-49.9	1,823	36.3	34.7-38.0	1,007	48.8	45.8-51.9	816	27.4	25.5-29.3
2007	2,907	58.8	56.6-60.9	1,511	72.7	69.0-76.5	1,396	48.9	46.4-51.6	1,907	37.9	36.2-39.7	1,004	48.7	45.7-51.8	903	30.7	28.6-32.8
2008	2,817	56.7	54.6-58.9	1,416	67.8	64.3-71.4	1,401	49.2	46.6-51.9	1,880	37.4	35.7-39.1	1,013	48.7	45.7-51.8	867	29.4	27.4-31.4
2009	2,842	56.8	54.7-58.9	1,426	67.9	64.4-71.5	1,416	49.2	46.6-51.9	1,747	34.5	32.9-36.1	887	42.6	39.8-45.5	860	28.9	26.9-30.9
2010	2,672	53.4	51.4-55.5	1,364	64.4	61.0-68.0	1,308	45.5	43.0-48.1	1,823	35.9	34.2-37.6	945	44.9	42.1-47.9	878	29.6	27.6-31.7
2011	2,750	54.0	52.0-56.1	1,379	63.9	60.6-67.4	1,371	47.0	44.5-49.6	1,732	33.4	31.8-35.0	908	42.3	39.6-45.2	824	26.9	25.0-28.8
2012	2,687	52.2	50.2-54.2	1,310	59.6	56.4-63.0	1,377	47.4	44.9-50.0	1,703	32.8	31.2-34.4	922	42.1	39.4-45.0	781	26.2	24.3-28.1
2013	2,744	52.4	50.5-54.5	1,356	60.7	57.5-64.1	1,388	46.8	44.3-49.4	1,752	33.2	31.6-34.8	920	41.7	39.0-44.5	832	27.4	25.5-29.4
2014	2,654	49.8	47.9-51.7	1,294	56.4	53.4-59.6	1,360	45.1	42.7-47.6	1,527	28.2	26.8-29.7	797	35.1	32.7-37.7	730	23.2	21.5-25.0
2015	2,592	48.1	46.2-50.0	1,332	57.6	54.5-60.9	1,260	41.3	39.0-43.7	1,550	28.2	26.8-29.7	804	34.9	32.5-37.4	746	23.4	21.7-25.2
2016	2,627	48.2	46.4-50.1	1,257	53.6	50.6-56.7	1,370	44.6	42.3-47.1	1,439	26.0	24.6-27.4	740	32.0	29.7-34.5	699	21.7	20.1-23.4
2017	2,692	49.0	47.1-50.9	1,344	56.9	53.9-60.1	1,348	43.2	40.9-45.6	1,399	24.8	23.5-26.2	726	30.4	28.2-32.8	673	20.6	19.0-22.2
2018	2,537	45.5	43.7-47.3	1,178	48.5	45.7-51.4	1,359	43.6	41.2-46.0	1,257	22.1	20.9-23.4	626	26.0	24.0-28.2	631	19.2	17.7-20.8
2019	2,559	45.1	43.4-46.9	1,197	48.4	45.7-51.3	1,362	42.8	40.5-45.2	1,257	21.8	20.6-23.1	635	26.0	24.0-28.1	622	18.7	17.2-20.2
2020	2,068	36.6	35.0-38.3	1,010	41.6	39.1-44.3	1,058	33.0	31.0-35.1	1,096	19.1	18.0-20.3	577	23.6	21.7-25.6	519	15.9	14.5-17.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	386	6.6	6.0-7.4	199	8.0	6.9-9.3	187	5.7	4.9-6.6	146	2.5	2.1-3.0	81	3.2	2.5-4.1	65	2.0	1.5-2.6
1977	387	6.6	6.0-7.4	172	6.9	5.8-8.0	215	6.5	5.6-7.4	142	2.4	2.0-2.8	72	2.9	2.3-3.7	70	2.0	1.5-2.5
1978	354	6.6	5.9-7.3	168	7.2	6.1-8.4	186	6.2	5.3-7.2	113	1.9	1.6-2.3	69	2.9	2.3-3.8	44	1.3	0.9-1.7
1979	391	7.3	6.5-8.0	194	8.0	6.9-9.2	197	6.8	5.8-7.8	140	2.6	2.1-3.0	75	3.2	2.5-4.0	65	2.0	1.6-2.6
1980	413	7.7	6.9-8.5	204	8.6	7.4-9.9	209	7.2	6.2-8.3	136	2.5	2.0-2.9	70	3.0	2.3-3.8	66	2.1	1.6-2.7
1981	372	7.0	6.3-7.8	185	7.9	6.8-9.2	187	6.4	5.4-7.4	156	3.0	2.5-3.5	87	3.9	3.1-4.8	69	2.3	1.7-2.9
1982	381	7.2	6.5-8.0	206	9.2	7.9-10.6	175	5.8	4.9-6.8	129	2.3	1.9-2.8	73	3.2	2.5-4.0	56	1.7	1.3-2.2
1983	433	8.2	7.4-9.1	217	9.6	8.4-11.1	216	7.3	6.3-8.4	133	2.5	2.1-3.0	77	3.5	2.8-4.4	56	1.9	1.4-2.4
1984	471	8.9	8.0-9.7	239	10.4	9.1-11.8	232	7.9	6.9-9.1	158	2.9	2.4-3.4	94	4.1	3.3-5.0	64	2.0	1.5-2.6
1985	423	8.1	7.3-9.0	241	10.6	9.3-12.1	182	6.4	5.5-7.5	165	3.1	2.7-3.7	88	4.0	3.2-4.9	77	2.6	2.0-3.3
1986	435	8.3	7.5-9.2	212	9.5	8.2-10.9	223	7.7	6.6-8.8	139	2.6	2.1-3.0	72	3.2	2.5-4.0	67	2.2	1.7-2.8
1987	426	8.3	7.5-9.2	228	10.6	9.2-12.1	198	6.8	5.9-7.9	141	2.7	2.2-3.2	83	3.8	3.0-4.7	58	2.0	1.5-2.6
1988	395	7.8	7.0-8.6	213	10.0	8.7-11.5	182	6.2	5.3-7.2	125	2.4	2.0-2.9	73	3.4	2.7-4.3	52	1.7	1.3-2.3
1989	398	7.9	7.1-8.7	203	9.6	8.3-11.0	195	6.7	5.7-7.7	105	2.1	1.7-2.6	67	3.3	2.5-4.2	38	1.4	1.0-2.0
1990	395	8.0	7.2-8.8	214	10.1	8.8-11.6	181	6.5	5.6-7.6	115	2.3	1.9-2.8	65	3.2	2.4-4.0	50	1.8	1.3-2.4
1991	415	8.3	7.5-9.1	227	10.7	9.3-12.2	188	6.5	5.6-7.6	112	2.2	1.8-2.6	58	2.8	2.1-3.7	54	1.8	1.3-2.3
1992	354	7.2	6.4-8.0	182	8.7	7.5-10.1	172	6.0	5.1-7.0	94	1.8	1.5-2.3	58	2.7	2.0-3.5	36	1.2	0.8-1.6
1993	364	7.4	6.6-8.2	206	10.0	8.7-11.5	158	5.5	4.7-6.5	84	1.7	1.3-2.1	49	2.5	1.8-3.3	35	1.2	0.8-1.7
1994	436	8.9	8.1-9.8	247	11.9	10.5-13.5	189	6.8	5.8-7.8	111	2.2	1.8-2.7	72	3.5	2.7-4.4	39	1.4	1.0-1.9
1995	432	8.9	8.1-9.8	252	12.1	10.6-13.7	180	6.6	5.6-7.7	115	2.4	2.0-2.9	65	3.2	2.5-4.1	50	1.8	1.3-2.4
1996	461	9.5	8.6-10.4	243	11.8	10.3-13.4	218	8.0	7.0-9.2	112	2.3	1.9-2.7	57	2.7	2.1-3.5	55	1.9	1.4-2.5
1997	496	10.2	9.3-11.2	270	13.4	11.8-15.1	226	8.3	7.2-9.4	109	2.3	1.9-2.7	60	2.9	2.2-3.8	49	1.8	1.3-2.4
1998	509	10.5	9.6-11.4	270	13.0	11.5-14.6	239	8.7	7.6-9.9	126	2.5	2.1-3.0	72	3.6	2.8-4.5	54	1.8	1.4-2.4
1999	504	10.2	9.3-11.1	276	13.1	11.5-14.7	228	8.3	7.2-9.5	106	2.2	1.8-2.6	64	3.1	2.4-3.9	42	1.4	1.0-2.0
2000	602	12.2	11.2-13.2	316	15.0	13.4-16.8	286	10.5	9.3-11.9	124	2.5	2.1-3.0	72	3.5	2.8-4.5	52	1.8	1.4-2.4
2001	611	12.5	11.5-13.5	329	15.6	13.9-17.4	282	10.3	9.1-11.6	112	2.2	1.8-2.7	60	2.9	2.2-3.8	52	1.8	1.3-2.4
2002	619	12.5	11.5-13.5	336	15.8	14.1-17.6	283	10.4	9.2-11.7	99	2.0	1.6-2.4	50	2.4	1.8-3.2	49	1.7	1.3-2.3
2003	544	11.0	10.1-12.0	284	13.3	11.8-15.0	260	9.6	8.5-10.9	107	2.1	1.7-2.6	50	2.4	1.8-3.2	57	2.0	1.5-2.6
2004	750	15.3	14.2-16.4	396	18.5	16.8-20.5	354	13.3	11.9-14.8	99	1.9	1.6-2.4	53	2.5	1.9-3.3	46	1.6	1.1-2.1
2005	828	16.9	15.7-18.1	439	20.5	18.6-22.5	389	14.7	13.3-16.3	96	1.9	1.5-2.3	49	2.4	1.8-3.2	47	1.6	1.2-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,003	20.6	19.3-21.9	554	25.8	23.6-28.0	449	17.0	15.4-18.6	110	2.2	1.8-2.7	67	3.3	2.5-4.2	43	1.5	1.1-2.1
2007	812	16.7	15.5-17.9	424	19.7	17.9-21.7	388	14.7	13.3-16.3	106	2.0	1.7-2.5	63	3.0	2.3-3.9	43	1.4	1.0-2.0
2008	835	17.1	16.0-18.4	456	21.4	19.4-23.4	379	14.4	12.9-15.9	100	2.0	1.6-2.4	57	2.7	2.1-3.5	43	1.5	1.1-2.1
2009	915	18.5	17.3-19.7	493	22.8	20.8-25.0	422	15.6	14.1-17.2	97	2.0	1.6-2.4	63	3.0	2.3-3.8	34	1.2	0.8-1.7
2010	792	16.1	15.0-17.3	427	19.8	18.0-21.8	365	13.6	12.2-15.2	101	2.0	1.6-2.4	59	2.7	2.0-3.5	42	1.4	1.0-1.9
2011	949	18.9	17.7-20.2	513	23.2	21.2-25.4	436	16.2	14.6-17.8	82	1.6	1.2-1.9	43	2.0	1.4-2.7	39	1.2	0.9-1.7
2012	820	16.1	15.0-17.3	448	19.9	18.1-21.9	372	13.5	12.1-15.0	124	2.3	1.9-2.8	69	3.1	2.4-3.9	55	1.8	1.3-2.3
2013	908	17.6	16.5-18.8	516	22.7	20.8-24.8	392	14.1	12.7-15.7	101	1.9	1.5-2.3	64	2.8	2.1-3.6	37	1.2	0.9-1.7
2014	875	16.9	15.7-18.0	467	20.7	18.8-22.7	408	14.4	13.0-15.9	98	1.8	1.5-2.2	67	3.0	2.3-3.8	31	0.9	0.6-1.3
2015	853	16.3	15.2-17.5	477	20.9	19.0-22.9	376	13.1	11.8-14.6	66	1.3	1.0-1.7	40	1.8	1.3-2.5	26	1.0	0.6-1.4
2016	882	16.8	15.7-18.0	490	21.1	19.2-23.1	392	13.9	12.5-15.4	77	1.4	1.1-1.8	43	1.8	1.3-2.5	34	1.1	0.7-1.6
2017	853	16.3	15.2-17.4	497	21.2	19.3-23.1	356	12.7	11.4-14.2	87	1.5	1.2-1.9	51	2.1	1.6-2.8	36	1.1	0.7-1.5
2018	839	15.5	14.5-16.6	478	19.8	18.0-21.6	361	12.4	11.1-13.8	67	1.2	0.9-1.5	44	1.8	1.3-2.5	23	0.7	0.4-1.1
2019	902	16.9	15.8-18.1	501	20.9	19.1-22.8	401	14.3	12.9-15.8	71	1.3	1.0-1.6	46	1.9	1.4-2.6	25	0.8	0.5-1.2
2020	684	12.7	11.7-13.7	380	15.8	14.2-17.4	304	10.6	9.4-11.9	63	1.2	0.9-1.5	32	1.4	0.9-1.9	31	1.0	0.7-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3,513	106.3	102.6-110.0	1,353	38.7	36.5-40.9
1977	3,324	102.5	98.8-106.3	1,339	39.3	37.2-41.6
1978	3,356	104.7	101.0-108.5	1,332	39.6	37.4-41.9
1979	3,280	104.4	100.7-108.3	1,199	36.6	34.5-38.9
1980	3,229	104.4	100.7-108.3	1,314	39.8	37.6-42.1
1981	3,364	110.1	106.2-114.1	1,273	39.3	37.1-41.7
1982	3,280	108.4	104.5-112.4	1,190	37.0	34.8-39.2
1983	3,269	108.2	104.3-112.2	1,119	35.2	33.0-37.4
1984	3,477	116.0	112.0-120.2	1,186	37.7	35.5-40.0
1985	3,442	115.3	111.2-119.4	1,153	36.5	34.3-38.8
1986	3,337	113.1	109.1-117.2	1,102	35.0	32.9-37.3
1987	3,437	116.9	112.8-121.1	1,069	34.3	32.2-36.5
1988	3,441	119.0	114.9-123.2	1,162	37.3	35.1-39.7
1989	3,384	119.9	115.7-124.2	1,097	36.4	34.2-38.7
1990	3,611	130.8	126.4-135.3	1,031	34.6	32.4-36.9
1991	3,646	133.6	129.2-138.2	986	33.5	31.4-35.8
1992	3,638	134.7	130.2-139.3	977	33.3	31.2-35.5
1993	3,421	127.2	122.8-131.6	987	33.6	31.5-35.9
1994	3,767	139.3	134.8-144.0	987	34.0	31.8-36.2
1995	3,661	136.7	132.2-141.4	1,030	35.7	33.5-38.0
1996	3,840	144.4	139.8-149.1	1,051	36.9	34.6-39.3
1997	3,672	137.8	133.3-142.4	904	32.0	29.9-34.2
1998	3,729	139.7	135.2-144.3	842	29.6	27.6-31.7
1999	3,733	139.0	134.5-143.6	818	28.1	26.2-30.2
2000	3,628	134.6	130.2-139.1	812	27.6	25.7-29.6
2001	3,472	128.2	123.9-132.6	828	28.3	26.3-30.3
2002	3,404	125.5	121.2-129.9	762	25.7	23.9-27.7
2003	3,320	123.5	119.2-127.8	742	25.3	23.5-27.3
2004	3,288	122.8	118.5-127.1	763	25.9	24.1-27.9
2005	3,212	120.7	116.5-125.0	720	24.7	22.9-26.6
2006	3,284	125.2	120.9-129.7	711	24.7	22.9-26.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	3,421	131.4	127.0-136.0	663	22.9	21.1-24.7
2008	3,438	131.1	126.7-135.7	659	22.7	21.0-24.6
2009	3,488	132.5	128.0-137.1	650	22.7	20.9-24.6
2010	3,337	126.8	122.4-131.3	641	22.0	20.3-23.8
2011	3,424	128.7	124.3-133.2	597	20.3	18.6-22.0
2012	3,287	123.1	118.8-127.5	606	20.6	19.0-22.4
2013	3,472	129.3	124.9-133.8	599	19.9	18.3-21.7
2014	3,533	130.2	125.8-134.7	567	18.7	17.1-20.4
2015	3,530	130.7	126.3-135.2	565	19.1	17.5-20.8
2016	3,482	127.8	123.4-132.2	565	18.1	16.6-19.8
2017	3,669	135.6	131.1-140.2	538	17.3	15.8-18.9
2018	3,666	134.5	130.0-139.1	550	17.6	16.1-19.2
2019	3,766	137.2	132.6-141.8	557	17.9	16.4-19.6
2020	3,128	115.8	111.6-120.1	487	16.0	14.5-17.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	383	12.7	11.4-14.1	129	3.9	3.2-4.7
1977	339	11.2	10.0-12.6	128	3.9	3.2-4.7
1978	342	12.3	10.9-13.7	126	4.1	3.4-4.9
1979	357	13.0	11.6-14.5	103	3.4	2.7-4.2
1980	319	12.0	10.7-13.5	105	3.3	2.7-4.0
1981	310	11.3	10.0-12.7	113	4.0	3.2-4.8
1982	311	11.9	10.5-13.3	109	3.7	3.0-4.5
1983	324	12.3	10.9-13.7	90	3.0	2.4-3.8
1984	303	11.5	10.2-13.0	77	2.7	2.1-3.4
1985	304	11.7	10.4-13.2	81	2.9	2.3-3.6
1986	269	10.3	9.0-11.7	66	2.3	1.7-2.9
1987	322	12.3	11.0-13.8	70	2.3	1.8-2.9
1988	295	11.4	10.1-12.9	77	2.9	2.2-3.6
1989	291	11.4	10.1-12.8	64	2.4	1.8-3.1
1990	322	12.9	11.5-14.4	89	3.3	2.7-4.1
1991	330	13.5	12.0-15.1	69	2.6	2.0-3.3
1992	328	12.9	11.5-14.4	62	2.3	1.8-3.0
1993	302	12.3	11.0-13.9	65	2.5	1.9-3.2
1994	347	14.1	12.7-15.7	95	3.7	3.0-4.5
1995	345	13.9	12.5-15.5	87	3.4	2.7-4.2
1996	339	13.8	12.4-15.4	96	3.6	2.9-4.4
1997	314	12.5	11.1-14.0	85	3.4	2.7-4.2
1998	287	11.5	10.2-12.9	75	2.9	2.3-3.6
1999	280	11.0	9.8-12.4	94	3.7	3.0-4.5
2000	260	10.3	9.1-11.7	81	3.0	2.4-3.7
2001	218	8.8	7.6-10.0	64	2.4	1.9-3.1
2002	245	10.0	8.7-11.3	71	2.7	2.1-3.4
2003	235	9.2	8.1-10.5	78	3.0	2.4-3.7
2004	223	8.7	7.6-10.0	69	2.7	2.1-3.4
2005	248	10.1	8.8-11.4	77	2.8	2.2-3.5
2006	219	8.7	7.6-10.0	76	2.9	2.3-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	215	9.0	7.8-10.3	66	2.6	2.0-3.3
2008	231	9.4	8.2-10.7	61	2.3	1.7-3.0
2009	222	9.1	7.9-10.4	59	2.2	1.7-2.9
2010	177	7.5	6.4-8.7	64	2.4	1.8-3.1
2011	188	7.6	6.6-8.8	68	2.6	2.0-3.3
2012	207	8.4	7.3-9.7	62	2.3	1.8-3.0
2013	204	8.2	7.0-9.4	61	2.1	1.6-2.7
2014	194	7.8	6.7-9.1	75	2.6	2.0-3.3
2015	193	7.7	6.6-8.9	61	2.2	1.7-2.9
2016	199	8.0	6.9-9.2	62	2.3	1.8-3.0
2017	202	8.1	7.0-9.3	57	2.2	1.6-2.8
2018	188	7.4	6.3-8.6	53	1.9	1.4-2.6
2019	162	6.5	5.5-7.7	31	1.1	0.8-1.7
2020	148	5.8	4.9-6.9	36	1.3	0.9-1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals**

Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	853	24.2	22.6-26.0	248	6.6	5.8-7.5
1977	749	21.7	20.1-23.4	198	5.4	4.7-6.3
1978	683	19.6	18.1-21.2	184	5.0	4.3-5.9
1979	755	22.4	20.8-24.2	169	4.6	3.9-5.4
1980	789	23.9	22.1-25.7	196	5.4	4.6-6.2
1981	734	22.3	20.6-24.0	171	4.8	4.1-5.6
1982	732	22.7	21.0-24.5	164	4.7	4.0-5.5
1983	752	23.0	21.3-24.8	165	4.6	3.9-5.4
1984	684	21.2	19.6-22.9	163	4.7	4.0-5.5
1985	700	21.9	20.3-23.7	130	3.8	3.1-4.5
1986	665	21.5	19.9-23.3	138	3.9	3.3-4.6
1987	666	21.7	20.0-23.4	153	4.5	3.8-5.4
1988	664	21.9	20.2-23.7	125	3.7	3.0-4.4
1989	636	21.3	19.7-23.1	130	3.9	3.3-4.7
1990	607	21.3	19.6-23.2	130	3.9	3.2-4.6
1991	697	24.8	23.0-26.8	144	4.7	3.9-5.5
1992	687	24.8	22.9-26.8	137	4.4	3.7-5.2
1993	657	23.5	21.7-25.5	146	4.7	3.9-5.5
1994	730	26.2	24.3-28.2	146	4.8	4.0-5.7
1995	730	26.8	24.8-28.8	155	5.1	4.3-6.0
1996	716	26.2	24.2-28.2	141	4.7	3.9-5.5
1997	766	28.3	26.3-30.4	148	5.0	4.2-5.8
1998	808	30.1	28.1-32.3	145	5.0	4.2-5.9
1999	813	29.7	27.6-31.8	153	5.2	4.4-6.1
2000	746	27.3	25.4-29.4	132	4.2	3.5-5.0
2001	807	29.6	27.6-31.7	125	4.2	3.5-5.0
2002	809	29.4	27.3-31.5	133	4.5	3.7-5.3
2003	808	29.3	27.3-31.4	140	4.7	4.0-5.6
2004	848	31.1	29.0-33.3	125	4.2	3.5-5.1
2005	825	30.2	28.2-32.4	144	5.0	4.2-5.9
2006	827	30.3	28.2-32.4	144	4.9	4.1-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals**

Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	776	28.4	26.4-30.6	152	5.2	4.4-6.1
2008	935	34.5	32.3-36.8	149	5.1	4.3-6.0
2009	852	31.6	29.4-33.8	125	4.3	3.5-5.1
2010	827	30.4	28.3-32.6	177	6.0	5.1-7.0
2011	891	32.3	30.2-34.5	165	5.6	4.7-6.5
2012	827	29.5	27.4-31.6	152	5.0	4.2-5.9
2013	917	32.6	30.5-34.9	159	5.3	4.5-6.2
2014	862	30.1	28.1-32.2	151	5.1	4.3-6.0
2015	913	32.3	30.2-34.5	188	6.0	5.2-7.0
2016	916	32.1	30.0-34.3	194	6.4	5.5-7.4
2017	889	31.0	29.0-33.2	174	5.5	4.7-6.4
2018	847	29.6	27.5-31.7	144	4.5	3.8-5.3
2019	812	28.3	26.4-30.4	173	5.6	4.8-6.5
2020	723	25.0	23.2-27.0	174	5.4	4.6-6.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	567	16.2	14.9-17.7	381	10.7	9.6-11.9
1977	507	15.6	14.2-17.1	395	11.2	10.1-12.5
1978	530	16.4	15.0-18.0	335	9.7	8.7-10.9
1979	529	16.9	15.4-18.5	340	9.8	8.7-11.0
1980	523	16.5	15.0-18.0	354	10.4	9.3-11.6
1981	548	17.5	16.0-19.2	336	10.0	8.9-11.2
1982	554	18.2	16.6-19.9	314	9.5	8.4-10.6
1983	540	17.5	16.0-19.2	319	9.6	8.5-10.8
1984	520	17.0	15.5-18.6	328	10.0	8.9-11.1
1985	556	18.6	17.0-20.3	296	9.4	8.3-10.6
1986	476	16.0	14.5-17.5	316	9.9	8.8-11.1
1987	516	17.9	16.3-19.6	304	9.7	8.6-10.9
1988	481	16.6	15.0-18.2	342	10.8	9.6-12.1
1989	474	16.4	14.9-18.0	269	8.6	7.6-9.8
1990	489	17.7	16.1-19.4	283	9.3	8.2-10.5
1991	527	18.9	17.2-20.6	312	10.5	9.4-11.8
1992	483	17.9	16.3-19.6	311	10.7	9.5-12.0
1993	460	17.1	15.6-18.8	265	8.8	7.8-10.0
1994	485	17.7	16.1-19.4	274	9.4	8.3-10.6
1995	542	20.3	18.6-22.2	277	9.7	8.5-10.9
1996	482	18.0	16.4-19.7	280	10.0	8.8-11.2
1997	484	18.1	16.5-19.8	289	10.2	9.0-11.5
1998	439	16.5	14.9-18.1	248	8.7	7.6-9.9
1999	435	16.2	14.7-17.8	257	8.9	7.9-10.1
2000	417	15.2	13.8-16.8	265	9.1	8.0-10.3
2001	463	17.1	15.6-18.8	239	8.3	7.3-9.4
2002	456	16.3	14.9-17.9	263	9.1	8.0-10.2
2003	396	14.5	13.1-16.1	269	9.3	8.2-10.5
2004	425	15.7	14.2-17.3	265	9.0	8.0-10.2
2005	378	14.1	12.7-15.6	244	8.7	7.6-9.8
2006	376	13.9	12.5-15.4	246	8.6	7.5-9.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	399	15.0	13.5-16.6	230	8.0	6.9-9.1
2008	388	14.3	12.9-15.8	232	8.0	7.0-9.1
2009	395	15.0	13.5-16.6	216	7.2	6.2-8.3
2010	373	13.9	12.5-15.4	223	7.6	6.6-8.7
2011	394	14.4	13.0-16.0	236	7.8	6.8-8.9
2012	341	12.3	11.0-13.7	233	8.0	7.0-9.1
2013	361	13.2	11.9-14.7	230	7.7	6.7-8.8
2014	368	13.2	11.9-14.7	201	6.5	5.6-7.5
2015	373	13.6	12.2-15.1	195	6.6	5.7-7.6
2016	342	12.4	11.1-13.8	213	6.9	6.0-7.9
2017	352	12.3	11.0-13.7	218	7.1	6.2-8.2
2018	279	9.9	8.7-11.2	193	6.3	5.4-7.3
2019	290	10.5	9.3-11.9	201	6.4	5.5-7.4
2020	275	10.2	9.0-11.5	154	5.0	4.2-5.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,580	68.0	64.5-71.6	563	26.3	24.1-28.7
1977	1,576	67.6	64.1-71.2	572	26.2	24.0-28.6
1978	1,556	67.7	64.2-71.3	591	27.4	25.2-29.8
1979	1,705	75.3	71.6-79.1	533	25.8	23.5-28.1
1980	1,822	80.8	77.0-84.7	552	26.8	24.5-29.2
1981	1,728	76.8	73.1-80.6	556	27.1	24.9-29.6
1982	1,693	76.6	72.8-80.4	519	25.4	23.2-27.7
1983	1,725	77.1	73.4-80.9	587	28.7	26.4-31.2
1984	1,748	79.2	75.4-83.1	612	30.1	27.7-32.7
1985	1,822	81.8	78.0-85.8	520	24.9	22.8-27.2
1986	1,847	85.1	81.2-89.1	507	24.7	22.5-27.0
1987	1,837	84.2	80.3-88.2	524	25.8	23.6-28.2
1988	1,733	82.1	78.2-86.1	583	29.1	26.8-31.7
1989	1,736	82.4	78.5-86.4	548	27.9	25.6-30.4
1990	2,128	101.8	97.5-106.3	568	29.3	26.9-31.9
1991	2,431	115.8	111.2-120.6	534	27.8	25.5-30.3
1992	2,999	143.1	138.0-148.4	596	31.0	28.5-33.6
1993	2,837	136.1	131.0-141.2	573	30.0	27.6-32.6
1994	2,974	143.1	137.9-148.3	606	32.2	29.7-34.9
1995	2,681	129.8	124.9-134.9	596	31.6	29.1-34.2
1996	2,725	133.5	128.5-138.6	559	29.8	27.4-32.4
1997	2,917	142.9	137.8-148.2	494	26.5	24.2-28.9
1998	2,929	142.4	137.3-147.7	531	28.0	25.7-30.5
1999	3,007	145.6	140.5-151.0	431	22.5	20.4-24.8
2000	3,457	166.1	160.6-171.8	484	24.9	22.8-27.3
2001	3,228	154.1	148.8-159.5	463	23.7	21.6-26.0
2002	3,176	150.7	145.5-156.0	463	23.7	21.6-25.9
2003	2,836	134.8	129.9-139.9	452	22.9	20.8-25.1
2004	2,799	131.9	127.0-136.9	424	21.5	19.5-23.6
2005	2,856	134.4	129.5-139.5	403	20.4	18.4-22.5
2006	3,219	152.8	147.6-158.2	382	19.3	17.4-21.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	3,330	156.4	151.1-161.9	392	19.8	17.8-21.8
2008	3,158	146.7	141.6-151.9	377	19.0	17.1-21.0
2009	3,179	145.8	140.7-151.0	393	19.7	17.8-21.7
2010	2,928	132.5	127.7-137.5	426	21.0	19.0-23.1
2011	3,126	139.0	134.1-144.0	391	18.8	16.9-20.7
2012	2,677	116.2	111.8-120.8	339	16.1	14.4-17.9
2013	2,609	111.8	107.5-116.2	327	15.3	13.7-17.1
2014	2,497	104.2	100.1-108.5	364	16.7	15.0-18.6
2015	2,495	102.7	98.7-106.9	358	15.9	14.2-17.6
2016	2,581	104.9	100.9-109.1	382	16.7	15.1-18.5
2017	2,721	109.2	105.1-113.4	389	16.8	15.2-18.6
2018	2,726	108.7	104.6-112.9	313	13.2	11.8-14.8
2019	2,970	116.4	112.2-120.8	334	13.9	12.4-15.5
2020	2,469	96.3	92.5-100.2	258	10.6	9.3-12.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Testis**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	76	2.8	2.2-3.6	16	0.6	0.3-1.0
1977	72	2.8	2.1-3.5	16	0.6	0.3-1.0
1978	70	2.8	2.2-3.6	15	0.6	0.3-1.0
1979	80	3.4	2.6-4.2	16	0.6	0.3-1.0
1980	96	4.0	3.2-5.0	16	0.7	0.4-1.1
1981	98	3.8	3.1-4.7	4	0.1	0.0-0.4
1982	108	4.3	3.5-5.3	7	0.3	0.1-0.6
1983	101	3.9	3.2-4.8	7	0.3	0.1-0.6
1984	94	3.9	3.1-4.8	10	0.4	0.2-0.8
1985	97	3.9	3.1-4.8	14	0.5	0.3-0.9
1986	109	4.3	3.5-5.3	9	0.4	0.2-0.8
1987	107	4.3	3.5-5.2	7	0.3	0.1-0.6
1988	90	3.6	2.9-4.5	7	0.2	0.1-0.5
1989	103	4.2	3.4-5.2	10	0.4	0.2-0.8
1990	118	4.9	4.0-5.9	8	0.3	0.1-0.7
1991	121	4.9	4.1-5.9	9	0.4	0.2-0.8
1992	115	5.0	4.1-6.0	9	0.4	0.2-0.8
1993	118	4.8	4.0-5.8	9	0.4	0.2-0.8
1994	117	4.8	4.0-5.8	9	0.4	0.2-0.8
1995	107	4.4	3.6-5.4	7	0.3	0.1-0.7
1996	108	4.4	3.6-5.3	5	0.2	0.1-0.5
1997	102	4.2	3.4-5.1	6	0.2	0.1-0.5
1998	100	4.1	3.4-5.1	7	0.3	0.1-0.6
1999	146	5.9	4.9-6.9	4	0.2	0.1-0.5
2000	128	5.2	4.3-6.2	2	0.1	0.0-0.3
2001	120	4.8	4.0-5.8	6	0.3	0.1-0.6
2002	127	5.2	4.3-6.2	3	0.1	0.0-0.4
2003	139	5.8	4.8-6.8	8	0.4	0.2-0.7
2004	146	6.0	5.1-7.1	8	0.3	0.1-0.7
2005	150	6.1	5.2-7.2	7	0.3	0.1-0.6
2006	150	6.1	5.2-7.2	11	0.5	0.2-0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Testis**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	161	6.7	5.7-7.8	1	0.0	0.0-0.2
2008	148	6.3	5.3-7.4	5	0.2	0.1-0.5
2009	170	7.1	6.1-8.3	8	0.3	0.1-0.7
2010	162	6.8	5.8-8.0	14	0.6	0.3-1.0
2011	167	6.9	5.9-8.1	7	0.3	0.1-0.6
2012	175	7.3	6.2-8.5	7	0.3	0.1-0.6
2013	160	6.6	5.6-7.7	6	0.2	0.1-0.5
2014	154	6.4	5.4-7.5	2	0.1	0.0-0.3
2015	162	6.5	5.5-7.7	5	0.2	0.1-0.5
2016	160	6.6	5.6-7.8	4	0.2	0.0-0.5
2017	172	7.0	6.0-8.1	5	0.2	0.1-0.5
2018	162	7.0	5.9-8.2	6	0.2	0.1-0.5
2019	172	7.3	6.2-8.5	1	0.0	0.0-0.2
2020	172	7.6	6.5-8.8	6	0.3	0.1-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,177	18.7	17.7-19.9	843	33.8	31.4-36.3	334	9.0	8.0-10.0	365	6.3	5.7-7.0	249	11.2	9.7-12.7	116	3.4	2.8-4.1
1977	1,266	20.5	19.4-21.7	893	36.2	33.8-38.8	373	10.3	9.2-11.4	384	6.5	5.9-7.2	267	11.9	10.4-13.4	117	3.3	2.7-4.0
1978	1,318	21.8	20.6-23.1	951	40.0	37.4-42.7	367	10.0	9.0-11.1	353	6.0	5.4-6.7	234	10.5	9.1-12.0	119	3.4	2.8-4.1
1979	1,328	22.3	21.1-23.6	926	39.7	37.1-42.5	402	11.4	10.2-12.6	346	5.9	5.3-6.6	242	11.1	9.7-12.6	104	2.9	2.3-3.5
1980	1,391	23.4	22.1-24.7	1,038	44.6	41.9-47.5	353	9.8	8.8-10.9	358	6.1	5.5-6.8	249	11.3	9.9-12.9	109	3.0	2.5-3.7
1981	1,354	23.0	21.8-24.3	986	42.9	40.2-45.7	368	10.4	9.4-11.6	301	5.2	4.6-5.8	202	9.3	8.0-10.7	99	2.7	2.2-3.3
1982	1,251	21.5	20.3-22.7	887	38.8	36.2-41.5	364	10.5	9.4-11.6	317	5.4	4.8-6.1	226	10.5	9.1-12.0	91	2.5	2.0-3.1
1983	1,389	23.9	22.6-25.2	977	43.8	41.0-46.7	412	11.6	10.5-12.8	284	4.9	4.3-5.5	189	9.0	7.7-10.4	95	2.6	2.1-3.2
1984	1,260	21.6	20.4-22.8	910	39.8	37.2-42.5	350	10.0	8.9-11.1	282	4.9	4.3-5.5	186	8.7	7.4-10.1	96	2.7	2.2-3.3
1985	1,134	20.0	18.8-21.2	802	36.4	33.8-39.1	332	9.5	8.5-10.6	292	5.1	4.5-5.8	201	9.8	8.5-11.3	91	2.5	2.0-3.1
1986	1,209	21.6	20.4-22.9	855	39.2	36.6-42.0	354	10.6	9.5-11.8	260	4.5	4.0-5.1	164	7.9	6.7-9.3	96	2.6	2.1-3.2
1987	1,234	21.8	20.6-23.1	880	40.0	37.4-42.8	354	10.4	9.3-11.6	243	4.3	3.8-4.9	172	8.4	7.2-9.8	71	2.1	1.6-2.6
1988	1,188	21.3	20.0-22.5	829	38.0	35.4-40.8	359	10.5	9.4-11.6	271	4.7	4.2-5.4	169	8.1	6.9-9.5	102	2.8	2.3-3.4
1989	1,117	20.7	19.5-21.9	792	37.7	35.1-40.5	325	10.0	8.9-11.1	277	5.0	4.4-5.7	175	8.8	7.5-10.3	102	2.9	2.4-3.6
1990	1,143	21.3	20.1-22.6	794	38.2	35.6-41.0	349	10.5	9.4-11.7	249	4.6	4.0-5.2	159	8.1	6.8-9.4	90	2.7	2.1-3.3
1991	1,210	23.0	21.7-24.4	865	42.1	39.3-45.1	345	10.8	9.7-12.1	226	4.2	3.7-4.8	148	7.5	6.4-8.9	78	2.3	1.8-2.8
1992	1,191	22.8	21.5-24.2	837	41.1	38.4-44.1	354	11.4	10.2-12.7	233	4.3	3.8-4.9	152	7.9	6.7-9.2	81	2.4	1.9-3.0
1993	1,099	21.3	20.0-22.6	785	38.6	35.9-41.4	314	10.2	9.0-11.4	273	5.2	4.6-5.8	187	9.6	8.3-11.1	86	2.6	2.1-3.3
1994	1,210	23.5	22.2-24.9	839	41.5	38.7-44.4	371	12.1	10.8-13.4	258	4.8	4.3-5.5	175	9.1	7.8-10.6	83	2.5	2.0-3.1
1995	1,176	23.1	21.8-24.5	843	42.1	39.3-45.1	333	10.8	9.7-12.1	288	5.5	4.9-6.2	185	9.6	8.3-11.1	103	3.0	2.5-3.7
1996	1,036	20.5	19.3-21.8	742	37.4	34.7-40.2	294	9.5	8.4-10.7	239	4.6	4.0-5.2	146	7.7	6.5-9.1	93	2.8	2.2-3.4
1997	1,185	23.7	22.4-25.1	851	42.5	39.7-45.5	334	10.9	9.8-12.2	233	4.5	3.9-5.1	154	8.1	6.9-9.5	79	2.5	2.0-3.1
1998	1,224	24.4	23.0-25.8	851	42.2	39.4-45.2	373	12.2	11.0-13.5	253	4.9	4.3-5.5	160	8.2	7.0-9.6	93	2.8	2.2-3.4
1999	1,119	22.1	20.8-23.4	786	38.9	36.2-41.7	333	11.0	9.8-12.3	240	4.5	4.0-5.2	166	8.6	7.3-10.0	74	2.2	1.7-2.8
2000	1,197	23.5	22.1-24.8	839	41.3	38.5-44.1	358	11.8	10.6-13.1	228	4.3	3.8-4.9	154	7.8	6.6-9.1	74	2.2	1.7-2.8
2001	1,130	22.2	20.9-23.5	810	39.5	36.8-42.3	320	10.6	9.4-11.8	234	4.4	3.9-5.1	165	8.3	7.0-9.6	69	2.1	1.6-2.7
2002	1,087	21.2	20.0-22.5	787	38.4	35.8-41.2	300	9.8	8.7-11.1	208	3.9	3.4-4.4	136	6.8	5.7-8.1	72	2.1	1.6-2.7
2003	1,155	22.6	21.3-24.0	832	40.5	37.8-43.4	323	10.5	9.4-11.8	192	3.6	3.1-4.1	131	6.6	5.5-7.8	61	1.8	1.4-2.4
2004	1,185	23.2	21.9-24.5	841	40.5	37.8-43.3	344	11.1	10.0-12.4	225	4.2	3.7-4.8	157	7.7	6.6-9.0	68	2.0	1.6-2.6
2005	1,155	22.8	21.5-24.2	810	39.1	36.4-41.9	345	11.5	10.3-12.8	225	4.3	3.7-4.9	143	7.2	6.0-8.4	82	2.5	1.9-3.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,121	22.2	20.9-23.5	836	40.6	37.9-43.5	285	9.6	8.5-10.8	240	4.6	4.0-5.2	156	7.9	6.7-9.2	84	2.5	2.0-3.2
2007	1,181	23.5	22.2-24.9	858	42.0	39.2-44.9	323	10.9	9.7-12.2	242	4.6	4.0-5.2	161	8.0	6.8-9.3	81	2.4	1.9-3.1
2008	1,169	23.2	21.9-24.6	891	42.8	40.0-45.7	278	9.5	8.4-10.7	227	4.4	3.8-5.0	159	7.9	6.7-9.2	68	2.2	1.7-2.8
2009	1,139	22.5	21.2-23.9	823	39.6	36.9-42.4	316	10.6	9.4-11.8	215	4.0	3.5-4.6	143	7.1	5.9-8.3	72	2.1	1.6-2.7
2010	1,098	21.6	20.3-22.9	835	39.8	37.1-42.6	263	8.9	7.9-10.1	241	4.5	3.9-5.1	170	8.1	6.9-9.5	71	2.1	1.6-2.7
2011	1,104	21.5	20.2-22.8	812	38.1	35.5-40.9	292	9.8	8.7-11.0	221	4.1	3.6-4.7	150	7.1	6.0-8.4	71	2.2	1.7-2.8
2012	1,121	21.5	20.2-22.8	824	38.1	35.5-40.8	297	9.7	8.6-11.0	234	4.3	3.7-4.9	159	7.4	6.3-8.6	75	2.2	1.7-2.8
2013	1,095	20.8	19.5-22.1	810	36.6	34.1-39.2	285	9.3	8.2-10.5	221	3.9	3.4-4.4	156	7.2	6.1-8.4	65	1.8	1.4-2.3
2014	1,117	20.7	19.5-22.0	825	36.7	34.2-39.4	292	9.2	8.2-10.4	243	4.3	3.7-4.8	170	7.7	6.6-9.0	73	2.0	1.6-2.6
2015	1,216	22.5	21.2-23.8	942	41.3	38.7-44.1	274	8.9	7.9-10.1	230	4.0	3.5-4.6	154	6.8	5.8-8.0	76	2.2	1.7-2.8
2016	1,129	20.6	19.4-21.9	864	37.4	34.9-40.0	265	8.5	7.5-9.6	227	4.0	3.5-4.5	154	6.8	5.8-8.0	73	2.1	1.7-2.7
2017	1,147	20.6	19.4-21.9	843	35.6	33.2-38.1	304	9.7	8.6-10.9	216	3.7	3.2-4.2	159	6.8	5.8-8.0	57	1.6	1.2-2.1
2018	1,136	20.2	19.0-21.4	851	35.5	33.1-38.0	285	8.9	7.8-10.0	206	3.5	3.0-4.0	139	5.9	4.9-6.9	67	1.9	1.4-2.4
2019	1,144	20.3	19.1-21.6	822	33.8	31.5-36.2	322	10.1	9.0-11.4	214	3.5	3.0-4.0	152	6.2	5.3-7.3	62	1.8	1.3-2.3
2020	929	16.4	15.3-17.5	695	28.4	26.3-30.6	234	7.5	6.5-8.5	157	2.6	2.2-3.1	90	3.7	3.0-4.5	67	1.9	1.5-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	463	7.5	6.8-8.2	286	10.9	9.6-12.3	177	5.1	4.3-5.9	241	3.9	3.4-4.4	158	6.4	5.4-7.5	83	2.2	1.8-2.8
1977	457	7.6	6.9-8.3	295	11.7	10.3-13.1	162	4.7	4.0-5.6	233	3.8	3.3-4.4	150	6.1	5.1-7.2	83	2.2	1.8-2.8
1978	448	7.4	6.7-8.2	272	11.0	9.7-12.5	176	4.9	4.2-5.7	221	3.7	3.2-4.2	144	5.7	4.8-6.8	77	2.3	1.8-2.9
1979	432	7.3	6.6-8.1	268	11.1	9.8-12.6	164	4.6	3.9-5.3	227	3.9	3.4-4.5	143	6.1	5.1-7.3	84	2.4	1.9-3.0
1980	414	7.0	6.4-7.8	258	10.4	9.1-11.7	156	4.8	4.0-5.6	192	3.2	2.8-3.7	117	5.0	4.1-6.0	75	2.1	1.6-2.6
1981	417	7.2	6.5-8.0	240	10.2	9.0-11.7	177	5.2	4.4-6.1	208	3.5	3.1-4.1	120	5.0	4.1-6.0	88	2.5	2.0-3.1
1982	463	8.3	7.6-9.2	274	12.1	10.6-13.6	189	5.8	4.9-6.7	209	3.6	3.1-4.2	124	5.5	4.6-6.6	85	2.4	1.9-3.0
1983	453	8.1	7.3-8.9	281	12.1	10.7-13.6	172	5.3	4.5-6.3	190	3.3	2.8-3.8	116	5.0	4.1-6.1	74	2.1	1.7-2.7
1984	483	8.7	7.9-9.5	301	13.1	11.7-14.8	182	5.5	4.7-6.4	199	3.5	3.1-4.1	122	5.4	4.5-6.5	77	2.3	1.8-2.9
1985	517	9.3	8.5-10.2	314	13.7	12.2-15.3	203	6.3	5.4-7.3	182	3.2	2.8-3.7	104	4.6	3.7-5.6	78	2.4	1.9-3.0
1986	494	8.8	8.0-9.6	297	13.0	11.5-14.6	197	5.8	5.0-6.8	205	3.6	3.1-4.1	126	5.6	4.6-6.7	79	2.2	1.8-2.8
1987	547	10.0	9.1-10.9	326	14.4	12.8-16.1	221	6.9	6.0-7.9	189	3.4	2.9-4.0	117	5.4	4.4-6.5	72	2.2	1.7-2.7
1988	456	8.7	7.9-9.5	284	13.1	11.6-14.7	172	5.7	4.9-6.7	176	3.3	2.8-3.8	106	4.9	4.0-5.9	70	2.2	1.7-2.9
1989	485	9.3	8.5-10.2	298	13.9	12.3-15.6	187	6.1	5.2-7.0	193	3.6	3.1-4.1	114	5.4	4.5-6.5	79	2.4	1.9-3.0
1990	529	10.2	9.4-11.2	298	13.9	12.4-15.6	231	7.8	6.8-8.9	185	3.5	3.0-4.1	102	4.9	4.0-6.0	83	2.6	2.0-3.2
1991	534	10.5	9.6-11.4	304	14.4	12.8-16.2	230	7.7	6.7-8.8	162	3.1	2.6-3.6	91	4.3	3.4-5.3	71	2.3	1.8-2.9
1992	534	10.7	9.8-11.6	316	15.2	13.6-17.0	218	7.4	6.4-8.5	152	2.9	2.5-3.4	86	4.3	3.4-5.3	66	2.1	1.6-2.7
1993	549	11.0	10.1-12.0	355	17.0	15.3-18.9	194	6.6	5.7-7.7	169	3.3	2.8-3.9	116	5.6	4.6-6.7	53	1.7	1.3-2.3
1994	595	12.1	11.2-13.2	367	18.0	16.2-20.0	228	7.8	6.8-8.9	195	3.8	3.3-4.4	120	5.9	4.9-7.0	75	2.4	1.8-3.0
1995	535	10.9	10.0-11.9	320	15.6	13.9-17.4	215	7.6	6.6-8.8	183	3.7	3.1-4.2	113	5.7	4.7-6.8	70	2.3	1.8-2.9
1996	625	12.8	11.8-13.8	370	18.3	16.4-20.2	255	8.9	7.8-10.1	176	3.5	3.0-4.1	98	4.9	4.0-6.0	78	2.5	1.9-3.1
1997	605	12.5	11.5-13.6	354	17.5	15.7-19.5	251	9.1	8.0-10.3	184	3.7	3.2-4.3	109	5.5	4.5-6.6	75	2.5	2.0-3.2
1998	645	13.3	12.3-14.4	389	18.9	17.1-20.9	256	9.2	8.1-10.5	174	3.5	3.0-4.0	95	4.8	3.9-5.8	79	2.7	2.1-3.4
1999	664	13.6	12.6-14.7	419	20.1	18.2-22.2	245	8.6	7.5-9.8	164	3.3	2.8-3.8	106	5.2	4.2-6.3	58	1.9	1.4-2.4
2000	685	13.9	12.9-15.0	402	19.2	17.4-21.2	283	10.0	8.8-11.3	158	3.1	2.7-3.7	101	4.9	4.0-6.0	57	1.9	1.4-2.4
2001	676	13.8	12.8-14.9	421	19.9	18.1-21.9	255	9.0	7.9-10.2	164	3.3	2.8-3.8	96	4.6	3.7-5.6	68	2.3	1.8-2.9
2002	680	13.9	12.8-15.0	423	19.9	18.1-21.9	257	8.9	7.9-10.1	170	3.3	2.8-3.9	106	5.1	4.2-6.2	64	2.1	1.6-2.7
2003	744	15.1	14.1-16.3	479	22.5	20.5-24.6	265	9.3	8.2-10.6	149	2.9	2.5-3.5	91	4.4	3.5-5.4	58	1.8	1.4-2.4
2004	757	15.5	14.4-16.6	476	22.4	20.4-24.5	281	10.2	9.0-11.5	163	3.2	2.7-3.8	100	4.8	3.9-5.9	63	2.1	1.6-2.7
2005	747	15.3	14.2-16.4	461	21.8	19.8-23.9	286	10.3	9.1-11.6	133	2.6	2.2-3.1	77	3.7	2.9-4.6	56	1.8	1.3-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	798	16.5	15.4-17.7	507	23.9	21.9-26.1	291	10.8	9.5-12.1	160	3.2	2.7-3.7	93	4.6	3.7-5.6	67	2.2	1.7-2.8
2007	764	15.8	14.7-17.0	480	22.9	20.8-25.0	284	10.4	9.2-11.7	163	3.2	2.7-3.7	101	4.8	3.9-5.8	62	2.0	1.5-2.6
2008	813	16.6	15.5-17.8	523	24.3	22.3-26.5	290	10.4	9.2-11.7	145	2.8	2.4-3.3	87	4.2	3.3-5.2	58	1.9	1.4-2.4
2009	723	14.9	13.8-16.0	436	20.3	18.5-22.4	287	10.5	9.3-11.8	181	3.6	3.1-4.2	125	6.0	5.0-7.1	56	1.9	1.4-2.5
2010	746	15.2	14.1-16.4	471	21.4	19.5-23.4	275	10.2	9.0-11.6	156	3.0	2.5-3.5	98	4.5	3.6-5.5	58	1.9	1.4-2.5
2011	751	15.3	14.2-16.4	469	21.1	19.2-23.1	282	10.3	9.1-11.6	138	2.6	2.2-3.0	92	4.1	3.3-5.1	46	1.4	1.0-1.9
2012	774	15.6	14.5-16.8	503	22.5	20.6-24.6	271	10.0	8.8-11.3	136	2.6	2.2-3.1	91	4.2	3.4-5.2	45	1.4	1.0-2.0
2013	795	15.6	14.5-16.7	511	22.4	20.4-24.4	284	10.0	8.9-11.3	135	2.5	2.1-3.0	75	3.4	2.7-4.3	60	1.8	1.3-2.3
2014	783	15.2	14.1-16.3	514	21.9	20.1-24.0	269	9.6	8.5-10.9	153	2.8	2.4-3.3	91	4.0	3.2-4.9	62	2.0	1.5-2.5
2015	807	15.4	14.4-16.6	512	21.5	19.7-23.5	295	10.2	9.0-11.5	179	3.3	2.8-3.8	118	5.1	4.2-6.2	61	1.8	1.4-2.4
2016	798	15.3	14.3-16.4	510	21.5	19.7-23.5	288	10.3	9.1-11.6	126	2.3	1.9-2.7	84	3.6	2.8-4.4	42	1.3	0.9-1.8
2017	853	16.1	15.0-17.3	542	22.3	20.5-24.3	311	11.0	9.7-12.3	145	2.7	2.2-3.1	91	3.9	3.1-4.8	54	1.8	1.3-2.4
2018	806	15.4	14.3-16.5	544	22.8	20.9-24.8	262	9.2	8.1-10.5	135	2.4	2.0-2.8	96	4.0	3.2-4.9	39	1.1	0.8-1.6
2019	867	16.6	15.5-17.8	587	24.3	22.3-26.4	280	10.2	9.0-11.5	129	2.3	1.9-2.7	80	3.2	2.6-4.0	49	1.5	1.1-2.0
2020	681	13.0	12.0-14.0	443	18.4	16.7-20.2	238	8.3	7.3-9.5	128	2.3	1.9-2.7	82	3.6	2.8-4.4	46	1.3	1.0-1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	351	5.8	5.2-6.4	189	7.0	6.0-8.1	162	4.9	4.1-5.7	273	4.6	4.0-5.2	152	5.9	5.0-6.9	121	3.7	3.0-4.4
1977	347	6.1	5.4-6.8	191	7.5	6.4-8.7	156	5.0	4.2-5.9	264	4.4	3.9-5.0	137	5.2	4.3-6.1	127	3.9	3.2-4.6
1978	351	6.3	5.6-7.0	175	7.0	6.0-8.2	176	5.6	4.8-6.6	262	4.4	3.9-5.0	135	5.4	4.5-6.4	127	3.6	3.0-4.4
1979	336	6.0	5.3-6.7	177	7.3	6.2-8.5	159	5.1	4.3-6.0	242	4.1	3.6-4.7	134	5.5	4.6-6.5	108	3.1	2.6-3.8
1980	333	6.0	5.3-6.7	166	6.9	5.9-8.1	167	5.3	4.5-6.2	268	4.5	4.0-5.1	144	5.7	4.8-6.8	124	3.7	3.0-4.4
1981	347	6.5	5.8-7.3	194	8.5	7.3-9.8	153	4.9	4.1-5.8	250	4.4	3.9-5.1	130	5.5	4.5-6.5	120	3.7	3.0-4.4
1982	342	6.3	5.6-7.0	185	7.9	6.8-9.2	157	5.1	4.3-6.0	255	4.6	4.0-5.2	125	5.3	4.4-6.4	130	4.0	3.3-4.8
1983	318	6.1	5.5-6.9	166	7.2	6.1-8.4	152	5.2	4.4-6.2	232	4.2	3.7-4.8	127	5.6	4.7-6.7	105	3.1	2.6-3.8
1984	344	6.6	5.9-7.3	161	7.0	5.9-8.2	183	6.3	5.4-7.3	232	4.1	3.6-4.7	115	4.9	4.0-5.9	117	3.7	3.0-4.4
1985	356	6.9	6.2-7.7	194	8.7	7.5-10.1	162	5.5	4.6-6.4	239	4.4	3.9-5.0	119	5.3	4.4-6.4	120	3.7	3.1-4.5
1986	333	6.6	5.9-7.3	188	8.6	7.4-9.9	145	5.2	4.3-6.1	244	4.6	4.0-5.2	135	6.3	5.3-7.5	109	3.3	2.7-4.0
1987	362	7.1	6.4-7.9	183	8.3	7.1-9.6	179	6.0	5.2-7.1	230	4.4	3.9-5.1	117	5.3	4.4-6.4	113	3.8	3.1-4.6
1988	367	7.3	6.5-8.1	190	8.7	7.5-10.1	177	6.2	5.3-7.3	169	3.2	2.7-3.7	90	4.0	3.2-5.0	79	2.5	2.0-3.2
1989	375	7.7	6.9-8.6	202	9.6	8.3-11.0	173	6.2	5.3-7.3	187	3.6	3.1-4.2	98	4.6	3.7-5.6	89	2.8	2.2-3.4
1990	374	7.8	7.0-8.7	191	9.1	7.9-10.6	183	6.9	5.9-8.0	194	3.9	3.4-4.5	106	5.1	4.2-6.2	88	3.0	2.4-3.8
1991	359	7.6	6.8-8.4	180	8.6	7.4-10.0	179	6.6	5.7-7.7	210	4.3	3.7-5.0	105	5.0	4.1-6.1	105	3.8	3.1-4.6
1992	352	7.4	6.7-8.3	192	9.0	7.8-10.4	160	6.0	5.1-7.1	173	3.5	3.0-4.1	91	4.5	3.6-5.5	82	2.8	2.2-3.5
1993	365	7.6	6.8-8.4	180	8.5	7.3-9.9	185	6.9	5.9-8.0	172	3.5	3.0-4.0	95	4.6	3.7-5.6	77	2.7	2.1-3.4
1994	394	8.4	7.6-9.3	197	9.5	8.2-11.0	197	7.4	6.4-8.6	184	3.8	3.3-4.4	94	4.6	3.7-5.6	90	3.2	2.6-4.0
1995	417	8.7	7.9-9.6	226	10.7	9.3-12.2	191	7.1	6.1-8.2	199	4.1	3.5-4.7	106	5.2	4.2-6.2	93	3.3	2.6-4.0
1996	414	8.8	8.0-9.7	212	10.1	8.8-11.6	202	7.7	6.6-8.8	187	3.9	3.4-4.6	94	4.6	3.7-5.7	93	3.3	2.7-4.1
1997	375	8.1	7.3-8.9	182	8.8	7.5-10.2	193	7.5	6.5-8.7	209	4.4	3.8-5.0	90	4.5	3.6-5.5	119	4.3	3.6-5.2
1998	431	9.2	8.3-10.1	229	10.8	9.4-12.3	202	8.0	6.9-9.2	215	4.5	3.9-5.1	104	5.1	4.1-6.1	111	4.0	3.3-4.8
1999	351	7.4	6.7-8.3	180	8.4	7.2-9.8	171	6.5	5.5-7.5	213	4.4	3.8-5.0	111	5.3	4.4-6.4	102	3.6	2.9-4.4
2000	365	7.6	6.9-8.5	174	8.2	7.0-9.5	191	7.2	6.2-8.3	185	3.9	3.3-4.5	93	4.4	3.6-5.4	92	3.4	2.8-4.3
2001	339	7.1	6.4-7.9	170	8.0	6.8-9.3	169	6.4	5.5-7.5	172	3.5	3.0-4.1	95	4.5	3.6-5.5	77	2.7	2.1-3.4
2002	353	7.4	6.6-8.2	181	8.4	7.3-9.8	172	6.6	5.6-7.6	165	3.4	2.9-4.0	87	4.1	3.3-5.1	78	2.9	2.3-3.6
2003	329	6.8	6.1-7.6	169	7.8	6.6-9.0	160	6.0	5.1-7.1	162	3.3	2.8-3.9	95	4.4	3.6-5.4	67	2.4	1.8-3.1
2004	333	7.1	6.3-7.9	186	8.8	7.6-10.2	147	5.7	4.8-6.8	147	3.0	2.5-3.6	76	3.6	2.8-4.5	71	2.6	2.0-3.3
2005	358	7.6	6.8-8.4	200	9.4	8.1-10.8	158	6.1	5.2-7.2	191	3.9	3.4-4.5	107	5.1	4.2-6.1	84	3.0	2.4-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	378	8.1	7.3-9.0	198	9.3	8.0-10.7	180	7.1	6.1-8.3	188	3.9	3.4-4.5	107	5.1	4.2-6.1	81	2.9	2.3-3.7
2007	319	6.9	6.2-7.7	166	7.9	6.7-9.2	153	6.1	5.2-7.2	205	4.2	3.7-4.8	109	5.1	4.2-6.2	96	3.5	2.8-4.3
2008	346	7.4	6.6-8.2	172	8.1	6.9-9.4	174	6.8	5.8-8.0	157	3.2	2.7-3.8	86	4.0	3.2-5.0	71	2.6	2.0-3.3
2009	354	7.4	6.6-8.2	187	8.5	7.3-9.9	167	6.3	5.4-7.4	195	4.0	3.5-4.7	106	5.0	4.1-6.1	89	3.4	2.7-4.2
2010	320	6.9	6.2-7.8	173	8.2	7.0-9.5	147	6.0	5.1-7.1	185	3.7	3.2-4.3	91	4.1	3.3-5.1	94	3.4	2.7-4.2
2011	311	6.5	5.8-7.3	166	7.6	6.5-8.9	145	5.7	4.8-6.7	192	4.0	3.4-4.6	102	4.8	3.9-5.8	90	3.3	2.6-4.1
2012	334	6.9	6.2-7.7	170	7.7	6.5-8.9	164	6.2	5.3-7.3	183	3.6	3.1-4.1	81	3.6	2.8-4.5	102	3.6	2.9-4.4
2013	328	6.8	6.1-7.7	177	7.9	6.8-9.2	151	5.8	4.9-6.9	170	3.3	2.8-3.9	85	3.9	3.1-4.9	85	2.8	2.2-3.5
2014	372	7.5	6.7-8.3	215	9.4	8.1-10.7	157	6.0	5.1-7.1	210	4.1	3.5-4.7	108	4.7	3.8-5.6	102	3.6	2.9-4.4
2015	365	7.4	6.6-8.2	186	8.2	7.0-9.5	179	6.7	5.7-7.8	193	3.7	3.2-4.3	96	4.1	3.3-5.0	97	3.4	2.7-4.2
2016	361	7.3	6.6-8.2	201	8.8	7.6-10.1	160	6.1	5.2-7.2	188	3.5	3.0-4.1	104	4.5	3.6-5.4	84	2.8	2.2-3.5
2017	371	7.6	6.8-8.4	202	8.9	7.7-10.2	169	6.4	5.4-7.5	219	4.2	3.7-4.8	118	5.0	4.2-6.1	101	3.6	2.9-4.4
2018	318	6.4	5.7-7.2	185	7.9	6.8-9.1	133	5.2	4.3-6.2	187	3.5	3.0-4.1	109	4.5	3.7-5.5	78	2.6	2.1-3.3
2019	319	6.4	5.7-7.2	176	7.5	6.4-8.7	143	5.4	4.5-6.4	192	3.7	3.1-4.2	100	4.2	3.4-5.1	92	3.2	2.5-3.9
2020	295	6.1	5.4-6.9	163	7.0	6.0-8.2	132	5.3	4.4-6.4	191	3.7	3.1-4.2	107	4.4	3.6-5.4	84	2.9	2.3-3.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	193	3.4	3.0-4.0	54	2.2	1.6-2.9	139	4.4	3.7-5.3	50	0.8	0.6-1.1	21	0.8	0.5-1.3	29	0.8	0.6-1.2
1977	223	3.9	3.4-4.5	62	2.4	1.8-3.1	161	5.2	4.4-6.2	47	0.8	0.6-1.0	25	0.9	0.6-1.4	22	0.6	0.4-0.9
1978	230	4.3	3.7-4.9	62	2.6	2.0-3.4	168	5.7	4.9-6.7	37	0.7	0.5-0.9	11	0.5	0.2-0.9	26	0.8	0.5-1.2
1979	203	3.9	3.3-4.5	79	3.4	2.6-4.2	124	4.4	3.6-5.3	33	0.6	0.4-0.8	15	0.6	0.4-1.1	18	0.5	0.3-0.8
1980	203	3.9	3.4-4.6	56	2.5	1.9-3.3	147	5.1	4.3-6.1	40	0.7	0.5-1.0	15	0.7	0.4-1.1	25	0.7	0.4-1.1
1981	191	3.6	3.1-4.2	56	2.4	1.8-3.1	135	4.7	3.9-5.7	38	0.7	0.5-0.9	12	0.5	0.2-0.9	26	0.8	0.5-1.1
1982	198	3.9	3.4-4.5	61	2.8	2.1-3.6	137	4.9	4.0-5.8	46	0.8	0.6-1.0	15	0.6	0.3-1.0	31	0.8	0.6-1.2
1983	233	4.7	4.1-5.4	58	2.5	1.9-3.3	175	6.7	5.7-7.8	27	0.5	0.3-0.7	7	0.3	0.1-0.6	20	0.6	0.4-1.0
1984	261	5.2	4.6-5.9	74	3.4	2.6-4.3	187	6.8	5.9-8.0	37	0.6	0.4-0.9	13	0.6	0.3-1.0	24	0.7	0.4-1.0
1985	263	5.1	4.5-5.8	79	3.4	2.7-4.3	184	6.5	5.5-7.5	44	0.8	0.6-1.1	25	1.2	0.7-1.7	19	0.6	0.3-0.9
1986	227	4.4	3.9-5.1	57	2.5	1.9-3.2	170	6.2	5.3-7.3	29	0.5	0.4-0.8	4	0.2	0.0-0.5	25	0.8	0.5-1.1
1987	210	4.3	3.7-4.9	65	3.0	2.3-3.9	145	5.4	4.5-6.4	30	0.5	0.4-0.8	10	0.5	0.2-0.9	20	0.5	0.3-0.9
1988	215	4.3	3.7-4.9	72	3.3	2.6-4.2	143	5.3	4.4-6.3	30	0.5	0.4-0.8	11	0.5	0.2-0.9	19	0.6	0.3-0.9
1989	188	4.0	3.4-4.6	51	2.3	1.7-3.0	137	5.6	4.6-6.6	28	0.5	0.3-0.8	7	0.3	0.1-0.7	21	0.6	0.4-1.0
1990	227	4.7	4.1-5.4	56	2.6	2.0-3.4	171	6.6	5.7-7.8	31	0.6	0.4-0.8	8	0.4	0.2-0.8	23	0.7	0.4-1.1
1991	247	5.3	4.6-6.0	65	3.0	2.3-3.9	182	7.4	6.3-8.6	30	0.6	0.4-0.8	8	0.4	0.2-0.8	22	0.7	0.4-1.1
1992	260	5.5	4.9-6.3	64	3.1	2.4-4.0	196	7.7	6.6-8.9	26	0.5	0.3-0.8	10	0.5	0.3-1.0	16	0.5	0.3-0.8
1993	283	6.0	5.4-6.8	77	3.7	2.9-4.6	206	8.3	7.2-9.5	26	0.5	0.3-0.7	9	0.4	0.2-0.8	17	0.5	0.3-0.8
1994	293	6.3	5.6-7.0	83	3.9	3.1-4.9	210	8.4	7.3-9.7	31	0.6	0.4-0.9	12	0.6	0.3-1.1	19	0.6	0.4-1.0
1995	311	6.5	5.8-7.3	89	4.2	3.3-5.1	222	8.7	7.6-10.0	23	0.4	0.3-0.7	7	0.3	0.1-0.7	16	0.5	0.3-0.8
1996	306	6.6	5.8-7.4	70	3.4	2.6-4.3	236	9.6	8.4-10.9	31	0.6	0.4-0.9	11	0.5	0.3-1.0	20	0.6	0.4-1.0
1997	347	7.3	6.6-8.1	93	4.3	3.5-5.3	254	10.1	8.9-11.4	30	0.6	0.4-0.9	9	0.4	0.2-0.8	21	0.7	0.4-1.1
1998	375	7.9	7.1-8.7	105	4.9	4.0-5.9	270	10.6	9.4-12.0	23	0.5	0.3-0.7	8	0.4	0.2-0.8	15	0.5	0.3-0.8
1999	415	8.7	7.9-9.6	97	4.3	3.5-5.3	318	12.7	11.3-14.2	22	0.4	0.3-0.7	10	0.5	0.2-0.9	12	0.4	0.2-0.7
2000	404	8.4	7.6-9.3	102	4.6	3.7-5.5	302	12.1	10.7-13.5	32	0.6	0.4-0.9	8	0.4	0.2-0.8	24	0.8	0.5-1.2
2001	471	9.8	9.0-10.8	123	5.7	4.7-6.8	348	13.8	12.4-15.4	26	0.5	0.3-0.7	10	0.5	0.2-0.9	16	0.5	0.3-0.8
2002	487	10.3	9.4-11.2	142	6.4	5.4-7.5	345	14.0	12.5-15.5	33	0.6	0.4-0.9	11	0.5	0.3-0.9	22	0.7	0.4-1.1
2003	465	9.7	8.9-10.7	111	5.0	4.1-6.0	354	14.1	12.7-15.7	23	0.4	0.3-0.7	9	0.4	0.2-0.8	14	0.4	0.2-0.7
2004	533	11.2	10.3-12.2	136	6.2	5.2-7.4	397	15.9	14.4-17.6	21	0.4	0.2-0.6	10	0.5	0.2-0.9	11	0.3	0.2-0.6
2005	720	15.3	14.2-16.4	182	8.3	7.1-9.6	538	21.9	20.1-23.9	37	0.7	0.5-1.0	16	0.8	0.4-1.3	21	0.7	0.4-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	699	14.8	13.7-16.0	174	7.7	6.6-9.0	525	21.5	19.7-23.5	41	0.8	0.6-1.1	13	0.7	0.4-1.1	28	1.0	0.6-1.4
2007	782	16.7	15.6-18.0	181	8.2	7.0-9.5	601	24.8	22.9-26.9	39	0.8	0.6-1.1	11	0.5	0.3-1.0	28	1.0	0.6-1.4
2008	941	19.8	18.6-21.2	227	10.1	8.8-11.5	714	29.3	27.1-31.5	26	0.5	0.3-0.8	14	0.7	0.4-1.2	12	0.4	0.2-0.7
2009	1,077	22.7	21.3-24.1	276	12.2	10.7-13.7	801	32.8	30.5-35.2	31	0.6	0.4-0.9	17	0.8	0.5-1.3	14	0.5	0.3-0.8
2010	1,024	21.6	20.3-23.0	248	11.1	9.8-12.6	776	31.9	29.6-34.2	31	0.6	0.4-0.8	14	0.6	0.3-1.1	17	0.5	0.3-0.9
2011	1,015	21.4	20.0-22.7	216	9.6	8.3-11.0	799	32.5	30.3-34.9	27	0.5	0.3-0.8	10	0.5	0.2-0.9	17	0.5	0.3-0.9
2012	1,107	23.1	21.8-24.6	277	12.0	10.6-13.5	830	33.9	31.5-36.3	26	0.5	0.3-0.8	13	0.6	0.3-1.0	13	0.5	0.2-0.8
2013	1,197	25.2	23.7-26.7	319	13.9	12.4-15.6	878	36.2	33.8-38.8	38	0.7	0.5-1.0	18	0.8	0.5-1.3	20	0.6	0.4-1.0
2014	1,163	24.1	22.7-25.5	292	12.6	11.2-14.2	871	35.3	33.0-37.8	34	0.6	0.4-0.9	12	0.5	0.3-0.9	22	0.7	0.4-1.1
2015	1,097	22.8	21.4-24.2	283	12.0	10.7-13.6	814	33.3	30.9-35.7	29	0.5	0.3-0.7	12	0.5	0.2-0.8	17	0.5	0.3-0.8
2016	1,109	23.0	21.7-24.5	263	11.2	9.8-12.6	846	34.7	32.3-37.2	32	0.6	0.4-0.8	12	0.5	0.3-1.0	20	0.6	0.4-1.0
2017	1,060	21.8	20.5-23.2	274	11.5	10.2-13.0	786	31.9	29.6-34.3	28	0.5	0.3-0.8	7	0.3	0.1-0.7	21	0.7	0.4-1.0
2018	953	20.0	18.7-21.3	242	10.4	9.1-11.8	711	29.5	27.3-31.8	26	0.4	0.3-0.6	4	0.2	0.0-0.4	22	0.7	0.4-1.0
2019	1,008	21.1	19.8-22.5	273	11.6	10.2-13.1	735	30.5	28.3-32.9	30	0.5	0.3-0.7	15	0.6	0.3-1.0	15	0.4	0.2-0.7
2020	802	16.9	15.7-18.2	234	10.3	9.0-11.7	568	23.5	21.5-25.5	25	0.5	0.3-0.7	12	0.5	0.3-0.9	13	0.4	0.2-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	218	3.7	3.2-4.2	116	4.3	3.6-5.2	102	3.2	2.6-3.9	78	1.4	1.1-1.8	51	2.0	1.5-2.7	27	0.8	0.5-1.3
1977	208	3.6	3.1-4.2	119	4.5	3.7-5.4	89	3.0	2.3-3.7	88	1.6	1.3-2.0	49	1.9	1.4-2.5	39	1.3	0.9-1.8
1978	213	3.7	3.2-4.2	112	4.2	3.4-5.1	101	3.3	2.6-4.0	78	1.4	1.1-1.7	43	1.7	1.2-2.3	35	1.1	0.8-1.6
1979	198	3.8	3.3-4.4	100	4.3	3.5-5.2	98	3.4	2.7-4.2	62	1.1	0.9-1.5	37	1.5	1.1-2.1	25	0.9	0.5-1.3
1980	199	3.7	3.2-4.3	103	4.4	3.5-5.3	96	3.2	2.5-3.9	72	1.3	1.0-1.6	35	1.6	1.1-2.2	37	1.1	0.8-1.5
1981	185	3.5	3.0-4.0	101	4.2	3.4-5.1	84	2.8	2.2-3.6	64	1.2	1.0-1.6	33	1.5	1.0-2.1	31	1.0	0.7-1.5
1982	210	4.0	3.4-4.6	120	5.1	4.2-6.1	90	3.1	2.5-3.9	51	0.9	0.6-1.2	26	1.0	0.6-1.5	25	0.8	0.5-1.2
1983	192	3.7	3.2-4.3	113	4.9	4.0-5.9	79	2.7	2.1-3.4	62	1.1	0.8-1.4	36	1.5	1.0-2.1	26	0.8	0.5-1.2
1984	187	3.6	3.1-4.2	102	4.3	3.5-5.3	85	3.1	2.4-3.8	63	1.2	0.9-1.6	34	1.5	1.0-2.1	29	0.9	0.6-1.4
1985	211	4.1	3.5-4.7	128	5.4	4.5-6.5	83	2.9	2.3-3.6	58	1.1	0.8-1.4	30	1.2	0.8-1.7	28	1.0	0.6-1.4
1986	210	4.1	3.5-4.7	130	5.5	4.6-6.6	80	2.9	2.2-3.6	56	1.0	0.8-1.4	38	1.6	1.1-2.2	18	0.6	0.3-1.0
1987	174	3.5	3.0-4.1	89	3.9	3.1-4.9	85	3.2	2.5-4.0	40	0.8	0.5-1.1	24	1.0	0.7-1.6	16	0.5	0.3-0.9
1988	198	4.0	3.4-4.6	112	4.9	4.0-5.9	86	3.1	2.5-3.9	39	0.7	0.5-1.0	22	1.0	0.6-1.6	17	0.6	0.3-0.9
1989	181	3.8	3.2-4.4	105	4.6	3.7-5.6	76	3.0	2.4-3.8	56	1.2	0.9-1.5	32	1.5	1.0-2.1	24	0.9	0.6-1.3
1990	164	3.4	2.9-4.0	97	4.4	3.6-5.4	67	2.5	1.9-3.2	55	1.1	0.8-1.4	38	1.7	1.2-2.4	17	0.6	0.3-0.9
1991	194	4.2	3.6-4.8	102	4.7	3.8-5.8	92	3.8	3.0-4.7	43	0.9	0.6-1.2	25	1.2	0.8-1.8	18	0.6	0.4-1.0
1992	183	3.7	3.2-4.3	109	4.8	3.9-5.8	74	2.8	2.2-3.5	39	0.8	0.6-1.1	22	0.9	0.6-1.4	17	0.7	0.4-1.1
1993	157	3.3	2.8-3.9	82	3.9	3.1-4.8	75	2.8	2.2-3.6	34	0.7	0.5-0.9	19	0.8	0.5-1.3	15	0.5	0.3-0.8
1994	175	3.7	3.2-4.4	103	4.8	3.9-5.8	72	2.9	2.2-3.6	41	0.8	0.6-1.1	18	0.9	0.5-1.4	23	0.8	0.5-1.3
1995	182	4.0	3.4-4.6	94	4.3	3.4-5.2	88	3.7	2.9-4.6	31	0.7	0.5-1.0	17	0.8	0.5-1.3	14	0.6	0.3-1.0
1996	144	3.1	2.6-3.6	76	3.5	2.7-4.4	68	2.7	2.1-3.5	31	0.6	0.4-0.9	16	0.7	0.4-1.2	15	0.5	0.3-0.8
1997	162	3.5	3.0-4.1	94	4.3	3.5-5.3	68	2.9	2.2-3.7	37	0.7	0.5-1.0	18	0.8	0.5-1.3	19	0.7	0.4-1.1
1998	183	3.9	3.3-4.5	97	4.3	3.5-5.3	86	3.5	2.8-4.3	35	0.7	0.5-1.0	17	0.8	0.4-1.2	18	0.7	0.4-1.1
1999	163	3.4	2.9-4.0	78	3.4	2.6-4.2	85	3.4	2.7-4.2	21	0.4	0.3-0.7	10	0.5	0.2-0.9	11	0.4	0.2-0.7
2000	206	4.4	3.8-5.0	108	4.8	3.9-5.8	98	4.1	3.3-5.0	36	0.7	0.5-1.0	19	0.9	0.5-1.4	17	0.6	0.4-1.0
2001	180	3.7	3.2-4.3	91	3.9	3.1-4.8	89	3.5	2.8-4.3	24	0.5	0.3-0.7	13	0.6	0.3-1.0	11	0.4	0.2-0.7
2002	201	4.2	3.6-4.8	106	4.6	3.8-5.6	95	3.8	3.0-4.6	29	0.6	0.4-0.9	21	1.0	0.6-1.5	8	0.3	0.1-0.6
2003	183	4.0	3.4-4.6	93	4.2	3.4-5.2	90	3.8	3.1-4.7	24	0.5	0.3-0.7	11	0.5	0.3-0.9	13	0.5	0.3-0.9
2004	174	3.7	3.1-4.3	91	4.0	3.2-4.9	83	3.3	2.7-4.2	23	0.5	0.3-0.7	13	0.6	0.3-1.0	10	0.4	0.2-0.7
2005	182	3.9	3.3-4.5	105	4.8	3.9-5.8	77	3.0	2.4-3.8	23	0.5	0.3-0.7	14	0.7	0.4-1.1	9	0.3	0.1-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	188	4.1	3.5-4.7	106	4.9	4.0-5.9	82	3.3	2.6-4.1	15	0.3	0.2-0.5	12	0.6	0.3-1.0	3	0.1	0.0-0.4
2007	192	4.0	3.5-4.7	84	3.7	3.0-4.7	108	4.3	3.5-5.2	32	0.7	0.4-0.9	17	0.8	0.5-1.3	15	0.5	0.3-0.9
2008	175	3.7	3.2-4.3	83	3.8	3.0-4.7	92	3.7	3.0-4.6	33	0.6	0.4-0.9	22	1.0	0.6-1.5	11	0.4	0.2-0.7
2009	179	3.9	3.3-4.5	104	4.8	3.9-5.8	75	3.1	2.4-3.9	23	0.5	0.3-0.7	15	0.7	0.4-1.2	8	0.3	0.1-0.6
2010	179	3.7	3.2-4.4	93	4.0	3.2-5.0	86	3.5	2.8-4.3	22	0.4	0.3-0.7	15	0.6	0.3-1.0	7	0.3	0.1-0.6
2011	183	3.9	3.3-4.5	100	4.3	3.5-5.3	83	3.5	2.7-4.3	32	0.7	0.4-0.9	17	0.8	0.5-1.3	15	0.5	0.3-0.9
2012	161	3.5	3.0-4.1	81	3.6	2.9-4.5	80	3.5	2.7-4.3	20	0.4	0.2-0.6	11	0.5	0.2-0.9	9	0.3	0.1-0.6
2013	179	3.7	3.2-4.3	77	3.4	2.7-4.3	102	3.9	3.2-4.8	22	0.4	0.3-0.7	13	0.6	0.3-1.0	9	0.3	0.1-0.7
2014	176	3.8	3.2-4.4	96	4.3	3.4-5.2	80	3.3	2.6-4.1	25	0.5	0.3-0.8	10	0.5	0.2-0.8	15	0.5	0.3-0.9
2015	181	3.8	3.3-4.5	105	4.6	3.7-5.6	76	3.1	2.4-3.9	24	0.5	0.3-0.7	13	0.6	0.3-1.0	11	0.4	0.2-0.7
2016	186	3.9	3.4-4.6	95	4.2	3.4-5.2	91	3.7	3.0-4.6	22	0.4	0.2-0.6	13	0.5	0.3-0.9	9	0.3	0.1-0.6
2017	153	3.2	2.7-3.8	90	3.8	3.1-4.7	63	2.6	2.0-3.4	13	0.3	0.1-0.5	6	0.3	0.1-0.6	7	0.3	0.1-0.6
2018	124	2.6	2.1-3.1	71	3.2	2.5-4.0	53	2.0	1.5-2.7	13	0.2	0.1-0.4	12	0.5	0.3-0.9	1	0.0	0.0-0.2
2019	141	3.1	2.6-3.6	76	3.4	2.7-4.3	65	2.7	2.0-3.4	11	0.2	0.1-0.4	8	0.3	0.1-0.6	3	0.1	0.0-0.4
2020	143	3.1	2.6-3.7	81	3.5	2.8-4.4	62	2.8	2.1-3.6	17	0.3	0.2-0.5	8	0.3	0.1-0.6	9	0.3	0.1-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	733	12.0	11.1-12.9	370	14.7	13.2-16.3	363	10.3	9.2-11.4	398	6.5	5.9-7.2	217	8.9	7.7-10.2	181	5.0	4.3-5.8
1977	695	11.6	10.7-12.5	350	13.7	12.3-15.3	345	10.0	9.0-11.2	354	5.8	5.2-6.5	169	6.8	5.7-7.9	185	5.2	4.4-6.0
1978	699	12.0	11.1-13.0	351	14.2	12.7-15.8	348	10.3	9.2-11.5	371	6.3	5.6-6.9	198	8.1	6.9-9.3	173	5.0	4.3-5.8
1979	746	12.9	12.0-13.9	347	14.8	13.3-16.5	399	11.6	10.4-12.8	364	6.4	5.7-7.1	181	7.9	6.7-9.2	183	5.3	4.5-6.2
1980	814	14.3	13.3-15.3	395	16.6	15.0-18.4	419	12.4	11.2-13.7	373	6.4	5.7-7.1	189	7.8	6.7-9.0	184	5.3	4.6-6.2
1981	803	14.1	13.1-15.2	381	16.2	14.6-18.0	422	12.5	11.3-13.8	375	6.5	5.8-7.2	172	7.5	6.4-8.8	203	5.8	5.0-6.7
1982	875	15.7	14.6-16.8	402	17.4	15.7-19.2	473	14.2	12.9-15.6	404	7.2	6.5-8.0	190	8.5	7.3-9.8	214	6.3	5.5-7.2
1983	870	16.1	15.0-17.2	447	20.3	18.4-22.4	423	12.7	11.5-14.1	434	7.9	7.1-8.7	214	9.7	8.4-11.1	220	6.5	5.6-7.4
1984	880	16.0	14.9-17.1	447	19.9	18.1-21.9	433	13.0	11.8-14.4	376	6.8	6.1-7.5	194	8.8	7.5-10.1	182	5.2	4.5-6.1
1985	972	18.0	16.9-19.2	522	23.8	21.8-26.0	450	13.6	12.3-15.0	445	8.0	7.3-8.8	240	10.9	9.5-12.4	205	6.0	5.2-6.9
1986	1,028	19.3	18.1-20.6	542	24.2	22.2-26.4	486	15.0	13.7-16.5	393	7.3	6.6-8.1	216	10.0	8.7-11.4	177	5.2	4.5-6.1
1987	1,038	20.1	18.8-21.3	561	26.1	24.0-28.5	477	15.0	13.6-16.4	411	7.6	6.9-8.4	214	9.8	8.5-11.3	197	5.8	5.0-6.7
1988	956	18.5	17.3-19.7	516	23.8	21.7-26.0	440	14.0	12.7-15.5	447	8.3	7.6-9.2	218	10.3	8.9-11.8	229	6.7	5.8-7.7
1989	987	20.0	18.7-21.3	578	27.5	25.2-29.8	409	13.5	12.2-15.0	420	8.1	7.3-8.9	238	11.4	10.0-13.0	182	5.7	4.9-6.6
1990	1,105	22.1	20.8-23.4	635	30.0	27.6-32.4	470	15.4	14.0-16.9	428	8.2	7.5-9.1	214	10.2	8.8-11.6	214	6.7	5.8-7.7
1991	1,231	24.8	23.4-26.3	704	33.7	31.2-36.3	527	17.6	16.1-19.2	414	8.3	7.5-9.1	222	10.8	9.4-12.4	192	6.0	5.2-7.0
1992	1,129	23.0	21.6-24.4	630	29.9	27.6-32.4	499	17.1	15.5-18.7	424	8.2	7.5-9.1	217	10.6	9.2-12.1	207	6.6	5.7-7.6
1993	1,113	22.9	21.6-24.3	650	31.2	28.8-33.7	463	15.9	14.5-17.5	399	7.8	7.1-8.7	213	10.5	9.1-12.0	186	6.0	5.2-7.0
1994	1,308	27.2	25.8-28.8	762	36.3	33.8-39.0	546	19.2	17.6-21.0	458	9.2	8.3-10.1	236	11.6	10.1-13.2	222	7.2	6.3-8.3
1995	1,234	25.6	24.2-27.1	694	33.4	30.9-36.0	540	18.9	17.3-20.6	429	8.6	7.8-9.5	223	11.1	9.7-12.6	206	6.7	5.8-7.8
1996	1,200	24.9	23.5-26.4	668	32.4	30.0-35.0	532	19.0	17.3-20.7	436	8.8	7.9-9.6	233	11.8	10.3-13.4	203	6.7	5.8-7.8
1997	1,148	23.7	22.3-25.1	603	29.3	27.0-31.8	545	18.9	17.3-20.6	445	8.9	8.1-9.8	229	11.4	10.0-13.0	216	7.1	6.2-8.2
1998	1,074	22.2	20.9-23.5	581	27.8	25.6-30.1	493	17.6	16.0-19.3	430	8.6	7.8-9.5	204	10.0	8.7-11.5	226	7.5	6.5-8.5
1999	1,096	22.4	21.1-23.8	563	26.9	24.7-29.3	533	18.8	17.2-20.5	400	8.0	7.2-8.8	194	9.6	8.3-11.1	206	6.8	5.9-7.8
2000	1,090	22.2	20.9-23.5	555	26.2	24.0-28.4	535	18.8	17.2-20.5	397	7.8	7.1-8.7	197	9.6	8.3-11.1	200	6.6	5.7-7.6
2001	1,059	21.4	20.1-22.7	571	27.1	24.9-29.4	488	17.1	15.6-18.7	381	7.4	6.7-8.2	214	10.4	9.0-11.9	167	5.2	4.4-6.1
2002	1,050	21.3	20.0-22.6	585	27.5	25.3-29.8	465	16.2	14.8-17.8	338	6.6	5.9-7.4	176	8.5	7.3-9.8	162	5.3	4.5-6.2
2003	1,154	23.3	22.0-24.7	594	27.9	25.7-30.3	560	19.7	18.0-21.4	326	6.4	5.7-7.1	166	7.9	6.8-9.2	160	5.1	4.3-6.0
2004	1,128	22.8	21.5-24.2	585	27.6	25.4-29.9	543	19.1	17.5-20.8	354	6.9	6.2-7.7	178	8.6	7.4-10.0	176	5.7	4.9-6.7
2005	1,165	23.7	22.3-25.1	582	27.4	25.2-29.7	583	20.8	19.1-22.6	350	6.9	6.2-7.7	179	8.7	7.4-10.0	171	5.6	4.8-6.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,170	24.0	22.6-25.4	648	30.5	28.2-33.0	522	18.6	17.0-20.3	358	7.1	6.4-7.9	187	9.1	7.8-10.5	171	5.6	4.8-6.5
2007	1,116	23.0	21.6-24.4	580	27.5	25.3-29.8	536	19.4	17.8-21.2	338	6.8	6.1-7.6	180	8.7	7.4-10.1	158	5.4	4.6-6.4
2008	1,179	24.0	22.6-25.4	635	29.6	27.4-32.1	544	19.6	18.0-21.4	313	6.1	5.5-6.8	178	8.5	7.3-9.8	135	4.4	3.6-5.2
2009	1,226	25.0	23.6-26.5	617	29.1	26.8-31.5	609	21.8	20.1-23.7	314	6.1	5.5-6.9	172	8.1	6.9-9.4	142	4.6	3.9-5.5
2010	1,181	23.8	22.5-25.3	624	28.9	26.6-31.3	557	19.7	18.1-21.5	290	5.7	5.1-6.4	150	7.3	6.1-8.5	140	4.6	3.8-5.4
2011	1,182	23.7	22.3-25.1	600	27.2	25.1-29.5	582	20.7	19.0-22.5	302	5.8	5.1-6.5	154	7.1	6.0-8.4	148	4.8	4.0-5.7
2012	1,183	23.6	22.2-25.0	652	29.5	27.3-31.9	531	19.0	17.4-20.7	293	5.5	4.9-6.2	162	7.5	6.4-8.7	131	4.0	3.3-4.8
2013	1,220	24.0	22.6-25.4	661	29.6	27.3-31.9	559	19.4	17.8-21.2	276	5.2	4.6-5.8	147	6.7	5.6-7.9	129	4.0	3.3-4.8
2014	1,195	23.2	21.8-24.5	643	28.5	26.3-30.8	552	19.0	17.4-20.7	298	5.5	4.9-6.2	159	7.1	6.0-8.3	139	4.2	3.5-5.0
2015	1,172	22.6	21.3-23.9	632	27.2	25.1-29.5	540	18.7	17.1-20.5	255	4.8	4.2-5.4	152	6.8	5.7-8.0	103	3.3	2.7-4.1
2016	1,189	22.5	21.2-23.8	637	27.1	25.0-29.3	552	18.8	17.2-20.5	295	5.2	4.6-5.9	160	6.8	5.8-8.0	135	4.0	3.3-4.8
2017	1,226	23.3	22.0-24.6	643	27.4	25.3-29.6	583	19.9	18.3-21.6	296	5.3	4.7-6.0	152	6.5	5.5-7.7	144	4.4	3.7-5.2
2018	1,184	22.4	21.1-23.8	618	26.2	24.2-28.4	566	19.4	17.8-21.2	242	4.3	3.7-4.8	136	5.8	4.8-6.8	106	3.1	2.5-3.8
2019	1,247	23.2	21.9-24.5	678	28.2	26.1-30.4	569	19.2	17.6-20.9	260	4.5	4.0-5.2	154	6.5	5.5-7.6	106	3.1	2.5-3.8
2020	1,008	18.9	17.7-20.1	525	21.9	20.1-23.9	483	16.4	14.9-18.0	248	4.3	3.8-4.9	138	5.7	4.8-6.7	110	3.2	2.6-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	243	3.9	3.4-4.5	123	4.9	4.1-5.9	120	3.3	2.7-4.0	151	2.5	2.1-3.0	75	3.2	2.5-4.1	76	2.1	1.6-2.6
1977	259	4.2	3.7-4.8	124	5.1	4.2-6.1	135	3.6	3.0-4.3	162	2.5	2.2-3.0	81	3.2	2.5-4.0	81	2.1	1.7-2.6
1978	253	4.2	3.7-4.7	123	5.1	4.2-6.1	130	3.5	2.9-4.2	149	2.4	2.0-2.9	66	2.7	2.0-3.4	83	2.2	1.8-2.8
1979	266	4.6	4.0-5.2	139	5.9	5.0-7.1	127	3.6	3.0-4.3	173	2.9	2.5-3.4	71	2.9	2.2-3.7	102	2.8	2.3-3.5
1980	241	4.2	3.7-4.8	114	5.1	4.2-6.1	127	3.6	3.0-4.3	167	2.8	2.4-3.3	78	3.3	2.6-4.2	89	2.5	2.0-3.1
1981	302	5.1	4.5-5.7	147	6.3	5.3-7.5	155	4.3	3.6-5.1	169	2.9	2.5-3.4	86	3.9	3.1-4.8	83	2.3	1.8-2.9
1982	268	4.6	4.1-5.2	123	5.4	4.5-6.5	145	4.2	3.5-4.9	159	2.7	2.3-3.2	70	3.0	2.3-3.9	89	2.6	2.1-3.2
1983	276	4.8	4.3-5.5	140	6.3	5.3-7.5	136	3.9	3.3-4.7	160	2.8	2.3-3.2	68	3.0	2.3-3.8	92	2.5	2.0-3.1
1984	233	4.1	3.6-4.7	111	5.2	4.2-6.3	122	3.4	2.8-4.2	157	2.7	2.3-3.2	75	3.4	2.6-4.2	82	2.3	1.8-2.9
1985	254	4.5	3.9-5.1	137	6.1	5.1-7.3	117	3.4	2.8-4.1	145	2.5	2.1-2.9	82	3.8	3.0-4.7	63	1.8	1.3-2.3
1986	255	4.5	4.0-5.1	129	5.8	4.8-6.9	126	3.6	3.0-4.4	144	2.5	2.1-3.0	68	3.0	2.3-3.9	76	2.2	1.7-2.8
1987	248	4.4	3.9-5.0	137	6.4	5.3-7.6	111	3.2	2.6-3.9	154	2.8	2.4-3.3	88	4.1	3.3-5.1	66	1.9	1.4-2.4
1988	247	4.5	4.0-5.1	128	6.2	5.1-7.4	119	3.5	2.9-4.3	141	2.5	2.1-2.9	74	3.5	2.7-4.4	67	1.9	1.4-2.4
1989	252	4.8	4.2-5.5	109	5.2	4.2-6.2	143	4.6	3.8-5.5	150	2.7	2.3-3.2	72	3.3	2.6-4.2	78	2.3	1.8-2.9
1990	266	5.0	4.4-5.7	132	6.2	5.1-7.3	134	4.3	3.6-5.1	139	2.5	2.1-3.0	69	3.3	2.5-4.2	70	2.1	1.6-2.7
1991	240	4.6	4.0-5.2	112	5.3	4.4-6.4	128	4.2	3.5-5.0	127	2.4	2.0-2.8	59	2.8	2.2-3.7	68	2.1	1.6-2.7
1992	273	5.2	4.6-5.9	147	7.2	6.1-8.5	126	4.0	3.3-4.8	149	2.8	2.4-3.3	74	3.7	2.9-4.6	75	2.4	1.9-3.0
1993	259	5.1	4.5-5.8	119	6.1	5.0-7.3	140	4.6	3.8-5.4	177	3.4	2.9-3.9	90	4.4	3.6-5.5	87	2.7	2.2-3.4
1994	238	4.8	4.2-5.4	113	5.5	4.5-6.6	125	4.3	3.5-5.1	154	3.0	2.6-3.5	73	3.5	2.8-4.5	81	2.7	2.1-3.3
1995	248	5.0	4.4-5.7	119	5.9	4.9-7.0	129	4.4	3.7-5.3	149	2.9	2.5-3.5	71	3.6	2.8-4.5	78	2.5	2.0-3.2
1996	286	5.7	5.1-6.5	139	7.0	5.9-8.3	147	5.0	4.2-5.9	151	3.0	2.5-3.5	63	3.2	2.4-4.1	88	2.9	2.3-3.6
1997	239	4.8	4.2-5.5	120	6.0	5.0-7.2	119	4.0	3.3-4.8	133	2.7	2.2-3.2	66	3.3	2.6-4.2	67	2.3	1.7-2.9
1998	293	5.9	5.2-6.6	145	7.1	6.0-8.4	148	5.1	4.3-6.0	139	2.8	2.3-3.3	69	3.4	2.6-4.3	70	2.3	1.8-2.9
1999	280	5.6	5.0-6.3	119	5.8	4.8-7.0	161	5.5	4.7-6.5	138	2.7	2.3-3.2	63	3.1	2.4-4.0	75	2.5	1.9-3.1
2000	278	5.5	4.9-6.2	130	6.3	5.3-7.5	148	5.1	4.3-6.0	168	3.3	2.8-3.8	72	3.5	2.7-4.4	96	3.1	2.5-3.8
2001	301	6.0	5.3-6.7	154	7.5	6.4-8.8	147	5.0	4.2-5.8	136	2.6	2.2-3.1	70	3.4	2.7-4.3	66	2.1	1.6-2.7
2002	307	6.1	5.4-6.8	161	7.8	6.6-9.1	146	5.0	4.2-5.9	143	2.8	2.3-3.3	71	3.5	2.8-4.5	72	2.4	1.8-3.0
2003	323	6.4	5.7-7.1	170	8.2	7.0-9.5	153	5.3	4.5-6.3	133	2.5	2.1-3.0	59	2.9	2.2-3.7	74	2.3	1.8-2.9
2004	309	6.2	5.5-6.9	155	7.4	6.3-8.6	154	5.3	4.5-6.3	121	2.3	1.9-2.8	56	2.8	2.1-3.6	65	2.1	1.6-2.7
2005	338	6.7	6.0-7.5	174	8.3	7.1-9.6	164	5.6	4.8-6.6	136	2.7	2.2-3.1	66	3.2	2.5-4.1	70	2.3	1.7-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	334	6.8	6.1-7.6	162	7.8	6.7-9.1	172	6.1	5.2-7.1	126	2.5	2.1-3.0	82	4.0	3.2-5.0	44	1.5	1.1-2.1
2007	317	6.4	5.7-7.1	174	8.3	7.1-9.7	143	4.9	4.2-5.9	123	2.4	2.0-2.9	68	3.3	2.6-4.2	55	1.8	1.3-2.3
2008	342	6.9	6.2-7.7	199	9.5	8.2-11.0	143	5.0	4.2-6.0	138	2.7	2.3-3.2	68	3.3	2.6-4.2	70	2.4	1.8-3.0
2009	382	7.6	6.9-8.4	209	9.9	8.6-11.4	173	5.9	5.1-6.9	117	2.3	1.9-2.7	66	3.2	2.5-4.0	51	1.6	1.2-2.2
2010	374	7.4	6.7-8.2	203	9.4	8.1-10.8	171	6.0	5.1-7.0	148	2.9	2.4-3.4	82	4.0	3.1-4.9	66	2.2	1.7-2.8
2011	407	8.0	7.2-8.9	234	10.7	9.4-12.2	173	6.1	5.2-7.1	118	2.3	1.9-2.7	53	2.5	1.9-3.3	65	2.2	1.7-2.8
2012	383	7.4	6.7-8.2	223	10.0	8.8-11.5	160	5.5	4.6-6.4	123	2.4	1.9-2.8	58	2.7	2.1-3.5	65	2.1	1.6-2.7
2013	404	7.6	6.9-8.4	204	9.0	7.8-10.4	200	6.7	5.7-7.7	147	2.7	2.3-3.2	85	3.8	3.0-4.7	62	1.9	1.5-2.5
2014	450	8.6	7.8-9.4	265	11.6	10.2-13.1	185	6.3	5.4-7.3	146	2.7	2.3-3.2	71	3.1	2.4-4.0	75	2.3	1.8-3.0
2015	411	7.7	7.0-8.5	214	9.3	8.1-10.7	197	6.5	5.6-7.5	135	2.5	2.1-3.0	77	3.4	2.7-4.2	58	1.8	1.4-2.4
2016	413	7.6	6.9-8.4	260	10.9	9.6-12.4	153	5.0	4.3-5.9	159	2.8	2.4-3.3	91	3.9	3.1-4.8	68	2.1	1.6-2.7
2017	444	8.1	7.3-8.9	238	9.9	8.6-11.2	206	6.7	5.8-7.7	136	2.4	2.0-2.9	71	3.1	2.4-3.9	65	2.0	1.5-2.5
2018	381	6.9	6.2-7.6	204	8.4	7.3-9.7	177	5.7	4.9-6.7	113	1.9	1.6-2.3	63	2.5	1.9-3.2	50	1.5	1.1-2.0
2019	408	7.4	6.7-8.2	218	8.9	7.7-10.2	190	6.3	5.4-7.3	136	2.3	1.9-2.7	68	2.8	2.1-3.5	68	2.0	1.5-2.5
2020	331	6.0	5.4-6.7	179	7.3	6.3-8.5	152	5.1	4.3-6.0	125	2.1	1.7-2.5	79	3.2	2.5-4.0	46	1.3	0.9-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	689	11.9	11.0-12.9	383	15.8	14.2-17.6	306	9.1	8.0-10.2	500	8.7	8.0-9.6	263	11.2	9.8-12.7	237	7.0	6.1-8.0
1977	640	11.2	10.3-12.1	348	14.5	12.9-16.2	292	8.8	7.7-9.9	458	8.0	7.3-8.8	254	10.8	9.5-12.3	204	6.1	5.3-7.1
1978	686	12.1	11.2-13.1	394	16.8	15.1-18.7	292	8.9	7.8-10.0	463	8.1	7.4-8.9	250	10.7	9.4-12.1	213	6.4	5.6-7.4
1979	646	11.5	10.6-12.5	329	13.8	12.3-15.5	317	9.9	8.8-11.1	467	8.3	7.6-9.2	238	10.2	8.9-11.7	229	7.2	6.2-8.2
1980	741	13.4	12.5-14.5	379	16.6	14.9-18.4	362	11.4	10.2-12.7	436	7.8	7.1-8.6	252	11.2	9.8-12.7	184	5.6	4.7-6.5
1981	655	11.9	11.0-12.9	362	16.0	14.3-17.7	293	9.1	8.0-10.2	450	8.2	7.4-9.0	245	10.8	9.5-12.3	205	6.4	5.5-7.4
1982	692	12.6	11.7-13.6	383	17.1	15.4-18.9	309	9.5	8.4-10.7	474	8.7	7.9-9.6	264	12.0	10.5-13.5	210	6.5	5.6-7.5
1983	648	12.2	11.3-13.2	328	14.8	13.2-16.5	320	10.7	9.5-12.1	419	7.8	7.1-8.6	224	10.4	9.0-11.9	195	6.1	5.2-7.1
1984	688	12.7	11.8-13.8	371	17.3	15.5-19.2	317	9.9	8.8-11.1	458	8.4	7.6-9.2	241	11.2	9.8-12.8	217	6.5	5.6-7.5
1985	650	11.9	11.0-12.9	353	16.1	14.4-17.9	297	9.2	8.1-10.4	407	7.4	6.7-8.2	204	9.4	8.1-10.8	203	6.1	5.3-7.1
1986	699	13.2	12.2-14.3	354	15.9	14.3-17.7	345	11.5	10.2-12.8	405	7.5	6.8-8.3	202	9.2	7.9-10.6	203	6.3	5.5-7.3
1987	560	10.6	9.7-11.5	266	12.5	11.0-14.1	294	9.3	8.3-10.5	358	6.6	5.9-7.3	172	8.0	6.8-9.3	186	5.9	5.0-6.8
1988	599	11.6	10.7-12.6	313	15.0	13.4-16.8	286	9.4	8.3-10.6	353	6.7	6.0-7.5	192	9.2	7.9-10.6	161	5.1	4.3-6.0
1989	679	13.1	12.1-14.2	362	17.3	15.6-19.3	317	10.4	9.2-11.6	366	6.9	6.2-7.7	192	9.2	8.0-10.7	174	5.4	4.6-6.3
1990	690	13.7	12.7-14.8	361	17.3	15.5-19.2	329	11.3	10.0-12.6	361	7.0	6.3-7.8	181	9.0	7.7-10.4	180	5.9	5.0-6.9
1991	627	12.8	11.8-13.8	350	17.1	15.3-19.0	277	9.5	8.3-10.7	355	7.0	6.3-7.8	208	10.4	9.0-12.0	147	4.8	4.0-5.7
1992	642	12.9	11.9-14.0	352	17.3	15.5-19.2	290	10.0	8.8-11.2	324	6.4	5.7-7.2	178	9.0	7.7-10.4	146	4.8	4.0-5.7
1993	601	12.3	11.3-13.3	293	14.3	12.7-16.1	308	11.0	9.8-12.4	336	6.6	5.9-7.4	190	9.3	8.0-10.7	146	4.9	4.2-5.9
1994	698	14.1	13.1-15.2	357	17.7	15.9-19.7	341	11.5	10.3-12.9	387	7.6	6.9-8.4	197	9.7	8.4-11.2	190	6.1	5.2-7.0
1995	619	12.7	11.7-13.8	318	15.7	14.0-17.6	301	10.6	9.4-11.9	378	7.6	6.8-8.4	199	10.0	8.7-11.5	179	6.0	5.1-6.9
1996	628	12.9	11.9-14.0	349	17.2	15.4-19.1	279	9.9	8.7-11.2	340	6.8	6.1-7.5	165	8.1	6.9-9.5	175	5.9	5.1-6.9
1997	668	13.7	12.7-14.8	355	17.6	15.8-19.5	313	11.1	9.9-12.5	346	7.0	6.3-7.8	199	10.0	8.7-11.5	147	4.9	4.1-5.8
1998	657	13.4	12.4-14.5	363	17.6	15.8-19.5	294	10.3	9.1-11.6	362	7.3	6.6-8.1	205	10.0	8.7-11.5	157	5.3	4.5-6.3
1999	736	14.9	13.9-16.1	385	18.7	16.9-20.7	351	12.2	10.9-13.6	370	7.3	6.6-8.1	203	9.9	8.6-11.4	167	5.6	4.7-6.5
2000	674	13.7	12.6-14.7	343	16.6	14.9-18.5	331	11.6	10.3-12.9	329	6.5	5.8-7.3	149	7.2	6.1-8.4	180	6.0	5.1-7.0
2001	703	14.2	13.1-15.3	358	17.3	15.5-19.2	345	12.0	10.7-13.4	344	6.7	6.0-7.5	176	8.5	7.3-9.9	168	5.4	4.6-6.3
2002	637	12.8	11.8-13.9	360	17.2	15.4-19.0	277	9.6	8.5-10.9	342	6.7	6.0-7.4	178	8.6	7.4-10.0	164	5.4	4.6-6.3
2003	725	14.7	13.6-15.8	373	17.8	16.0-19.7	352	12.4	11.1-13.9	309	6.1	5.4-6.8	162	7.8	6.6-9.1	147	4.9	4.2-5.9
2004	694	13.9	12.9-15.0	366	17.6	15.8-19.5	328	11.4	10.2-12.8	354	6.9	6.2-7.7	185	9.0	7.8-10.4	169	5.6	4.8-6.5
2005	722	14.7	13.7-15.9	407	19.3	17.5-21.3	315	11.3	10.1-12.7	345	6.8	6.1-7.6	193	9.3	8.0-10.7	152	5.1	4.3-6.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
White Individuals
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	698	14.4	13.4-15.6	381	18.4	16.6-20.4	317	11.5	10.2-12.9	344	6.8	6.1-7.6	177	8.6	7.4-10.0	167	5.6	4.8-6.6
2007	742	15.5	14.4-16.7	414	19.8	17.9-21.8	328	12.3	10.9-13.7	325	6.5	5.8-7.3	178	8.7	7.4-10.1	147	5.3	4.4-6.2
2008	739	15.4	14.3-16.5	409	19.8	17.9-21.8	330	12.2	10.9-13.6	347	7.0	6.3-7.8	171	8.3	7.1-9.7	176	6.2	5.3-7.2
2009	809	16.6	15.5-17.8	434	20.6	18.7-22.7	375	13.5	12.1-15.0	316	6.3	5.6-7.1	143	6.9	5.8-8.1	173	5.9	5.0-6.9
2010	836	17.0	15.9-18.3	470	22.1	20.1-24.2	366	13.1	11.7-14.6	327	6.4	5.7-7.2	178	8.5	7.3-9.9	149	5.0	4.2-6.0
2011	853	16.9	15.8-18.1	467	21.4	19.5-23.5	386	13.7	12.3-15.2	340	6.5	5.8-7.3	197	9.3	8.0-10.7	143	4.5	3.8-5.4
2012	792	15.7	14.6-16.8	460	20.8	19.0-22.9	332	11.9	10.6-13.3	311	5.9	5.3-6.7	176	8.2	7.0-9.5	135	4.3	3.6-5.2
2013	838	16.5	15.4-17.7	487	22.1	20.2-24.2	351	12.3	11.0-13.7	342	6.6	5.9-7.3	196	9.0	7.8-10.4	146	4.9	4.1-5.8
2014	892	17.6	16.4-18.8	463	20.8	18.9-22.8	429	15.3	13.8-16.9	293	5.5	4.9-6.2	148	6.8	5.7-8.0	145	4.7	3.9-5.5
2015	912	17.6	16.5-18.8	523	23.0	21.1-25.1	389	13.4	12.0-14.8	330	6.1	5.5-6.8	182	8.2	7.0-9.5	148	4.7	3.9-5.5
2016	869	16.5	15.4-17.7	512	22.1	20.2-24.1	357	12.2	11.0-13.6	322	5.9	5.3-6.6	160	7.0	6.0-8.2	162	5.3	4.5-6.2
2017	814	15.6	14.5-16.7	463	19.8	18.0-21.7	351	12.4	11.1-13.8	298	5.3	4.7-5.9	153	6.5	5.5-7.7	145	4.4	3.7-5.3
2018	792	15.3	14.3-16.5	460	19.9	18.1-21.8	332	11.5	10.3-12.9	276	4.9	4.3-5.6	143	5.9	5.0-7.0	133	4.2	3.5-5.0
2019	726	13.8	12.8-14.9	412	17.5	15.9-19.4	314	10.7	9.5-12.0	312	5.6	5.0-6.3	170	7.3	6.2-8.5	142	4.4	3.7-5.2
2020	706	13.5	12.5-14.6	403	17.2	15.6-19.0	303	10.6	9.4-12.0	256	4.6	4.0-5.2	151	6.4	5.4-7.5	105	3.3	2.7-4.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
All Invasive Malignant Tumors

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	35,773	399.1	394.9-403.3	17,586	473.2	465.9-480.7	18,187	360.7	355.4-366.1	19,253	217.2	214.0-220.3	10,040	277.4	271.7-283.2	9,213	180.9	177.1-184.6
1977	36,141	398.2	394.0-402.4	17,753	472.3	465.0-479.7	18,388	359.6	354.3-364.9	19,718	219.9	216.7-223.0	10,324	281.8	276.1-287.6	9,394	182.0	178.3-185.7
1978	36,664	400.8	396.6-405.0	17,975	475.6	468.3-483.0	18,689	361.9	356.7-367.3	19,927	219.0	215.9-222.1	10,361	281.7	276.0-287.5	9,566	182.0	178.3-185.7
1979	37,490	404.9	400.7-409.1	18,351	479.6	472.3-487.0	19,139	366.0	360.8-371.4	20,008	218.1	215.0-221.2	10,323	278.0	272.4-283.7	9,685	182.4	178.7-186.1
1980	38,893	414.6	410.4-418.8	19,439	499.8	492.4-507.2	19,454	368.2	362.9-373.5	20,562	220.9	217.8-224.0	10,715	284.0	278.3-289.7	9,847	182.9	179.2-186.6
1981	39,511	418.7	414.5-423.0	19,566	503.3	495.9-510.8	19,945	374.9	369.6-380.2	20,580	219.0	216.0-222.1	10,724	283.5	277.8-289.2	9,856	181.1	177.5-184.8
1982	40,300	422.2	418.0-426.4	19,920	506.0	498.7-513.5	20,380	379.0	373.7-384.3	20,850	220.1	217.1-223.2	10,783	283.7	278.0-289.4	10,067	183.1	179.4-186.7
1983	41,470	431.8	427.6-436.1	20,448	515.1	507.7-522.6	21,022	389.5	384.2-395.0	21,092	220.3	217.3-223.3	10,891	284.3	278.7-289.9	10,201	183.5	179.9-187.2
1984	41,960	434.0	429.8-438.2	20,449	510.3	502.9-517.6	21,511	395.3	390.0-400.8	21,310	221.0	218.0-224.1	10,826	280.1	274.5-285.7	10,484	187.4	183.8-191.1
1985	43,405	446.6	442.3-450.8	21,241	525.2	517.8-532.6	22,164	406.5	401.0-412.0	21,636	223.1	220.1-226.2	11,029	283.5	278.0-289.1	10,607	188.5	184.9-192.2
1986	43,480	444.9	440.7-449.2	21,188	522.3	514.9-529.7	22,292	406.7	401.2-412.2	22,000	225.3	222.3-228.4	11,282	287.9	282.3-293.5	10,718	189.2	185.6-192.9
1987	44,308	449.4	445.1-453.7	21,429	523.8	516.5-531.2	22,879	413.3	407.9-418.8	21,927	222.9	219.9-225.9	11,161	283.2	277.7-288.7	10,766	187.9	184.3-191.5
1988	44,445	447.0	442.8-451.2	21,486	521.2	514.0-528.6	22,959	412.2	406.7-417.6	22,220	223.3	220.4-226.3	11,354	284.4	279.0-289.9	10,866	188.1	184.5-191.8
1989	44,808	446.9	442.7-451.1	21,992	527.7	520.4-535.0	22,816	406.5	401.1-411.9	21,864	218.4	215.5-221.3	11,092	275.4	270.0-280.8	10,772	185.5	182.0-189.1
1990	47,089	464.4	460.2-468.7	23,248	547.0	539.7-554.3	23,841	421.0	415.6-426.5	21,869	215.5	212.6-218.4	11,055	271.6	266.3-276.9	10,814	183.0	179.6-186.6
1991	47,804	467.5	463.3-471.8	23,986	558.5	551.2-565.8	23,818	418.4	413.0-423.9	22,298	218.0	215.1-220.9	11,310	275.2	270.0-280.6	10,988	184.9	181.4-188.4
1992	49,764	481.5	477.2-485.8	25,678	590.1	582.7-597.6	24,086	419.2	413.8-424.6	22,511	217.6	214.8-220.5	11,202	270.9	265.7-276.1	11,309	188.2	184.7-191.7
1993	49,863	479.0	474.8-483.2	25,350	573.5	566.3-580.8	24,513	423.4	418.0-428.8	22,289	213.3	210.5-216.2	11,105	265.1	260.0-270.3	11,184	184.0	180.5-187.5
1994	50,316	481.5	477.2-485.7	25,362	571.4	564.2-578.6	24,954	429.1	423.7-434.6	23,050	219.0	216.2-221.9	11,657	275.7	270.5-280.9	11,393	186.0	182.6-189.5
1995	50,768	484.0	479.8-488.2	25,138	558.8	551.8-565.9	25,630	440.8	435.4-446.4	22,750	214.7	211.9-217.5	11,416	268.1	263.1-273.2	11,334	184.2	180.8-187.7
1996	52,517	499.0	494.7-503.3	26,386	583.0	575.9-590.2	26,131	448.1	442.6-453.6	22,600	212.1	209.4-214.9	11,351	263.1	258.2-268.1	11,249	182.1	178.7-185.5
1997	53,713	506.7	502.4-511.0	26,987	590.2	583.0-597.3	26,726	454.7	449.2-460.3	22,202	206.3	203.6-209.0	11,167	255.5	250.7-260.4	11,035	176.5	173.2-179.9
1998	55,039	514.8	510.5-519.1	27,454	591.9	584.8-599.0	27,585	467.4	461.9-473.0	21,751	200.0	197.4-202.7	10,852	242.6	238.0-247.3	10,899	173.0	169.7-176.3
1999	55,889	518.3	514.0-522.6	28,069	600.8	593.7-607.9	27,820	466.0	460.5-471.6	22,340	203.2	200.5-205.9	11,153	247.9	243.3-252.6	11,187	175.8	172.5-179.1
2000	57,428	527.5	523.1-531.8	29,006	616.0	608.9-623.2	28,422	471.1	465.6-476.7	22,276	200.1	197.5-202.8	11,071	244.3	239.7-248.9	11,205	173.6	170.3-176.9
2001	57,145	521.8	517.5-526.1	28,825	605.2	598.2-612.2	28,320	468.7	463.2-474.3	22,024	196.4	193.8-199.0	10,955	238.8	234.3-243.4	11,069	170.8	167.6-174.1
2002	58,176	525.9	521.7-530.2	29,954	620.5	613.5-627.6	28,222	463.4	457.9-468.9	22,012	194.2	191.6-196.8	10,917	234.5	230.1-239.0	11,095	170.1	166.9-173.3
2003	57,993	519.1	514.9-523.4	29,418	601.1	594.2-608.0	28,575	465.3	459.8-470.8	21,476	187.2	184.7-189.7	10,525	222.7	218.4-227.0	10,951	165.9	162.7-169.1
2004	58,364	518.7	514.4-522.9	29,584	597.6	590.7-604.5	28,780	467.1	461.6-472.6	21,435	185.3	182.8-187.8	10,588	220.1	215.9-224.4	10,847	163.3	160.2-166.4
2005	59,135	522.5	518.3-526.8	30,025	599.7	592.9-606.6	29,110	470.1	464.6-475.6	21,045	180.4	178.0-182.9	10,423	214.7	210.6-218.9	10,622	158.6	155.6-161.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
All Invasive Malignant Tumors

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	60,851	534.3	530.1-538.6	31,288	617.4	610.5-624.4	29,563	477.0	471.4-482.6	21,126	180.0	177.6-182.5	10,459	212.3	208.2-216.5	10,667	159.1	156.0-162.2
2007	61,563	534.2	530.0-538.5	31,857	617.3	610.5-624.2	29,706	476.8	471.3-482.4	20,927	175.8	173.4-178.2	10,511	209.0	205.0-213.1	10,416	154.0	151.0-157.1
2008	62,656	537.8	533.5-542.1	32,038	612.0	605.2-618.8	30,618	487.7	482.1-493.4	21,127	174.9	172.5-177.3	10,629	208.0	204.0-212.1	10,498	153.2	150.2-156.2
2009	62,472	529.4	525.2-533.6	31,993	599.6	592.9-606.3	30,479	480.5	474.9-486.1	20,713	169.8	167.4-172.2	10,530	203.0	199.1-206.9	10,183	147.9	144.9-150.8
2010	61,284	514.0	509.9-518.2	31,073	573.1	566.6-579.6	30,211	473.9	468.4-479.5	20,729	167.5	165.2-169.9	10,261	194.3	190.5-198.1	10,468	149.8	146.9-152.8
2011	63,040	523.4	519.2-527.6	32,175	584.6	578.1-591.1	30,865	480.8	475.2-486.4	20,649	164.6	162.3-166.9	10,302	191.6	187.8-195.3	10,347	146.4	143.5-149.4
2012	61,715	506.7	502.6-510.8	31,067	556.2	549.9-562.6	30,648	474.8	469.2-480.3	21,101	166.8	164.5-169.2	10,719	197.5	193.7-201.3	10,382	145.7	142.9-148.7
2013	63,218	514.2	510.1-518.4	31,584	556.8	550.6-563.2	31,634	488.2	482.6-493.9	20,707	160.9	158.7-163.2	10,537	189.1	185.4-192.8	10,170	141.1	138.3-144.0
2014	63,153	507.5	503.4-511.6	31,528	546.7	540.5-553.0	31,625	483.8	478.2-489.4	20,176	154.9	152.7-157.1	10,356	183.1	179.5-186.8	9,820	135.3	132.5-138.1
2015	63,987	509.9	505.8-514.1	31,851	544.4	538.2-550.6	32,136	490.1	484.5-495.8	20,166	152.8	150.7-155.0	10,208	178.3	174.8-181.9	9,958	135.1	132.4-137.9
2016	64,516	509.1	505.0-513.2	32,734	551.6	545.4-557.8	31,782	482.0	476.4-487.6	20,203	152.2	150.0-154.3	10,221	175.9	172.4-179.4	9,982	135.8	133.0-138.6
2017	65,350	510.4	506.3-514.5	33,172	551.3	545.2-557.5	32,178	484.8	479.2-490.5	19,973	147.6	145.5-149.7	10,187	172.0	168.6-175.5	9,786	130.3	127.6-133.0
2018	64,982	501.7	497.7-505.8	33,209	542.9	536.9-548.9	31,773	474.2	468.7-479.8	19,916	145.7	143.6-147.8	10,123	168.7	165.4-172.1	9,793	129.9	127.2-132.6
2019	66,644	509.8	505.8-513.9	34,168	550.1	544.1-556.2	32,476	483.3	477.7-488.9	19,460	140.6	138.6-142.6	9,899	162.3	159.0-165.6	9,561	124.9	122.3-127.6
2020	59,801	454.2	450.4-458.1	30,626	489.3	483.7-495.0	29,175	431.1	425.8-436.4	19,174	137.6	135.6-139.6	9,732	158.6	155.4-161.9	9,442	122.7	120.1-125.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Oral cavity and pharynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	989	10.7	10.0-11.4	671	16.7	15.4-18.0	318	6.2	5.5-6.9	396	4.3	3.8-4.7	285	7.1	6.3-8.1	111	2.1	1.7-2.5
1977	981	10.4	9.7-11.1	658	16.0	14.8-17.4	323	6.2	5.5-6.9	396	4.2	3.8-4.7	270	6.8	6.0-7.7	126	2.4	2.0-2.8
1978	1,010	10.6	10.0-11.3	665	16.3	15.0-17.7	345	6.5	5.8-7.3	470	5.0	4.6-5.5	327	8.3	7.4-9.3	143	2.7	2.3-3.2
1979	1,055	10.9	10.2-11.6	674	16.1	14.8-17.4	381	7.1	6.4-7.8	413	4.3	3.9-4.8	257	6.2	5.4-7.0	156	2.9	2.5-3.4
1980	1,019	10.5	9.9-11.2	665	15.9	14.7-17.2	354	6.5	5.8-7.2	401	4.1	3.7-4.6	258	6.3	5.5-7.2	143	2.6	2.1-3.0
1981	1,060	10.9	10.2-11.6	725	17.1	15.9-18.5	335	6.1	5.5-6.8	395	4.0	3.6-4.4	271	6.5	5.7-7.4	124	2.2	1.8-2.6
1982	1,112	11.5	10.8-12.2	740	17.8	16.4-19.1	372	6.9	6.2-7.6	417	4.3	3.9-4.7	276	6.9	6.1-7.8	141	2.5	2.1-3.0
1983	1,049	10.7	10.0-11.3	690	16.1	14.9-17.5	359	6.7	6.0-7.4	398	4.0	3.6-4.5	261	6.2	5.5-7.1	137	2.4	2.0-2.9
1984	1,061	10.8	10.2-11.5	714	17.0	15.7-18.3	347	6.3	5.6-7.0	389	4.0	3.6-4.4	249	6.1	5.3-6.9	140	2.5	2.1-3.0
1985	1,059	10.7	10.1-11.4	699	16.5	15.2-17.8	360	6.4	5.8-7.2	377	3.8	3.4-4.2	240	5.8	5.1-6.6	137	2.5	2.1-2.9
1986	1,037	10.6	9.9-11.2	685	16.3	15.0-17.6	352	6.3	5.6-7.0	409	4.1	3.7-4.6	266	6.3	5.5-7.1	143	2.5	2.1-3.0
1987	1,036	10.5	9.9-11.2	652	15.3	14.1-16.6	384	6.8	6.1-7.6	382	3.9	3.5-4.3	241	5.9	5.2-6.8	141	2.4	2.1-2.9
1988	1,032	10.4	9.7-11.0	649	15.1	13.9-16.3	383	6.9	6.2-7.6	399	4.0	3.6-4.4	261	6.5	5.7-7.4	138	2.3	2.0-2.8
1989	999	9.9	9.3-10.5	662	15.0	13.8-16.2	337	6.0	5.4-6.7	332	3.3	3.0-3.7	201	4.8	4.1-5.5	131	2.3	1.9-2.7
1990	983	9.8	9.2-10.4	622	14.1	12.9-15.2	361	6.5	5.8-7.2	345	3.4	3.0-3.7	220	5.0	4.4-5.8	125	2.1	1.7-2.5
1991	1,013	10.1	9.5-10.7	645	14.6	13.5-15.8	368	6.5	5.8-7.2	343	3.4	3.0-3.8	209	4.9	4.2-5.6	134	2.3	1.9-2.7
1992	1,008	9.9	9.2-10.5	639	14.3	13.2-15.5	369	6.4	5.8-7.2	302	2.9	2.6-3.2	182	4.1	3.5-4.7	120	2.0	1.6-2.4
1993	1,006	9.8	9.2-10.5	663	14.9	13.8-16.1	343	5.9	5.3-6.5	296	2.8	2.5-3.2	175	3.9	3.3-4.6	121	2.0	1.7-2.4
1994	1,006	9.8	9.2-10.4	661	14.7	13.6-15.9	345	6.0	5.4-6.7	305	2.9	2.6-3.3	211	4.7	4.1-5.4	94	1.6	1.3-1.9
1995	937	9.1	8.5-9.7	570	12.5	11.5-13.6	367	6.3	5.6-6.9	322	3.1	2.8-3.4	214	4.9	4.2-5.6	108	1.8	1.5-2.2
1996	1,030	9.9	9.3-10.5	666	14.3	13.2-15.4	364	6.3	5.7-7.0	283	2.7	2.4-3.0	184	4.2	3.6-4.8	99	1.6	1.3-2.0
1997	1,086	10.4	9.8-11.0	703	15.1	13.9-16.2	383	6.5	5.9-7.2	272	2.6	2.3-2.9	177	3.9	3.3-4.5	95	1.5	1.2-1.9
1998	1,054	10.0	9.4-10.6	665	14.2	13.1-15.3	389	6.6	6.0-7.3	303	2.8	2.5-3.1	190	4.1	3.6-4.8	113	1.8	1.5-2.1
1999	1,028	9.6	9.1-10.2	695	14.6	13.5-15.7	333	5.5	4.9-6.2	261	2.4	2.1-2.7	167	3.5	3.0-4.1	94	1.5	1.2-1.8
2000	1,040	9.6	9.1-10.2	677	14.1	13.1-15.3	363	6.0	5.4-6.7	236	2.1	1.9-2.4	147	3.2	2.7-3.7	89	1.4	1.1-1.7
2001	1,079	9.9	9.3-10.5	688	13.9	12.9-15.0	391	6.4	5.8-7.1	274	2.5	2.2-2.8	172	3.6	3.1-4.2	102	1.5	1.2-1.8
2002	1,110	10.1	9.5-10.7	725	14.5	13.5-15.6	385	6.3	5.6-6.9	251	2.2	2.0-2.5	158	3.3	2.8-3.8	93	1.4	1.1-1.7
2003	1,116	10.0	9.4-10.6	736	14.6	13.6-15.7	380	6.2	5.6-6.8	240	2.1	1.9-2.4	149	3.0	2.5-3.5	91	1.4	1.1-1.7
2004	1,170	10.4	9.8-11.0	795	15.6	14.5-16.7	375	6.0	5.4-6.7	272	2.3	2.1-2.6	179	3.5	3.0-4.1	93	1.4	1.1-1.7
2005	1,121	9.9	9.3-10.5	718	13.8	12.8-14.8	403	6.6	6.0-7.3	241	2.1	1.8-2.3	161	3.1	2.7-3.7	80	1.2	0.9-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Oral cavity and pharynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,214	10.6	10.0-11.2	850	16.1	15.1-17.3	364	5.9	5.3-6.6	249	2.1	1.9-2.4	161	3.1	2.7-3.7	88	1.3	1.0-1.6
2007	1,305	11.2	10.6-11.9	906	16.8	15.7-18.0	399	6.4	5.8-7.1	234	2.0	1.7-2.2	159	3.0	2.5-3.5	75	1.1	0.9-1.4
2008	1,281	10.9	10.3-11.5	870	16.0	14.9-17.1	411	6.5	5.9-7.2	250	2.0	1.8-2.3	182	3.3	2.8-3.8	68	1.0	0.8-1.3
2009	1,344	11.1	10.5-11.8	912	16.2	15.2-17.3	432	6.7	6.0-7.4	247	2.0	1.7-2.3	164	3.1	2.6-3.6	83	1.1	0.9-1.4
2010	1,392	11.5	10.8-12.1	943	16.7	15.6-17.8	449	6.9	6.3-7.6	262	2.1	1.9-2.4	170	3.1	2.6-3.6	92	1.3	1.0-1.6
2011	1,407	11.4	10.8-12.0	981	16.9	15.8-18.0	426	6.5	5.8-7.1	258	2.0	1.8-2.3	168	3.0	2.5-3.5	90	1.3	1.0-1.6
2012	1,428	11.5	10.9-12.1	983	17.0	15.9-18.1	445	6.6	6.0-7.3	279	2.2	1.9-2.5	197	3.5	3.0-4.0	82	1.1	0.8-1.3
2013	1,484	11.9	11.3-12.5	1,048	18.1	17.0-19.2	436	6.4	5.8-7.1	277	2.1	1.9-2.4	191	3.2	2.8-3.8	86	1.1	0.9-1.4
2014	1,519	12.0	11.4-12.6	1,048	17.6	16.5-18.7	471	7.0	6.4-7.7	226	1.7	1.5-2.0	159	2.7	2.3-3.2	67	0.9	0.7-1.2
2015	1,618	12.7	12.1-13.3	1,167	19.4	18.3-20.6	451	6.8	6.2-7.5	281	2.1	1.9-2.4	189	3.2	2.8-3.7	92	1.2	1.0-1.6
2016	1,706	13.1	12.4-13.7	1,188	19.4	18.3-20.6	518	7.5	6.9-8.3	309	2.3	2.0-2.5	218	3.5	3.1-4.1	91	1.2	0.9-1.5
2017	1,630	12.5	11.9-13.2	1,134	18.6	17.5-19.7	496	7.1	6.5-7.8	304	2.2	2.0-2.5	210	3.4	2.9-3.9	94	1.3	1.0-1.6
2018	1,661	12.6	12.0-13.3	1,155	18.5	17.5-19.7	506	7.3	6.6-8.0	312	2.3	2.0-2.5	211	3.4	2.9-3.9	101	1.3	1.1-1.7
2019	1,773	13.4	12.8-14.1	1,232	19.7	18.5-20.8	541	7.8	7.2-8.6	319	2.3	2.0-2.6	224	3.5	3.1-4.0	95	1.2	1.0-1.5
2020	1,651	12.2	11.6-12.9	1,173	18.4	17.3-19.5	478	6.8	6.1-7.4	310	2.2	2.0-2.5	211	3.2	2.8-3.7	99	1.3	1.1-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Esophagus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	362	4.0	3.6-4.4	276	7.2	6.3-8.1	86	1.7	1.3-2.1	330	3.6	3.2-4.1	252	6.6	5.8-7.5	78	1.5	1.2-1.9
1977	362	4.0	3.6-4.4	255	6.7	5.9-7.7	107	2.0	1.7-2.5	323	3.5	3.1-3.9	232	6.0	5.2-6.9	91	1.7	1.4-2.1
1978	353	3.8	3.4-4.2	251	6.6	5.8-7.5	102	1.9	1.6-2.3	371	4.0	3.6-4.5	271	7.1	6.3-8.1	100	1.9	1.5-2.3
1979	375	4.0	3.6-4.4	257	6.4	5.6-7.2	118	2.2	1.8-2.6	325	3.5	3.1-3.9	225	5.8	5.0-6.6	100	1.8	1.5-2.3
1980	375	3.9	3.5-4.3	260	6.6	5.8-7.5	115	2.1	1.7-2.5	339	3.5	3.1-3.9	228	5.7	4.9-6.5	111	2.0	1.6-2.4
1981	376	3.9	3.5-4.3	259	6.4	5.6-7.3	117	2.1	1.7-2.5	355	3.7	3.3-4.1	262	6.7	5.8-7.6	93	1.7	1.4-2.1
1982	376	3.8	3.4-4.2	251	6.0	5.3-6.9	125	2.2	1.8-2.6	354	3.6	3.3-4.0	238	5.9	5.2-6.8	116	2.0	1.7-2.4
1983	396	4.0	3.6-4.4	280	6.8	6.0-7.7	116	2.0	1.7-2.4	368	3.8	3.4-4.3	276	7.1	6.2-8.0	92	1.6	1.3-2.0
1984	402	4.0	3.7-4.5	275	6.7	5.9-7.6	127	2.2	1.8-2.7	371	3.7	3.4-4.1	256	6.4	5.6-7.3	115	2.0	1.6-2.4
1985	373	3.8	3.4-4.2	280	6.8	6.0-7.6	93	1.6	1.3-2.0	359	3.6	3.2-4.0	265	6.4	5.6-7.3	94	1.6	1.3-2.0
1986	424	4.3	3.9-4.7	299	7.1	6.3-8.0	125	2.2	1.8-2.6	370	3.7	3.3-4.1	264	6.2	5.5-7.1	106	1.8	1.5-2.2
1987	399	4.0	3.6-4.4	276	6.5	5.8-7.4	123	2.1	1.7-2.5	391	3.9	3.5-4.3	284	6.9	6.0-7.7	107	1.8	1.5-2.2
1988	417	4.1	3.7-4.6	299	7.0	6.2-7.9	118	2.0	1.6-2.4	424	4.2	3.8-4.6	317	7.5	6.7-8.4	107	1.8	1.5-2.2
1989	414	4.0	3.6-4.4	300	6.9	6.1-7.7	114	1.9	1.6-2.3	405	4.0	3.6-4.4	301	7.1	6.3-7.9	104	1.7	1.4-2.1
1990	466	4.5	4.1-5.0	320	7.2	6.4-8.1	146	2.4	2.0-2.8	395	3.8	3.4-4.2	287	6.4	5.7-7.2	108	1.8	1.4-2.1
1991	450	4.3	3.9-4.7	310	6.9	6.1-7.7	140	2.3	2.0-2.8	431	4.1	3.8-4.5	311	7.1	6.3-7.9	120	2.0	1.6-2.3
1992	514	4.9	4.5-5.4	351	7.8	6.9-8.6	163	2.6	2.2-3.1	460	4.4	4.0-4.8	334	7.5	6.7-8.4	126	2.1	1.7-2.5
1993	501	4.8	4.3-5.2	357	7.9	7.1-8.8	144	2.3	2.0-2.7	455	4.3	3.9-4.7	335	7.5	6.7-8.3	120	1.9	1.6-2.3
1994	514	4.9	4.5-5.3	368	8.2	7.3-9.1	146	2.3	2.0-2.7	460	4.3	3.9-4.7	325	7.2	6.4-8.0	135	2.2	1.8-2.6
1995	494	4.7	4.3-5.1	354	7.9	7.1-8.8	140	2.2	1.9-2.6	502	4.8	4.4-5.2	378	8.5	7.6-9.4	124	1.9	1.6-2.3
1996	532	5.0	4.6-5.5	373	8.1	7.3-9.0	159	2.5	2.1-3.0	474	4.5	4.1-4.9	337	7.5	6.7-8.4	137	2.2	1.8-2.6
1997	481	4.5	4.1-4.9	339	7.4	6.6-8.3	142	2.3	1.9-2.7	477	4.4	4.0-4.9	368	8.1	7.3-8.9	109	1.7	1.4-2.0
1998	545	5.1	4.6-5.5	411	8.8	7.9-9.7	134	2.2	1.8-2.6	485	4.5	4.1-4.9	366	7.8	7.1-8.7	119	1.9	1.6-2.3
1999	533	4.9	4.5-5.3	386	8.2	7.4-9.1	147	2.3	1.9-2.7	499	4.6	4.2-5.0	367	7.9	7.1-8.7	132	2.0	1.7-2.4
2000	583	5.3	4.9-5.8	426	9.0	8.2-9.9	157	2.4	2.0-2.8	504	4.6	4.2-5.0	366	7.8	7.0-8.7	138	2.1	1.8-2.5
2001	536	4.8	4.4-5.3	402	8.3	7.5-9.1	134	2.1	1.7-2.4	539	4.8	4.4-5.3	412	8.6	7.8-9.5	127	1.9	1.6-2.3
2002	585	5.2	4.8-5.7	441	9.0	8.2-9.9	144	2.2	1.8-2.6	507	4.5	4.1-4.9	384	7.9	7.1-8.7	123	1.8	1.5-2.2
2003	607	5.4	4.9-5.8	462	9.3	8.5-10.2	145	2.1	1.8-2.5	567	5.0	4.6-5.4	421	8.6	7.8-9.5	146	2.1	1.8-2.5
2004	626	5.5	5.1-5.9	479	9.6	8.8-10.6	147	2.2	1.8-2.6	586	5.1	4.7-5.6	452	9.0	8.2-9.9	134	1.9	1.6-2.3
2005	682	5.9	5.5-6.4	526	10.4	9.5-11.4	156	2.4	2.0-2.8	604	5.2	4.8-5.6	471	9.5	8.6-10.4	133	1.9	1.6-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Esophagus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	702	6.1	5.6-6.5	550	10.7	9.8-11.7	152	2.3	2.0-2.7	553	4.8	4.4-5.2	436	8.6	7.8-9.5	117	1.8	1.5-2.2
2007	663	5.6	5.2-6.1	512	9.8	8.9-10.7	151	2.3	1.9-2.7	585	4.9	4.5-5.4	434	8.4	7.7-9.3	151	2.2	1.9-2.6
2008	702	5.9	5.5-6.4	549	10.4	9.6-11.4	153	2.3	1.9-2.7	575	4.8	4.4-5.2	460	8.7	7.9-9.5	115	1.7	1.4-2.0
2009	736	6.1	5.7-6.6	576	10.6	9.8-11.6	160	2.3	2.0-2.7	567	4.6	4.2-5.0	455	8.4	7.6-9.2	112	1.6	1.3-1.9
2010	678	5.5	5.1-5.9	514	9.3	8.5-10.2	164	2.3	2.0-2.7	585	4.8	4.4-5.2	470	8.7	7.9-9.5	115	1.6	1.3-1.9
2011	651	5.2	4.8-5.6	486	8.6	7.8-9.4	165	2.4	2.0-2.8	623	5.0	4.6-5.4	477	8.6	7.8-9.4	146	2.0	1.7-2.4
2012	640	5.1	4.7-5.5	511	9.0	8.2-9.8	129	1.8	1.5-2.1	584	4.6	4.2-5.0	458	8.2	7.5-9.0	126	1.7	1.4-2.0
2013	683	5.3	4.9-5.7	535	9.2	8.4-10.0	148	2.0	1.7-2.4	589	4.5	4.2-4.9	457	7.9	7.2-8.7	132	1.8	1.5-2.2
2014	686	5.2	4.8-5.6	536	8.9	8.1-9.7	150	2.1	1.7-2.4	576	4.5	4.1-4.9	475	8.3	7.5-9.1	101	1.4	1.1-1.7
2015	744	5.7	5.3-6.1	570	9.4	8.7-10.3	174	2.5	2.1-2.9	559	4.2	3.8-4.6	436	7.3	6.6-8.0	123	1.6	1.4-2.0
2016	737	5.5	5.1-6.0	567	9.3	8.6-10.2	170	2.3	2.0-2.7	599	4.5	4.1-4.8	474	7.8	7.1-8.6	125	1.7	1.4-2.0
2017	678	5.0	4.7-5.4	525	8.5	7.7-9.2	153	2.1	1.8-2.5	589	4.3	3.9-4.7	479	7.7	7.0-8.4	110	1.5	1.2-1.8
2018	715	5.3	4.9-5.7	564	9.1	8.3-9.9	151	2.0	1.7-2.4	565	4.1	3.8-4.5	437	7.0	6.4-7.7	128	1.7	1.4-2.1
2019	733	5.3	4.9-5.7	574	9.0	8.3-9.8	159	2.0	1.7-2.4	559	3.9	3.6-4.3	446	6.9	6.3-7.6	113	1.4	1.2-1.7
2020	674	4.8	4.5-5.2	534	8.4	7.7-9.2	140	1.8	1.5-2.1	561	4.0	3.7-4.4	451	7.3	6.6-8.0	110	1.4	1.1-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Stomach

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	962	11.1	10.4-11.8	554	15.3	14.0-16.7	408	8.1	7.3-8.9	762	8.9	8.3-9.6	418	11.9	10.7-13.2	344	6.8	6.1-7.6
1977	955	10.9	10.2-11.7	563	15.7	14.4-17.1	392	7.6	6.9-8.4	749	8.6	8.0-9.3	443	12.6	11.4-13.9	306	6.0	5.3-6.7
1978	943	10.6	9.9-11.3	530	14.6	13.3-16.0	413	7.9	7.1-8.7	698	8.0	7.4-8.6	390	10.9	9.8-12.1	308	5.9	5.3-6.6
1979	965	10.6	9.9-11.3	572	15.5	14.2-16.9	393	7.3	6.6-8.1	728	8.1	7.5-8.7	420	11.5	10.4-12.7	308	5.7	5.1-6.4
1980	920	10.0	9.4-10.7	557	14.8	13.5-16.2	363	6.8	6.1-7.5	731	8.1	7.5-8.7	419	11.5	10.4-12.8	312	5.8	5.2-6.5
1981	1,012	10.9	10.2-11.6	610	16.0	14.7-17.4	402	7.4	6.7-8.2	701	7.6	7.0-8.2	413	11.1	10.0-12.2	288	5.3	4.7-5.9
1982	989	10.5	9.8-11.1	592	15.6	14.3-17.0	397	7.1	6.4-7.9	702	7.5	7.0-8.1	433	11.6	10.5-12.8	269	4.8	4.3-5.5
1983	973	10.3	9.6-10.9	590	15.3	14.1-16.7	383	6.8	6.1-7.5	687	7.3	6.8-7.9	412	10.9	9.8-12.1	275	4.9	4.3-5.5
1984	961	10.0	9.4-10.7	586	14.9	13.6-16.2	375	6.5	5.9-7.2	725	7.5	7.0-8.1	426	10.7	9.7-11.8	299	5.2	4.6-5.8
1985	982	10.1	9.5-10.8	617	15.7	14.4-17.0	365	6.4	5.7-7.1	650	6.8	6.3-7.3	378	10.0	9.0-11.1	272	4.7	4.2-5.3
1986	961	9.8	9.2-10.5	604	15.4	14.1-16.7	357	6.1	5.5-6.8	700	7.2	6.7-7.8	425	10.8	9.8-12.0	275	4.7	4.1-5.3
1987	969	9.9	9.2-10.5	582	14.6	13.4-15.9	387	6.6	5.9-7.3	676	6.9	6.4-7.4	404	10.5	9.4-11.6	272	4.6	4.1-5.2
1988	935	9.5	8.9-10.1	534	13.3	12.1-14.5	401	6.8	6.2-7.6	646	6.6	6.1-7.1	364	9.2	8.3-10.3	282	4.7	4.2-5.3
1989	894	8.9	8.3-9.5	595	14.7	13.5-16.0	299	4.9	4.4-5.5	676	6.8	6.3-7.3	398	10.2	9.2-11.3	278	4.6	4.0-5.1
1990	996	9.9	9.2-10.5	632	15.1	13.9-16.4	364	6.0	5.4-6.7	619	6.2	5.7-6.7	390	9.7	8.7-10.7	229	3.7	3.3-4.3
1991	893	8.7	8.1-9.3	560	13.3	12.2-14.5	333	5.5	4.9-6.1	596	5.8	5.4-6.3	385	9.6	8.6-10.6	211	3.4	3.0-3.9
1992	924	8.9	8.4-9.5	584	13.9	12.8-15.1	340	5.4	4.9-6.0	616	6.0	5.5-6.5	388	9.6	8.6-10.6	228	3.6	3.1-4.1
1993	962	9.2	8.7-9.8	587	13.8	12.7-15.0	375	5.9	5.3-6.5	598	5.7	5.3-6.2	362	8.9	8.0-9.9	236	3.7	3.2-4.2
1994	922	8.8	8.2-9.4	566	13.2	12.1-14.4	356	5.6	5.0-6.2	614	5.9	5.4-6.4	363	8.7	7.8-9.6	251	3.9	3.4-4.4
1995	907	8.6	8.0-9.1	550	12.6	11.6-13.8	357	5.7	5.1-6.3	581	5.5	5.1-6.0	330	7.9	7.1-8.8	251	3.9	3.4-4.4
1996	897	8.4	7.9-9.0	552	12.7	11.6-13.8	345	5.3	4.8-5.9	551	5.2	4.7-5.6	320	7.5	6.7-8.4	231	3.5	3.1-4.0
1997	926	8.6	8.1-9.2	564	12.6	11.6-13.7	362	5.6	5.0-6.2	521	4.8	4.4-5.3	292	6.7	5.9-7.5	229	3.5	3.0-4.0
1998	931	8.6	8.0-9.1	595	13.0	12.0-14.1	336	5.2	4.6-5.8	505	4.6	4.2-5.0	287	6.4	5.7-7.3	218	3.3	2.9-3.8
1999	885	8.1	7.6-8.7	561	12.3	11.3-13.3	324	5.0	4.5-5.6	494	4.5	4.1-4.9	300	6.6	5.9-7.4	194	2.9	2.5-3.4
2000	902	8.1	7.6-8.6	561	12.2	11.2-13.3	341	5.0	4.5-5.6	498	4.4	4.1-4.9	283	6.4	5.6-7.1	215	3.2	2.8-3.6
2001	860	7.7	7.2-8.3	550	11.7	10.8-12.8	310	4.8	4.3-5.4	473	4.2	3.8-4.6	271	5.9	5.2-6.7	202	3.0	2.6-3.4
2002	857	7.6	7.1-8.1	524	11.1	10.1-12.1	333	5.1	4.5-5.6	441	3.9	3.5-4.2	255	5.5	4.9-6.3	186	2.7	2.3-3.1
2003	906	8.0	7.5-8.5	565	11.7	10.7-12.7	341	5.2	4.6-5.7	425	3.7	3.3-4.0	243	5.2	4.5-5.9	182	2.7	2.3-3.1
2004	861	7.5	7.0-8.0	524	10.8	9.9-11.8	337	5.1	4.5-5.6	470	4.0	3.7-4.4	277	5.9	5.2-6.6	193	2.8	2.4-3.2
2005	800	6.9	6.4-7.4	495	9.9	9.0-10.8	305	4.6	4.1-5.1	419	3.6	3.2-3.9	240	5.0	4.3-5.6	179	2.6	2.2-3.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Stomach

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	799	6.9	6.4-7.4	492	9.8	8.9-10.7	307	4.7	4.1-5.2	371	3.1	2.8-3.5	212	4.2	3.7-4.9	159	2.3	2.0-2.7
2007	837	7.2	6.7-7.7	544	10.6	9.7-11.5	293	4.5	4.0-5.0	367	3.1	2.8-3.5	218	4.3	3.7-4.9	149	2.2	1.9-2.6
2008	775	6.5	6.0-7.0	505	9.8	9.0-10.7	270	4.0	3.6-4.6	394	3.2	2.9-3.5	240	4.7	4.1-5.3	154	2.1	1.8-2.5
2009	749	6.3	5.8-6.8	483	9.3	8.5-10.2	266	3.8	3.4-4.3	349	2.9	2.6-3.2	216	4.2	3.7-4.8	133	1.9	1.6-2.2
2010	820	6.8	6.3-7.3	531	9.9	9.0-10.8	289	4.3	3.8-4.8	340	2.8	2.5-3.1	204	3.9	3.4-4.5	136	1.9	1.6-2.3
2011	884	7.2	6.7-7.7	554	10.2	9.4-11.1	330	4.8	4.3-5.4	320	2.5	2.3-2.8	194	3.6	3.1-4.2	126	1.7	1.4-2.1
2012	847	6.8	6.4-7.3	533	9.6	8.8-10.4	314	4.7	4.1-5.2	355	2.8	2.5-3.1	212	3.9	3.4-4.5	143	2.0	1.7-2.4
2013	924	7.4	6.9-7.9	589	10.5	9.7-11.4	335	5.0	4.4-5.5	354	2.7	2.5-3.0	214	3.8	3.3-4.4	140	1.9	1.6-2.3
2014	846	6.7	6.3-7.2	567	10.0	9.2-10.9	279	4.1	3.6-4.7	338	2.7	2.4-3.0	215	3.8	3.3-4.4	123	1.8	1.5-2.1
2015	847	6.6	6.1-7.1	541	9.3	8.5-10.2	306	4.4	3.9-5.0	303	2.4	2.1-2.7	187	3.3	2.8-3.8	116	1.7	1.4-2.0
2016	887	6.9	6.5-7.4	545	9.3	8.5-10.1	342	5.1	4.5-5.7	314	2.4	2.1-2.7	197	3.3	2.9-3.8	117	1.6	1.3-2.0
2017	919	7.1	6.7-7.6	570	9.6	8.8-10.5	349	5.1	4.5-5.7	304	2.3	2.0-2.6	170	2.8	2.4-3.3	134	1.9	1.6-2.2
2018	878	6.8	6.4-7.3	537	9.1	8.3-9.9	341	5.0	4.5-5.6	310	2.3	2.0-2.6	187	3.2	2.7-3.7	123	1.6	1.3-2.0
2019	892	6.6	6.2-7.1	571	9.3	8.5-10.1	321	4.4	3.9-4.9	308	2.4	2.1-2.7	188	3.3	2.8-3.8	120	1.6	1.3-1.9
2020	811	6.2	5.7-6.6	534	8.8	8.0-9.6	277	4.1	3.6-4.6	294	2.1	1.9-2.4	172	2.8	2.4-3.2	122	1.7	1.4-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Colorectal

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	5,931	67.5	65.7-69.2	2,914	80.5	77.5-83.7	3,017	59.0	56.9-61.1	3,006	34.9	33.6-36.2	1,433	41.2	38.9-43.5	1,573	30.9	29.4-32.5
1977	6,067	67.8	66.0-69.5	2,944	80.1	77.0-83.2	3,123	59.9	57.8-62.0	3,015	34.7	33.5-36.0	1,438	41.3	39.0-43.6	1,577	30.5	29.0-32.1
1978	6,166	68.5	66.8-70.3	3,006	81.3	78.3-84.5	3,160	60.0	57.9-62.1	3,198	36.1	34.9-37.4	1,550	43.6	41.4-46.0	1,648	31.4	29.8-32.9
1979	6,228	68.1	66.4-69.9	2,978	79.5	76.5-82.6	3,250	60.5	58.4-62.7	3,152	35.1	33.9-36.4	1,510	42.7	40.5-45.1	1,642	30.6	29.1-32.1
1980	6,448	69.3	67.6-71.0	3,236	84.5	81.4-87.6	3,212	58.8	56.8-60.9	3,115	34.2	33.0-35.4	1,511	41.5	39.3-43.8	1,604	29.5	28.0-31.0
1981	6,602	70.5	68.8-72.2	3,313	87.5	84.3-90.6	3,289	59.5	57.4-61.6	3,164	34.4	33.2-35.6	1,543	42.1	39.9-44.4	1,621	29.3	27.9-30.8
1982	6,599	69.6	67.9-71.3	3,269	84.3	81.2-87.4	3,330	59.5	57.4-61.5	3,181	34.2	33.0-35.4	1,565	42.8	40.6-45.1	1,616	28.8	27.4-30.2
1983	6,751	70.6	68.9-72.3	3,374	86.8	83.8-90.0	3,377	59.8	57.8-61.9	3,125	33.1	31.9-34.3	1,526	41.3	39.1-43.5	1,599	28.0	26.7-29.5
1984	6,811	70.7	69.0-72.4	3,325	84.6	81.6-87.6	3,486	61.2	59.2-63.3	3,260	34.3	33.1-35.5	1,609	42.9	40.8-45.2	1,651	28.9	27.5-30.3
1985	7,053	72.5	70.8-74.3	3,548	89.4	86.3-92.5	3,505	61.1	59.1-63.2	3,174	33.2	32.0-34.4	1,541	40.9	38.8-43.1	1,633	28.2	26.8-29.6
1986	6,784	69.3	67.7-71.0	3,345	84.2	81.3-87.3	3,439	59.5	57.5-61.5	3,168	32.8	31.6-33.9	1,566	41.4	39.2-43.6	1,602	27.4	26.1-28.8
1987	6,739	68.3	66.6-69.9	3,343	83.8	80.9-86.8	3,396	58.3	56.3-60.3	3,154	32.3	31.2-33.5	1,578	41.3	39.2-43.4	1,576	26.6	25.3-28.0
1988	6,494	65.0	63.4-66.6	3,202	78.8	75.9-81.7	3,292	55.7	53.8-57.6	3,135	31.8	30.7-32.9	1,570	40.1	38.1-42.3	1,565	26.3	25.0-27.7
1989	6,437	64.0	62.5-65.6	3,232	79.2	76.4-82.1	3,205	53.9	52.0-55.8	2,908	29.1	28.1-30.2	1,406	35.7	33.8-37.7	1,502	24.9	23.7-26.2
1990	6,425	62.9	61.4-64.5	3,176	76.3	73.5-79.1	3,249	53.7	51.8-55.6	2,818	27.9	26.8-28.9	1,437	36.2	34.3-38.2	1,381	22.6	21.4-23.8
1991	6,380	62.1	60.5-63.6	3,175	75.6	72.9-78.3	3,205	52.8	51.0-54.7	2,862	28.0	27.0-29.1	1,383	34.3	32.4-36.2	1,479	23.9	22.7-25.2
1992	6,281	60.4	58.9-62.0	3,174	74.9	72.2-77.6	3,107	50.5	48.8-52.4	2,725	26.3	25.4-27.4	1,320	32.4	30.6-34.3	1,405	22.5	21.3-23.7
1993	6,374	60.7	59.2-62.2	3,094	71.8	69.2-74.4	3,280	52.9	51.0-54.7	2,788	26.6	25.6-27.6	1,356	33.0	31.2-34.9	1,432	22.5	21.3-23.7
1994	6,536	61.9	60.4-63.4	3,228	75.0	72.4-77.7	3,308	53.2	51.4-55.1	2,840	26.9	26.0-28.0	1,349	32.8	31.0-34.7	1,491	23.2	22.0-24.4
1995	6,436	60.5	59.0-62.0	3,196	73.3	70.8-76.0	3,240	51.5	49.7-53.3	2,789	26.1	25.2-27.1	1,332	32.1	30.3-33.9	1,457	22.5	21.4-23.7
1996	6,297	58.9	57.4-60.4	3,180	72.5	69.9-75.1	3,117	49.3	47.6-51.1	2,649	24.6	23.7-25.6	1,316	31.1	29.4-32.8	1,333	20.4	19.3-21.6
1997	6,633	61.5	60.0-63.0	3,305	73.7	71.2-76.3	3,328	52.5	50.7-54.3	2,516	23.2	22.3-24.1	1,248	28.9	27.3-30.6	1,268	19.3	18.2-20.4
1998	6,961	63.9	62.4-65.4	3,448	75.8	73.3-78.4	3,513	55.2	53.3-57.0	2,518	22.9	22.0-23.8	1,217	27.5	25.9-29.1	1,301	19.6	18.5-20.7
1999	7,017	63.8	62.3-65.3	3,490	76.2	73.7-78.8	3,527	54.7	52.9-56.6	2,478	22.3	21.4-23.2	1,216	27.4	25.9-29.0	1,262	18.7	17.7-19.8
2000	7,013	63.2	61.7-64.7	3,480	75.0	72.5-77.5	3,533	54.4	52.6-56.3	2,497	22.2	21.3-23.0	1,256	28.0	26.5-29.6	1,241	18.0	17.0-19.0
2001	6,678	59.7	58.3-61.2	3,298	70.2	67.8-72.6	3,380	51.8	50.0-53.6	2,411	21.2	20.4-22.1	1,138	25.1	23.7-26.7	1,273	18.6	17.5-19.6
2002	6,615	58.6	57.2-60.1	3,302	69.1	66.7-71.5	3,313	50.4	48.7-52.2	2,412	21.0	20.2-21.9	1,174	25.4	24.0-26.9	1,238	17.9	16.9-19.0
2003	6,314	55.2	53.9-56.6	3,068	63.4	61.1-65.7	3,246	49.0	47.3-50.8	2,198	18.9	18.1-19.7	1,083	23.1	21.7-24.5	1,115	16.0	15.1-17.0
2004	6,200	53.8	52.5-55.2	2,979	60.8	58.6-63.0	3,221	48.6	46.9-50.3	2,146	18.3	17.5-19.0	1,030	21.4	20.1-22.8	1,116	15.7	14.7-16.6
2005	5,870	50.5	49.2-51.8	2,948	59.5	57.3-61.7	2,922	43.8	42.2-45.5	1,974	16.6	15.9-17.4	962	20.1	18.8-21.4	1,012	14.2	13.3-15.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Colorectal

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	5,594	47.9	46.7-49.2	2,770	55.3	53.2-57.4	2,824	42.2	40.6-43.8	1,930	16.2	15.4-16.9	921	19.0	17.8-20.3	1,009	14.1	13.2-15.0
2007	5,730	48.6	47.4-49.9	2,820	55.2	53.2-57.3	2,910	43.5	41.9-45.1	1,922	15.9	15.2-16.6	944	18.9	17.7-20.2	978	13.6	12.8-14.5
2008	5,560	46.6	45.3-47.8	2,737	52.9	50.9-54.9	2,823	41.8	40.2-43.4	1,890	15.4	14.7-16.2	911	18.0	16.8-19.2	979	13.4	12.6-14.3
2009	5,212	43.3	42.1-44.5	2,594	49.4	47.5-51.3	2,618	38.3	36.8-39.8	1,821	14.8	14.1-15.5	896	17.5	16.4-18.7	925	12.8	12.0-13.7
2010	5,302	43.6	42.4-44.8	2,593	48.5	46.6-50.5	2,709	39.5	38.0-41.1	1,800	14.3	13.6-15.0	878	16.7	15.6-17.8	922	12.4	11.6-13.3
2011	5,117	42.0	40.8-43.2	2,512	46.7	44.8-48.6	2,605	38.2	36.6-39.7	1,775	13.9	13.2-14.5	873	16.2	15.1-17.3	902	12.0	11.2-12.8
2012	4,910	39.9	38.7-41.0	2,523	45.9	44.1-47.8	2,387	35.0	33.5-36.5	1,735	13.5	12.8-14.2	839	15.5	14.4-16.6	896	11.9	11.1-12.8
2013	4,934	39.7	38.5-40.8	2,543	45.5	43.7-47.4	2,391	35.0	33.5-36.5	1,714	13.3	12.6-13.9	907	16.5	15.4-17.6	807	10.6	9.9-11.5
2014	4,875	38.7	37.6-39.8	2,500	44.2	42.5-46.1	2,375	34.2	32.8-35.7	1,667	12.7	12.0-13.3	854	15.2	14.2-16.3	813	10.7	9.9-11.5
2015	4,910	39.1	38.0-40.2	2,486	43.6	41.8-45.4	2,424	35.3	33.8-36.8	1,673	12.6	12.0-13.3	859	15.1	14.1-16.2	814	10.7	9.9-11.5
2016	4,971	39.3	38.2-40.5	2,546	44.2	42.4-46.0	2,425	35.1	33.7-36.6	1,569	11.8	11.2-12.5	772	13.4	12.4-14.4	797	10.6	9.8-11.4
2017	4,907	38.7	37.6-39.8	2,506	43.3	41.6-45.1	2,401	34.8	33.3-36.3	1,696	12.7	12.0-13.3	841	14.5	13.5-15.5	855	11.2	10.4-12.0
2018	4,703	36.8	35.7-38.0	2,446	41.9	40.2-43.6	2,257	32.4	31.0-33.9	1,540	11.5	10.9-12.2	777	13.4	12.4-14.4	763	10.1	9.3-10.9
2019	4,788	37.3	36.2-38.5	2,496	42.3	40.6-44.1	2,292	33.0	31.6-34.5	1,544	11.2	10.6-11.8	805	13.5	12.5-14.5	739	9.2	8.5-9.9
2020	4,341	33.9	32.8-34.9	2,296	38.8	37.2-40.5	2,045	29.6	28.3-31.0	1,510	11.0	10.5-11.6	783	13.0	12.1-14.0	727	9.4	8.7-10.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Colon excluding rectum

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	4,136	47.5	46.0-49.0	1,919	53.9	51.4-56.6	2,217	43.4	41.6-45.2	2,355	27.4	26.2-28.5	1,057	30.4	28.5-32.4	1,298	25.5	24.2-27.0
1977	4,237	47.8	46.4-49.3	1,944	54.4	51.8-57.0	2,293	44.1	42.3-45.9	2,442	28.2	27.1-29.4	1,132	32.6	30.6-34.6	1,310	25.4	24.0-26.8
1978	4,336	48.5	47.1-50.0	1,999	55.2	52.6-57.8	2,337	44.4	42.6-46.2	2,563	29.0	27.8-30.1	1,186	33.5	31.5-35.5	1,377	26.2	24.8-27.6
1979	4,419	48.8	47.4-50.3	2,008	54.8	52.3-57.4	2,411	45.0	43.2-46.9	2,597	28.9	27.8-30.0	1,211	34.2	32.2-36.3	1,386	25.8	24.4-27.2
1980	4,584	49.6	48.2-51.1	2,217	58.6	56.1-61.3	2,367	43.4	41.7-45.2	2,542	27.9	26.8-29.0	1,212	33.4	31.4-35.4	1,330	24.4	23.1-25.8
1981	4,730	50.8	49.3-52.3	2,275	60.7	58.1-63.4	2,455	44.5	42.8-46.4	2,663	29.0	27.9-30.2	1,279	35.1	33.1-37.2	1,384	25.1	23.8-26.5
1982	4,642	49.3	47.9-50.8	2,175	57.0	54.5-59.6	2,467	44.0	42.3-45.8	2,625	28.2	27.1-29.3	1,269	34.7	32.7-36.8	1,356	24.2	22.9-25.5
1983	4,759	50.1	48.7-51.6	2,242	59.0	56.5-61.6	2,517	44.7	42.9-46.5	2,618	27.7	26.7-28.8	1,251	34.0	32.0-36.0	1,367	24.0	22.7-25.3
1984	4,894	51.1	49.6-52.5	2,237	57.8	55.3-60.4	2,657	46.5	44.8-48.4	2,735	28.8	27.7-29.9	1,321	35.2	33.2-37.2	1,414	24.7	23.4-26.0
1985	5,034	52.0	50.5-53.5	2,422	61.7	59.1-64.3	2,612	45.4	43.7-47.2	2,665	27.8	26.8-28.9	1,269	33.6	31.7-35.6	1,396	24.1	22.8-25.4
1986	4,902	50.3	48.9-51.7	2,299	58.7	56.2-61.3	2,603	45.0	43.2-46.7	2,689	27.8	26.7-28.9	1,305	34.6	32.7-36.7	1,384	23.7	22.4-25.0
1987	4,782	48.6	47.2-50.0	2,275	57.9	55.4-60.4	2,507	42.8	41.1-44.5	2,661	27.3	26.2-28.3	1,322	34.6	32.6-36.6	1,339	22.7	21.5-23.9
1988	4,649	46.6	45.3-48.0	2,158	53.7	51.3-56.1	2,491	42.0	40.3-43.7	2,695	27.3	26.3-28.4	1,330	34.0	32.1-35.9	1,365	22.9	21.7-24.2
1989	4,544	45.4	44.1-46.8	2,199	55.1	52.7-57.5	2,345	39.3	37.7-40.9	2,468	24.7	23.7-25.7	1,162	29.5	27.8-31.3	1,306	21.7	20.5-22.9
1990	4,538	44.6	43.3-45.9	2,140	52.3	50.0-54.7	2,398	39.5	37.9-41.2	2,447	24.2	23.2-25.2	1,252	31.7	29.9-33.6	1,195	19.5	18.4-20.7
1991	4,596	44.7	43.4-46.0	2,198	53.0	50.7-55.4	2,398	39.2	37.7-40.9	2,416	23.7	22.8-24.7	1,151	28.6	26.9-30.4	1,265	20.4	19.3-21.6
1992	4,485	43.2	42.0-44.5	2,135	51.0	48.8-53.3	2,350	38.1	36.5-39.7	2,312	22.4	21.5-23.3	1,088	26.9	25.3-28.6	1,224	19.6	18.5-20.7
1993	4,607	43.9	42.7-45.2	2,117	49.7	47.6-51.9	2,490	39.9	38.3-41.5	2,390	22.8	21.9-23.8	1,155	28.3	26.6-30.0	1,235	19.4	18.3-20.6
1994	4,739	44.9	43.6-46.2	2,253	53.0	50.7-55.3	2,486	39.8	38.2-41.4	2,437	23.1	22.2-24.0	1,132	27.7	26.1-29.4	1,305	20.3	19.2-21.4
1995	4,665	43.7	42.5-45.0	2,229	51.5	49.3-53.7	2,436	38.4	36.8-39.9	2,355	22.1	21.2-23.0	1,096	26.6	25.0-28.2	1,259	19.4	18.3-20.5
1996	4,498	41.9	40.7-43.2	2,161	50.0	47.8-52.2	2,337	36.7	35.2-38.2	2,245	20.9	20.0-21.8	1,101	26.1	24.6-27.7	1,144	17.5	16.5-18.5
1997	4,884	45.2	43.9-46.5	2,307	51.8	49.7-54.0	2,577	40.3	38.8-42.0	2,175	20.1	19.2-20.9	1,051	24.4	22.9-25.9	1,124	17.0	16.0-18.1
1998	5,069	46.4	45.2-47.7	2,424	53.9	51.7-56.1	2,645	41.3	39.7-42.9	2,131	19.4	18.6-20.2	1,013	22.9	21.5-24.4	1,118	16.7	15.8-17.8
1999	5,058	45.8	44.5-47.1	2,393	52.6	50.5-54.8	2,665	40.9	39.3-42.5	2,073	18.6	17.8-19.4	997	22.5	21.2-24.0	1,076	15.9	14.9-16.9
2000	5,097	45.7	44.5-47.0	2,402	52.2	50.2-54.4	2,695	41.0	39.5-42.6	2,106	18.6	17.9-19.5	1,039	23.3	21.9-24.8	1,067	15.4	14.5-16.4
2001	4,843	43.1	41.9-44.4	2,294	49.2	47.1-51.2	2,549	38.6	37.1-40.2	2,040	17.9	17.2-18.7	945	20.9	19.6-22.3	1,095	15.9	15.0-16.9
2002	4,792	42.3	41.1-43.5	2,295	48.4	46.4-50.4	2,497	37.7	36.2-39.2	1,991	17.3	16.6-18.1	954	20.7	19.4-22.1	1,037	14.9	14.0-15.9
2003	4,578	39.9	38.7-41.1	2,123	44.2	42.3-46.2	2,455	36.7	35.2-38.2	1,860	16.0	15.3-16.7	894	19.1	17.9-20.4	966	13.8	13.0-14.8
2004	4,452	38.5	37.4-39.7	2,025	41.7	39.9-43.6	2,427	36.2	34.7-37.7	1,773	15.0	14.3-15.8	833	17.4	16.2-18.6	940	13.1	12.3-14.0
2005	4,283	36.7	35.6-37.9	2,063	41.9	40.1-43.8	2,220	33.0	31.6-34.4	1,623	13.6	13.0-14.3	771	16.2	15.0-17.4	852	11.9	11.1-12.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Colon excluding rectum

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	4,117	35.1	34.0-36.2	1,965	39.6	37.8-41.4	2,152	31.6	30.3-33.0	1,602	13.4	12.7-14.1	750	15.5	14.4-16.7	852	11.8	11.0-12.7
2007	4,196	35.5	34.4-36.6	1,975	39.0	37.3-40.8	2,221	32.7	31.3-34.1	1,618	13.3	12.7-14.0	772	15.6	14.5-16.7	846	11.7	10.9-12.6
2008	4,036	33.7	32.6-34.7	1,882	36.6	34.9-38.3	2,154	31.7	30.3-33.1	1,540	12.6	11.9-13.2	728	14.4	13.4-15.5	812	11.1	10.3-11.9
2009	3,719	30.7	29.7-31.7	1,747	33.4	31.8-35.0	1,972	28.4	27.1-29.7	1,498	12.2	11.6-12.8	718	14.1	13.1-15.2	780	10.8	10.0-11.6
2010	3,808	31.2	30.2-32.2	1,767	33.5	31.9-35.1	2,041	29.3	28.0-30.6	1,466	11.6	11.0-12.2	691	13.2	12.2-14.2	775	10.3	9.6-11.1
2011	3,650	29.7	28.8-30.7	1,686	31.6	30.1-33.2	1,964	28.2	26.9-29.5	1,452	11.3	10.7-11.9	686	12.8	11.8-13.8	766	10.1	9.4-10.9
2012	3,537	28.5	27.6-29.5	1,708	31.3	29.8-32.8	1,829	26.3	25.1-27.6	1,426	11.1	10.5-11.7	660	12.2	11.3-13.2	766	10.2	9.4-11.0
2013	3,467	27.6	26.7-28.6	1,701	30.6	29.1-32.1	1,766	25.3	24.1-26.6	1,388	10.7	10.1-11.3	712	13.0	12.1-14.0	676	8.9	8.2-9.6
2014	3,448	27.2	26.3-28.2	1,662	29.7	28.2-31.2	1,786	25.3	24.1-26.6	1,334	10.1	9.6-10.7	667	12.0	11.1-12.9	667	8.7	8.0-9.4
2015	3,497	27.9	26.9-28.8	1,665	29.7	28.2-31.2	1,832	26.3	25.1-27.6	1,361	10.2	9.7-10.8	673	11.8	10.9-12.8	688	8.9	8.3-9.7
2016	3,569	28.1	27.1-29.1	1,715	29.9	28.5-31.4	1,854	26.5	25.2-27.8	1,223	9.2	8.7-9.7	577	10.0	9.2-10.9	646	8.5	7.8-9.2
2017	3,513	27.5	26.5-28.4	1,699	29.5	28.0-30.9	1,814	25.9	24.6-27.2	1,340	10.0	9.4-10.6	643	11.1	10.2-12.0	697	9.0	8.3-9.8
2018	3,308	25.8	24.8-26.7	1,628	28.1	26.7-29.6	1,680	23.7	22.5-24.9	1,193	8.9	8.4-9.4	573	9.9	9.1-10.8	620	8.1	7.5-8.8
2019	3,386	26.0	25.1-27.0	1,667	28.1	26.7-29.5	1,719	24.3	23.1-25.5	1,200	8.7	8.2-9.2	604	10.2	9.4-11.1	596	7.4	6.8-8.0
2020	3,060	23.7	22.8-24.6	1,544	26.2	24.8-27.5	1,516	21.6	20.4-22.8	1,146	8.3	7.8-8.8	576	9.6	8.8-10.4	570	7.3	6.7-8.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Rectum & rectosigmoid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,795	20.0	19.1-21.0	995	26.6	24.9-28.4	800	15.6	14.5-16.7	651	7.5	6.9-8.1	376	10.8	9.6-12.0	275	5.4	4.8-6.1
1977	1,830	19.9	19.0-20.9	1,000	25.7	24.0-27.4	830	15.8	14.8-17.0	573	6.5	6.0-7.1	306	8.7	7.7-9.8	267	5.1	4.5-5.8
1978	1,830	20.0	19.1-20.9	1,007	26.2	24.5-27.9	823	15.6	14.5-16.7	635	7.2	6.6-7.8	364	10.2	9.1-11.4	271	5.2	4.6-5.8
1979	1,809	19.3	18.4-20.2	970	24.7	23.1-26.4	839	15.5	14.5-16.6	555	6.2	5.7-6.8	299	8.6	7.6-9.6	256	4.8	4.2-5.4
1980	1,864	19.7	18.8-20.6	1,019	25.8	24.2-27.6	845	15.4	14.4-16.5	573	6.3	5.8-6.8	299	8.2	7.2-9.2	274	5.1	4.5-5.7
1981	1,872	19.7	18.8-20.6	1,038	26.7	25.0-28.5	834	14.9	13.9-16.0	501	5.3	4.9-5.8	264	7.0	6.1-7.9	237	4.2	3.7-4.8
1982	1,957	20.2	19.3-21.2	1,094	27.2	25.6-29.0	863	15.4	14.4-16.5	556	6.0	5.5-6.5	296	8.1	7.2-9.2	260	4.6	4.1-5.2
1983	1,992	20.5	19.6-21.4	1,132	27.8	26.1-29.6	860	15.2	14.1-16.2	507	5.4	4.9-5.9	275	7.3	6.4-8.3	232	4.1	3.6-4.6
1984	1,917	19.6	18.7-20.5	1,088	26.7	25.1-28.5	829	14.7	13.7-15.7	525	5.5	5.1-6.0	288	7.7	6.8-8.7	237	4.2	3.6-4.7
1985	2,019	20.5	19.6-21.5	1,126	27.7	26.0-29.4	893	15.7	14.7-16.8	509	5.3	4.9-5.8	272	7.3	6.5-8.3	237	4.1	3.6-4.6
1986	1,882	19.0	18.2-19.9	1,046	25.5	23.9-27.2	836	14.5	13.5-15.6	479	5.0	4.5-5.4	261	6.7	5.9-7.6	218	3.8	3.3-4.3
1987	1,957	19.7	18.8-20.6	1,068	25.9	24.3-27.6	889	15.4	14.4-16.5	493	5.1	4.6-5.5	256	6.7	5.9-7.6	237	4.0	3.5-4.5
1988	1,845	18.3	17.5-19.2	1,044	25.1	23.5-26.7	801	13.7	12.8-14.7	440	4.5	4.1-4.9	240	6.1	5.4-7.0	200	3.4	2.9-3.9
1989	1,893	18.6	17.8-19.5	1,033	24.1	22.6-25.7	860	14.6	13.7-15.7	440	4.4	4.0-4.9	244	6.2	5.4-7.0	196	3.3	2.8-3.7
1990	1,887	18.4	17.5-19.2	1,036	24.0	22.5-25.5	851	14.2	13.2-15.2	371	3.7	3.3-4.1	185	4.5	3.8-5.2	186	3.1	2.6-3.5
1991	1,784	17.3	16.5-18.2	977	22.6	21.1-24.1	807	13.6	12.6-14.5	446	4.3	3.9-4.7	232	5.7	4.9-6.5	214	3.5	3.0-4.0
1992	1,796	17.2	16.4-18.0	1,039	23.9	22.4-25.4	757	12.5	11.6-13.4	413	4.0	3.6-4.4	232	5.5	4.8-6.3	181	2.9	2.5-3.4
1993	1,767	16.8	16.0-17.6	977	22.1	20.7-23.5	790	13.0	12.1-13.9	398	3.8	3.4-4.2	201	4.7	4.1-5.4	197	3.1	2.7-3.5
1994	1,797	17.0	16.2-17.8	975	22.0	20.6-23.5	822	13.4	12.5-14.4	403	3.9	3.5-4.3	217	5.1	4.5-5.9	186	2.9	2.5-3.4
1995	1,771	16.8	16.0-17.6	967	21.8	20.4-23.3	804	13.1	12.2-14.1	434	4.1	3.7-4.5	236	5.5	4.8-6.3	198	3.1	2.7-3.6
1996	1,799	17.0	16.2-17.8	1,019	22.5	21.1-24.0	780	12.7	11.8-13.6	404	3.8	3.4-4.2	215	5.0	4.3-5.7	189	2.9	2.5-3.4
1997	1,749	16.3	15.5-17.1	998	21.9	20.5-23.3	751	12.1	11.3-13.0	341	3.2	2.8-3.5	197	4.6	3.9-5.3	144	2.2	1.9-2.6
1998	1,892	17.5	16.7-18.3	1,024	21.9	20.6-23.4	868	13.9	13.0-14.9	387	3.5	3.2-3.9	204	4.5	3.9-5.2	183	2.8	2.4-3.3
1999	1,959	18.0	17.2-18.8	1,097	23.6	22.2-25.1	862	13.9	12.9-14.8	405	3.7	3.3-4.0	219	4.9	4.3-5.6	186	2.8	2.4-3.3
2000	1,916	17.5	16.7-18.3	1,078	22.7	21.4-24.1	838	13.4	12.5-14.3	391	3.5	3.2-3.9	217	4.7	4.1-5.4	174	2.6	2.2-3.0
2001	1,835	16.6	15.8-17.3	1,004	21.0	19.7-22.4	831	13.2	12.3-14.1	371	3.3	3.0-3.6	193	4.2	3.6-4.9	178	2.6	2.3-3.1
2002	1,823	16.3	15.6-17.1	1,007	20.7	19.4-22.0	816	12.7	11.8-13.6	421	3.7	3.4-4.1	220	4.7	4.1-5.4	201	3.0	2.6-3.5
2003	1,736	15.4	14.6-16.1	945	19.1	17.9-20.4	791	12.3	11.5-13.2	338	2.9	2.6-3.3	189	4.0	3.4-4.6	149	2.2	1.9-2.6
2004	1,748	15.3	14.6-16.1	954	19.0	17.8-20.3	794	12.4	11.6-13.3	373	3.2	2.9-3.6	197	4.0	3.5-4.6	176	2.5	2.1-2.9
2005	1,587	13.8	13.1-14.5	885	17.6	16.4-18.8	702	10.9	10.1-11.7	351	3.0	2.7-3.3	191	3.9	3.4-4.5	160	2.3	1.9-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Rectum & rectosigmoid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,477	12.9	12.2-13.5	805	15.7	14.6-16.8	672	10.6	9.8-11.4	328	2.8	2.5-3.1	171	3.5	3.0-4.0	157	2.3	1.9-2.7
2007	1,534	13.1	12.5-13.8	845	16.2	15.1-17.3	689	10.8	10.0-11.7	304	2.6	2.3-2.9	172	3.4	2.9-3.9	132	1.9	1.6-2.3
2008	1,524	12.9	12.2-13.6	855	16.3	15.2-17.5	669	10.1	9.3-10.9	350	2.9	2.6-3.2	183	3.6	3.1-4.1	167	2.3	2.0-2.7
2009	1,493	12.6	12.0-13.3	847	16.0	14.9-17.1	646	9.9	9.1-10.7	323	2.6	2.3-2.9	178	3.4	2.9-3.9	145	2.0	1.7-2.4
2010	1,494	12.4	11.7-13.0	826	15.0	14.0-16.1	668	10.2	9.4-11.1	334	2.7	2.4-3.0	187	3.5	3.0-4.0	147	2.1	1.8-2.5
2011	1,467	12.3	11.6-12.9	826	15.0	14.0-16.1	641	10.0	9.2-10.8	323	2.6	2.3-2.9	187	3.4	2.9-3.9	136	1.9	1.6-2.3
2012	1,373	11.3	10.7-12.0	815	14.7	13.6-15.7	558	8.6	7.9-9.4	309	2.4	2.2-2.7	179	3.3	2.8-3.8	130	1.8	1.5-2.1
2013	1,467	12.0	11.4-12.7	842	14.9	13.9-16.0	625	9.7	8.9-10.5	326	2.6	2.3-2.9	195	3.5	3.0-4.0	131	1.8	1.5-2.2
2014	1,427	11.5	10.9-12.1	838	14.6	13.6-15.6	589	8.9	8.2-9.7	333	2.6	2.3-2.9	187	3.2	2.8-3.7	146	2.0	1.7-2.4
2015	1,413	11.2	10.6-11.9	821	13.9	12.9-14.9	592	9.0	8.2-9.8	312	2.4	2.1-2.7	186	3.3	2.8-3.8	126	1.7	1.4-2.1
2016	1,402	11.2	10.6-11.8	831	14.3	13.3-15.3	571	8.6	7.9-9.4	346	2.6	2.4-3.0	195	3.4	2.9-3.9	151	2.1	1.8-2.5
2017	1,394	11.2	10.6-11.8	807	13.9	12.9-14.9	587	8.9	8.2-9.7	356	2.7	2.4-3.0	198	3.4	2.9-3.9	158	2.1	1.8-2.5
2018	1,395	11.1	10.5-11.7	818	13.8	12.8-14.8	577	8.8	8.0-9.5	347	2.7	2.4-3.0	204	3.5	3.0-4.0	143	1.9	1.6-2.3
2019	1,402	11.3	10.7-12.0	829	14.2	13.2-15.3	573	8.8	8.0-9.6	344	2.5	2.2-2.8	201	3.3	2.8-3.8	143	1.8	1.5-2.2
2020	1,281	10.2	9.6-10.8	752	12.7	11.7-13.7	529	8.1	7.3-8.8	364	2.7	2.4-3.0	207	3.5	3.0-4.0	157	2.1	1.7-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Liver / intrahepatic bile duct

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	234	2.6	2.3-3.0	137	3.6	3.0-4.3	97	1.9	1.5-2.3	176	2.0	1.7-2.3	107	2.9	2.3-3.5	69	1.3	1.0-1.7
1977	278	2.9	2.6-3.3	186	4.6	3.9-5.4	92	1.7	1.4-2.1	237	2.5	2.2-2.9	139	3.4	2.8-4.1	98	1.9	1.5-2.3
1978	265	2.9	2.5-3.2	152	3.9	3.3-4.6	113	2.1	1.7-2.5	236	2.5	2.2-2.9	138	3.5	2.9-4.2	98	1.8	1.5-2.2
1979	252	2.7	2.4-3.1	132	3.3	2.7-3.9	120	2.3	1.9-2.8	199	2.2	1.9-2.5	108	2.7	2.2-3.3	91	1.7	1.4-2.1
1980	250	2.7	2.3-3.0	150	4.0	3.4-4.8	100	1.8	1.5-2.2	217	2.3	2.0-2.6	114	3.0	2.5-3.6	103	1.9	1.5-2.3
1981	254	2.7	2.4-3.1	148	3.9	3.2-4.6	106	1.9	1.6-2.3	236	2.5	2.2-2.9	131	3.5	2.9-4.2	105	1.9	1.6-2.3
1982	260	2.7	2.4-3.1	158	4.0	3.4-4.7	102	1.8	1.5-2.2	220	2.3	2.0-2.7	140	3.7	3.1-4.4	80	1.4	1.1-1.8
1983	276	2.9	2.5-3.2	186	4.5	3.9-5.3	90	1.7	1.3-2.1	231	2.4	2.1-2.7	153	3.9	3.2-4.6	78	1.4	1.1-1.7
1984	265	2.7	2.4-3.1	152	3.8	3.2-4.5	113	2.0	1.7-2.4	233	2.4	2.1-2.8	136	3.4	2.8-4.1	97	1.7	1.4-2.1
1985	256	2.7	2.3-3.0	162	4.2	3.5-4.9	94	1.7	1.4-2.1	239	2.5	2.2-2.8	149	3.8	3.2-4.5	90	1.6	1.3-2.0
1986	265	2.7	2.4-3.1	169	4.1	3.5-4.8	96	1.7	1.4-2.1	253	2.6	2.3-2.9	156	3.8	3.2-4.5	97	1.7	1.4-2.1
1987	281	2.8	2.5-3.2	176	4.2	3.6-4.9	105	1.9	1.5-2.3	281	2.9	2.5-3.2	168	4.2	3.6-5.0	113	2.0	1.6-2.4
1988	330	3.3	3.0-3.7	212	5.1	4.4-5.9	118	2.0	1.7-2.4	281	2.8	2.5-3.2	179	4.3	3.7-5.1	102	1.7	1.4-2.1
1989	312	3.1	2.7-3.4	196	4.5	3.9-5.2	116	2.0	1.6-2.4	257	2.5	2.2-2.9	156	3.7	3.1-4.4	101	1.7	1.4-2.1
1990	348	3.4	3.1-3.8	222	5.2	4.5-6.0	126	2.1	1.7-2.5	312	3.1	2.7-3.4	194	4.8	4.1-5.6	118	1.9	1.6-2.3
1991	342	3.3	3.0-3.7	209	4.7	4.1-5.4	133	2.2	1.9-2.6	310	3.0	2.6-3.3	188	4.3	3.7-5.0	122	2.0	1.7-2.4
1992	391	3.8	3.4-4.2	228	5.5	4.8-6.3	163	2.7	2.3-3.2	324	3.1	2.8-3.5	195	4.7	4.0-5.4	129	2.1	1.8-2.5
1993	412	3.9	3.5-4.3	258	6.0	5.3-6.8	154	2.5	2.1-2.9	401	3.8	3.4-4.2	240	5.6	4.9-6.4	161	2.5	2.2-3.0
1994	384	3.6	3.3-4.0	250	5.8	5.1-6.5	134	2.1	1.8-2.6	363	3.4	3.1-3.8	210	5.0	4.3-5.7	153	2.5	2.1-2.9
1995	445	4.2	3.8-4.6	287	6.5	5.7-7.3	158	2.5	2.2-3.0	398	3.7	3.4-4.1	251	5.8	5.1-6.6	147	2.3	1.9-2.7
1996	399	3.7	3.4-4.1	244	5.5	4.8-6.2	155	2.5	2.1-2.9	380	3.5	3.2-3.9	224	5.1	4.4-5.8	156	2.4	2.0-2.8
1997	463	4.4	4.0-4.8	300	6.5	5.8-7.3	163	2.6	2.2-3.1	386	3.6	3.3-4.0	243	5.4	4.7-6.1	143	2.2	1.9-2.6
1998	499	4.6	4.2-5.0	335	7.2	6.5-8.1	164	2.6	2.2-3.0	369	3.4	3.1-3.8	219	4.8	4.2-5.4	150	2.3	2.0-2.8
1999	489	4.5	4.1-4.9	320	6.9	6.1-7.7	169	2.7	2.3-3.1	413	3.8	3.4-4.2	250	5.4	4.7-6.1	163	2.5	2.1-2.9
2000	486	4.4	4.0-4.8	307	6.5	5.8-7.2	179	2.8	2.4-3.3	406	3.7	3.3-4.0	246	5.3	4.6-6.0	160	2.4	2.0-2.8
2001	508	4.6	4.2-5.0	359	7.5	6.8-8.4	149	2.3	2.0-2.7	414	3.7	3.4-4.1	269	5.7	5.1-6.5	145	2.2	1.8-2.6
2002	526	4.7	4.3-5.1	370	7.6	6.9-8.4	156	2.4	2.0-2.8	469	4.1	3.8-4.5	305	6.4	5.7-7.2	164	2.4	2.1-2.8
2003	531	4.7	4.3-5.2	340	6.8	6.1-7.6	191	3.0	2.6-3.5	411	3.6	3.3-4.0	261	5.4	4.7-6.0	150	2.2	1.9-2.6
2004	577	5.1	4.7-5.5	416	8.2	7.4-9.0	161	2.5	2.1-2.9	466	4.1	3.7-4.4	288	5.8	5.1-6.5	178	2.6	2.2-3.0
2005	612	5.4	4.9-5.8	426	8.3	7.5-9.1	186	2.9	2.5-3.4	481	4.1	3.8-4.5	313	6.2	5.5-7.0	168	2.5	2.1-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Liver / intrahepatic bile duct

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	624	5.4	4.9-5.8	456	8.7	7.9-9.6	168	2.6	2.2-3.0	505	4.3	3.9-4.7	307	6.1	5.4-6.8	198	2.9	2.5-3.4
2007	685	5.8	5.4-6.3	486	9.1	8.3-10.0	199	3.1	2.6-3.5	484	4.1	3.7-4.4	322	6.2	5.6-6.9	162	2.3	2.0-2.7
2008	738	6.2	5.7-6.6	519	9.5	8.7-10.4	219	3.3	2.9-3.8	581	4.8	4.4-5.2	373	7.0	6.3-7.8	208	3.0	2.6-3.4
2009	729	6.0	5.6-6.5	510	9.2	8.4-10.1	219	3.2	2.8-3.7	554	4.5	4.2-4.9	357	6.6	6.0-7.4	197	2.8	2.4-3.2
2010	692	5.6	5.2-6.0	525	9.3	8.5-10.1	167	2.5	2.1-2.9	554	4.4	4.1-4.8	348	6.3	5.7-7.0	206	3.0	2.6-3.4
2011	757	6.0	5.6-6.5	536	9.1	8.3-9.9	221	3.3	2.9-3.8	609	4.8	4.4-5.2	376	6.6	5.9-7.3	233	3.3	2.8-3.7
2012	828	6.4	6.0-6.9	610	10.2	9.4-11.1	218	3.2	2.8-3.7	650	5.1	4.7-5.5	423	7.4	6.7-8.1	227	3.2	2.8-3.7
2013	914	7.0	6.6-7.5	654	10.9	10.0-11.8	260	3.7	3.3-4.3	685	5.2	4.8-5.7	466	8.0	7.3-8.8	219	3.0	2.6-3.4
2014	949	7.2	6.8-7.7	693	11.4	10.5-12.3	256	3.7	3.2-4.2	635	4.8	4.4-5.2	438	7.2	6.6-8.0	197	2.8	2.4-3.2
2015	928	6.9	6.5-7.4	674	10.8	10.0-11.7	254	3.6	3.1-4.1	656	4.9	4.5-5.3	429	7.1	6.5-7.9	227	3.0	2.6-3.5
2016	1,005	7.5	7.0-8.0	728	11.5	10.7-12.5	277	3.9	3.5-4.5	690	5.1	4.7-5.5	455	7.3	6.7-8.1	235	3.2	2.8-3.6
2017	1,007	7.3	6.8-7.8	729	11.4	10.5-12.3	278	3.8	3.4-4.3	641	4.7	4.3-5.1	453	7.3	6.6-8.0	188	2.6	2.2-3.1
2018	926	6.7	6.2-7.1	659	10.2	9.4-11.1	267	3.7	3.2-4.2	675	4.8	4.5-5.2	455	7.2	6.5-7.9	220	2.9	2.5-3.3
2019	1,007	7.2	6.8-7.7	699	10.7	9.9-11.6	308	4.3	3.8-4.8	634	4.5	4.2-4.9	428	6.7	6.0-7.4	206	2.8	2.4-3.2
2020	963	6.8	6.4-7.3	659	10.1	9.3-10.9	304	4.1	3.6-4.6	669	4.7	4.4-5.1	430	6.6	6.0-7.3	239	3.1	2.7-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Pancreas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,052	11.8	11.1-12.6	566	15.1	13.8-16.5	486	9.5	8.7-10.4	1,023	11.5	10.8-12.3	521	14.1	12.9-15.5	502	9.7	8.9-10.6
1977	1,111	12.5	11.8-13.3	587	16.3	15.0-17.8	524	10.0	9.2-10.9	1,085	12.2	11.5-13.0	568	15.5	14.2-16.9	517	10.0	9.1-10.9
1978	1,115	12.3	11.6-13.1	575	15.2	13.9-16.5	540	10.2	9.4-11.1	1,086	11.9	11.2-12.7	565	15.0	13.7-16.4	521	9.8	9.0-10.7
1979	1,219	13.3	12.6-14.1	644	17.1	15.7-18.5	575	10.7	9.8-11.6	1,173	12.8	12.0-13.5	615	16.5	15.1-17.9	558	10.3	9.4-11.1
1980	1,169	12.5	11.8-13.3	576	14.8	13.6-16.2	593	10.9	10.0-11.8	1,100	11.9	11.2-12.6	554	14.6	13.4-16.0	546	10.0	9.1-10.9
1981	1,137	12.2	11.4-12.9	573	14.8	13.6-16.2	564	10.3	9.5-11.2	1,102	11.8	11.1-12.5	554	14.4	13.2-15.7	548	10.0	9.2-10.9
1982	1,130	11.9	11.2-12.6	554	14.3	13.1-15.6	576	10.2	9.4-11.1	1,037	10.9	10.2-11.6	533	13.8	12.6-15.0	504	8.9	8.2-9.8
1983	1,087	11.4	10.7-12.1	536	13.6	12.4-14.9	551	9.8	9.0-10.7	1,129	11.8	11.1-12.5	549	13.7	12.6-15.0	580	10.2	9.4-11.1
1984	1,130	11.7	11.0-12.4	559	14.4	13.1-15.7	571	9.9	9.1-10.8	1,089	11.3	10.6-12.0	524	13.6	12.4-14.8	565	9.8	9.0-10.7
1985	1,232	12.6	11.9-13.3	580	14.3	13.1-15.5	652	11.3	10.4-12.2	1,126	11.5	10.9-12.3	528	13.4	12.2-14.6	598	10.4	9.6-11.3
1986	1,167	12.0	11.3-12.7	574	14.2	13.0-15.5	593	10.3	9.4-11.1	1,133	11.6	10.9-12.3	560	13.7	12.5-14.9	573	9.8	9.0-10.7
1987	1,206	12.1	11.5-12.8	571	14.3	13.1-15.6	635	10.7	9.9-11.6	1,132	11.5	10.8-12.2	554	14.0	12.8-15.2	578	9.8	9.0-10.6
1988	1,246	12.5	11.9-13.3	607	15.0	13.8-16.3	639	10.7	9.9-11.6	1,180	11.8	11.1-12.5	568	14.2	13.0-15.4	612	10.2	9.4-11.0
1989	1,164	11.5	10.8-12.2	573	13.8	12.7-15.1	591	9.9	9.1-10.8	1,089	10.8	10.2-11.5	547	13.4	12.3-14.6	542	9.1	8.3-9.9
1990	1,232	12.2	11.5-12.9	592	14.3	13.1-15.6	640	10.6	9.8-11.4	1,144	11.3	10.6-11.9	544	13.1	12.0-14.3	600	9.9	9.1-10.7
1991	1,290	12.6	11.9-13.3	618	14.5	13.4-15.8	672	11.0	10.2-11.9	1,148	11.2	10.6-11.9	541	13.0	11.9-14.2	607	9.9	9.1-10.7
1992	1,241	12.0	11.4-12.7	591	14.1	12.9-15.3	650	10.6	9.8-11.5	1,153	11.1	10.5-11.8	536	12.9	11.8-14.0	617	9.9	9.1-10.7
1993	1,256	12.0	11.3-12.7	617	14.3	13.1-15.5	639	10.3	9.5-11.1	1,217	11.6	10.9-12.3	581	13.5	12.4-14.6	636	10.2	9.4-11.0
1994	1,345	12.7	12.0-13.4	672	15.3	14.2-16.6	673	10.6	9.8-11.4	1,256	11.9	11.2-12.5	624	14.5	13.3-15.7	632	9.9	9.1-10.7
1995	1,341	12.6	11.9-13.3	614	13.7	12.7-14.9	727	11.5	10.6-12.3	1,246	11.7	11.0-12.3	578	12.9	11.8-14.0	668	10.4	9.6-11.3
1996	1,276	12.0	11.3-12.6	647	14.6	13.5-15.8	629	10.0	9.2-10.8	1,192	11.1	10.5-11.8	576	13.2	12.1-14.4	616	9.6	8.9-10.4
1997	1,334	12.4	11.7-13.1	646	14.3	13.2-15.5	688	10.8	10.0-11.6	1,287	11.9	11.2-12.5	641	14.5	13.4-15.7	646	9.9	9.2-10.8
1998	1,412	13.0	12.3-13.7	677	14.8	13.7-16.0	735	11.5	10.7-12.4	1,249	11.4	10.8-12.1	614	13.7	12.6-14.8	635	9.8	9.0-10.6
1999	1,386	12.6	11.9-13.3	654	14.1	13.1-15.3	732	11.3	10.5-12.1	1,294	11.7	11.1-12.4	632	13.7	12.7-14.9	662	10.1	9.4-11.0
2000	1,392	12.5	11.9-13.2	655	14.1	13.0-15.2	737	11.3	10.5-12.2	1,277	11.4	10.8-12.1	594	13.0	12.0-14.1	683	10.3	9.5-11.1
2001	1,420	12.7	12.0-13.3	650	13.7	12.7-14.8	770	11.7	10.9-12.6	1,277	11.3	10.7-12.0	585	12.6	11.6-13.6	692	10.4	9.6-11.2
2002	1,491	13.2	12.5-13.9	701	14.6	13.5-15.7	790	12.0	11.2-12.9	1,319	11.6	11.0-12.3	633	13.2	12.2-14.3	686	10.2	9.5-11.0
2003	1,502	13.1	12.4-13.8	726	14.9	13.9-16.1	776	11.7	10.9-12.5	1,290	11.2	10.6-11.8	591	12.3	11.3-13.3	699	10.3	9.5-11.1
2004	1,479	12.8	12.1-13.5	718	14.6	13.5-15.7	761	11.4	10.6-12.3	1,309	11.3	10.7-11.9	648	13.2	12.2-14.3	661	9.8	9.1-10.6
2005	1,517	13.0	12.4-13.7	747	14.9	13.8-16.0	770	11.3	10.5-12.2	1,342	11.4	10.8-12.1	631	12.7	11.8-13.8	711	10.3	9.6-11.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Pancreas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,540	13.1	12.5-13.8	736	14.6	13.6-15.7	804	11.9	11.1-12.8	1,344	11.4	10.8-12.0	661	13.1	12.1-14.1	683	9.8	9.1-10.6
2007	1,607	13.5	12.9-14.2	785	15.3	14.2-16.4	822	12.0	11.2-12.9	1,344	11.2	10.6-11.8	660	12.9	11.9-14.0	684	9.9	9.2-10.7
2008	1,679	14.0	13.3-14.7	827	15.9	14.8-17.1	852	12.4	11.5-13.2	1,388	11.5	10.9-12.1	667	12.9	12.0-14.0	721	10.2	9.4-11.0
2009	1,679	13.7	13.1-14.4	871	16.4	15.3-17.5	808	11.6	10.8-12.5	1,395	11.3	10.7-11.9	698	13.2	12.2-14.2	697	9.9	9.1-10.6
2010	1,722	14.0	13.3-14.7	825	15.3	14.3-16.4	897	12.8	12.0-13.7	1,441	11.4	10.9-12.1	702	12.9	12.0-13.9	739	10.2	9.5-11.0
2011	1,755	14.0	13.4-14.7	894	16.2	15.2-17.4	861	12.1	11.3-13.0	1,411	11.1	10.6-11.7	688	12.6	11.7-13.6	723	9.9	9.2-10.7
2012	1,780	14.1	13.4-14.8	916	16.6	15.5-17.7	864	12.1	11.3-13.0	1,479	11.7	11.1-12.3	771	14.0	13.0-15.1	708	9.7	9.0-10.5
2013	1,830	14.2	13.6-14.9	920	16.2	15.2-17.3	910	12.5	11.7-13.4	1,512	11.7	11.1-12.3	757	13.4	12.5-14.4	755	10.2	9.5-11.0
2014	1,924	14.8	14.2-15.5	985	16.9	15.8-18.0	939	13.1	12.2-14.0	1,499	11.4	10.8-12.0	766	13.3	12.3-14.3	733	9.8	9.1-10.6
2015	1,884	14.3	13.6-15.0	941	15.9	14.9-17.0	943	12.9	12.1-13.8	1,442	10.8	10.3-11.4	734	12.6	11.7-13.5	708	9.3	8.6-10.0
2016	1,998	15.3	14.6-16.0	1,035	17.5	16.4-18.6	963	13.5	12.6-14.4	1,533	11.4	10.9-12.0	757	12.7	11.8-13.7	776	10.4	9.6-11.2
2017	2,052	15.2	14.5-15.9	1,080	17.8	16.7-18.9	972	13.1	12.2-14.0	1,555	11.4	10.8-12.0	791	13.1	12.2-14.1	764	10.0	9.3-10.8
2018	2,037	15.0	14.4-15.7	1,042	17.0	16.0-18.1	995	13.3	12.5-14.2	1,569	11.3	10.7-11.9	801	13.0	12.1-13.9	768	9.9	9.2-10.7
2019	2,124	15.4	14.8-16.1	1,091	17.6	16.6-18.7	1,033	13.6	12.7-14.5	1,596	11.5	10.9-12.0	837	13.5	12.5-14.4	759	9.7	9.0-10.4
2020	2,050	14.7	14.0-15.4	1,088	17.2	16.1-18.2	962	12.5	11.7-13.4	1,599	11.4	10.8-12.0	812	12.9	12.0-13.9	787	10.1	9.3-10.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	504	5.2	4.8-5.7	432	10.4	9.4-11.5	72	1.3	1.0-1.7	175	1.9	1.6-2.2	145	3.8	3.1-4.5	30	0.6	0.4-0.8
1977	510	5.3	4.8-5.8	417	10.1	9.1-11.2	93	1.7	1.4-2.1	169	1.8	1.5-2.1	141	3.6	3.0-4.3	28	0.5	0.3-0.7
1978	548	5.6	5.1-6.1	458	10.9	9.9-12.0	90	1.7	1.3-2.0	185	1.9	1.7-2.2	145	3.7	3.1-4.4	40	0.7	0.5-1.0
1979	567	5.7	5.2-6.2	445	10.3	9.3-11.4	122	2.2	1.8-2.7	209	2.2	1.9-2.5	170	4.2	3.6-5.0	39	0.7	0.5-1.0
1980	578	5.9	5.4-6.4	473	11.2	10.1-12.3	105	2.0	1.6-2.4	184	1.9	1.6-2.2	148	3.7	3.1-4.4	36	0.7	0.5-0.9
1981	629	6.3	5.8-6.9	514	12.0	10.9-13.1	115	2.1	1.7-2.5	184	1.8	1.6-2.1	155	3.9	3.2-4.6	29	0.5	0.3-0.7
1982	562	5.7	5.2-6.2	453	10.5	9.5-11.6	109	2.0	1.6-2.4	191	1.9	1.6-2.2	152	3.6	3.0-4.3	39	0.7	0.5-0.9
1983	570	5.8	5.3-6.3	457	10.6	9.6-11.6	113	2.1	1.8-2.6	183	1.8	1.6-2.1	139	3.4	2.8-4.0	44	0.8	0.6-1.0
1984	572	5.7	5.2-6.2	449	10.2	9.3-11.3	123	2.3	1.9-2.7	162	1.6	1.4-1.9	126	3.1	2.5-3.7	36	0.6	0.4-0.9
1985	588	5.8	5.4-6.3	465	10.5	9.5-11.5	123	2.2	1.9-2.7	178	1.7	1.5-2.0	137	3.2	2.6-3.8	41	0.7	0.5-1.0
1986	572	5.7	5.2-6.2	442	10.2	9.2-11.2	130	2.3	1.9-2.8	167	1.7	1.4-2.0	132	3.2	2.7-3.8	35	0.6	0.4-0.8
1987	609	6.0	5.5-6.5	487	11.2	10.2-12.3	122	2.1	1.8-2.6	176	1.7	1.5-2.0	139	3.4	2.8-4.0	37	0.6	0.4-0.9
1988	544	5.3	4.9-5.8	414	9.3	8.4-10.3	130	2.3	1.9-2.8	172	1.7	1.5-2.0	136	3.3	2.7-3.9	36	0.6	0.4-0.9
1989	607	6.0	5.5-6.5	471	10.7	9.7-11.8	136	2.4	2.0-2.9	171	1.7	1.4-1.9	118	2.7	2.2-3.3	53	0.9	0.7-1.2
1990	609	5.9	5.4-6.4	456	9.9	9.0-10.9	153	2.7	2.3-3.2	153	1.5	1.3-1.8	121	2.9	2.4-3.5	32	0.5	0.4-0.8
1991	552	5.4	4.9-5.9	441	9.8	8.9-10.7	111	1.9	1.6-2.3	159	1.5	1.3-1.8	123	2.7	2.3-3.3	36	0.6	0.4-0.8
1992	565	5.5	5.0-6.0	430	9.5	8.6-10.4	135	2.4	2.0-2.8	205	1.9	1.7-2.2	157	3.6	3.1-4.3	48	0.8	0.6-1.1
1993	543	5.2	4.8-5.7	394	8.8	7.9-9.7	149	2.6	2.2-3.0	168	1.6	1.4-1.9	131	3.0	2.5-3.6	37	0.6	0.4-0.9
1994	525	5.0	4.6-5.5	400	8.8	7.9-9.7	125	2.2	1.8-2.6	169	1.6	1.3-1.8	127	2.9	2.4-3.5	42	0.7	0.5-0.9
1995	544	5.2	4.8-5.7	413	8.9	8.1-9.9	131	2.3	1.9-2.8	165	1.6	1.3-1.8	127	2.9	2.4-3.4	38	0.6	0.4-0.9
1996	482	4.6	4.2-5.0	371	8.0	7.2-8.8	111	2.0	1.6-2.4	165	1.6	1.3-1.8	122	2.7	2.2-3.2	43	0.7	0.5-1.0
1997	493	4.7	4.3-5.2	382	8.2	7.4-9.1	111	2.0	1.6-2.4	141	1.3	1.1-1.6	104	2.3	1.9-2.8	37	0.6	0.4-0.9
1998	467	4.4	4.0-4.8	337	7.1	6.4-8.0	130	2.2	1.9-2.7	151	1.4	1.2-1.6	112	2.4	2.0-2.9	39	0.6	0.4-0.9
1999	522	4.9	4.5-5.3	415	8.6	7.8-9.5	107	1.8	1.5-2.2	155	1.4	1.2-1.7	116	2.5	2.0-3.0	39	0.6	0.5-0.9
2000	500	4.6	4.2-5.0	392	8.2	7.4-9.0	108	1.8	1.5-2.2	140	1.3	1.1-1.5	113	2.4	2.0-2.9	27	0.4	0.3-0.7
2001	471	4.3	3.9-4.7	367	7.5	6.7-8.3	104	1.7	1.4-2.1	138	1.2	1.0-1.5	115	2.4	2.0-2.9	23	0.4	0.2-0.5
2002	463	4.2	3.8-4.6	360	7.4	6.6-8.2	103	1.8	1.4-2.1	146	1.3	1.1-1.5	115	2.4	2.0-2.9	31	0.5	0.3-0.7
2003	450	4.0	3.7-4.4	350	7.0	6.2-7.7	100	1.6	1.3-2.0	129	1.1	0.9-1.3	96	2.0	1.6-2.4	33	0.5	0.3-0.7
2004	453	4.0	3.6-4.4	363	7.1	6.3-7.8	90	1.5	1.2-1.8	128	1.1	0.9-1.3	95	2.0	1.6-2.4	33	0.5	0.3-0.7
2005	510	4.4	4.1-4.8	397	7.6	6.9-8.4	113	1.8	1.5-2.2	144	1.3	1.1-1.5	103	2.1	1.7-2.5	41	0.7	0.5-0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	486	4.2	3.9-4.6	365	7.1	6.4-7.9	121	2.0	1.6-2.3	139	1.2	1.0-1.4	103	2.0	1.7-2.5	36	0.6	0.4-0.8
2007	440	3.7	3.4-4.1	353	6.7	6.0-7.4	87	1.3	1.1-1.6	126	1.0	0.9-1.3	106	2.0	1.6-2.4	20	0.3	0.2-0.5
2008	494	4.2	3.8-4.6	390	7.2	6.5-8.0	104	1.6	1.3-2.0	112	0.9	0.8-1.1	87	1.6	1.3-2.0	25	0.4	0.2-0.6
2009	456	3.8	3.5-4.2	354	6.6	5.9-7.3	102	1.6	1.3-1.9	128	1.0	0.9-1.2	93	1.8	1.4-2.2	35	0.5	0.3-0.7
2010	437	3.6	3.3-4.0	342	6.2	5.5-6.9	95	1.5	1.2-1.8	142	1.1	1.0-1.4	106	2.0	1.6-2.4	36	0.5	0.4-0.7
2011	441	3.5	3.2-3.9	358	6.2	5.5-6.9	83	1.3	1.0-1.6	132	1.1	0.9-1.3	96	1.7	1.4-2.1	36	0.5	0.4-0.7
2012	455	3.6	3.3-4.0	356	6.3	5.6-7.0	99	1.4	1.2-1.8	117	0.9	0.8-1.1	91	1.6	1.3-2.0	26	0.4	0.2-0.6
2013	470	3.6	3.3-4.0	373	6.2	5.6-6.9	97	1.5	1.2-1.8	153	1.1	1.0-1.3	124	2.1	1.7-2.5	29	0.4	0.3-0.6
2014	443	3.4	3.1-3.7	347	5.7	5.1-6.4	96	1.4	1.1-1.7	137	1.1	0.9-1.3	108	1.8	1.5-2.2	29	0.4	0.3-0.6
2015	465	3.6	3.2-3.9	370	6.1	5.5-6.8	95	1.4	1.1-1.7	114	0.8	0.7-1.0	87	1.5	1.2-1.8	27	0.4	0.2-0.5
2016	418	3.2	2.9-3.5	336	5.6	5.0-6.2	82	1.2	0.9-1.5	125	1.0	0.8-1.2	100	1.7	1.4-2.1	25	0.4	0.2-0.5
2017	442	3.3	3.0-3.6	338	5.4	4.9-6.1	104	1.5	1.2-1.9	109	0.8	0.6-0.9	84	1.3	1.0-1.6	25	0.3	0.2-0.5
2018	428	3.1	2.8-3.4	336	5.2	4.7-5.9	92	1.3	1.0-1.6	125	0.9	0.7-1.1	93	1.5	1.2-1.8	32	0.4	0.3-0.6
2019	413	3.0	2.7-3.3	334	5.1	4.5-5.7	79	1.1	0.9-1.5	102	0.7	0.6-0.9	81	1.2	1.0-1.6	21	0.3	0.2-0.4
2020	373	2.7	2.4-3.0	296	4.6	4.0-5.1	77	1.1	0.8-1.3	100	0.7	0.6-0.9	80	1.3	1.0-1.6	20	0.2	0.1-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Lung and bronchus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	5,325	57.3	55.7-58.9	3,856	98.8	95.5-102.1	1,469	28.3	26.8-29.8	4,289	46.5	45.1-47.9	3,163	82.6	79.6-85.7	1,126	21.7	20.4-23.0
1977	5,339	56.7	55.2-58.3	3,810	96.5	93.3-99.8	1,529	29.3	27.8-30.8	4,533	48.6	47.2-50.1	3,304	85.3	82.2-88.4	1,229	23.4	22.1-24.8
1978	5,468	57.5	56.0-59.1	3,921	98.7	95.4-101.9	1,547	29.1	27.7-30.6	4,552	48.0	46.6-49.5	3,282	83.7	80.8-86.8	1,270	23.7	22.4-25.1
1979	5,626	58.8	57.3-60.4	3,902	97.7	94.5-101.0	1,724	32.3	30.8-33.9	4,516	47.3	45.9-48.7	3,191	81.0	78.0-84.0	1,325	24.5	23.2-25.9
1980	5,957	61.3	59.8-63.0	4,167	102.4	99.1-105.7	1,790	32.9	31.4-34.5	4,892	50.6	49.2-52.1	3,484	86.9	83.9-90.0	1,408	25.7	24.4-27.1
1981	6,013	61.8	60.2-63.4	4,099	101.1	97.9-104.4	1,914	35.3	33.7-36.9	4,939	50.8	49.4-52.3	3,483	87.0	84.0-90.1	1,456	26.6	25.2-28.0
1982	6,254	63.6	62.0-65.2	4,206	102.7	99.4-106.0	2,048	37.2	35.6-38.9	5,074	52.0	50.6-53.5	3,425	85.4	82.4-88.4	1,649	29.9	28.4-31.4
1983	6,542	66.1	64.5-67.7	4,286	103.4	100.2-106.7	2,256	40.6	38.9-42.3	5,253	53.2	51.8-54.7	3,531	87.2	84.2-90.3	1,722	30.8	29.3-32.3
1984	6,490	65.3	63.7-66.9	4,211	100.7	97.5-103.9	2,279	41.2	39.5-42.9	5,249	52.9	51.4-54.3	3,460	83.9	81.0-86.8	1,789	31.9	30.4-33.5
1985	6,810	68.2	66.5-69.8	4,343	103.6	100.4-106.9	2,467	44.4	42.6-46.2	5,592	56.1	54.6-57.6	3,649	88.7	85.7-91.8	1,943	34.4	32.9-36.0
1986	6,763	67.7	66.1-69.3	4,268	102.1	99.0-105.4	2,495	44.6	42.9-46.5	5,625	56.2	54.7-57.7	3,612	87.6	84.7-90.7	2,013	35.7	34.2-37.4
1987	6,953	69.0	67.4-70.7	4,321	102.0	98.9-105.3	2,632	46.7	44.9-48.6	5,603	55.7	54.3-57.2	3,586	86.5	83.6-89.5	2,017	35.2	33.7-36.9
1988	7,139	70.2	68.5-71.9	4,359	102.7	99.6-105.9	2,780	48.7	46.9-50.6	5,792	57.0	55.6-58.6	3,619	86.4	83.5-89.3	2,173	37.8	36.2-39.5
1989	7,258	70.9	69.3-72.6	4,361	101.2	98.1-104.4	2,897	50.5	48.7-52.5	5,851	57.3	55.8-58.8	3,613	85.0	82.1-87.9	2,238	38.8	37.1-40.4
1990	7,309	70.7	69.1-72.4	4,413	101.9	98.8-105.0	2,896	49.8	48.0-51.7	5,988	58.0	56.5-59.5	3,619	84.5	81.7-87.5	2,369	40.2	38.6-41.9
1991	7,412	71.5	69.9-73.2	4,339	99.6	96.5-102.7	3,073	52.7	50.8-54.6	6,180	59.5	58.1-61.1	3,720	86.5	83.7-89.4	2,460	41.7	40.0-43.4
1992	7,461	71.3	69.7-73.0	4,290	97.3	94.3-100.3	3,171	53.8	51.9-55.7	6,226	59.5	58.0-61.0	3,616	83.7	80.9-86.6	2,610	43.5	41.9-45.3
1993	7,466	71.0	69.3-72.6	4,184	94.0	91.1-97.0	3,282	55.4	53.5-57.4	6,038	57.3	55.8-58.7	3,488	79.6	76.9-82.4	2,550	42.5	40.9-44.2
1994	7,799	73.7	72.0-75.3	4,370	97.8	94.8-100.8	3,429	57.7	55.8-59.7	6,240	58.7	57.3-60.2	3,656	83.1	80.4-85.9	2,584	42.5	40.9-44.2
1995	7,754	72.9	71.3-74.5	4,339	95.9	93.0-98.8	3,415	57.1	55.1-59.0	6,246	58.6	57.1-60.0	3,598	81.1	78.4-83.9	2,648	43.6	41.9-45.3
1996	7,862	73.8	72.1-75.4	4,403	96.9	94.1-99.9	3,459	57.8	55.8-59.7	6,137	57.4	56.0-58.9	3,501	78.3	75.7-81.0	2,636	43.4	41.7-45.1
1997	7,908	73.7	72.1-75.3	4,304	94.1	91.3-97.0	3,604	59.9	57.9-61.9	6,037	56.0	54.6-57.4	3,358	74.5	71.9-77.1	2,679	43.5	41.8-45.2
1998	8,027	74.1	72.5-75.8	4,311	92.6	89.8-95.4	3,716	61.5	59.5-63.5	6,052	55.6	54.2-57.0	3,400	73.8	71.3-76.4	2,652	42.8	41.1-44.4
1999	8,106	74.3	72.6-75.9	4,359	93.2	90.4-96.0	3,747	61.3	59.3-63.3	6,019	54.9	53.5-56.3	3,295	71.3	68.9-73.8	2,724	43.7	42.1-45.4
2000	8,260	74.9	73.3-76.6	4,361	92.3	89.6-95.1	3,899	63.0	61.0-65.0	6,180	55.7	54.3-57.1	3,362	71.9	69.5-74.4	2,818	44.6	42.9-46.3
2001	8,069	72.8	71.2-74.4	4,154	87.5	84.8-90.2	3,915	63.2	61.2-65.2	6,077	54.4	53.0-55.8	3,301	70.6	68.2-73.0	2,776	43.8	42.2-45.5
2002	8,287	73.9	72.3-75.5	4,355	90.5	87.8-93.3	3,932	62.9	60.9-64.9	6,199	54.9	53.6-56.3	3,294	69.2	66.8-71.6	2,905	45.6	43.9-47.3
2003	8,367	74.0	72.4-75.6	4,379	89.6	87.0-92.3	3,988	63.3	61.4-65.3	6,013	52.8	51.4-54.1	3,206	66.4	64.1-68.7	2,807	43.5	41.9-45.2
2004	8,388	73.5	71.9-75.1	4,230	85.6	83.0-88.2	4,158	65.3	63.3-67.3	6,029	52.5	51.2-53.8	3,089	63.1	60.9-65.4	2,940	45.2	43.6-46.9
2005	8,416	73.3	71.8-74.9	4,265	85.6	83.0-88.2	4,151	64.8	62.8-66.8	5,985	51.8	50.5-53.2	3,118	63.4	61.2-65.7	2,867	43.7	42.1-45.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Lung and bronchus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	8,657	75.2	73.6-76.8	4,241	84.2	81.7-86.8	4,416	69.3	67.2-71.4	6,027	51.9	50.6-53.2	3,176	63.7	61.5-66.0	2,851	43.7	42.1-45.4
2007	8,597	73.5	71.9-75.1	4,310	84.3	81.8-86.9	4,287	66.2	64.2-68.3	6,023	51.1	49.8-52.4	3,098	60.9	58.8-63.1	2,925	44.2	42.6-45.9
2008	8,883	75.0	73.5-76.6	4,426	85.4	82.9-88.0	4,457	68.1	66.1-70.2	5,986	49.9	48.6-51.2	3,069	59.5	57.4-61.6	2,917	43.4	41.8-45.1
2009	8,737	72.7	71.2-74.3	4,440	84.1	81.6-86.6	4,297	64.7	62.7-66.7	5,818	48.2	47.0-49.5	3,037	57.9	55.9-60.1	2,781	41.3	39.8-42.9
2010	8,641	71.1	69.6-72.7	4,245	78.8	76.4-81.2	4,396	66.2	64.2-68.3	5,772	47.2	46.0-48.5	2,928	54.9	52.9-56.9	2,844	41.8	40.3-43.4
2011	8,802	71.4	69.9-72.9	4,345	79.5	77.1-81.9	4,457	65.9	63.9-67.9	5,740	46.3	45.1-47.5	2,921	54.0	52.1-56.1	2,819	40.8	39.3-42.4
2012	8,713	69.6	68.1-71.1	4,297	77.4	75.1-79.8	4,416	64.3	62.4-66.3	5,918	47.1	45.9-48.4	3,071	56.0	54.0-58.0	2,847	40.8	39.3-42.4
2013	8,799	69.4	67.9-70.9	4,280	75.5	73.2-77.9	4,519	65.1	63.2-67.1	5,655	44.1	42.9-45.3	2,863	50.6	48.8-52.6	2,792	39.3	37.8-40.8
2014	8,564	66.2	64.8-67.7	4,174	72.4	70.2-74.7	4,390	62.0	60.1-63.9	5,434	42.0	40.9-43.1	2,783	48.8	47.0-50.7	2,651	37.0	35.6-38.5
2015	9,053	69.4	67.9-70.8	4,299	73.7	71.5-76.0	4,754	66.8	64.9-68.8	5,420	41.2	40.1-42.3	2,755	47.5	45.8-49.4	2,665	36.5	35.1-38.0
2016	8,887	66.8	65.4-68.3	4,288	71.8	69.7-74.1	4,599	63.8	61.9-65.7	5,121	38.5	37.4-39.6	2,573	43.9	42.1-45.6	2,548	34.5	33.2-35.9
2017	9,235	68.5	67.1-69.9	4,350	71.5	69.3-73.7	4,885	66.8	64.9-68.7	4,921	36.2	35.1-37.2	2,460	40.9	39.2-42.5	2,461	32.7	31.3-34.0
2018	9,105	66.3	65.0-67.8	4,301	69.3	67.2-71.5	4,804	64.6	62.7-66.5	4,915	35.7	34.7-36.7	2,477	40.5	38.9-42.2	2,438	32.2	30.9-33.5
2019	8,988	64.3	62.9-65.6	4,206	66.5	64.5-68.6	4,782	63.2	61.4-65.1	4,766	33.8	32.8-34.8	2,365	37.9	36.3-39.5	2,401	30.9	29.6-32.2
2020	8,049	56.8	55.5-58.1	3,728	58.0	56.1-59.9	4,321	56.3	54.6-58.0	4,447	31.1	30.2-32.1	2,225	35.0	33.5-36.5	2,222	28.4	27.2-29.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Melanoma of the skin

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	536	6.0	5.5-6.5	274	6.6	5.9-7.5	262	5.5	4.8-6.2	233	2.6	2.3-3.0	130	3.2	2.7-3.9	103	2.1	1.7-2.6
1977	689	7.6	7.0-8.2	373	9.0	8.1-10.0	316	6.6	5.8-7.3	246	2.8	2.4-3.2	159	4.0	3.4-4.7	87	1.8	1.5-2.3
1978	657	7.3	6.7-7.9	335	8.2	7.3-9.2	322	6.8	6.0-7.6	219	2.5	2.1-2.8	139	3.5	2.9-4.1	80	1.6	1.3-2.0
1979	676	7.5	6.9-8.1	361	8.8	7.9-9.8	315	6.5	5.8-7.3	236	2.6	2.3-3.0	133	3.3	2.8-3.9	103	2.1	1.7-2.5
1980	832	9.1	8.5-9.7	458	11.2	10.2-12.3	374	7.6	6.8-8.4	251	2.7	2.4-3.1	152	3.6	3.1-4.3	99	2.0	1.6-2.5
1981	809	8.8	8.2-9.4	423	10.2	9.2-11.3	386	7.8	7.0-8.6	223	2.4	2.1-2.7	132	3.2	2.6-3.8	91	1.8	1.4-2.2
1982	846	9.0	8.4-9.7	412	9.8	8.8-10.8	434	8.6	7.8-9.5	253	2.7	2.4-3.1	157	3.8	3.2-4.5	96	1.8	1.5-2.3
1983	916	9.8	9.2-10.5	506	12.1	11.1-13.3	410	8.4	7.6-9.2	244	2.6	2.3-2.9	136	3.2	2.7-3.9	108	2.2	1.8-2.6
1984	818	8.7	8.1-9.3	447	10.4	9.4-11.5	371	7.4	6.7-8.3	254	2.7	2.4-3.0	144	3.5	2.9-4.1	110	2.1	1.7-2.6
1985	915	9.7	9.1-10.4	491	11.5	10.5-12.6	424	8.4	7.6-9.3	252	2.6	2.3-3.0	151	3.7	3.1-4.3	101	1.9	1.5-2.3
1986	976	10.3	9.6-10.9	525	12.3	11.3-13.5	451	9.0	8.2-9.9	273	2.9	2.5-3.3	158	3.8	3.2-4.5	115	2.2	1.8-2.7
1987	914	9.5	8.9-10.2	501	11.6	10.6-12.7	413	8.1	7.3-8.9	297	3.1	2.8-3.5	179	4.2	3.6-4.9	118	2.3	1.9-2.8
1988	880	9.1	8.5-9.7	503	11.7	10.7-12.8	377	7.2	6.5-8.0	291	3.0	2.6-3.3	187	4.4	3.8-5.1	104	1.9	1.5-2.3
1989	929	9.6	9.0-10.2	524	12.0	11.0-13.1	405	7.7	7.0-8.6	312	3.2	2.9-3.6	199	4.8	4.1-5.6	113	2.1	1.7-2.5
1990	970	9.7	9.1-10.4	521	11.7	10.7-12.7	449	8.5	7.7-9.3	297	3.0	2.7-3.4	181	4.2	3.6-4.9	116	2.2	1.8-2.6
1991	896	9.0	8.4-9.6	521	11.7	10.7-12.8	375	7.0	6.3-7.7	298	3.0	2.6-3.3	187	4.4	3.7-5.0	111	2.0	1.6-2.4
1992	928	9.2	8.6-9.8	541	12.1	11.1-13.2	387	7.1	6.4-7.8	295	2.9	2.6-3.3	172	3.9	3.4-4.6	123	2.2	1.8-2.6
1993	931	9.1	8.5-9.7	533	11.6	10.7-12.7	398	7.2	6.5-8.0	290	2.8	2.5-3.2	183	4.2	3.6-4.9	107	1.9	1.5-2.3
1994	1,018	10.0	9.4-10.6	597	13.1	12.0-14.2	421	7.7	7.0-8.5	294	2.9	2.5-3.2	185	4.2	3.6-4.8	109	1.9	1.5-2.3
1995	1,114	10.9	10.3-11.6	666	14.5	13.4-15.6	448	8.2	7.4-9.0	311	3.0	2.7-3.4	183	4.2	3.6-4.8	128	2.2	1.8-2.6
1996	1,193	11.6	11.0-12.3	685	14.9	13.8-16.0	508	9.2	8.4-10.1	319	3.1	2.7-3.4	197	4.3	3.7-4.9	122	2.1	1.8-2.6
1997	1,278	12.3	11.7-13.0	752	16.1	15.0-17.4	526	9.5	8.7-10.4	303	2.9	2.6-3.2	204	4.6	3.9-5.2	99	1.6	1.3-2.0
1998	1,361	13.1	12.4-13.8	777	16.5	15.4-17.8	584	10.7	9.8-11.6	292	2.7	2.4-3.1	187	4.0	3.5-4.6	105	1.8	1.4-2.1
1999	1,401	13.3	12.6-14.1	811	17.1	16.0-18.4	590	10.5	9.7-11.4	284	2.6	2.3-3.0	167	3.6	3.1-4.2	117	2.0	1.6-2.4
2000	1,558	14.7	14.0-15.5	895	18.8	17.6-20.1	663	11.9	11.0-12.8	304	2.8	2.5-3.1	196	4.1	3.5-4.7	108	1.8	1.5-2.2
2001	1,729	16.2	15.5-17.0	974	20.2	19.0-21.5	755	13.4	12.5-14.4	356	3.2	2.9-3.6	229	4.8	4.2-5.5	127	2.1	1.7-2.5
2002	1,708	15.9	15.1-16.6	988	20.3	19.0-21.6	720	12.7	11.8-13.7	313	2.8	2.5-3.2	168	3.5	3.0-4.0	145	2.4	2.0-2.8
2003	1,774	16.3	15.5-17.1	1,037	21.1	19.8-22.4	737	12.8	11.9-13.8	312	2.8	2.5-3.1	191	3.9	3.4-4.5	121	1.9	1.6-2.3
2004	2,112	19.3	18.5-20.2	1,215	24.4	23.0-25.8	897	15.7	14.7-16.8	322	2.8	2.5-3.2	197	4.0	3.4-4.6	125	2.0	1.7-2.4
2005	2,315	21.2	20.3-22.1	1,355	27.1	25.7-28.6	960	16.8	15.8-18.0	335	2.9	2.6-3.3	209	4.2	3.6-4.8	126	2.0	1.7-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Melanoma of the skin

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	2,487	22.5	21.6-23.4	1,433	28.3	26.8-29.8	1,054	18.4	17.3-19.6	340	3.0	2.7-3.3	221	4.4	3.8-5.0	119	1.8	1.5-2.2
2007	2,408	21.7	20.8-22.6	1,392	27.3	25.9-28.8	1,016	17.8	16.7-19.0	327	2.8	2.5-3.1	235	4.6	4.1-5.3	92	1.4	1.1-1.8
2008	2,639	23.5	22.6-24.5	1,525	29.6	28.2-31.2	1,114	19.3	18.2-20.6	345	2.9	2.6-3.3	228	4.4	3.8-5.0	117	1.8	1.5-2.2
2009	2,814	24.8	23.9-25.7	1,653	31.6	30.1-33.2	1,161	20.0	18.9-21.3	350	3.0	2.7-3.3	206	4.0	3.5-4.6	144	2.2	1.9-2.6
2010	2,698	23.6	22.7-24.6	1,547	29.3	27.8-30.8	1,151	19.5	18.4-20.8	347	2.9	2.6-3.2	212	4.0	3.5-4.6	135	2.1	1.7-2.5
2011	3,000	25.6	24.6-26.5	1,753	32.4	30.9-34.0	1,247	20.6	19.4-21.9	368	3.0	2.7-3.4	241	4.5	3.9-5.1	127	1.9	1.6-2.3
2012	2,923	24.9	24.0-25.9	1,660	30.3	28.8-31.8	1,263	21.3	20.1-22.5	394	3.2	2.8-3.5	250	4.7	4.1-5.3	144	2.0	1.7-2.4
2013	3,246	27.5	26.5-28.5	1,868	33.5	32.0-35.1	1,378	23.4	22.1-24.7	379	3.1	2.8-3.4	252	4.7	4.1-5.3	127	1.8	1.5-2.2
2014	3,226	27.2	26.2-28.2	1,891	33.8	32.3-35.4	1,335	22.3	21.1-23.6	336	2.6	2.4-3.0	217	3.8	3.3-4.4	119	1.7	1.4-2.0
2015	3,186	26.2	25.3-27.2	1,883	33.1	31.6-34.7	1,303	21.4	20.2-22.6	293	2.3	2.0-2.6	193	3.4	2.9-3.9	100	1.4	1.2-1.8
2016	3,212	26.2	25.3-27.2	1,908	33.0	31.5-34.5	1,304	21.4	20.2-22.7	322	2.4	2.2-2.7	205	3.6	3.1-4.1	117	1.6	1.3-1.9
2017	3,290	26.9	25.9-27.9	1,947	33.9	32.3-35.5	1,343	22.0	20.7-23.3	290	2.2	2.0-2.5	188	3.3	2.8-3.8	102	1.4	1.1-1.7
2018	3,302	26.3	25.4-27.3	1,908	32.2	30.7-33.7	1,394	22.2	20.9-23.5	308	2.2	2.0-2.5	186	3.0	2.6-3.5	122	1.6	1.3-1.9
2019	3,369	26.8	25.8-27.8	1,982	33.0	31.5-34.5	1,387	22.5	21.2-23.8	273	2.1	1.8-2.3	173	3.0	2.5-3.5	100	1.4	1.1-1.7
2020	2,911	23.0	22.1-23.9	1,651	27.4	26.1-28.8	1,260	19.9	18.7-21.1	302	2.3	2.0-2.6	204	3.5	3.0-4.1	98	1.3	1.0-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Female breast

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	5,064	102.9	100.0-105.8	1,957	38.9	37.2-40.7
1977	5,094	102.5	99.7-105.5	2,012	39.7	37.9-41.5
1978	5,213	104.0	101.1-106.9	2,027	39.2	37.5-41.0
1979	5,169	102.1	99.3-105.0	1,997	38.6	36.9-40.4
1980	5,333	104.3	101.4-107.2	1,986	37.7	36.0-39.4
1981	5,386	105.0	102.1-107.9	1,994	37.5	35.8-39.2
1982	5,506	105.9	103.1-108.9	2,022	37.7	36.1-39.5
1983	5,878	112.9	110.0-115.9	2,059	38.2	36.6-40.0
1984	6,032	114.8	111.8-117.8	2,126	39.1	37.4-40.8
1985	6,384	121.9	118.8-125.0	2,162	39.8	38.1-41.6
1986	6,553	123.9	120.8-127.0	2,195	39.9	38.2-41.6
1987	6,776	127.1	124.0-130.3	2,196	39.4	37.7-41.1
1988	6,900	128.3	125.2-131.5	2,165	38.9	37.3-40.6
1989	6,769	125.3	122.3-128.4	2,158	38.7	37.0-40.4
1990	7,175	131.7	128.6-134.8	2,125	37.3	35.7-39.0
1991	7,082	128.7	125.7-131.8	2,054	36.0	34.4-37.6
1992	7,127	128.9	125.9-132.0	2,043	35.3	33.8-36.9
1993	7,148	128.3	125.3-131.4	2,026	34.5	33.0-36.0
1994	7,237	129.7	126.7-132.7	2,020	34.3	32.8-35.9
1995	7,591	136.0	133.0-139.2	1,933	32.8	31.4-34.4
1996	8,043	143.3	140.1-146.5	1,962	32.7	31.2-34.2
1997	8,197	144.7	141.5-147.9	1,846	30.6	29.2-32.0
1998	8,472	148.1	145.0-151.3	1,864	30.4	29.1-31.9
1999	8,377	145.0	141.9-148.2	1,740	28.0	26.7-29.4
2000	8,456	144.5	141.4-147.6	1,759	28.0	26.7-29.3
2001	8,310	141.3	138.2-144.4	1,708	27.1	25.8-28.4
2002	8,308	140.3	137.2-143.4	1,684	26.3	25.0-27.6
2003	8,253	137.9	134.9-141.0	1,678	26.0	24.8-27.3
2004	8,117	134.9	131.9-137.9	1,535	23.6	22.4-24.8
2005	8,252	135.9	132.9-138.9	1,506	23.0	21.8-24.2
2006	8,363	138.2	135.2-141.3	1,444	22.0	20.9-23.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
White Individuals
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	8,349	136.4	133.4-139.4	1,504	22.6	21.5-23.8
2008	8,675	140.9	137.9-144.0	1,435	21.4	20.3-22.6
2009	8,876	142.0	138.9-145.0	1,440	21.1	20.0-22.3
2010	8,308	132.6	129.6-135.5	1,493	21.5	20.4-22.7
2011	8,764	139.2	136.2-142.2	1,453	20.8	19.7-22.0
2012	8,677	137.3	134.3-140.3	1,395	19.8	18.7-20.9
2013	8,966	141.2	138.2-144.3	1,397	19.9	18.8-21.0
2014	9,166	143.7	140.6-146.8	1,246	17.5	16.5-18.5
2015	9,243	144.7	141.6-147.8	1,317	18.3	17.3-19.4
2016	8,916	139.0	136.0-142.1	1,376	19.4	18.3-20.5
2017	9,176	143.0	139.9-146.1	1,289	17.6	16.6-18.6
2018	9,283	143.3	140.2-146.4	1,361	18.6	17.6-19.7
2019	9,592	148.4	145.3-151.6	1,272	17.4	16.4-18.5
2020	8,502	131.3	128.4-134.4	1,299	17.6	16.6-18.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Cervix uteri

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	600	12.5	11.5-13.5	234	4.6	4.1-5.3
1977	523	10.7	9.8-11.7	191	3.7	3.2-4.3
1978	520	10.6	9.7-11.6	223	4.4	3.8-5.0
1979	510	10.5	9.6-11.5	215	4.3	3.7-4.9
1980	552	10.9	10.0-11.9	209	4.2	3.6-4.8
1981	457	9.0	8.2-9.9	217	4.1	3.6-4.7
1982	465	9.4	8.6-10.3	191	3.6	3.1-4.1
1983	479	9.6	8.7-10.5	167	3.2	2.7-3.7
1984	518	10.5	9.6-11.4	164	3.2	2.7-3.7
1985	479	9.5	8.7-10.4	161	3.1	2.7-3.7
1986	436	8.7	7.9-9.6	182	3.5	3.0-4.0
1987	499	9.8	8.9-10.7	181	3.5	3.0-4.1
1988	513	10.2	9.3-11.1	170	3.2	2.7-3.7
1989	483	9.5	8.6-10.4	151	2.8	2.3-3.3
1990	515	10.1	9.2-11.0	168	3.0	2.6-3.5
1991	500	9.6	8.7-10.5	141	2.6	2.2-3.1
1992	458	8.8	8.0-9.6	163	3.0	2.6-3.5
1993	489	9.3	8.5-10.2	144	2.6	2.2-3.1
1994	467	8.8	8.0-9.7	157	2.9	2.4-3.3
1995	554	10.5	9.7-11.5	117	2.1	1.8-2.6
1996	539	10.2	9.3-11.1	147	2.7	2.3-3.2
1997	487	9.2	8.4-10.1	155	2.7	2.3-3.2
1998	448	8.5	7.7-9.3	122	2.2	1.8-2.6
1999	440	8.3	7.5-9.1	144	2.5	2.1-3.0
2000	406	7.6	6.9-8.4	121	2.1	1.7-2.5
2001	433	8.0	7.2-8.8	129	2.2	1.8-2.6
2002	372	6.9	6.2-7.7	132	2.3	1.9-2.7
2003	412	7.7	7.0-8.5	113	1.9	1.5-2.3
2004	349	6.6	5.9-7.3	122	2.1	1.7-2.5
2005	399	7.5	6.8-8.3	120	2.0	1.7-2.4
2006	382	7.2	6.5-8.0	113	1.9	1.6-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Cervix uteri

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	393	7.3	6.6-8.1	118	2.0	1.6-2.4
2008	374	7.1	6.3-7.8	99	1.6	1.3-2.0
2009	405	7.6	6.9-8.5	91	1.5	1.2-1.9
2010	390	7.6	6.8-8.4	113	1.8	1.5-2.2
2011	315	5.9	5.2-6.6	110	1.8	1.5-2.2
2012	350	6.5	5.8-7.2	139	2.4	2.0-2.8
2013	366	7.0	6.3-7.8	113	1.9	1.5-2.3
2014	353	6.7	6.0-7.5	116	2.0	1.6-2.4
2015	362	7.1	6.3-7.9	101	1.6	1.3-1.9
2016	333	6.7	5.9-7.5	120	2.0	1.6-2.4
2017	369	7.2	6.5-8.1	114	1.8	1.5-2.2
2018	310	6.1	5.4-6.9	93	1.5	1.2-1.9
2019	320	6.7	5.9-7.5	105	1.8	1.4-2.2
2020	290	5.7	5.0-6.5	87	1.4	1.1-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,518	29.4	27.9-31.0	285	5.5	4.9-6.2
1977	1,391	26.4	25.0-27.8	286	5.4	4.8-6.1
1978	1,239	23.2	21.9-24.6	256	4.8	4.2-5.4
1979	1,359	25.5	24.1-26.9	290	5.3	4.7-5.9
1980	1,358	25.7	24.3-27.1	292	5.2	4.6-5.9
1981	1,466	27.3	25.9-28.7	283	5.1	4.5-5.8
1982	1,422	26.1	24.7-27.5	300	5.4	4.8-6.0
1983	1,353	25.0	23.7-26.4	280	4.8	4.3-5.5
1984	1,400	25.7	24.3-27.1	291	5.1	4.5-5.7
1985	1,330	24.1	22.8-25.5	281	4.9	4.3-5.5
1986	1,353	24.7	23.3-26.0	274	4.6	4.1-5.2
1987	1,438	26.1	24.7-27.5	271	4.6	4.0-5.2
1988	1,316	23.8	22.5-25.1	275	4.6	4.1-5.2
1989	1,330	23.7	22.4-25.0	276	4.6	4.1-5.2
1990	1,386	24.7	23.4-26.1	239	3.9	3.5-4.5
1991	1,403	25.5	24.2-27.0	271	4.4	3.9-5.0
1992	1,481	26.3	25.0-27.7	281	4.6	4.1-5.2
1993	1,417	25.1	23.8-26.5	293	4.7	4.2-5.3
1994	1,526	26.7	25.4-28.1	289	4.7	4.1-5.2
1995	1,525	26.9	25.6-28.3	298	4.7	4.2-5.3
1996	1,623	28.4	27.0-29.8	276	4.4	3.9-4.9
1997	1,572	27.5	26.2-29.0	273	4.4	3.8-4.9
1998	1,724	30.0	28.6-31.5	272	4.3	3.8-4.9
1999	1,649	28.3	27.0-29.8	248	3.8	3.3-4.3
2000	1,652	28.0	26.7-29.4	277	4.3	3.8-4.8
2001	1,806	30.7	29.3-32.1	269	4.1	3.6-4.6
2002	1,834	30.7	29.3-32.1	289	4.5	4.0-5.0
2003	1,821	30.2	28.8-31.6	304	4.6	4.1-5.2
2004	1,841	30.1	28.7-31.5	293	4.4	3.9-5.0
2005	1,898	30.8	29.5-32.3	291	4.4	3.9-5.0
2006	1,864	29.9	28.6-31.3	317	4.7	4.2-5.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
White Individuals
Corpus uteris and uterus, not otherwise specified**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1,946	31.2	29.8-32.6	298	4.3	3.9-4.9
2008	2,067	32.7	31.3-34.2	306	4.5	4.0-5.1
2009	1,962	30.6	29.3-32.1	304	4.5	4.0-5.0
2010	2,003	31.0	29.7-32.5	303	4.3	3.8-4.8
2011	2,081	31.6	30.3-33.1	335	4.9	4.4-5.5
2012	2,177	32.5	31.1-34.0	343	4.8	4.3-5.4
2013	2,312	34.4	32.9-35.9	375	5.1	4.6-5.7
2014	2,283	33.6	32.2-35.0	360	4.9	4.4-5.4
2015	2,233	32.5	31.1-34.0	320	4.4	3.9-4.9
2016	2,313	34.0	32.5-35.4	373	5.0	4.5-5.6
2017	2,291	33.1	31.7-34.6	371	5.0	4.5-5.6
2018	2,168	31.0	29.6-32.4	405	5.4	4.9-6.0
2019	2,239	31.4	30.1-32.9	412	5.4	4.9-6.0
2020	2,149	30.1	28.8-31.5	400	5.2	4.7-5.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Ovary

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	870	17.3	16.2-18.5	652	12.7	11.7-13.7
1977	852	16.6	15.5-17.8	554	10.7	9.8-11.6
1978	838	16.2	15.1-17.4	596	11.2	10.3-12.1
1979	877	16.9	15.8-18.1	571	10.8	9.9-11.7
1980	872	16.7	15.6-17.9	583	10.9	10.0-11.8
1981	946	18.1	16.9-19.3	624	11.4	10.5-12.4
1982	884	16.9	15.7-18.0	619	11.3	10.4-12.3
1983	952	18.0	16.8-19.2	613	11.0	10.1-11.9
1984	904	17.0	15.9-18.2	601	10.9	10.0-11.8
1985	929	17.7	16.6-18.9	588	10.6	9.8-11.5
1986	947	17.9	16.7-19.1	567	10.2	9.4-11.1
1987	933	17.3	16.2-18.5	611	10.9	10.1-11.8
1988	948	17.6	16.5-18.8	618	10.9	10.0-11.8
1989	952	17.5	16.3-18.6	623	10.9	10.0-11.8
1990	998	18.2	17.0-19.3	632	10.8	10.0-11.7
1991	1,003	18.2	17.1-19.4	655	11.2	10.4-12.1
1992	1,003	18.1	17.0-19.3	629	10.9	10.0-11.8
1993	1,000	17.7	16.6-18.9	624	10.5	9.6-11.3
1994	1,027	18.2	17.1-19.3	618	10.4	9.6-11.3
1995	1,028	18.2	17.0-19.3	640	10.7	9.8-11.5
1996	1,043	18.3	17.2-19.5	600	9.9	9.1-10.8
1997	999	17.6	16.5-18.7	650	10.7	9.9-11.6
1998	923	16.1	15.1-17.2	573	9.4	8.6-10.2
1999	965	16.5	15.5-17.6	582	9.4	8.7-10.2
2000	998	16.8	15.8-17.9	633	10.1	9.3-10.9
2001	986	16.6	15.5-17.7	636	10.0	9.2-10.8
2002	951	15.9	14.9-17.0	604	9.5	8.8-10.3
2003	973	16.1	15.1-17.2	627	9.7	9.0-10.5
2004	930	15.3	14.3-16.3	609	9.4	8.6-10.2
2005	889	14.6	13.6-15.6	653	10.0	9.2-10.8
2006	935	15.2	14.2-16.2	627	9.7	8.9-10.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Ovary

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	907	14.6	13.6-15.6	574	8.8	8.0-9.5
2008	904	14.4	13.5-15.4	615	9.1	8.4-9.9
2009	840	13.2	12.3-14.2	574	8.5	7.8-9.2
2010	895	14.1	13.2-15.1	557	8.3	7.6-9.0
2011	884	13.7	12.8-14.6	566	8.2	7.5-8.9
2012	828	12.9	12.0-13.8	544	7.8	7.1-8.5
2013	930	14.5	13.5-15.5	586	8.3	7.6-9.0
2014	785	12.1	11.2-13.0	516	7.2	6.6-7.9
2015	816	12.7	11.8-13.6	524	7.3	6.7-8.0
2016	824	12.6	11.7-13.6	508	7.1	6.5-7.8
2017	779	11.6	10.8-12.5	480	6.6	6.0-7.2
2018	749	11.7	10.8-12.6	487	6.5	6.0-7.2
2019	774	11.9	11.0-12.9	477	6.3	5.7-6.9
2020	656	10.0	9.2-10.9	504	6.7	6.1-7.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Prostate

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2,551	79.0	75.8-82.3	938	31.5	29.4-33.7
1977	2,621	79.0	75.9-82.3	928	30.8	28.7-32.9
1978	2,592	77.9	74.8-81.1	975	32.6	30.5-34.8
1979	2,663	79.6	76.5-82.9	994	32.5	30.4-34.7
1980	2,991	85.9	82.6-89.2	996	32.1	30.0-34.2
1981	3,048	87.6	84.4-90.9	952	30.6	28.6-32.7
1982	3,190	89.9	86.6-93.2	1,011	31.7	29.6-33.8
1983	3,371	93.3	90.0-96.6	1,093	33.6	31.6-35.8
1984	3,318	90.5	87.3-93.8	1,134	34.8	32.7-37.0
1985	3,503	94.0	90.8-97.4	1,103	33.6	31.5-35.7
1986	3,576	94.9	91.7-98.2	1,188	34.9	32.8-37.0
1987	3,924	102.7	99.4-106.1	1,163	34.1	32.1-36.2
1988	4,041	104.9	101.5-108.3	1,184	34.2	32.2-36.3
1989	4,260	108.6	105.3-112.1	1,218	35.1	33.1-37.2
1990	5,025	122.2	118.7-125.8	1,234	34.8	32.8-36.9
1991	6,052	143.6	139.8-147.4	1,354	37.4	35.3-39.5
1992	7,692	176.5	172.5-180.6	1,302	35.9	33.9-38.0
1993	7,380	166.3	162.4-170.3	1,329	36.1	34.2-38.2
1994	6,694	149.2	145.5-152.9	1,359	36.3	34.4-38.4
1995	6,636	145.1	141.6-148.7	1,290	34.3	32.4-36.3
1996	7,317	159.1	155.4-162.8	1,293	33.4	31.5-35.3
1997	7,671	165.4	161.6-169.2	1,304	33.0	31.2-34.9
1998	7,555	160.8	157.2-164.5	1,149	28.4	26.8-30.2
1999	7,990	168.7	165.0-172.5	1,166	28.3	26.7-30.0
2000	8,657	181.8	178.0-185.7	1,148	28.0	26.4-29.7
2001	8,707	180.4	176.6-184.3	1,149	27.4	25.8-29.1
2002	9,065	185.1	181.3-189.0	1,068	25.0	23.5-26.6
2003	8,471	171.0	167.3-174.7	1,083	24.8	23.3-26.3
2004	8,217	163.3	159.8-166.9	1,038	23.4	22.0-24.9
2005	8,332	163.1	159.6-166.7	961	21.2	19.9-22.6
2006	9,363	181.1	177.5-184.9	942	20.4	19.1-21.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	9,501	179.2	175.6-182.9	993	20.9	19.6-22.2
2008	9,207	169.8	166.3-173.3	941	19.3	18.1-20.6
2009	8,980	161.3	157.9-164.7	872	17.7	16.5-18.9
2010	8,512	149.7	146.4-152.9	945	18.6	17.4-19.8
2011	8,790	151.8	148.6-155.1	914	17.7	16.6-18.9
2012	7,522	126.5	123.6-129.4	883	17.0	15.9-18.1
2013	7,318	120.7	117.9-123.6	838	15.6	14.6-16.7
2014	7,157	115.3	112.5-118.0	902	16.5	15.4-17.6
2015	7,569	119.7	116.9-122.5	869	15.8	14.8-17.0
2016	7,836	122.1	119.3-124.9	926	16.4	15.3-17.5
2017	8,420	128.2	125.4-131.1	928	16.3	15.2-17.4
2018	8,938	134.5	131.7-137.4	928	15.8	14.8-16.8
2019	9,357	138.2	135.4-141.1	862	14.6	13.6-15.6
2020	8,058	117.9	115.3-120.6	916	15.4	14.4-16.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Testis

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	173	3.7	3.1-4.3	32	0.6	0.4-0.9
1977	169	3.6	3.0-4.2	28	0.5	0.4-0.8
1978	193	4.2	3.6-4.9	34	0.7	0.5-1.0
1979	223	4.8	4.2-5.5	31	0.7	0.5-1.0
1980	228	4.7	4.1-5.4	28	0.6	0.4-0.8
1981	218	4.5	3.9-5.1	28	0.5	0.4-0.8
1982	213	4.4	3.8-5.1	25	0.5	0.3-0.8
1983	238	4.9	4.2-5.5	23	0.5	0.3-0.8
1984	287	5.9	5.2-6.6	17	0.4	0.2-0.6
1985	255	5.2	4.6-5.9	20	0.4	0.3-0.7
1986	247	4.9	4.3-5.6	25	0.5	0.3-0.8
1987	274	5.5	4.8-6.2	20	0.4	0.2-0.6
1988	261	5.1	4.5-5.8	16	0.4	0.2-0.6
1989	260	5.1	4.5-5.8	21	0.4	0.3-0.6
1990	290	5.7	5.1-6.4	16	0.3	0.2-0.5
1991	268	5.3	4.7-6.0	13	0.2	0.1-0.4
1992	285	5.7	5.0-6.4	19	0.4	0.2-0.6
1993	303	6.1	5.4-6.8	23	0.5	0.3-0.8
1994	295	6.0	5.3-6.7	12	0.2	0.1-0.4
1995	308	6.3	5.6-7.1	15	0.3	0.2-0.5
1996	302	6.3	5.6-7.0	21	0.4	0.3-0.7
1997	325	6.8	6.1-7.6	6	0.1	0.1-0.3
1998	330	7.0	6.2-7.8	18	0.4	0.2-0.6
1999	324	6.9	6.2-7.7	13	0.3	0.1-0.5
2000	294	6.3	5.6-7.1	13	0.3	0.1-0.5
2001	316	6.8	6.1-7.6	13	0.3	0.2-0.5
2002	300	6.5	5.8-7.3	22	0.5	0.3-0.7
2003	339	7.5	6.7-8.3	11	0.2	0.1-0.4
2004	318	7.1	6.3-7.9	11	0.2	0.1-0.4
2005	339	7.6	6.8-8.4	7	0.1	0.1-0.3
2006	314	7.0	6.2-7.8	12	0.2	0.1-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Testis

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	339	7.7	6.9-8.6	6	0.1	0.0-0.3
2008	370	8.5	7.6-9.4	10	0.2	0.1-0.4
2009	336	7.6	6.8-8.5	7	0.1	0.1-0.3
2010	348	7.9	7.1-8.8	14	0.3	0.2-0.5
2011	342	7.7	6.9-8.6	9	0.2	0.1-0.3
2012	365	8.4	7.5-9.3	11	0.2	0.1-0.4
2013	318	7.4	6.6-8.2	7	0.1	0.1-0.3
2014	345	8.0	7.2-8.9	15	0.3	0.2-0.6
2015	337	7.7	6.9-8.6	13	0.3	0.1-0.5
2016	355	8.3	7.4-9.2	10	0.2	0.1-0.4
2017	331	7.9	7.0-8.8	9	0.2	0.1-0.3
2018	364	8.6	7.7-9.6	10	0.2	0.1-0.4
2019	315	7.6	6.8-8.5	26	0.5	0.3-0.8
2020	288	7.0	6.2-7.9	17	0.4	0.2-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Urinary bladder (incl. in situ)

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,907	21.6	20.6-22.6	1,406	39.3	37.2-41.6	501	9.7	8.9-10.6	585	7.0	6.5-7.6	424	13.3	12.0-14.7	161	3.2	2.8-3.8
1977	1,970	22.0	21.0-23.0	1,446	39.4	37.3-41.6	524	10.2	9.4-11.1	583	6.9	6.4-7.5	410	12.7	11.4-14.0	173	3.4	2.9-4.0
1978	2,070	22.9	21.9-24.0	1,504	41.1	38.9-43.3	566	10.9	10.1-11.9	595	6.9	6.4-7.5	419	12.9	11.6-14.3	176	3.4	2.9-3.9
1979	2,091	22.7	21.8-23.8	1,579	42.6	40.5-44.9	512	9.5	8.7-10.4	579	6.7	6.2-7.3	389	11.7	10.5-13.0	190	3.6	3.1-4.2
1980	2,201	23.6	22.6-24.6	1,622	42.9	40.7-45.1	579	10.7	9.8-11.6	555	6.2	5.7-6.7	392	11.6	10.4-12.9	163	3.0	2.6-3.5
1981	2,155	22.8	21.8-23.8	1,603	42.0	39.9-44.3	552	10.1	9.3-11.0	615	6.8	6.3-7.4	436	13.0	11.7-14.3	179	3.3	2.8-3.8
1982	2,326	24.4	23.4-25.4	1,701	43.9	41.8-46.2	625	11.3	10.4-12.2	564	6.2	5.7-6.7	397	11.5	10.3-12.7	167	3.0	2.5-3.5
1983	2,246	23.3	22.3-24.3	1,678	43.4	41.2-45.6	568	10.1	9.3-11.0	560	6.0	5.6-6.6	409	12.1	10.9-13.4	151	2.6	2.2-3.1
1984	2,261	23.2	22.3-24.2	1,636	41.6	39.5-43.7	625	11.1	10.2-12.0	570	6.2	5.7-6.7	389	11.6	10.4-12.9	181	3.1	2.7-3.6
1985	2,387	24.4	23.4-25.4	1,746	44.1	42.0-46.3	641	11.3	10.4-12.2	562	6.0	5.5-6.5	384	10.9	9.8-12.1	178	3.1	2.6-3.6
1986	2,472	24.9	24.0-26.0	1,821	45.1	42.9-47.3	651	11.4	10.5-12.3	574	6.1	5.6-6.6	403	11.9	10.7-13.2	171	2.9	2.5-3.4
1987	2,330	23.3	22.3-24.3	1,675	41.0	39.0-43.1	655	11.2	10.4-12.1	530	5.5	5.1-6.0	356	10.0	9.0-11.2	174	2.9	2.5-3.4
1988	2,341	23.4	22.4-24.4	1,704	41.8	39.8-44.0	637	10.9	10.1-11.8	558	5.7	5.2-6.2	374	10.1	9.1-11.3	184	3.0	2.6-3.5
1989	2,400	23.6	22.7-24.6	1,758	42.7	40.6-44.8	642	10.8	10.0-11.7	541	5.5	5.0-5.9	388	10.7	9.6-11.8	153	2.5	2.1-2.9
1990	2,502	24.5	23.6-25.5	1,794	43.1	41.0-45.2	708	12.0	11.1-12.9	550	5.5	5.1-6.0	361	9.9	8.8-11.0	189	3.0	2.6-3.5
1991	2,437	23.5	22.6-24.5	1,769	41.8	39.8-43.8	668	11.1	10.3-12.0	539	5.3	4.8-5.8	377	10.0	9.0-11.1	162	2.6	2.2-3.0
1992	2,457	23.6	22.7-24.6	1,771	41.8	39.8-43.8	686	11.4	10.6-12.3	566	5.5	5.0-5.9	338	8.9	7.9-9.9	228	3.6	3.1-4.1
1993	2,611	24.8	23.8-25.7	1,837	42.4	40.4-44.4	774	12.6	11.7-13.6	557	5.3	4.9-5.8	374	9.8	8.8-10.9	183	2.8	2.4-3.2
1994	2,485	23.6	22.6-24.5	1,791	41.2	39.3-43.2	694	11.4	10.5-12.3	572	5.4	5.0-5.9	383	9.7	8.7-10.8	189	2.8	2.5-3.3
1995	2,653	25.0	24.0-26.0	1,867	42.6	40.6-44.6	786	12.8	11.9-13.7	547	5.1	4.7-5.6	363	8.9	8.0-9.9	184	2.8	2.4-3.2
1996	2,865	26.8	25.8-27.8	2,108	47.7	45.7-49.9	757	12.3	11.4-13.2	601	5.6	5.1-6.0	401	10.0	9.0-11.0	200	3.0	2.6-3.5
1997	2,990	27.8	26.8-28.8	2,148	48.0	46.0-50.1	842	13.5	12.6-14.4	624	5.7	5.3-6.2	418	10.0	9.1-11.0	206	3.1	2.7-3.5
1998	3,093	28.5	27.5-29.5	2,264	49.9	47.8-52.0	829	13.4	12.5-14.4	579	5.2	4.8-5.7	409	9.9	8.9-10.9	170	2.5	2.1-2.9
1999	3,129	28.6	27.6-29.6	2,268	49.4	47.4-51.5	861	13.7	12.8-14.6	590	5.3	4.8-5.7	407	9.6	8.7-10.6	183	2.6	2.3-3.1
2000	3,141	28.3	27.4-29.3	2,271	49.0	46.9-51.0	870	13.5	12.6-14.5	601	5.3	4.9-5.7	415	9.4	8.5-10.4	186	2.7	2.3-3.1
2001	3,146	28.1	27.2-29.1	2,310	49.5	47.4-51.5	836	13.0	12.1-14.0	613	5.3	4.9-5.8	432	9.9	9.0-10.9	181	2.5	2.2-2.9
2002	3,314	29.4	28.4-30.4	2,474	52.1	50.0-54.2	840	13.0	12.2-14.0	609	5.3	4.9-5.7	407	9.1	8.2-10.1	202	2.9	2.5-3.4
2003	3,368	29.6	28.6-30.6	2,493	51.5	49.5-53.6	875	13.5	12.6-14.4	563	4.8	4.4-5.2	381	8.4	7.6-9.3	182	2.5	2.2-3.0
2004	3,477	30.3	29.3-31.4	2,578	53.1	51.0-55.2	899	13.8	12.9-14.8	605	5.1	4.7-5.6	419	9.0	8.1-9.9	186	2.6	2.2-3.0
2005	3,526	30.5	29.5-31.5	2,602	52.9	50.9-55.0	924	14.0	13.1-14.9	565	4.7	4.3-5.1	383	8.1	7.3-9.0	182	2.5	2.1-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Urinary bladder (incl. in situ)

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	3,449	29.5	28.5-30.5	2,554	51.4	49.4-53.4	895	13.6	12.7-14.5	576	4.8	4.4-5.2	394	8.3	7.5-9.1	182	2.5	2.2-2.9
2007	3,437	29.2	28.2-30.2	2,539	50.1	48.1-52.1	898	13.6	12.7-14.6	640	5.2	4.8-5.7	458	9.4	8.5-10.3	182	2.5	2.1-2.9
2008	3,566	29.7	28.8-30.7	2,688	52.3	50.4-54.4	878	12.9	12.1-13.8	649	5.2	4.8-5.6	446	9.0	8.2-9.9	203	2.7	2.3-3.1
2009	3,613	29.8	28.8-30.8	2,713	51.9	49.9-53.9	900	13.1	12.2-14.0	641	5.1	4.7-5.5	485	9.7	8.8-10.6	156	2.0	1.7-2.4
2010	3,375	27.5	26.5-28.4	2,552	48.1	46.2-50.0	823	11.9	11.1-12.8	666	5.2	4.8-5.6	467	9.0	8.2-9.9	199	2.6	2.2-3.0
2011	3,650	29.6	28.6-30.6	2,726	50.6	48.7-52.6	924	13.5	12.6-14.4	617	4.7	4.3-5.1	437	8.4	7.6-9.2	180	2.3	2.0-2.7
2012	3,751	29.7	28.8-30.7	2,782	51.0	49.1-53.0	969	13.6	12.8-14.6	641	4.9	4.5-5.3	443	8.3	7.6-9.1	198	2.5	2.2-2.9
2013	3,788	29.9	28.9-30.9	2,832	51.5	49.6-53.5	956	13.5	12.6-14.4	643	4.9	4.5-5.3	465	8.6	7.9-9.5	178	2.3	1.9-2.6
2014	3,703	28.6	27.6-29.5	2,772	49.0	47.2-50.9	931	12.8	11.9-13.6	680	5.0	4.7-5.5	461	8.4	7.6-9.2	219	2.8	2.5-3.3
2015	3,590	27.5	26.6-28.4	2,679	46.6	44.8-48.5	911	12.7	11.9-13.6	633	4.6	4.3-5.0	438	7.8	7.1-8.6	195	2.5	2.2-2.9
2016	3,748	28.3	27.3-29.2	2,822	48.2	46.4-50.1	926	12.6	11.7-13.4	664	4.9	4.5-5.3	476	8.5	7.7-9.3	188	2.3	2.0-2.7
2017	3,674	27.4	26.5-28.3	2,765	46.8	45.0-48.6	909	12.1	11.3-13.0	645	4.6	4.2-5.0	445	7.7	7.0-8.5	200	2.4	2.1-2.8
2018	3,651	26.7	25.8-27.6	2,741	45.3	43.6-47.1	910	12.0	11.2-12.9	647	4.6	4.2-4.9	464	7.9	7.2-8.6	183	2.3	1.9-2.7
2019	3,652	26.4	25.5-27.3	2,748	44.7	43.0-46.4	904	11.9	11.1-12.7	592	4.1	3.8-4.5	432	7.3	6.6-8.0	160	1.9	1.6-2.3
2020	3,343	23.8	23.0-24.6	2,538	40.6	39.0-42.2	805	10.4	9.7-11.2	612	4.2	3.9-4.6	449	7.6	6.9-8.3	163	2.0	1.7-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Kidney and renal pelvis

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	644	7.0	6.5-7.6	403	10.2	9.2-11.3	241	4.7	4.1-5.3	357	3.9	3.5-4.3	223	5.7	4.9-6.5	134	2.6	2.1-3.1
1977	703	7.6	7.1-8.2	433	10.8	9.8-12.0	270	5.2	4.6-5.8	400	4.4	4.0-4.9	243	6.3	5.5-7.1	157	3.0	2.6-3.5
1978	701	7.6	7.0-8.2	431	10.8	9.8-12.0	270	5.2	4.6-5.9	354	3.8	3.4-4.2	217	5.5	4.8-6.4	137	2.6	2.1-3.0
1979	717	7.7	7.1-8.2	437	10.8	9.7-11.9	280	5.2	4.6-5.8	409	4.4	4.0-4.8	255	6.4	5.6-7.2	154	2.9	2.5-3.4
1980	726	7.7	7.1-8.3	437	10.7	9.7-11.9	289	5.4	4.8-6.1	406	4.3	3.8-4.7	240	6.1	5.3-7.0	166	3.0	2.5-3.5
1981	799	8.5	7.9-9.2	474	11.9	10.8-13.1	325	6.1	5.5-6.8	385	4.1	3.7-4.5	238	6.0	5.2-6.9	147	2.7	2.3-3.1
1982	838	8.6	8.0-9.3	510	12.0	11.0-13.2	328	6.0	5.4-6.7	390	4.0	3.6-4.4	234	5.8	5.0-6.6	156	2.8	2.4-3.3
1983	903	9.3	8.7-10.0	553	13.2	12.1-14.4	350	6.4	5.7-7.1	398	4.1	3.7-4.6	234	5.7	5.0-6.5	164	3.0	2.5-3.5
1984	893	9.2	8.6-9.9	562	13.6	12.4-14.8	331	6.0	5.4-6.8	400	4.2	3.8-4.6	249	6.4	5.6-7.3	151	2.7	2.3-3.1
1985	939	9.6	9.0-10.2	571	13.6	12.5-14.9	368	6.6	5.9-7.3	432	4.5	4.0-4.9	246	6.2	5.4-7.0	186	3.3	2.8-3.8
1986	953	9.8	9.2-10.4	592	14.1	12.9-15.3	361	6.7	6.0-7.4	449	4.6	4.1-5.0	269	6.5	5.7-7.4	180	3.2	2.7-3.7
1987	1,026	10.4	9.8-11.1	607	14.3	13.2-15.5	419	7.5	6.8-8.3	463	4.7	4.3-5.2	287	7.1	6.3-8.1	176	3.0	2.6-3.5
1988	996	10.0	9.4-10.6	594	13.9	12.8-15.1	402	7.1	6.4-7.8	446	4.4	4.0-4.9	253	6.1	5.3-6.9	193	3.3	2.8-3.8
1989	980	9.8	9.2-10.5	613	14.5	13.3-15.7	367	6.4	5.8-7.2	433	4.3	3.9-4.8	262	6.4	5.6-7.3	171	2.9	2.5-3.4
1990	1,098	10.9	10.2-11.5	688	15.9	14.7-17.2	410	7.2	6.5-7.9	417	4.1	3.7-4.5	253	5.9	5.2-6.7	164	2.7	2.3-3.2
1991	1,120	11.0	10.4-11.7	677	15.4	14.2-16.6	443	7.8	7.1-8.6	464	4.5	4.1-5.0	279	6.6	5.8-7.5	185	3.0	2.6-3.5
1992	1,142	11.2	10.5-11.8	688	15.4	14.2-16.6	454	8.0	7.2-8.8	425	4.1	3.7-4.5	257	6.1	5.4-7.0	168	2.8	2.4-3.2
1993	1,177	11.4	10.7-12.0	701	15.6	14.4-16.8	476	8.2	7.5-9.0	410	3.9	3.6-4.3	250	5.7	5.0-6.4	160	2.6	2.2-3.0
1994	1,173	11.3	10.6-12.0	701	15.6	14.4-16.8	472	8.2	7.4-9.0	467	4.5	4.1-4.9	290	6.8	6.0-7.7	177	2.9	2.5-3.3
1995	1,213	11.6	11.0-12.3	722	15.7	14.6-16.9	491	8.4	7.7-9.2	445	4.2	3.8-4.6	250	5.8	5.1-6.6	195	3.2	2.7-3.7
1996	1,285	12.3	11.6-13.0	794	17.3	16.1-18.5	491	8.3	7.6-9.1	473	4.5	4.1-4.9	275	6.2	5.5-7.0	198	3.2	2.8-3.7
1997	1,372	13.0	12.3-13.7	864	18.8	17.5-20.1	508	8.5	7.8-9.3	449	4.2	3.8-4.6	283	6.4	5.6-7.2	166	2.7	2.3-3.1
1998	1,438	13.6	12.9-14.3	880	18.6	17.4-19.9	558	9.4	8.7-10.3	460	4.2	3.9-4.7	290	6.3	5.6-7.1	170	2.6	2.2-3.1
1999	1,513	14.1	13.4-14.9	928	19.6	18.3-20.9	585	9.8	9.0-10.7	384	3.5	3.2-3.9	237	5.2	4.5-5.9	147	2.2	1.9-2.7
2000	1,511	13.9	13.2-14.7	899	18.7	17.5-20.0	612	10.1	9.3-11.0	447	4.0	3.7-4.4	269	5.9	5.2-6.6	178	2.7	2.4-3.2
2001	1,609	14.8	14.0-15.5	998	20.6	19.4-22.0	611	10.1	9.3-10.9	474	4.2	3.9-4.6	278	6.0	5.3-6.7	196	3.1	2.7-3.6
2002	1,624	14.7	14.0-15.5	1,015	20.8	19.5-22.1	609	9.9	9.1-10.8	435	3.8	3.5-4.2	257	5.5	4.8-6.2	178	2.7	2.3-3.1
2003	1,681	15.1	14.4-15.8	1,050	20.9	19.7-22.2	631	10.2	9.4-11.1	455	4.0	3.6-4.4	270	5.5	4.9-6.2	185	2.7	2.3-3.1
2004	1,857	16.6	15.8-17.4	1,153	22.8	21.5-24.2	704	11.5	10.7-12.4	472	4.1	3.7-4.5	283	5.8	5.1-6.5	189	2.8	2.4-3.3
2005	1,860	16.6	15.8-17.3	1,139	22.6	21.3-23.9	721	11.8	10.9-12.7	469	4.0	3.7-4.4	302	6.1	5.4-6.8	167	2.5	2.1-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Kidney and renal pelvis

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	2,017	17.7	16.9-18.5	1,280	24.8	23.5-26.3	737	11.8	10.9-12.7	472	4.0	3.6-4.4	288	5.8	5.2-6.5	184	2.7	2.3-3.2
2007	2,117	18.4	17.7-19.3	1,351	25.9	24.5-27.3	766	12.5	11.6-13.4	449	3.8	3.4-4.2	279	5.5	4.9-6.2	170	2.5	2.1-2.9
2008	2,052	17.6	16.8-18.4	1,283	24.2	22.8-25.5	769	12.2	11.3-13.1	461	3.8	3.5-4.2	304	5.9	5.2-6.6	157	2.2	1.9-2.6
2009	2,027	17.3	16.6-18.1	1,260	23.5	22.2-24.8	767	12.2	11.4-13.2	440	3.6	3.3-4.0	281	5.3	4.7-6.0	159	2.3	1.9-2.7
2010	2,084	17.5	16.8-18.3	1,304	24.0	22.7-25.3	780	12.2	11.3-13.1	437	3.5	3.2-3.9	280	5.1	4.5-5.8	157	2.2	1.9-2.6
2011	2,160	18.1	17.3-18.9	1,408	25.5	24.1-26.9	752	11.8	11.0-12.7	475	3.8	3.4-4.2	292	5.5	4.8-6.1	183	2.5	2.2-3.0
2012	2,248	18.5	17.7-19.3	1,426	25.4	24.1-26.8	822	12.6	11.7-13.5	495	3.9	3.5-4.2	326	5.9	5.3-6.6	169	2.3	2.0-2.7
2013	2,227	18.2	17.5-19.0	1,436	25.2	23.9-26.6	791	12.2	11.3-13.1	444	3.5	3.1-3.8	291	5.2	4.6-5.9	153	2.1	1.8-2.5
2014	2,310	18.7	17.9-19.5	1,492	25.8	24.5-27.2	818	12.5	11.6-13.4	427	3.2	2.9-3.5	277	4.7	4.2-5.3	150	2.0	1.7-2.4
2015	2,341	18.7	17.9-19.5	1,514	26.0	24.6-27.4	827	12.3	11.4-13.2	450	3.4	3.0-3.7	283	4.9	4.3-5.5	167	2.2	1.9-2.6
2016	2,323	18.6	17.8-19.4	1,548	26.5	25.1-27.9	775	11.7	10.9-12.6	404	3.0	2.7-3.3	263	4.4	3.9-5.0	141	1.9	1.6-2.3
2017	2,365	18.8	18.0-19.6	1,500	25.4	24.1-26.8	865	12.9	12.0-13.9	409	3.1	2.8-3.4	251	4.3	3.7-4.8	158	2.1	1.8-2.5
2018	2,464	19.4	18.6-20.2	1,578	26.2	24.9-27.6	886	13.5	12.6-14.5	440	3.2	2.9-3.5	277	4.6	4.1-5.2	163	2.1	1.8-2.5
2019	2,493	19.3	18.5-20.1	1,646	27.0	25.7-28.4	847	12.5	11.7-13.5	404	2.9	2.6-3.2	252	4.0	3.5-4.5	152	2.0	1.7-2.3
2020	2,212	17.1	16.4-17.9	1,473	24.3	23.0-25.6	739	10.8	10.0-11.7	393	2.8	2.5-3.1	262	4.2	3.7-4.8	131	1.6	1.3-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Brain and other central nervous system

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	555	5.8	5.3-6.3	322	7.2	6.4-8.0	233	4.6	4.0-5.3	415	4.4	3.9-4.8	224	5.1	4.4-5.8	191	3.7	3.2-4.3
1977	570	5.9	5.4-6.4	324	7.3	6.5-8.1	246	4.7	4.1-5.4	428	4.4	4.0-4.8	239	5.4	4.7-6.2	189	3.6	3.1-4.1
1978	542	5.6	5.2-6.2	286	6.5	5.7-7.3	256	5.0	4.4-5.7	398	4.1	3.7-4.5	219	4.9	4.3-5.7	179	3.3	2.9-3.9
1979	617	6.4	5.9-6.9	354	8.0	7.1-8.9	263	5.0	4.4-5.7	404	4.1	3.7-4.6	229	5.1	4.5-5.9	175	3.3	2.8-3.9
1980	620	6.3	5.8-6.8	357	7.7	6.9-8.6	263	5.1	4.5-5.8	455	4.7	4.2-5.1	274	6.2	5.5-7.0	181	3.4	2.9-4.0
1981	614	6.4	5.9-6.9	342	7.8	7.0-8.7	272	5.2	4.5-5.8	421	4.3	3.9-4.7	238	5.5	4.8-6.3	183	3.4	2.9-4.0
1982	655	6.8	6.3-7.3	334	7.6	6.8-8.5	321	6.2	5.5-6.9	435	4.5	4.0-4.9	223	5.1	4.4-5.8	212	4.0	3.4-4.5
1983	631	6.5	6.0-7.1	316	7.1	6.3-8.0	315	6.1	5.4-6.8	425	4.3	3.9-4.7	225	5.1	4.4-5.8	200	3.6	3.1-4.1
1984	699	7.1	6.6-7.7	380	8.6	7.7-9.5	319	6.0	5.3-6.7	430	4.4	4.0-4.8	235	5.5	4.8-6.2	195	3.5	3.1-4.1
1985	660	6.8	6.2-7.3	354	7.9	7.1-8.8	306	5.8	5.1-6.5	413	4.3	3.9-4.7	206	4.8	4.2-5.6	207	3.9	3.4-4.5
1986	688	7.0	6.5-7.6	374	8.4	7.5-9.3	314	5.9	5.2-6.6	458	4.6	4.2-5.1	251	5.7	5.0-6.5	207	3.7	3.2-4.3
1987	739	7.5	7.0-8.1	390	8.7	7.8-9.6	349	6.5	5.8-7.2	458	4.6	4.2-5.1	234	5.3	4.6-6.1	224	4.0	3.5-4.5
1988	718	7.2	6.7-7.8	399	8.8	7.9-9.7	319	6.0	5.3-6.7	488	4.8	4.3-5.2	273	6.0	5.3-6.8	215	3.8	3.3-4.3
1989	710	7.1	6.6-7.7	409	8.9	8.0-9.8	301	5.7	5.1-6.4	457	4.5	4.1-5.0	254	5.7	5.0-6.4	203	3.6	3.2-4.2
1990	766	7.7	7.1-8.3	423	9.4	8.5-10.4	343	6.3	5.7-7.1	419	4.1	3.8-4.6	250	5.6	4.9-6.4	169	2.9	2.5-3.4
1991	720	7.2	6.7-7.7	394	8.8	7.9-9.7	326	6.1	5.4-6.8	445	4.5	4.0-4.9	262	6.0	5.3-6.8	183	3.4	2.9-3.9
1992	739	7.3	6.8-7.9	387	8.5	7.7-9.4	352	6.5	5.8-7.2	475	4.6	4.2-5.1	241	5.3	4.6-6.0	234	4.2	3.6-4.7
1993	740	7.3	6.8-7.9	411	8.9	8.0-9.8	329	6.0	5.4-6.7	455	4.4	4.0-4.9	248	5.4	4.8-6.1	207	3.6	3.1-4.2
1994	815	8.0	7.5-8.6	461	10.0	9.1-11.0	354	6.3	5.7-7.0	529	5.1	4.7-5.6	293	6.4	5.7-7.2	236	4.1	3.6-4.6
1995	857	8.4	7.9-9.0	423	9.0	8.2-10.0	434	7.9	7.1-8.7	467	4.5	4.1-5.0	253	5.5	4.8-6.2	214	3.7	3.2-4.3
1996	836	8.2	7.7-8.8	458	9.9	9.0-10.9	378	6.9	6.2-7.7	514	5.0	4.6-5.4	284	6.2	5.5-7.0	230	4.0	3.5-4.6
1997	791	7.7	7.2-8.3	442	9.4	8.5-10.3	349	6.3	5.6-7.0	490	4.7	4.3-5.1	272	5.8	5.1-6.5	218	3.8	3.3-4.3
1998	840	8.2	7.6-8.8	470	9.8	8.9-10.7	370	6.8	6.1-7.5	476	4.5	4.1-5.0	271	5.7	5.0-6.4	205	3.5	3.0-4.0
1999	783	7.6	7.1-8.2	415	8.8	7.9-9.7	368	6.6	5.9-7.3	483	4.6	4.2-5.0	269	5.6	5.0-6.4	214	3.7	3.2-4.2
2000	877	8.5	7.9-9.0	460	9.6	8.8-10.6	417	7.3	6.6-8.1	468	4.4	4.0-4.8	243	5.0	4.4-5.7	225	3.7	3.2-4.2
2001	815	7.8	7.3-8.4	444	9.3	8.4-10.2	371	6.6	5.9-7.3	462	4.3	3.9-4.7	242	5.0	4.4-5.7	220	3.7	3.2-4.2
2002	826	7.9	7.4-8.5	467	9.7	8.9-10.7	359	6.4	5.7-7.1	506	4.6	4.2-5.1	286	5.8	5.2-6.6	220	3.7	3.2-4.2
2003	850	8.2	7.6-8.7	441	9.0	8.2-9.9	409	7.4	6.7-8.2	457	4.2	3.8-4.6	256	5.2	4.5-5.8	201	3.4	3.0-4.0
2004	808	7.7	7.2-8.2	451	9.3	8.4-10.2	357	6.4	5.7-7.1	510	4.6	4.2-5.1	274	5.5	4.8-6.2	236	3.9	3.4-4.4
2005	839	8.0	7.5-8.6	482	9.9	9.0-10.8	357	6.4	5.7-7.1	507	4.6	4.2-5.0	278	5.6	4.9-6.3	229	3.8	3.3-4.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Brain and other central nervous system

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	819	7.7	7.2-8.3	478	9.6	8.7-10.5	341	6.1	5.4-6.8	473	4.2	3.8-4.6	284	5.6	4.9-6.3	189	3.1	2.6-3.5
2007	800	7.6	7.0-8.1	435	8.8	8.0-9.7	365	6.6	5.9-7.3	485	4.3	3.9-4.7	294	5.7	5.1-6.4	191	3.1	2.7-3.6
2008	822	7.7	7.2-8.3	462	9.3	8.5-10.2	360	6.4	5.7-7.1	481	4.2	3.8-4.6	283	5.5	4.9-6.2	198	3.1	2.7-3.6
2009	838	7.8	7.2-8.3	448	8.9	8.0-9.7	390	6.9	6.2-7.7	538	4.6	4.3-5.1	298	5.6	5.0-6.3	240	3.8	3.3-4.4
2010	809	7.5	7.0-8.0	446	8.9	8.0-9.7	363	6.3	5.7-7.0	515	4.5	4.1-4.9	285	5.5	4.8-6.1	230	3.6	3.2-4.2
2011	806	7.4	6.9-8.0	452	8.8	8.0-9.7	354	6.3	5.6-7.0	472	4.0	3.6-4.3	259	4.8	4.2-5.4	213	3.2	2.8-3.7
2012	844	7.6	7.1-8.2	467	8.9	8.1-9.7	377	6.6	5.9-7.4	570	4.8	4.4-5.2	322	6.0	5.3-6.7	248	3.9	3.4-4.4
2013	827	7.5	7.0-8.0	483	9.2	8.3-10.0	344	5.9	5.3-6.6	531	4.3	3.9-4.7	295	5.2	4.6-5.8	236	3.6	3.1-4.1
2014	905	8.2	7.6-8.8	494	9.3	8.5-10.2	411	7.2	6.5-8.0	552	4.5	4.1-4.9	311	5.4	4.8-6.1	241	3.7	3.3-4.3
2015	847	7.6	7.1-8.2	448	8.4	7.6-9.3	399	6.9	6.2-7.7	537	4.4	4.0-4.8	282	5.0	4.4-5.6	255	3.9	3.4-4.4
2016	830	7.4	6.8-7.9	485	9.0	8.2-9.9	345	5.9	5.2-6.6	566	4.6	4.2-5.0	333	5.8	5.1-6.5	233	3.5	3.1-4.1
2017	893	8.0	7.5-8.6	506	9.5	8.6-10.4	387	6.8	6.1-7.5	575	4.6	4.2-5.0	335	5.6	5.0-6.3	240	3.6	3.2-4.2
2018	804	7.1	6.6-7.6	445	8.1	7.3-8.9	359	6.2	5.5-6.9	575	4.6	4.2-5.0	321	5.5	4.9-6.2	254	3.9	3.4-4.4
2019	768	6.9	6.4-7.5	425	8.2	7.4-9.0	343	5.8	5.2-6.5	557	4.3	4.0-4.7	323	5.4	4.8-6.1	234	3.4	2.9-3.9
2020	795	7.0	6.5-7.5	440	8.0	7.2-8.8	355	6.1	5.4-6.8	561	4.4	4.1-4.8	327	5.6	5.0-6.3	234	3.4	2.9-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Thyroid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	300	3.4	3.0-3.8	101	2.5	2.0-3.0	199	4.2	3.6-4.8	52	0.6	0.5-0.8	20	0.6	0.3-0.9	32	0.6	0.4-0.9
1977	346	3.8	3.4-4.3	108	2.6	2.1-3.1	238	5.0	4.4-5.7	67	0.8	0.6-1.0	21	0.6	0.3-0.9	46	0.9	0.6-1.2
1978	330	3.6	3.2-4.0	96	2.3	1.9-2.9	234	4.7	4.1-5.4	58	0.7	0.5-0.9	20	0.6	0.3-0.9	38	0.7	0.5-1.0
1979	309	3.4	3.0-3.8	92	2.2	1.8-2.8	217	4.4	3.8-5.0	43	0.5	0.3-0.6	13	0.3	0.2-0.6	30	0.5	0.4-0.8
1980	336	3.6	3.2-4.0	106	2.5	2.0-3.0	230	4.5	4.0-5.2	69	0.8	0.6-1.0	33	0.9	0.6-1.2	36	0.7	0.5-0.9
1981	331	3.5	3.1-3.9	102	2.5	2.0-3.0	229	4.6	4.0-5.2	54	0.6	0.4-0.8	24	0.6	0.4-0.9	30	0.6	0.4-0.8
1982	314	3.4	3.0-3.8	99	2.3	1.8-2.8	215	4.4	3.8-5.0	51	0.6	0.4-0.7	26	0.7	0.4-1.0	25	0.5	0.3-0.7
1983	368	3.9	3.5-4.3	101	2.3	1.8-2.8	267	5.3	4.6-6.0	51	0.6	0.4-0.7	14	0.4	0.2-0.6	37	0.7	0.5-0.9
1984	389	4.1	3.7-4.5	112	2.6	2.2-3.2	277	5.5	4.8-6.2	64	0.7	0.5-0.9	24	0.7	0.4-1.0	40	0.7	0.5-0.9
1985	390	4.1	3.7-4.5	109	2.4	2.0-3.0	281	5.6	5.0-6.4	45	0.5	0.4-0.7	18	0.5	0.3-0.8	27	0.5	0.3-0.7
1986	403	4.2	3.8-4.6	119	2.7	2.2-3.2	284	5.6	5.0-6.3	52	0.6	0.4-0.7	24	0.6	0.4-1.0	28	0.5	0.3-0.7
1987	402	4.2	3.8-4.6	131	2.9	2.4-3.4	271	5.4	4.7-6.1	50	0.5	0.4-0.7	15	0.3	0.2-0.5	35	0.6	0.4-0.8
1988	393	4.1	3.7-4.5	102	2.3	1.9-2.8	291	5.8	5.1-6.5	48	0.5	0.4-0.6	18	0.4	0.2-0.7	30	0.5	0.3-0.7
1989	410	4.2	3.8-4.7	114	2.5	2.1-3.1	296	5.8	5.2-6.5	54	0.5	0.4-0.7	17	0.4	0.2-0.7	37	0.6	0.4-0.8
1990	414	4.3	3.9-4.7	129	2.9	2.4-3.5	285	5.5	4.9-6.2	46	0.4	0.3-0.6	21	0.5	0.3-0.7	25	0.4	0.3-0.6
1991	476	4.8	4.4-5.3	144	3.1	2.6-3.7	332	6.4	5.7-7.2	42	0.4	0.3-0.6	15	0.4	0.2-0.6	27	0.5	0.3-0.7
1992	483	4.9	4.4-5.3	135	3.0	2.5-3.5	348	6.6	5.9-7.4	55	0.5	0.4-0.7	16	0.4	0.2-0.6	39	0.6	0.4-0.8
1993	491	4.9	4.5-5.4	133	2.8	2.4-3.4	358	7.0	6.3-7.8	57	0.6	0.4-0.7	21	0.5	0.3-0.8	36	0.6	0.4-0.8
1994	546	5.5	5.0-6.0	141	3.1	2.6-3.6	405	7.9	7.1-8.7	50	0.5	0.4-0.6	23	0.6	0.4-0.8	27	0.4	0.3-0.6
1995	621	6.2	5.7-6.7	171	3.7	3.2-4.3	450	8.8	8.0-9.6	55	0.5	0.4-0.7	24	0.5	0.3-0.8	31	0.5	0.3-0.7
1996	681	6.9	6.4-7.4	182	3.8	3.3-4.4	499	9.8	9.0-10.7	81	0.8	0.6-1.0	27	0.6	0.4-0.9	54	0.9	0.7-1.2
1997	744	7.5	7.0-8.1	202	4.2	3.7-4.9	542	10.6	9.7-11.5	55	0.5	0.4-0.7	23	0.5	0.3-0.8	32	0.5	0.3-0.7
1998	746	7.5	7.0-8.1	211	4.4	3.8-5.1	535	10.4	9.6-11.4	55	0.5	0.4-0.7	17	0.4	0.2-0.6	38	0.6	0.4-0.8
1999	902	9.0	8.5-9.7	218	4.5	3.9-5.1	684	13.5	12.5-14.5	54	0.5	0.4-0.6	20	0.4	0.3-0.7	34	0.5	0.3-0.7
2000	1,036	10.4	9.7-11.0	267	5.5	4.9-6.2	769	15.1	14.0-16.2	48	0.4	0.3-0.6	17	0.4	0.2-0.6	31	0.5	0.3-0.7
2001	1,048	10.4	9.8-11.1	225	4.6	4.0-5.2	823	16.1	15.0-17.3	54	0.5	0.4-0.6	23	0.5	0.3-0.7	31	0.5	0.3-0.7
2002	1,085	10.7	10.1-11.4	280	5.7	5.0-6.4	805	15.7	14.6-16.8	47	0.4	0.3-0.6	18	0.4	0.2-0.6	29	0.5	0.3-0.7
2003	1,135	11.1	10.5-11.8	307	6.2	5.5-6.9	828	16.0	14.9-17.1	54	0.5	0.4-0.6	14	0.3	0.2-0.5	40	0.6	0.4-0.8
2004	1,252	12.3	11.6-13.0	302	6.0	5.3-6.7	950	18.4	17.2-19.6	47	0.4	0.3-0.5	20	0.4	0.3-0.7	27	0.4	0.2-0.6
2005	1,399	13.8	13.0-14.5	354	7.0	6.3-7.8	1,045	20.4	19.1-21.7	46	0.4	0.3-0.5	24	0.5	0.3-0.7	22	0.3	0.2-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Thyroid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,532	15.1	14.3-15.9	387	7.8	7.0-8.6	1,145	22.2	20.9-23.5	55	0.5	0.3-0.6	25	0.5	0.3-0.7	30	0.4	0.3-0.6
2007	1,665	16.2	15.4-17.0	436	8.7	7.9-9.6	1,229	23.5	22.1-24.8	54	0.5	0.3-0.6	26	0.5	0.3-0.7	28	0.4	0.3-0.6
2008	1,843	18.0	17.1-18.8	433	8.5	7.7-9.4	1,410	27.2	25.8-28.7	58	0.5	0.4-0.6	26	0.5	0.3-0.7	32	0.5	0.3-0.7
2009	1,915	18.7	17.8-19.6	494	9.8	8.9-10.7	1,421	27.4	26.0-28.9	65	0.5	0.4-0.7	26	0.5	0.3-0.7	39	0.5	0.4-0.7
2010	1,960	19.0	18.2-19.9	505	9.7	8.8-10.6	1,455	28.2	26.7-29.7	59	0.5	0.3-0.6	25	0.5	0.3-0.7	34	0.4	0.3-0.6
2011	2,086	20.3	19.4-21.2	566	10.9	10.0-11.9	1,520	29.4	27.9-31.0	74	0.6	0.5-0.8	39	0.8	0.5-1.0	35	0.5	0.3-0.7
2012	2,077	20.2	19.3-21.1	557	10.5	9.7-11.5	1,520	29.8	28.2-31.4	66	0.5	0.4-0.7	28	0.5	0.3-0.7	38	0.5	0.4-0.7
2013	2,138	20.7	19.8-21.7	599	11.3	10.4-12.3	1,539	30.1	28.5-31.7	64	0.5	0.4-0.6	28	0.5	0.3-0.7	36	0.5	0.3-0.7
2014	2,136	20.9	20.0-21.8	593	11.2	10.3-12.2	1,543	30.5	28.9-32.1	52	0.4	0.3-0.5	26	0.4	0.3-0.6	26	0.3	0.2-0.5
2015	2,149	21.3	20.3-22.2	578	11.1	10.1-12.1	1,571	31.4	29.8-33.1	68	0.5	0.4-0.7	33	0.6	0.4-0.8	35	0.5	0.3-0.7
2016	1,966	19.4	18.5-20.4	559	10.7	9.8-11.7	1,407	28.2	26.6-29.8	70	0.5	0.4-0.7	27	0.5	0.3-0.7	43	0.6	0.4-0.8
2017	1,882	18.8	17.9-19.7	509	9.9	9.0-10.9	1,373	27.6	26.0-29.1	66	0.5	0.4-0.6	34	0.6	0.4-0.9	32	0.4	0.3-0.6
2018	1,808	17.9	17.1-18.8	560	10.8	9.9-11.8	1,248	25.1	23.6-26.6	70	0.5	0.4-0.6	34	0.6	0.4-0.8	36	0.4	0.3-0.6
2019	1,959	19.5	18.6-20.4	594	11.2	10.2-12.2	1,365	27.8	26.3-29.4	72	0.5	0.4-0.7	35	0.6	0.4-0.9	37	0.5	0.3-0.6
2020	1,571	15.5	14.7-16.4	470	8.8	8.0-9.7	1,101	22.3	20.9-23.8	61	0.5	0.4-0.6	22	0.4	0.3-0.6	39	0.5	0.4-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Hodgkin lymphoma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	349	3.4	3.1-3.8	191	4.0	3.4-4.6	158	3.0	2.5-3.5	139	1.5	1.2-1.8	75	1.8	1.4-2.2	64	1.2	1.0-1.6
1977	329	3.3	3.0-3.7	183	3.9	3.3-4.5	146	2.8	2.3-3.3	137	1.5	1.2-1.8	79	1.9	1.5-2.4	58	1.1	0.8-1.4
1978	357	3.5	3.2-3.9	187	4.0	3.4-4.6	170	3.1	2.7-3.6	119	1.3	1.0-1.5	66	1.6	1.2-2.0	53	1.0	0.8-1.4
1979	377	3.7	3.4-4.2	213	4.6	3.9-5.2	164	3.0	2.6-3.5	108	1.1	0.9-1.4	60	1.4	1.1-1.8	48	0.9	0.7-1.2
1980	397	3.9	3.5-4.3	223	4.7	4.1-5.4	174	3.2	2.7-3.7	131	1.4	1.1-1.6	77	1.8	1.4-2.3	54	1.0	0.8-1.3
1981	379	3.7	3.4-4.1	199	4.1	3.6-4.8	180	3.4	2.9-3.9	112	1.1	0.9-1.4	65	1.5	1.1-1.9	47	0.9	0.6-1.2
1982	369	3.6	3.2-4.0	180	3.8	3.3-4.5	189	3.4	2.9-3.9	131	1.4	1.1-1.6	71	1.6	1.2-2.1	60	1.2	0.9-1.5
1983	354	3.5	3.2-3.9	199	4.2	3.6-4.8	155	2.9	2.4-3.4	90	0.9	0.7-1.1	52	1.1	0.8-1.5	38	0.7	0.5-0.9
1984	381	3.8	3.4-4.2	200	4.3	3.7-5.0	181	3.3	2.8-3.8	94	1.0	0.8-1.2	59	1.4	1.0-1.8	35	0.6	0.4-0.9
1985	395	3.9	3.5-4.3	202	4.2	3.6-4.8	193	3.6	3.1-4.2	97	1.0	0.8-1.2	54	1.2	0.9-1.5	43	0.8	0.6-1.0
1986	353	3.4	3.1-3.8	177	3.6	3.0-4.1	176	3.3	2.8-3.8	118	1.2	1.0-1.4	61	1.4	1.1-1.8	57	1.0	0.8-1.3
1987	346	3.4	3.0-3.8	178	3.7	3.1-4.3	168	3.1	2.7-3.7	100	1.0	0.8-1.2	62	1.3	1.0-1.7	38	0.7	0.5-1.0
1988	364	3.6	3.2-4.0	201	4.2	3.6-4.9	163	3.1	2.6-3.6	80	0.8	0.6-1.0	54	1.2	0.9-1.5	26	0.5	0.3-0.7
1989	376	3.7	3.3-4.1	206	4.3	3.7-4.9	170	3.2	2.7-3.7	93	1.0	0.8-1.2	63	1.4	1.1-1.9	30	0.6	0.4-0.8
1990	362	3.6	3.2-4.0	189	4.0	3.4-4.6	173	3.3	2.8-3.8	82	0.8	0.7-1.0	46	1.0	0.8-1.4	36	0.7	0.5-0.9
1991	352	3.5	3.1-3.9	189	3.9	3.3-4.5	163	3.1	2.6-3.6	67	0.7	0.5-0.9	36	0.8	0.6-1.1	31	0.6	0.4-0.8
1992	378	3.8	3.4-4.2	196	4.2	3.6-4.8	182	3.5	3.0-4.0	89	0.9	0.7-1.1	42	0.9	0.6-1.2	47	0.8	0.6-1.1
1993	348	3.5	3.2-3.9	169	3.5	3.0-4.1	179	3.5	3.0-4.1	68	0.7	0.5-0.8	36	0.8	0.5-1.1	32	0.6	0.4-0.8
1994	396	4.0	3.6-4.4	197	4.1	3.6-4.8	199	3.8	3.3-4.4	72	0.7	0.6-0.9	37	0.8	0.6-1.1	35	0.6	0.4-0.9
1995	332	3.4	3.0-3.8	183	3.8	3.3-4.4	149	2.9	2.5-3.5	69	0.7	0.5-0.9	33	0.7	0.5-1.0	36	0.6	0.4-0.9
1996	349	3.6	3.2-4.0	175	3.7	3.2-4.3	174	3.5	3.0-4.1	58	0.6	0.4-0.8	29	0.7	0.4-0.9	29	0.6	0.4-0.8
1997	320	3.3	3.0-3.7	174	3.7	3.2-4.3	146	3.0	2.5-3.5	59	0.6	0.4-0.7	35	0.7	0.5-1.0	24	0.4	0.3-0.6
1998	357	3.7	3.3-4.1	178	3.8	3.3-4.4	179	3.6	3.1-4.2	58	0.6	0.4-0.7	31	0.6	0.4-0.9	27	0.5	0.3-0.8
1999	340	3.5	3.2-3.9	191	4.1	3.5-4.7	149	3.0	2.5-3.5	62	0.6	0.5-0.8	29	0.6	0.4-0.9	33	0.6	0.4-0.8
2000	332	3.4	3.1-3.8	183	3.9	3.4-4.5	149	3.0	2.5-3.5	50	0.5	0.4-0.6	25	0.5	0.3-0.8	25	0.4	0.3-0.6
2001	358	3.7	3.4-4.2	187	4.0	3.4-4.6	171	3.6	3.0-4.2	61	0.6	0.5-0.8	32	0.7	0.5-1.0	29	0.5	0.3-0.8
2002	385	4.0	3.6-4.4	221	4.8	4.1-5.4	164	3.4	2.9-3.9	64	0.6	0.5-0.8	27	0.6	0.4-0.8	37	0.7	0.5-0.9
2003	333	3.4	3.1-3.8	189	4.0	3.4-4.6	144	2.9	2.4-3.4	46	0.5	0.3-0.6	21	0.4	0.3-0.7	25	0.5	0.3-0.7
2004	371	3.8	3.4-4.2	190	4.0	3.4-4.6	181	3.7	3.2-4.3	52	0.5	0.4-0.6	31	0.6	0.4-0.9	21	0.3	0.2-0.5
2005	372	3.9	3.5-4.3	200	4.2	3.7-4.9	172	3.6	3.1-4.2	47	0.4	0.3-0.6	20	0.4	0.3-0.6	27	0.5	0.3-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Hodgkin lymphoma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	338	3.5	3.1-3.9	187	3.9	3.4-4.5	151	3.0	2.6-3.6	56	0.5	0.4-0.7	29	0.6	0.4-0.8	27	0.5	0.3-0.7
2007	343	3.5	3.2-3.9	182	3.8	3.2-4.4	161	3.3	2.8-3.9	52	0.5	0.4-0.6	25	0.5	0.3-0.7	27	0.4	0.3-0.6
2008	345	3.5	3.2-3.9	191	4.0	3.4-4.6	154	3.1	2.6-3.6	51	0.5	0.3-0.6	28	0.6	0.4-0.8	23	0.4	0.2-0.5
2009	348	3.6	3.2-4.0	189	3.9	3.4-4.6	159	3.3	2.8-3.9	40	0.4	0.3-0.5	20	0.4	0.2-0.6	20	0.3	0.2-0.5
2010	333	3.3	3.0-3.7	192	4.0	3.4-4.6	141	2.8	2.3-3.3	44	0.4	0.3-0.6	23	0.5	0.3-0.7	21	0.4	0.2-0.6
2011	339	3.5	3.1-3.9	190	4.0	3.4-4.6	149	3.1	2.6-3.6	37	0.3	0.2-0.4	26	0.5	0.3-0.8	11	0.2	0.1-0.3
2012	359	3.8	3.4-4.2	220	4.7	4.1-5.4	139	2.9	2.4-3.5	47	0.4	0.3-0.5	20	0.4	0.2-0.6	27	0.4	0.3-0.6
2013	321	3.3	2.9-3.7	167	3.4	2.9-4.0	154	3.2	2.7-3.7	33	0.3	0.2-0.4	14	0.3	0.2-0.5	19	0.3	0.2-0.5
2014	372	3.8	3.4-4.2	196	4.1	3.5-4.7	176	3.5	2.9-4.0	34	0.3	0.2-0.4	15	0.3	0.1-0.5	19	0.3	0.2-0.5
2015	293	3.1	2.7-3.5	164	3.5	2.9-4.1	129	2.7	2.3-3.3	29	0.2	0.2-0.3	13	0.2	0.1-0.4	16	0.2	0.1-0.4
2016	347	3.6	3.2-4.0	175	3.6	3.1-4.2	172	3.5	3.0-4.1	40	0.3	0.2-0.5	23	0.4	0.3-0.6	17	0.3	0.2-0.5
2017	332	3.4	3.0-3.8	178	3.6	3.1-4.2	154	3.2	2.7-3.8	34	0.3	0.2-0.4	19	0.3	0.2-0.6	15	0.2	0.1-0.3
2018	307	3.0	2.7-3.4	179	3.6	3.0-4.1	128	2.5	2.1-3.0	43	0.4	0.2-0.5	26	0.5	0.3-0.7	17	0.3	0.1-0.4
2019	347	3.6	3.2-4.0	184	3.7	3.2-4.4	163	3.5	3.0-4.2	41	0.3	0.2-0.4	27	0.4	0.3-0.7	14	0.2	0.1-0.4
2020	282	3.0	2.7-3.4	156	3.4	2.9-4.0	126	2.7	2.2-3.2	24	0.2	0.1-0.3	16	0.3	0.2-0.5	8	0.1	0.1-0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Non-Hodgkin lymphomas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,023	11.2	10.6-12.0	532	13.4	12.2-14.6	491	9.6	8.8-10.5	572	6.3	5.8-6.9	302	7.8	6.9-8.7	270	5.2	4.6-5.9
1977	1,092	11.8	11.1-12.5	559	14.0	12.8-15.3	533	10.2	9.3-11.1	618	6.7	6.2-7.3	306	7.7	6.9-8.7	312	5.9	5.3-6.6
1978	1,208	12.9	12.2-13.7	641	15.6	14.4-17.0	567	10.8	9.9-11.7	618	6.8	6.2-7.3	312	8.2	7.3-9.2	306	5.8	5.1-6.5
1979	1,116	12.0	11.3-12.8	565	14.1	12.9-15.3	551	10.4	9.5-11.3	630	6.8	6.3-7.4	340	8.6	7.7-9.6	290	5.4	4.8-6.1
1980	1,206	12.8	12.0-13.5	623	15.1	13.8-16.3	583	10.9	10.0-11.8	646	6.9	6.4-7.5	314	7.8	6.9-8.8	332	6.2	5.5-6.9
1981	1,229	13.0	12.2-13.7	607	15.0	13.8-16.3	622	11.4	10.5-12.4	639	6.8	6.2-7.3	333	8.4	7.5-9.4	306	5.6	4.9-6.2
1982	1,252	13.1	12.3-13.8	651	15.8	14.6-17.2	601	11.1	10.2-12.0	653	6.9	6.4-7.5	338	8.7	7.8-9.7	315	5.6	5.0-6.3
1983	1,359	13.9	13.2-14.7	671	16.0	14.8-17.3	688	12.3	11.4-13.3	685	7.1	6.6-7.7	359	8.8	7.9-9.9	326	5.9	5.3-6.6
1984	1,408	14.6	13.8-15.4	739	18.3	17.0-19.8	669	12.0	11.1-13.0	691	7.1	6.6-7.7	374	9.6	8.6-10.7	317	5.6	5.0-6.2
1985	1,480	15.2	14.4-16.0	780	18.4	17.1-19.8	700	12.5	11.6-13.5	736	7.6	7.0-8.2	385	9.5	8.5-10.5	351	6.2	5.5-6.9
1986	1,548	15.8	15.0-16.7	833	19.9	18.5-21.4	715	12.8	11.9-13.8	783	8.1	7.5-8.7	428	10.7	9.7-11.8	355	6.1	5.5-6.8
1987	1,517	15.3	14.6-16.2	805	18.9	17.6-20.3	712	12.5	11.6-13.5	781	8.0	7.4-8.5	384	9.4	8.4-10.4	397	6.9	6.2-7.6
1988	1,638	16.6	15.8-17.5	838	19.6	18.3-21.1	800	14.4	13.4-15.4	792	8.0	7.5-8.6	398	9.8	8.8-10.8	394	6.8	6.1-7.5
1989	1,706	17.2	16.3-18.0	894	20.6	19.2-22.0	812	14.4	13.4-15.5	823	8.2	7.7-8.8	439	10.6	9.6-11.7	384	6.5	5.9-7.2
1990	1,790	17.7	16.9-18.6	951	21.8	20.4-23.3	839	14.5	13.5-15.6	855	8.5	7.9-9.1	434	10.5	9.5-11.6	421	7.0	6.4-7.8
1991	1,771	17.5	16.7-18.4	925	21.6	20.2-23.1	846	14.7	13.7-15.7	866	8.5	7.9-9.1	452	10.9	9.9-12.0	414	6.8	6.1-7.5
1992	1,764	17.3	16.5-18.1	925	20.9	19.6-22.4	839	14.4	13.5-15.5	911	8.9	8.3-9.5	459	11.0	10.0-12.1	452	7.4	6.8-8.2
1993	1,868	18.3	17.4-19.1	983	21.9	20.5-23.3	885	15.2	14.2-16.2	844	8.1	7.6-8.7	418	9.7	8.8-10.7	426	6.9	6.2-7.6
1994	1,969	19.0	18.2-19.9	1,059	23.6	22.2-25.1	910	15.3	14.3-16.4	961	9.2	8.6-9.8	492	11.5	10.5-12.6	469	7.6	6.9-8.3
1995	1,985	19.2	18.3-20.0	1,026	22.9	21.5-24.3	959	16.3	15.3-17.4	980	9.3	8.7-9.9	508	11.9	10.9-13.0	472	7.5	6.9-8.3
1996	2,149	20.6	19.7-21.5	1,127	24.9	23.4-26.4	1,022	17.3	16.2-18.4	990	9.3	8.7-9.9	508	11.7	10.7-12.7	482	7.7	7.0-8.4
1997	2,206	21.0	20.1-21.8	1,133	24.6	23.2-26.1	1,073	18.0	16.9-19.1	1,033	9.7	9.1-10.3	518	11.6	10.6-12.7	515	8.2	7.5-8.9
1998	2,244	21.1	20.2-22.0	1,125	24.3	22.8-25.7	1,119	18.7	17.6-19.8	952	8.8	8.2-9.3	458	10.1	9.2-11.1	494	7.7	7.0-8.4
1999	2,272	21.2	20.3-22.1	1,136	24.3	22.9-25.8	1,136	18.8	17.7-20.0	912	8.3	7.8-8.9	443	9.9	9.0-10.9	469	7.3	6.6-8.0
2000	2,294	21.2	20.3-22.1	1,173	24.9	23.4-26.3	1,121	18.3	17.2-19.4	957	8.6	8.1-9.2	469	10.3	9.4-11.3	488	7.5	6.8-8.2
2001	2,292	21.0	20.1-21.8	1,160	24.4	23.0-25.9	1,132	18.4	17.3-19.5	880	7.9	7.4-8.4	474	10.1	9.2-11.0	406	6.1	5.5-6.7
2002	2,361	21.5	20.6-22.4	1,247	25.9	24.5-27.4	1,114	18.1	17.0-19.2	885	7.8	7.3-8.4	478	10.2	9.3-11.2	407	6.1	5.6-6.8
2003	2,454	22.1	21.2-23.0	1,253	25.7	24.3-27.1	1,201	19.2	18.1-20.3	808	7.0	6.5-7.5	406	8.7	7.9-9.6	402	5.9	5.3-6.5
2004	2,520	22.6	21.7-23.5	1,344	27.3	25.8-28.8	1,176	19.0	17.9-20.2	790	6.8	6.3-7.3	383	8.0	7.2-8.8	407	5.9	5.3-6.5
2005	2,578	22.9	22.0-23.8	1,381	27.8	26.3-29.3	1,197	19.0	17.9-20.1	761	6.5	6.0-7.0	403	8.4	7.6-9.2	358	5.1	4.6-5.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Non-Hodgkin lymphomas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	2,697	23.8	22.9-24.7	1,387	27.6	26.2-29.1	1,310	20.6	19.5-21.8	849	7.2	6.7-7.7	430	8.8	7.9-9.6	419	5.9	5.3-6.5
2007	2,724	23.7	22.8-24.6	1,454	28.6	27.1-30.1	1,270	20.0	18.9-21.1	750	6.2	5.8-6.7	416	8.3	7.5-9.1	334	4.7	4.2-5.3
2008	2,731	23.6	22.7-24.5	1,444	28.1	26.6-29.6	1,287	20.2	19.1-21.3	817	6.8	6.3-7.3	423	8.4	7.6-9.2	394	5.7	5.1-6.3
2009	2,773	23.6	22.7-24.5	1,498	28.5	27.1-30.1	1,275	19.7	18.6-20.8	766	6.2	5.8-6.7	427	8.3	7.5-9.1	339	4.8	4.3-5.3
2010	2,786	23.4	22.6-24.3	1,474	27.9	26.4-29.4	1,312	19.8	18.7-20.9	792	6.4	5.9-6.8	414	7.9	7.2-8.8	378	5.2	4.7-5.8
2011	2,762	23.2	22.4-24.2	1,516	28.4	27.0-29.9	1,246	19.2	18.1-20.3	778	6.2	5.8-6.7	411	7.7	7.0-8.5	367	5.1	4.6-5.7
2012	2,796	23.1	22.2-24.0	1,513	28.0	26.5-29.4	1,283	19.1	18.1-20.3	749	6.0	5.6-6.4	413	7.8	7.1-8.7	336	4.5	4.0-5.1
2013	2,867	23.5	22.6-24.4	1,562	28.4	27.0-29.9	1,305	19.5	18.4-20.6	705	5.5	5.1-5.9	383	7.0	6.3-7.8	322	4.3	3.8-4.8
2014	2,957	23.8	22.9-24.7	1,625	29.1	27.7-30.6	1,332	19.4	18.3-20.5	753	5.7	5.3-6.2	420	7.6	6.9-8.4	333	4.3	3.8-4.8
2015	2,957	23.7	22.8-24.6	1,631	28.8	27.4-30.3	1,326	19.6	18.5-20.7	735	5.6	5.2-6.1	410	7.3	6.6-8.1	325	4.3	3.8-4.8
2016	3,003	24.0	23.1-24.9	1,588	27.8	26.4-29.3	1,415	20.9	19.8-22.1	752	5.6	5.2-6.1	403	7.1	6.4-7.8	349	4.6	4.1-5.1
2017	2,948	23.1	22.2-24.0	1,626	28.0	26.6-29.4	1,322	19.2	18.1-20.3	774	5.7	5.3-6.2	441	7.7	6.9-8.4	333	4.3	3.8-4.8
2018	2,866	22.4	21.6-23.3	1,514	25.7	24.4-27.1	1,352	19.9	18.7-21.0	672	4.9	4.5-5.3	387	6.7	6.0-7.4	285	3.7	3.3-4.2
2019	2,948	22.6	21.8-23.5	1,609	26.9	25.6-28.3	1,339	19.2	18.1-20.3	679	4.9	4.5-5.3	365	6.1	5.5-6.7	314	4.0	3.5-4.5
2020	2,682	20.7	19.8-21.5	1,465	24.5	23.2-25.8	1,217	17.5	16.5-18.6	640	4.6	4.2-5.0	336	5.6	5.0-6.3	304	3.8	3.4-4.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Myeloma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	395	4.4	4.0-4.8	199	5.3	4.6-6.2	196	3.8	3.2-4.3	258	3.0	2.6-3.3	136	3.8	3.1-4.5	122	2.4	2.0-2.9
1977	370	4.1	3.7-4.5	191	5.1	4.4-5.9	179	3.4	2.9-3.9	248	2.7	2.4-3.1	122	3.1	2.6-3.8	126	2.4	2.0-2.8
1978	392	4.3	3.9-4.8	181	4.9	4.2-5.7	211	4.0	3.5-4.6	251	2.8	2.4-3.1	128	3.4	2.8-4.1	123	2.3	1.9-2.7
1979	417	4.5	4.1-5.0	187	5.0	4.3-5.8	230	4.2	3.7-4.8	258	2.9	2.5-3.2	131	3.7	3.0-4.4	127	2.4	2.0-2.8
1980	409	4.3	3.9-4.8	211	5.5	4.8-6.4	198	3.6	3.1-4.2	311	3.3	3.0-3.7	150	4.1	3.4-4.8	161	2.9	2.5-3.4
1981	450	4.8	4.4-5.3	212	5.6	4.8-6.4	238	4.4	3.8-5.0	288	3.0	2.7-3.4	132	3.5	2.9-4.2	156	2.8	2.4-3.3
1982	445	4.6	4.2-5.1	237	6.0	5.2-6.8	208	3.7	3.2-4.2	286	3.0	2.7-3.4	142	3.8	3.2-4.5	144	2.5	2.1-3.0
1983	434	4.5	4.1-5.0	237	6.3	5.5-7.2	197	3.5	3.0-4.0	298	3.1	2.8-3.5	155	4.2	3.5-4.9	143	2.5	2.1-3.0
1984	483	5.0	4.6-5.5	256	6.4	5.6-7.3	227	4.0	3.5-4.5	330	3.4	3.0-3.8	164	4.1	3.5-4.9	166	2.9	2.5-3.4
1985	482	5.0	4.5-5.4	242	6.1	5.3-6.9	240	4.2	3.7-4.8	343	3.5	3.1-3.9	175	4.5	3.8-5.3	168	2.8	2.4-3.3
1986	444	4.5	4.1-5.0	244	6.1	5.3-6.9	200	3.5	3.0-4.0	343	3.5	3.1-3.9	169	4.4	3.8-5.2	174	2.9	2.5-3.4
1987	504	5.1	4.7-5.6	249	6.2	5.4-7.0	255	4.3	3.8-4.9	340	3.4	3.1-3.8	169	4.3	3.7-5.0	171	2.9	2.5-3.4
1988	475	4.8	4.4-5.3	251	6.2	5.4-7.0	224	3.8	3.3-4.4	361	3.6	3.3-4.0	199	5.2	4.4-6.0	162	2.7	2.3-3.1
1989	517	5.2	4.7-5.7	241	6.1	5.3-6.9	276	4.7	4.1-5.3	320	3.2	2.8-3.6	164	4.1	3.5-4.8	156	2.6	2.2-3.0
1990	544	5.3	4.9-5.8	282	6.7	6.0-7.6	262	4.4	3.9-5.0	322	3.2	2.8-3.5	163	4.1	3.5-4.8	159	2.6	2.2-3.0
1991	512	5.0	4.5-5.4	268	6.5	5.7-7.3	244	4.0	3.5-4.5	352	3.4	3.1-3.8	172	4.2	3.6-4.9	180	2.9	2.5-3.4
1992	557	5.3	4.9-5.8	281	6.7	5.9-7.5	276	4.5	4.0-5.1	353	3.4	3.0-3.8	185	4.6	3.9-5.3	168	2.6	2.3-3.1
1993	562	5.4	4.9-5.8	285	6.5	5.7-7.3	277	4.6	4.0-5.2	385	3.7	3.3-4.1	189	4.5	3.9-5.2	196	3.1	2.6-3.5
1994	564	5.4	4.9-5.8	296	6.9	6.1-7.7	268	4.3	3.8-4.9	411	3.9	3.5-4.3	183	4.4	3.7-5.1	228	3.6	3.2-4.2
1995	595	5.6	5.2-6.1	315	7.3	6.5-8.2	280	4.5	4.0-5.1	404	3.8	3.4-4.2	211	4.9	4.3-5.6	193	3.0	2.6-3.5
1996	578	5.4	5.0-5.9	300	6.9	6.1-7.7	278	4.5	3.9-5.0	385	3.6	3.2-4.0	198	4.8	4.1-5.5	187	2.9	2.5-3.4
1997	595	5.5	5.1-6.0	288	6.4	5.7-7.2	307	5.0	4.4-5.6	375	3.5	3.1-3.8	183	4.3	3.7-5.0	192	3.0	2.6-3.5
1998	662	6.1	5.6-6.6	337	7.3	6.6-8.2	325	5.2	4.6-5.8	387	3.5	3.2-3.9	192	4.3	3.7-5.0	195	3.0	2.6-3.4
1999	681	6.2	5.7-6.7	362	7.9	7.1-8.8	319	5.0	4.5-5.6	405	3.6	3.3-4.0	213	4.7	4.1-5.4	192	2.9	2.5-3.3
2000	680	6.1	5.7-6.6	364	7.8	7.0-8.6	316	5.0	4.4-5.6	410	3.6	3.3-4.0	218	4.8	4.2-5.5	192	2.9	2.5-3.4
2001	645	5.8	5.3-6.2	376	8.0	7.2-8.9	269	4.1	3.6-4.6	349	3.1	2.8-3.4	191	4.1	3.6-4.8	158	2.3	2.0-2.7
2002	717	6.4	5.9-6.8	404	8.5	7.7-9.3	313	4.8	4.3-5.4	394	3.4	3.1-3.8	218	4.8	4.1-5.4	176	2.6	2.2-3.0
2003	667	5.8	5.4-6.3	373	7.7	7.0-8.6	294	4.5	4.0-5.1	373	3.2	2.9-3.6	215	4.6	4.0-5.2	158	2.3	2.0-2.7
2004	715	6.2	5.8-6.7	398	8.0	7.3-8.9	317	4.8	4.3-5.4	364	3.1	2.8-3.5	200	4.2	3.6-4.8	164	2.4	2.0-2.8
2005	683	5.9	5.4-6.3	374	7.5	6.7-8.3	309	4.6	4.1-5.2	377	3.2	2.9-3.6	198	4.1	3.5-4.7	179	2.6	2.2-3.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Myeloma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	723	6.2	5.7-6.6	413	8.1	7.3-8.9	310	4.6	4.1-5.2	360	3.0	2.7-3.4	199	4.0	3.5-4.6	161	2.4	2.0-2.8
2007	790	6.7	6.2-7.2	430	8.4	7.6-9.3	360	5.4	4.9-6.0	374	3.1	2.8-3.4	193	3.8	3.3-4.4	181	2.6	2.2-3.0
2008	796	6.6	6.2-7.1	431	8.2	7.5-9.1	365	5.4	4.9-6.0	358	2.9	2.6-3.3	200	4.0	3.5-4.6	158	2.2	1.9-2.6
2009	836	6.9	6.4-7.4	446	8.4	7.6-9.3	390	5.7	5.1-6.3	380	3.1	2.7-3.4	203	3.9	3.4-4.5	177	2.5	2.1-2.9
2010	842	6.9	6.4-7.4	461	8.5	7.7-9.3	381	5.6	5.0-6.2	386	3.1	2.8-3.4	189	3.6	3.1-4.2	197	2.7	2.3-3.1
2011	878	7.1	6.6-7.6	499	9.2	8.4-10.1	379	5.4	4.9-6.0	363	2.8	2.5-3.1	197	3.6	3.1-4.2	166	2.2	1.9-2.6
2012	873	6.9	6.4-7.4	486	8.6	7.8-9.4	387	5.6	5.0-6.2	394	3.0	2.7-3.4	198	3.6	3.1-4.2	196	2.6	2.3-3.0
2013	938	7.4	6.9-7.9	543	9.5	8.7-10.4	395	5.7	5.1-6.3	344	2.7	2.4-3.0	181	3.3	2.8-3.8	163	2.2	1.9-2.6
2014	1,009	7.8	7.4-8.4	542	9.4	8.6-10.3	467	6.6	6.0-7.3	392	2.9	2.7-3.3	220	3.8	3.3-4.4	172	2.3	1.9-2.6
2015	940	7.3	6.8-7.8	528	9.1	8.3-9.9	412	5.9	5.3-6.5	384	2.9	2.6-3.2	221	3.9	3.4-4.5	163	2.1	1.8-2.5
2016	996	7.7	7.2-8.2	562	9.6	8.8-10.4	434	6.3	5.7-7.0	366	2.7	2.4-3.0	175	3.0	2.5-3.5	191	2.5	2.1-2.9
2017	954	7.2	6.8-7.7	560	9.3	8.6-10.2	394	5.5	4.9-6.1	382	2.8	2.5-3.1	206	3.5	3.0-4.0	176	2.2	1.9-2.6
2018	931	7.0	6.5-7.5	539	8.9	8.2-9.8	392	5.4	4.9-6.0	363	2.6	2.3-2.9	189	3.1	2.7-3.6	174	2.2	1.8-2.5
2019	995	7.4	6.9-7.9	560	9.1	8.3-9.9	435	6.0	5.4-6.6	346	2.4	2.2-2.7	207	3.4	3.0-3.9	139	1.7	1.4-2.0
2020	852	6.2	5.8-6.7	487	7.8	7.1-8.5	365	4.9	4.4-5.5	305	2.2	1.9-2.5	160	2.7	2.2-3.1	145	1.8	1.5-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Leukemias

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,183	13.2	12.4-14.0	672	18.1	16.7-19.6	511	10.0	9.2-10.9	727	8.2	7.6-8.9	415	11.7	10.5-13.0	312	6.1	5.5-6.9
1977	1,116	12.4	11.7-13.2	619	16.3	15.0-17.7	497	9.7	8.9-10.6	785	8.8	8.2-9.5	420	11.2	10.1-12.4	365	7.2	6.5-8.0
1978	1,156	13.0	12.2-13.7	638	17.0	15.7-18.5	518	10.2	9.3-11.1	740	8.2	7.6-8.8	386	10.3	9.3-11.5	354	6.8	6.1-7.6
1979	1,267	13.7	13.0-14.5	721	18.5	17.1-20.0	546	10.4	9.5-11.3	779	8.6	8.0-9.2	421	11.3	10.1-12.5	358	6.9	6.2-7.6
1980	1,269	13.7	13.0-14.5	692	18.0	16.6-19.4	577	11.0	10.1-12.0	820	8.9	8.3-9.5	446	11.7	10.5-12.9	374	7.0	6.3-7.7
1981	1,171	12.7	11.9-13.4	660	17.1	15.7-18.5	511	9.6	8.8-10.5	798	8.7	8.1-9.4	444	11.9	10.8-13.1	354	6.7	6.0-7.5
1982	1,208	13.0	12.3-13.8	682	17.6	16.3-19.1	526	9.9	9.1-10.8	821	8.9	8.3-9.5	464	12.5	11.3-13.7	357	6.7	6.0-7.4
1983	1,187	12.5	11.8-13.3	678	17.0	15.7-18.4	509	9.4	8.6-10.3	800	8.6	8.0-9.2	414	11.1	10.0-12.2	386	7.0	6.3-7.7
1984	1,279	13.4	12.7-14.2	684	17.0	15.6-18.3	595	10.8	9.9-11.7	753	7.9	7.4-8.5	364	9.2	8.3-10.3	389	7.1	6.4-7.8
1985	1,338	13.9	13.2-14.7	761	18.8	17.4-20.3	577	10.5	9.6-11.4	799	8.3	7.8-9.0	469	12.1	10.9-13.3	330	5.9	5.2-6.5
1986	1,244	13.0	12.3-13.7	691	17.3	16.0-18.7	553	10.0	9.2-10.9	789	8.3	7.7-8.9	451	11.5	10.4-12.7	338	6.0	5.4-6.7
1987	1,225	12.5	11.8-13.3	670	16.3	15.0-17.6	555	10.0	9.2-10.9	786	8.2	7.6-8.8	434	11.0	9.9-12.1	352	6.2	5.6-6.9
1988	1,244	12.6	11.9-13.4	699	17.0	15.7-18.3	545	9.8	9.0-10.7	800	8.1	7.6-8.7	461	11.7	10.6-12.8	339	5.9	5.2-6.5
1989	1,223	12.3	11.6-13.0	686	16.7	15.4-18.1	537	9.4	8.6-10.2	797	8.1	7.5-8.6	433	10.8	9.8-11.9	364	6.3	5.7-7.0
1990	1,371	13.8	13.1-14.6	753	17.8	16.5-19.1	618	10.9	10.0-11.8	790	7.9	7.4-8.5	405	9.9	8.9-11.0	385	6.6	5.9-7.3
1991	1,323	13.2	12.5-13.9	720	17.1	15.8-18.5	603	10.5	9.7-11.4	786	7.8	7.2-8.3	395	9.7	8.7-10.8	391	6.6	5.9-7.2
1992	1,353	13.4	12.7-14.1	768	18.2	16.9-19.5	585	10.1	9.3-11.0	842	8.2	7.7-8.8	464	11.3	10.2-12.4	378	6.3	5.7-7.0
1993	1,355	13.3	12.6-14.0	754	17.3	16.1-18.6	601	10.5	9.6-11.4	747	7.3	6.8-7.8	405	9.9	8.9-10.9	342	5.7	5.1-6.3
1994	1,471	14.2	13.5-15.0	818	18.7	17.4-20.1	653	11.0	10.2-11.9	839	8.1	7.5-8.6	476	11.3	10.3-12.4	363	5.9	5.3-6.6
1995	1,447	13.9	13.1-14.6	805	18.2	16.9-19.5	642	10.9	10.0-11.8	835	7.9	7.4-8.5	472	11.3	10.3-12.4	363	5.9	5.3-6.5
1996	1,503	14.3	13.6-15.1	873	19.4	18.1-20.7	630	10.5	9.7-11.4	895	8.5	7.9-9.0	496	11.4	10.4-12.5	399	6.5	5.9-7.2
1997	1,516	14.4	13.7-15.2	825	18.3	17.1-19.6	691	11.4	10.6-12.3	858	8.0	7.4-8.5	470	10.7	9.7-11.7	388	6.1	5.5-6.7
1998	1,529	14.4	13.7-15.2	885	19.2	18.0-20.6	644	10.7	9.9-11.6	828	7.7	7.2-8.2	454	10.1	9.2-11.1	374	5.9	5.3-6.5
1999	1,661	15.5	14.8-16.3	927	20.3	19.0-21.6	734	12.0	11.2-13.0	890	8.1	7.6-8.7	482	10.8	9.8-11.8	408	6.3	5.7-7.0
2000	1,775	16.4	15.6-17.2	1,008	21.9	20.5-23.3	767	12.3	11.4-13.2	893	8.1	7.6-8.7	506	11.2	10.2-12.2	387	6.0	5.4-6.6
2001	1,744	16.0	15.3-16.8	971	20.8	19.5-22.1	773	12.6	11.7-13.5	870	7.8	7.3-8.4	472	10.4	9.5-11.4	398	6.1	5.5-6.8
2002	1,649	15.1	14.3-15.8	943	19.8	18.6-21.2	706	11.5	10.6-12.4	866	7.7	7.2-8.2	483	10.4	9.5-11.4	383	5.8	5.2-6.4
2003	1,794	16.2	15.5-17.0	1,020	21.4	20.1-22.7	774	12.5	11.6-13.4	846	7.4	6.9-8.0	486	10.5	9.5-11.4	360	5.4	4.8-6.0
2004	1,856	16.7	15.9-17.5	1,036	21.3	20.0-22.7	820	13.2	12.3-14.1	906	7.9	7.4-8.4	506	10.6	9.7-11.6	400	6.0	5.4-6.6
2005	1,885	16.9	16.1-17.7	1,041	21.2	19.9-22.5	844	13.4	12.5-14.3	873	7.5	7.0-8.0	500	10.5	9.6-11.5	373	5.5	4.9-6.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
White Individuals
Leukemias

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,875	16.7	15.9-17.5	1,086	22.1	20.8-23.5	789	12.5	11.6-13.4	886	7.7	7.2-8.2	494	10.3	9.4-11.2	392	5.7	5.1-6.3
2007	1,945	17.1	16.4-17.9	1,132	22.5	21.2-23.8	813	13.1	12.2-14.0	853	7.3	6.8-7.8	487	9.9	9.0-10.8	366	5.4	4.8-6.0
2008	1,881	16.5	15.7-17.2	1,082	21.1	19.9-22.4	799	12.8	11.9-13.8	872	7.3	6.8-7.8	512	10.2	9.3-11.1	360	5.2	4.6-5.8
2009	2,017	17.4	16.6-18.2	1,188	23.0	21.6-24.3	829	13.0	12.1-14.0	905	7.7	7.2-8.2	546	10.9	10.0-11.8	359	5.3	4.7-5.9
2010	2,123	18.1	17.4-19.0	1,236	23.7	22.3-25.1	887	13.8	12.8-14.7	827	6.8	6.3-7.3	473	9.3	8.4-10.1	354	5.0	4.5-5.6
2011	2,256	19.2	18.4-20.0	1,309	24.9	23.5-26.3	947	14.6	13.6-15.6	883	7.2	6.7-7.7	504	9.5	8.7-10.4	379	5.4	4.9-6.1
2012	2,182	18.3	17.5-19.1	1,275	23.7	22.4-25.1	907	14.1	13.1-15.1	885	7.1	6.6-7.6	519	9.9	9.1-10.8	366	5.1	4.5-5.6
2013	2,350	19.6	18.8-20.4	1,377	25.4	24.0-26.8	973	15.0	14.0-16.0	863	6.8	6.4-7.3	495	9.3	8.4-10.1	368	5.1	4.5-5.7
2014	2,341	19.1	18.3-19.9	1,417	25.7	24.3-27.1	924	13.8	12.9-14.8	856	6.6	6.2-7.1	483	8.9	8.1-9.7	373	5.1	4.6-5.7
2015	2,337	19.1	18.3-19.9	1,361	24.5	23.2-25.9	976	14.7	13.8-15.7	857	6.6	6.1-7.1	498	9.0	8.2-9.8	359	4.8	4.3-5.3
2016	2,392	18.9	18.2-19.8	1,427	24.9	23.6-26.3	965	14.0	13.1-14.9	827	6.4	5.9-6.8	481	8.7	7.9-9.6	346	4.6	4.1-5.2
2017	2,296	18.4	17.6-19.2	1,377	24.1	22.8-25.5	919	13.8	12.8-14.7	796	6.0	5.5-6.4	454	8.0	7.3-8.8	342	4.4	3.9-4.9
2018	2,176	17.4	16.6-18.2	1,287	22.4	21.2-23.7	889	13.3	12.4-14.3	835	6.2	5.8-6.7	483	8.5	7.7-9.3	352	4.6	4.1-5.1
2019	2,206	17.1	16.4-17.9	1,321	22.4	21.2-23.7	885	12.8	11.9-13.7	817	6.1	5.7-6.5	483	8.3	7.5-9.1	334	4.4	3.9-5.0
2020	2,006	15.9	15.2-16.6	1,215	20.9	19.7-22.2	791	11.6	10.8-12.5	783	5.7	5.3-6.1	467	7.9	7.2-8.7	316	4.0	3.5-4.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	5,620	386.2	375.1-397.5	2,780	491.3	470.2-513.0	2,840	323.6	310.9-336.6	3,435	246.8	237.7-256.1	1,833	330.6	313.0-348.8	1,602	194.4	184.3-205.0
1977	5,969	401.2	390.1-412.6	2,951	520.8	499.2-543.0	3,018	331.8	319.2-344.7	3,498	241.5	232.7-250.4	1,911	343.2	325.5-361.4	1,587	179.9	170.5-189.6
1978	6,289	416.1	404.9-427.5	3,132	537.8	516.2-559.9	3,157	345.1	332.4-358.2	3,769	254.2	245.4-263.2	1,979	338.2	321.3-355.7	1,790	202.7	192.7-212.9
1979	6,528	420.2	409.2-431.4	3,260	549.6	528.2-571.5	3,268	345.2	332.7-358.0	3,904	261.1	252.2-270.2	2,110	366.3	348.6-384.7	1,794	198.3	188.6-208.3
1980	7,028	434.4	423.5-445.5	3,571	584.8	563.2-606.9	3,457	348.5	336.4-361.0	4,013	257.3	248.7-266.0	2,142	362.0	344.7-379.9	1,871	196.1	186.8-205.7
1981	6,862	412.1	401.7-422.7	3,506	555.8	535.1-577.0	3,356	330.4	318.8-342.3	4,167	258.6	250.2-267.2	2,219	364.2	347.1-381.9	1,948	197.6	188.4-207.0
1982	7,096	413.0	402.8-423.4	3,568	550.4	530.3-570.9	3,528	335.2	323.8-347.0	4,156	249.0	241.0-257.3	2,241	357.8	341.3-374.8	1,915	186.7	178.0-195.6
1983	7,216	402.6	392.8-412.6	3,593	534.6	515.1-554.5	3,623	330.2	319.2-341.6	4,378	252.0	244.2-260.1	2,358	358.8	342.8-375.3	2,020	190.0	181.5-198.8
1984	7,771	422.2	412.3-432.3	3,860	554.3	534.9-574.2	3,911	349.1	337.8-360.6	4,330	245.5	237.8-253.4	2,282	344.9	329.2-361.1	2,048	189.4	181.0-198.1
1985	8,024	425.3	415.5-435.2	3,995	560.9	541.7-580.6	4,029	350.3	339.2-361.6	4,343	240.5	233.0-248.2	2,316	343.7	328.2-359.6	2,027	182.8	174.6-191.2
1986	8,214	424.4	414.8-434.2	4,185	580.4	560.9-600.3	4,029	339.9	329.2-350.9	4,479	241.2	233.8-248.7	2,348	342.9	327.6-358.8	2,131	185.6	177.6-193.9
1987	8,251	413.9	404.5-423.4	4,116	550.8	532.3-569.8	4,135	339.2	328.6-350.0	4,430	233.5	226.3-240.8	2,334	338.3	323.1-354.0	2,096	177.6	169.9-185.6
1988	8,421	416.6	407.3-426.1	4,114	550.9	532.3-570.0	4,307	346.7	336.1-357.5	4,481	232.2	225.1-239.4	2,331	332.0	317.2-347.3	2,150	178.6	170.9-186.6
1989	8,708	419.5	410.3-428.8	4,399	572.0	553.3-591.1	4,309	338.4	328.1-348.9	4,477	225.9	219.1-233.0	2,363	328.3	313.7-343.2	2,114	171.1	163.7-178.8
1990	9,052	430.2	421.0-439.6	4,495	579.6	560.9-598.7	4,557	351.8	341.4-362.4	4,592	229.8	222.8-236.8	2,340	321.8	307.5-336.7	2,252	181.1	173.5-188.9
1991	9,661	449.8	440.5-459.3	4,815	606.2	587.2-625.5	4,846	367.2	356.7-377.9	4,555	223.0	216.2-229.8	2,360	320.7	306.5-335.4	2,195	171.9	164.6-179.4
1992	10,028	455.7	446.4-465.1	5,118	628.1	609.1-647.4	4,910	363.7	353.4-374.2	4,540	218.4	211.8-225.1	2,221	300.4	286.6-314.6	2,319	177.8	170.5-185.3
1993	10,050	447.0	438.0-456.2	5,258	627.4	608.9-646.3	4,792	348.0	338.1-358.2	4,519	212.5	206.1-219.0	2,291	302.0	288.3-316.0	2,228	167.6	160.6-174.8
1994	10,703	466.9	457.8-476.2	5,483	638.8	620.4-657.5	5,220	372.2	362.0-382.6	4,974	230.1	223.5-236.8	2,521	324.3	310.5-338.5	2,453	181.5	174.3-188.9
1995	10,680	455.1	446.2-464.1	5,605	630.8	613.0-649.1	5,075	354.8	345.0-364.9	4,973	225.1	218.7-231.6	2,447	305.8	292.6-319.3	2,526	183.3	176.1-190.6
1996	11,324	471.3	462.4-480.3	5,820	638.3	620.6-656.3	5,504	376.3	366.3-386.5	4,856	215.5	209.3-221.9	2,434	297.5	284.7-310.6	2,422	172.4	165.5-179.5
1997	11,400	470.0	461.2-479.0	5,844	639.0	621.4-656.9	5,556	374.8	364.9-384.9	4,835	211.3	205.2-217.5	2,410	291.7	279.2-304.6	2,425	169.4	162.7-176.4
1998	11,719	469.2	460.5-478.0	5,856	615.4	598.5-632.6	5,863	386.0	376.0-396.1	4,840	206.3	200.4-212.4	2,342	272.4	260.5-284.6	2,498	171.4	164.7-178.3
1999	11,886	465.8	457.2-474.4	6,140	630.8	614.0-647.9	5,746	369.9	360.3-379.7	4,884	203.1	197.3-209.0	2,335	268.6	256.9-280.7	2,549	170.1	163.5-176.9
2000	12,215	465.7	457.3-474.2	6,406	640.4	623.7-657.3	5,809	364.0	354.6-373.5	4,879	197.1	191.5-202.9	2,379	265.5	254.1-277.2	2,500	161.8	155.5-168.3
2001	12,296	457.6	449.3-465.9	6,468	627.8	611.6-644.3	5,828	357.9	348.7-367.3	4,923	195.2	189.6-200.8	2,311	255.0	243.9-266.4	2,612	165.5	159.1-172.0
2002	12,508	455.2	447.1-463.4	6,457	614.1	598.2-630.3	6,051	363.8	354.6-373.1	4,853	187.3	181.9-192.7	2,368	250.9	240.1-262.0	2,485	154.3	148.2-160.5
2003	12,800	456.3	448.3-464.5	6,602	612.3	596.6-628.2	6,198	365.6	356.5-374.9	4,727	178.1	172.9-183.3	2,265	231.8	221.6-242.2	2,462	149.5	143.6-155.6
2004	13,171	460.1	452.1-468.2	6,744	605.8	590.5-621.4	6,427	373.6	364.4-382.9	4,784	178.2	173.1-183.5	2,319	234.6	224.4-245.0	2,465	148.1	142.3-154.1
2005	13,360	459.3	451.4-467.3	6,844	603.6	588.6-619.0	6,516	373.7	364.6-383.0	4,781	173.8	168.9-179.0	2,237	220.0	210.3-229.9	2,544	149.8	144.0-155.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	13,662	462.5	454.6-470.5	7,051	611.3	596.3-626.6	6,611	373.6	364.6-382.8	4,606	165.4	160.6-170.4	2,152	210.2	200.8-219.8	2,454	142.6	136.9-148.4
2007	14,118	468.9	461.0-476.8	7,472	628.6	613.7-643.9	6,646	371.0	362.1-380.1	4,817	169.3	164.5-174.3	2,259	213.9	204.6-223.5	2,558	146.3	140.6-152.1
2008	14,401	467.7	460.0-475.6	7,395	604.7	590.2-619.4	7,006	384.0	375.0-393.2	4,698	161.6	156.9-166.4	2,185	203.0	194.1-212.3	2,513	140.3	134.8-145.9
2009	14,735	468.0	460.3-475.8	7,606	601.0	586.8-615.4	7,129	384.1	375.1-393.2	4,763	160.7	156.0-165.4	2,315	210.4	201.3-219.6	2,448	134.1	128.8-139.6
2010	14,821	459.1	451.6-466.7	7,622	587.8	574.0-601.9	7,199	379.6	370.7-388.6	4,848	159.8	155.2-164.4	2,345	209.6	200.6-218.8	2,503	133.9	128.7-139.4
2011	15,945	479.2	471.6-486.9	8,193	609.0	595.2-623.1	7,752	397.9	389.0-407.0	5,111	163.5	159.0-168.2	2,380	201.1	192.6-209.8	2,731	143.2	137.8-148.8
2012	15,345	450.9	443.7-458.3	7,639	553.3	540.3-566.5	7,706	388.1	379.4-397.0	5,068	157.3	152.9-161.8	2,376	194.4	186.2-202.8	2,692	137.4	132.2-142.8
2013	15,847	454.0	446.8-461.2	7,859	551.4	538.7-564.4	7,988	393.8	385.1-402.7	4,930	149.2	145.0-153.5	2,349	188.0	180.1-196.2	2,581	127.7	122.7-132.7
2014	15,949	446.2	439.2-453.3	7,831	531.1	518.9-543.6	8,118	393.3	384.7-402.1	5,119	150.6	146.4-154.9	2,390	184.3	176.6-192.2	2,729	132.4	127.4-137.5
2015	16,122	440.6	433.7-447.6	7,903	523.5	511.5-535.7	8,219	389.8	381.3-398.4	4,913	140.9	136.9-144.9	2,261	168.7	161.5-176.2	2,652	125.5	120.7-130.5
2016	16,677	446.8	439.9-453.7	8,251	534.6	522.6-546.7	8,426	392.4	383.9-401.0	5,082	141.1	137.2-145.1	2,403	171.3	164.3-178.7	2,679	123.7	119.0-128.6
2017	16,619	438.5	431.8-445.4	8,163	518.7	507.1-530.6	8,456	389.1	380.7-397.6	4,803	131.0	127.2-134.8	2,273	160.7	153.9-167.7	2,530	114.3	109.8-118.9
2018	16,577	428.6	421.9-435.3	8,128	503.0	491.7-514.5	8,449	382.2	373.9-390.6	5,055	134.5	130.8-138.4	2,303	158.7	152.1-165.6	2,752	121.4	116.9-126.1
2019	17,255	440.8	434.1-447.6	8,313	505.6	494.4-517.0	8,942	402.4	393.9-411.0	5,107	132.7	129.0-136.5	2,355	157.3	150.8-164.0	2,752	118.9	114.4-123.5
2020	14,642	367.3	361.3-373.5	7,135	425.4	415.2-435.7	7,507	331.9	324.3-339.7	4,676	118.7	115.3-122.3	2,133	138.3	132.3-144.5	2,543	108.3	104.1-112.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	251	15.0	13.1-17.1	171	25.2	21.1-29.8	80	8.1	6.4-10.2	111	7.2	5.8-8.7	75	12.6	9.5-16.2	36	3.7	2.6-5.2
1977	237	13.9	12.1-15.8	181	26.1	22.1-30.5	56	5.5	4.1-7.2	127	7.6	6.3-9.2	100	13.9	11.2-17.1	27	3.0	1.9-4.5
1978	268	14.7	13.0-16.7	185	24.7	21.0-28.9	83	7.8	6.2-9.8	133	7.7	6.4-9.3	100	15.1	11.9-18.8	33	3.1	2.1-4.3
1979	272	15.6	13.7-17.6	204	28.4	24.4-32.9	68	6.7	5.1-8.6	148	8.4	7.0-9.9	109	15.4	12.4-18.9	39	3.7	2.6-5.1
1980	277	15.0	13.2-16.9	196	26.3	22.5-30.5	81	7.3	5.8-9.1	140	7.8	6.5-9.3	114	16.0	12.9-19.6	26	2.3	1.5-3.4
1981	303	15.8	14.0-17.8	219	27.1	23.4-31.2	84	7.7	6.1-9.6	120	6.4	5.3-7.7	85	11.3	8.9-14.2	35	3.2	2.2-4.5
1982	292	15.0	13.3-16.9	200	25.4	21.8-29.4	92	8.0	6.4-9.9	152	8.0	6.7-9.4	116	15.3	12.4-18.6	36	3.1	2.1-4.3
1983	285	14.7	13.0-16.6	208	26.3	22.6-30.5	77	6.8	5.4-8.6	145	7.7	6.4-9.1	110	14.7	11.8-18.0	35	3.2	2.2-4.5
1984	334	16.1	14.4-18.0	226	26.5	23.0-30.3	108	8.9	7.3-10.8	121	6.1	5.0-7.3	92	11.8	9.3-14.7	29	2.4	1.6-3.5
1985	325	15.9	14.1-17.7	222	26.6	23.0-30.6	103	8.6	7.0-10.5	116	5.8	4.8-7.0	82	10.2	8.0-12.9	34	2.9	2.0-4.0
1986	322	15.0	13.4-16.8	230	26.3	22.8-30.1	92	7.3	5.8-9.0	135	6.9	5.7-8.2	97	12.5	9.9-15.4	38	3.3	2.3-4.6
1987	350	16.1	14.4-17.9	240	26.8	23.4-30.6	110	8.6	7.1-10.4	105	5.0	4.1-6.1	77	9.3	7.2-11.8	28	2.2	1.4-3.2
1988	306	13.7	12.1-15.3	203	22.6	19.4-26.2	103	7.7	6.2-9.3	133	6.0	5.0-7.2	96	10.7	8.6-13.3	37	2.9	2.0-3.9
1989	332	14.3	12.7-15.9	226	24.1	20.9-27.6	106	7.7	6.3-9.3	134	5.9	4.9-7.0	94	10.3	8.2-12.8	40	3.0	2.1-4.1
1990	307	13.6	12.1-15.3	214	23.9	20.6-27.6	93	7.0	5.6-8.6	101	4.6	3.7-5.7	70	8.5	6.4-11.1	31	2.3	1.5-3.3
1991	399	17.4	15.7-19.2	283	31.5	27.7-35.7	116	8.3	6.9-10.0	142	6.4	5.3-7.5	101	11.0	8.8-13.4	41	3.2	2.3-4.4
1992	369	15.6	14.0-17.3	242	25.8	22.4-29.5	127	9.0	7.5-10.7	118	5.0	4.1-6.0	84	9.0	7.0-11.2	34	2.4	1.7-3.4
1993	349	14.5	12.9-16.1	229	23.5	20.4-26.9	120	8.4	6.9-10.0	103	4.4	3.6-5.3	73	7.9	6.1-10.0	30	2.1	1.4-3.1
1994	336	13.6	12.2-15.2	232	23.4	20.3-26.8	104	7.1	5.8-8.6	119	5.0	4.1-5.9	89	9.6	7.6-12.0	30	2.1	1.4-2.9
1995	304	12.1	10.7-13.6	225	21.7	18.8-24.9	79	5.4	4.3-6.8	139	5.7	4.8-6.8	101	10.2	8.2-12.5	38	2.7	1.9-3.7
1996	333	13.1	11.7-14.6	219	21.8	18.8-25.0	114	7.5	6.2-9.0	108	4.4	3.6-5.4	76	7.8	6.1-9.9	32	2.2	1.5-3.2
1997	328	12.6	11.2-14.0	220	20.3	17.6-23.3	108	7.1	5.9-8.6	89	3.7	2.9-4.5	66	7.1	5.3-9.1	23	1.6	1.0-2.4
1998	296	11.0	9.8-12.4	195	17.6	15.1-20.4	101	6.5	5.3-7.9	97	3.8	3.1-4.6	70	6.9	5.3-8.8	27	1.8	1.2-2.6
1999	269	9.9	8.8-11.2	181	16.2	13.8-18.8	88	5.6	4.5-6.9	91	3.5	2.8-4.3	57	5.7	4.2-7.4	34	2.2	1.5-3.0
2000	280	10.3	9.1-11.6	186	17.1	14.6-19.9	94	5.8	4.7-7.1	101	3.9	3.1-4.7	71	6.8	5.3-8.7	30	1.9	1.3-2.8
2001	253	8.8	7.7-10.0	185	15.7	13.5-18.3	68	4.0	3.1-5.1	86	3.2	2.5-3.9	60	5.5	4.1-7.2	26	1.6	1.1-2.4
2002	293	10.3	9.1-11.6	188	16.2	13.8-18.8	105	6.3	5.2-7.7	76	2.7	2.1-3.4	54	4.6	3.4-6.1	22	1.4	0.8-2.1
2003	269	9.1	8.0-10.2	180	14.4	12.3-16.7	89	5.2	4.2-6.5	79	2.8	2.2-3.5	60	5.3	4.0-6.8	19	1.2	0.7-1.8
2004	297	9.8	8.7-11.0	194	15.6	13.4-18.1	103	5.8	4.7-7.0	102	3.5	2.8-4.2	78	6.6	5.1-8.3	24	1.4	0.9-2.1
2005	266	8.7	7.7-9.9	184	15.1	12.9-17.6	82	4.6	3.7-5.7	83	2.8	2.2-3.5	60	5.3	4.0-6.9	23	1.3	0.8-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	260	8.3	7.3-9.4	172	13.5	11.4-15.7	88	4.8	3.9-5.9	64	2.1	1.6-2.7	51	4.1	3.0-5.5	13	0.7	0.4-1.2
2007	306	9.8	8.7-11.0	203	15.7	13.5-18.2	103	5.8	4.7-7.0	82	2.7	2.1-3.3	61	5.1	3.8-6.7	21	1.2	0.7-1.8
2008	279	8.7	7.7-9.8	181	13.3	11.4-15.5	98	5.3	4.3-6.4	76	2.5	1.9-3.1	54	4.4	3.2-5.8	22	1.2	0.8-1.9
2009	279	8.5	7.5-9.6	179	13.0	11.1-15.1	100	5.3	4.3-6.4	71	2.2	1.7-2.8	49	3.7	2.7-5.0	22	1.2	0.7-1.8
2010	297	8.8	7.8-9.8	192	13.7	11.7-15.8	105	5.4	4.4-6.5	67	2.1	1.6-2.6	38	2.9	2.0-4.0	29	1.5	1.0-2.2
2011	287	8.2	7.3-9.3	203	13.7	11.8-15.8	84	4.3	3.4-5.3	99	2.9	2.4-3.6	68	4.8	3.7-6.1	31	1.6	1.1-2.3
2012	287	8.1	7.2-9.1	176	11.9	10.2-13.9	111	5.5	4.5-6.6	74	2.2	1.7-2.8	50	3.6	2.6-4.8	24	1.2	0.8-1.9
2013	275	7.7	6.8-8.7	184	12.1	10.4-14.1	91	4.5	3.6-5.5	88	2.6	2.1-3.2	61	4.4	3.3-5.7	27	1.4	0.9-2.0
2014	281	7.5	6.7-8.5	179	11.2	9.6-13.1	102	4.9	4.0-6.0	65	1.8	1.4-2.3	40	2.8	1.9-3.8	25	1.2	0.8-1.8
2015	277	7.5	6.6-8.4	178	11.6	9.9-13.5	99	4.6	3.7-5.7	70	1.9	1.5-2.4	47	3.0	2.2-4.1	23	1.1	0.7-1.7
2016	312	8.3	7.4-9.3	203	13.0	11.2-15.0	109	5.1	4.1-6.1	59	1.6	1.2-2.0	38	2.5	1.8-3.5	21	0.9	0.6-1.4
2017	310	8.2	7.3-9.2	191	12.1	10.4-14.0	119	5.5	4.5-6.6	72	1.9	1.5-2.4	50	3.3	2.4-4.3	22	1.0	0.6-1.6
2018	318	8.1	7.2-9.1	202	12.4	10.7-14.4	116	5.2	4.2-6.2	72	1.9	1.5-2.4	50	3.2	2.3-4.3	22	1.0	0.6-1.6
2019	275	7.0	6.1-7.8	170	10.1	8.6-11.8	105	4.7	3.8-5.7	76	2.0	1.5-2.5	58	3.6	2.7-4.7	18	0.8	0.5-1.2
2020	281	7.0	6.2-7.9	177	10.5	9.0-12.2	104	4.6	3.7-5.6	67	1.7	1.3-2.1	41	2.5	1.8-3.5	26	1.0	0.7-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	202	12.5	10.7-14.5	144	21.7	18.0-26.0	58	6.2	4.6-8.1	204	13.0	11.2-15.1	155	24.7	20.5-29.4	49	5.3	3.8-7.2
1977	220	13.4	11.6-15.5	155	23.9	19.8-28.5	65	6.6	5.1-8.5	196	11.9	10.2-13.8	145	22.6	18.6-27.2	51	5.0	3.7-6.7
1978	247	15.1	13.1-17.2	178	27.1	22.9-31.9	69	7.1	5.5-9.1	179	10.5	9.0-12.3	124	18.0	14.7-21.7	55	5.5	4.1-7.3
1979	237	13.9	12.1-15.9	168	24.9	20.9-29.4	69	6.7	5.2-8.6	215	12.8	11.0-14.7	161	24.9	20.7-29.6	54	5.2	3.9-6.8
1980	258	14.8	12.9-16.8	194	28.3	24.1-33.0	64	6.1	4.6-7.8	213	12.4	10.7-14.4	148	22.2	18.4-26.6	65	6.2	4.8-8.0
1981	262	14.0	12.3-15.9	186	24.7	21.1-28.8	76	6.9	5.4-8.6	212	11.3	9.7-12.9	157	20.6	17.3-24.3	55	4.9	3.7-6.5
1982	239	12.9	11.2-14.7	176	24.5	20.7-28.7	63	5.5	4.2-7.0	213	11.8	10.2-13.5	155	22.3	18.6-26.4	58	5.2	3.9-6.7
1983	247	12.7	11.1-14.4	175	22.4	18.9-26.2	72	6.3	4.9-8.0	211	11.1	9.6-12.7	151	19.8	16.6-23.5	60	5.4	4.1-7.0
1984	263	13.3	11.7-15.1	179	22.9	19.4-26.9	84	7.1	5.7-8.9	207	10.8	9.3-12.5	145	19.6	16.3-23.4	62	5.4	4.1-6.9
1985	240	12.1	10.6-13.8	168	21.5	18.1-25.3	72	6.1	4.7-7.7	217	10.9	9.4-12.5	163	20.5	17.2-24.1	54	4.6	3.4-6.0
1986	254	12.7	11.1-14.4	190	24.7	21.0-28.7	64	5.3	4.0-6.7	218	11.0	9.5-12.6	152	19.9	16.6-23.7	66	5.5	4.3-7.1
1987	225	11.0	9.6-12.6	146	18.5	15.4-21.9	79	6.3	5.0-7.9	179	8.8	7.5-10.2	131	16.4	13.5-19.6	48	4.0	2.9-5.3
1988	254	12.2	10.7-13.9	181	22.7	19.2-26.5	73	5.8	4.5-7.3	202	10.0	8.6-11.5	149	19.4	16.2-23.1	53	4.2	3.2-5.6
1989	257	12.1	10.6-13.7	180	21.9	18.6-25.6	77	6.0	4.7-7.5	184	8.7	7.4-10.1	134	16.0	13.3-19.2	50	4.0	3.0-5.3
1990	230	10.7	9.3-12.3	158	19.0	15.9-22.4	72	5.7	4.4-7.1	191	9.0	7.7-10.4	123	15.3	12.5-18.4	68	5.2	4.0-6.6
1991	211	9.6	8.3-11.0	144	17.3	14.4-20.6	67	4.9	3.8-6.2	177	8.0	6.8-9.3	120	14.1	11.5-17.1	57	4.2	3.2-5.5
1992	223	10.1	8.8-11.6	153	17.9	14.9-21.2	70	5.3	4.1-6.7	148	6.6	5.6-7.8	100	11.6	9.3-14.3	48	3.6	2.6-4.8
1993	236	10.2	8.9-11.6	159	17.1	14.4-20.2	77	5.6	4.4-7.0	182	8.0	6.9-9.3	134	15.6	12.9-18.8	48	3.5	2.6-4.7
1994	233	10.2	8.9-11.6	152	17.4	14.5-20.6	81	5.9	4.7-7.3	177	7.8	6.7-9.0	122	14.1	11.5-17.0	55	4.0	3.0-5.2
1995	186	8.0	6.8-9.2	123	13.2	10.8-15.9	63	4.6	3.5-5.9	172	7.5	6.4-8.8	119	13.2	10.8-16.0	53	3.9	2.9-5.1
1996	195	8.4	7.3-9.7	122	13.1	10.8-15.8	73	5.3	4.2-6.7	166	7.1	6.1-8.3	109	11.9	9.7-14.5	57	4.1	3.1-5.3
1997	189	7.9	6.8-9.1	132	13.9	11.5-16.7	57	4.0	3.0-5.2	150	6.4	5.4-7.6	99	10.9	8.8-13.5	51	3.6	2.7-4.8
1998	183	7.4	6.3-8.6	126	12.8	10.6-15.4	57	3.9	2.9-5.0	147	6.0	5.1-7.1	95	9.8	7.8-12.2	52	3.6	2.7-4.8
1999	194	7.5	6.5-8.7	135	13.6	11.3-16.2	59	3.7	2.8-4.8	131	5.1	4.3-6.1	91	9.1	7.2-11.3	40	2.6	1.9-3.6
2000	187	7.5	6.5-8.7	102	10.6	8.5-12.9	85	5.6	4.5-7.0	166	6.6	5.6-7.6	103	10.6	8.6-13.0	63	4.1	3.1-5.2
2001	174	6.4	5.5-7.5	118	11.4	9.3-13.7	56	3.4	2.6-4.4	127	4.8	4.0-5.8	92	9.2	7.3-11.4	35	2.2	1.5-3.1
2002	157	5.9	5.0-6.9	98	9.4	7.5-11.6	59	3.6	2.8-4.7	132	4.9	4.1-5.9	86	8.3	6.6-10.4	46	2.8	2.1-3.8
2003	185	6.8	5.8-7.9	124	11.8	9.7-14.2	61	3.7	2.9-4.8	128	4.6	3.9-5.5	80	7.2	5.6-9.1	48	2.9	2.2-3.9
2004	174	6.2	5.3-7.2	109	9.5	7.7-11.6	65	4.0	3.1-5.1	139	5.1	4.2-6.0	86	8.0	6.3-10.0	53	3.2	2.4-4.2
2005	155	5.3	4.4-6.2	92	7.7	6.1-9.5	63	3.6	2.8-4.6	108	3.9	3.2-4.7	67	5.9	4.5-7.6	41	2.5	1.8-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	130	4.7	3.9-5.6	91	8.9	7.0-11.0	39	2.3	1.6-3.2	114	4.1	3.3-4.9	71	7.0	5.4-8.9	43	2.5	1.8-3.3
2007	139	4.6	3.8-5.4	100	8.2	6.6-10.0	39	2.2	1.6-3.0	98	3.3	2.7-4.0	67	5.6	4.3-7.2	31	1.8	1.2-2.5
2008	148	4.9	4.1-5.7	102	8.9	7.1-10.9	46	2.5	1.8-3.3	120	4.0	3.3-4.8	74	6.4	4.9-8.1	46	2.6	1.9-3.4
2009	148	4.7	4.0-5.6	102	8.4	6.7-10.3	46	2.5	1.8-3.4	102	3.3	2.7-4.0	81	6.8	5.3-8.6	21	1.2	0.7-1.8
2010	127	4.0	3.3-4.8	81	6.5	5.1-8.1	46	2.5	1.8-3.3	110	3.5	2.9-4.3	74	6.3	4.8-7.9	36	1.9	1.3-2.7
2011	156	4.8	4.1-5.6	105	8.2	6.6-10.0	51	2.7	2.0-3.5	92	2.8	2.3-3.5	64	4.9	3.7-6.3	28	1.5	1.0-2.2
2012	128	3.8	3.2-4.5	83	6.3	4.9-7.9	45	2.2	1.6-2.9	88	2.6	2.0-3.2	59	4.2	3.1-5.5	29	1.4	0.9-2.1
2013	128	3.8	3.2-4.6	89	7.0	5.5-8.6	39	1.9	1.4-2.6	88	2.7	2.1-3.3	55	4.1	3.0-5.4	33	1.7	1.2-2.4
2014	128	3.7	3.1-4.5	91	6.8	5.4-8.5	37	1.8	1.3-2.5	99	3.0	2.4-3.6	64	5.0	3.8-6.5	35	1.7	1.2-2.4
2015	124	3.4	2.8-4.1	79	5.6	4.4-7.1	45	2.1	1.5-2.8	72	2.1	1.6-2.6	47	3.5	2.5-4.7	25	1.2	0.7-1.7
2016	149	4.1	3.4-4.8	96	6.7	5.4-8.3	53	2.5	1.8-3.2	94	2.6	2.1-3.2	64	4.7	3.5-6.0	30	1.4	0.9-2.0
2017	128	3.4	2.8-4.1	84	5.4	4.3-6.8	44	2.0	1.5-2.7	80	2.1	1.7-2.7	52	3.7	2.7-4.9	28	1.2	0.8-1.8
2018	123	3.2	2.7-3.9	73	4.8	3.8-6.1	50	2.2	1.6-2.9	74	1.9	1.5-2.4	54	3.5	2.6-4.6	20	0.9	0.6-1.4
2019	122	3.0	2.5-3.6	82	5.1	4.0-6.3	40	1.7	1.2-2.3	75	1.8	1.4-2.3	50	3.3	2.4-4.4	25	0.9	0.6-1.4
2020	103	2.6	2.1-3.2	67	4.3	3.3-5.5	36	1.6	1.1-2.2	79	1.9	1.5-2.4	48	2.9	2.1-3.9	31	1.3	0.8-1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	243	18.6	16.1-21.4	138	26.3	21.4-31.9	105	13.8	11.1-16.9	189	14.3	12.1-16.6	117	23.0	18.4-28.3	72	8.9	6.9-11.4
1977	271	19.8	17.3-22.5	149	27.0	22.3-32.4	122	15.1	12.3-18.2	212	16.2	13.8-18.7	124	23.8	19.2-29.1	88	11.4	8.9-14.2
1978	244	17.7	15.3-20.2	138	23.8	19.5-28.6	106	13.5	10.9-16.5	209	15.4	13.2-17.9	106	19.3	15.3-23.9	103	13.0	10.4-15.9
1979	253	18.4	16.0-21.0	143	26.0	21.3-31.3	110	13.7	11.1-16.7	180	12.9	10.9-15.0	99	17.5	13.8-21.8	81	10.0	7.8-12.5
1980	316	21.9	19.3-24.6	173	30.0	25.1-35.4	143	16.8	14.0-19.9	208	15.1	13.0-17.5	114	21.6	17.4-26.5	94	11.3	9.0-13.9
1981	283	18.7	16.4-21.2	174	29.0	24.2-34.3	109	12.5	10.1-15.2	240	16.2	14.0-18.5	149	25.2	20.8-30.2	91	10.7	8.5-13.2
1982	282	18.3	16.1-20.7	164	28.7	23.9-33.9	118	12.4	10.2-14.9	208	14.5	12.5-16.8	130	25.6	20.9-31.0	78	8.5	6.7-10.7
1983	283	17.0	15.0-19.2	145	22.5	18.6-26.8	138	13.7	11.4-16.3	214	13.0	11.2-15.0	119	18.8	15.3-22.9	95	9.5	7.6-11.7
1984	264	15.5	13.6-17.6	154	24.6	20.5-29.3	110	10.4	8.4-12.6	179	10.9	9.2-12.7	102	18.1	14.4-22.4	77	7.1	5.6-9.0
1985	288	16.9	14.9-19.1	153	22.3	18.5-26.5	135	13.4	11.2-16.0	184	10.6	9.0-12.3	97	14.5	11.5-18.1	87	8.2	6.5-10.1
1986	308	17.2	15.2-19.4	184	28.0	23.6-32.9	124	11.2	9.3-13.5	209	12.2	10.5-14.1	124	19.1	15.5-23.3	85	8.2	6.5-10.2
1987	319	17.9	15.9-20.1	185	25.6	21.7-30.0	134	12.9	10.8-15.3	200	11.5	9.9-13.3	107	15.6	12.4-19.2	93	9.0	7.2-11.0
1988	306	16.9	14.9-18.9	176	25.6	21.6-30.0	130	11.8	9.8-14.1	215	12.2	10.5-14.0	121	18.2	14.8-22.0	94	8.8	7.0-10.8
1989	310	16.8	14.9-18.9	162	21.7	18.1-25.7	148	13.5	11.4-15.9	217	11.7	10.2-13.5	121	17.0	13.8-20.6	96	8.5	6.9-10.5
1990	323	16.8	14.9-18.8	180	25.1	21.1-29.4	143	12.1	10.2-14.3	211	11.5	10.0-13.3	113	15.8	12.7-19.3	98	8.9	7.2-10.9
1991	296	14.8	13.1-16.7	156	20.4	17.0-24.2	140	11.5	9.7-13.6	205	10.4	8.9-12.0	122	16.2	13.1-19.7	83	7.0	5.5-8.7
1992	325	16.0	14.2-17.9	163	20.3	17.0-24.1	162	13.2	11.2-15.4	229	11.7	10.2-13.4	102	13.2	10.5-16.3	127	10.6	8.8-12.6
1993	333	16.3	14.5-18.2	176	22.5	19.0-26.4	157	12.5	10.6-14.6	193	9.5	8.2-11.0	98	12.9	10.2-15.9	95	7.6	6.1-9.3
1994	369	17.7	15.9-19.7	199	25.1	21.4-29.2	170	13.3	11.4-15.5	245	11.9	10.5-13.6	141	18.5	15.3-22.1	104	8.3	6.8-10.1
1995	326	15.4	13.7-17.2	166	20.4	17.1-24.0	160	12.2	10.4-14.2	213	10.2	8.8-11.7	102	12.6	10.1-15.5	111	8.6	7.0-10.3
1996	339	15.4	13.8-17.2	193	23.5	20.0-27.3	146	10.7	9.0-12.6	216	10.2	8.9-11.7	122	15.8	12.8-19.1	94	7.1	5.7-8.7
1997	332	15.1	13.4-16.8	185	23.0	19.5-26.9	147	10.7	9.0-12.6	195	8.6	7.4-9.9	113	13.3	10.8-16.2	82	5.8	4.6-7.2
1998	342	14.8	13.2-16.5	184	20.9	17.7-24.4	158	11.1	9.4-13.0	205	9.1	7.9-10.5	113	13.4	10.8-16.2	92	6.6	5.3-8.1
1999	318	13.5	12.0-15.1	171	19.7	16.7-23.2	147	10.1	8.5-11.8	201	8.7	7.5-10.1	107	13.0	10.4-15.9	94	6.5	5.2-7.9
2000	338	14.1	12.6-15.8	174	19.7	16.6-23.0	164	11.0	9.3-12.8	195	8.2	7.0-9.4	108	12.5	10.1-15.3	87	5.8	4.6-7.1
2001	326	13.5	12.1-15.1	159	17.5	14.7-20.7	167	11.1	9.5-13.0	167	7.0	6.0-8.2	93	10.1	8.0-12.5	74	5.1	4.0-6.3
2002	324	12.6	11.3-14.1	162	16.9	14.2-20.0	162	10.2	8.7-11.9	189	7.6	6.5-8.8	101	11.6	9.3-14.3	88	5.6	4.5-6.9
2003	327	12.5	11.1-13.9	168	16.6	14.0-19.5	159	9.9	8.4-11.6	174	6.7	5.7-7.7	93	8.8	7.0-11.0	81	5.1	4.1-6.4
2004	316	12.1	10.8-13.6	153	16.2	13.5-19.1	163	9.9	8.5-11.6	174	6.7	5.8-7.8	84	8.9	7.0-11.1	90	5.5	4.4-6.8
2005	409	15.4	13.9-17.0	211	21.3	18.4-24.6	198	12.0	10.3-13.8	212	8.2	7.1-9.4	122	13.2	10.8-15.9	90	5.5	4.4-6.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	368	13.4	12.1-14.9	202	19.5	16.8-22.6	166	9.8	8.4-11.4	188	7.2	6.1-8.3	107	11.1	8.9-13.5	81	4.9	3.9-6.1
2007	336	12.0	10.7-13.3	173	16.6	14.1-19.4	163	9.4	8.0-11.0	178	6.5	5.6-7.6	86	8.1	6.4-10.1	92	5.5	4.4-6.7
2008	307	10.9	9.7-12.2	163	15.6	13.2-18.4	144	8.1	6.9-9.6	180	6.4	5.4-7.4	93	8.9	7.1-11.0	87	5.0	4.0-6.1
2009	362	12.5	11.2-13.8	203	17.7	15.2-20.5	159	9.1	7.7-10.6	159	5.7	4.9-6.7	85	8.5	6.7-10.5	74	4.2	3.3-5.3
2010	348	11.7	10.5-13.0	184	15.7	13.4-18.3	164	9.1	7.7-10.6	159	5.6	4.7-6.5	92	8.8	7.0-10.8	67	3.7	2.9-4.7
2011	376	12.1	10.8-13.4	190	15.6	13.4-18.1	186	9.8	8.4-11.4	171	5.7	4.9-6.6	98	8.6	6.9-10.6	73	3.9	3.1-4.9
2012	385	12.0	10.8-13.3	212	16.6	14.4-19.1	173	8.9	7.6-10.4	165	5.4	4.6-6.3	85	6.9	5.4-8.6	80	4.3	3.4-5.4
2013	381	11.5	10.4-12.8	204	15.9	13.7-18.3	177	8.9	7.6-10.3	172	5.4	4.6-6.2	91	7.3	5.8-9.1	81	4.1	3.2-5.1
2014	383	11.1	10.0-12.3	209	15.3	13.2-17.7	174	8.4	7.2-9.8	189	5.7	4.9-6.6	111	8.4	6.9-10.2	78	4.0	3.2-5.0
2015	407	11.6	10.5-12.8	219	15.5	13.4-17.8	188	9.0	7.8-10.4	149	4.3	3.6-5.1	90	7.0	5.6-8.6	59	2.7	2.1-3.5
2016	405	11.3	10.2-12.5	225	15.7	13.6-18.0	180	8.6	7.4-10.0	196	5.5	4.7-6.3	119	8.2	6.8-9.9	77	3.6	2.8-4.5
2017	386	10.8	9.8-12.0	199	14.2	12.2-16.4	187	8.8	7.6-10.2	124	3.5	2.9-4.2	65	4.5	3.5-5.8	59	2.8	2.1-3.6
2018	381	10.3	9.3-11.4	205	13.9	12.0-16.1	176	8.0	6.9-9.3	147	4.0	3.3-4.7	87	6.1	4.8-7.6	60	2.7	2.0-3.5
2019	408	10.7	9.7-11.8	203	13.0	11.2-15.0	205	9.2	8.0-10.6	184	4.9	4.2-5.7	111	7.7	6.3-9.4	73	3.2	2.5-4.0
2020	334	8.5	7.6-9.5	175	10.9	9.3-12.7	159	6.8	5.7-7.9	133	3.5	2.9-4.2	70	4.5	3.5-5.8	63	2.7	2.1-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	642	48.2	44.2-52.5	273	51.5	44.7-58.9	369	46.1	41.1-51.4	384	31.2	27.8-34.8	168	35.1	29.2-41.8	216	28.9	24.8-33.3
1977	780	59.9	55.4-64.8	321	64.8	56.9-73.4	459	57.0	51.5-63.0	375	29.5	26.3-32.9	174	36.1	30.4-42.5	201	25.4	21.8-29.5
1978	802	60.3	55.8-65.1	335	65.7	57.8-74.3	467	57.4	51.9-63.2	462	36.5	32.9-40.3	202	41.7	35.3-48.7	260	33.5	29.2-38.1
1979	844	60.5	56.2-65.1	370	68.9	61.1-77.3	474	55.7	50.5-61.2	476	35.4	32.0-39.0	206	40.9	34.8-47.7	270	32.3	28.3-36.7
1980	889	60.6	56.3-65.0	392	70.2	62.5-78.5	497	55.2	50.2-60.6	463	32.7	29.6-36.1	197	36.4	30.9-42.6	266	30.7	26.9-34.8
1981	866	57.6	53.6-61.9	375	67.0	59.6-75.1	491	52.7	47.9-57.8	496	35.1	31.9-38.6	201	38.8	33.0-45.2	295	33.3	29.4-37.5
1982	873	54.9	51.1-58.9	398	65.0	58.0-72.5	475	48.9	44.4-53.7	468	30.5	27.6-33.5	208	35.4	30.2-41.2	260	27.6	24.2-31.3
1983	871	53.0	49.3-56.8	383	64.0	56.9-71.5	488	47.6	43.3-52.1	443	27.2	24.5-30.0	203	31.6	26.9-36.8	240	24.3	21.2-27.8
1984	930	55.5	51.8-59.4	430	66.1	59.3-73.4	500	49.1	44.8-53.8	483	29.6	26.8-32.5	211	35.3	30.2-41.0	272	26.7	23.5-30.2
1985	987	57.0	53.3-60.9	447	69.9	62.8-77.4	540	50.1	45.9-54.7	492	30.1	27.3-33.0	201	33.6	28.6-39.1	291	28.4	25.1-32.0
1986	1,042	57.9	54.3-61.8	473	69.5	62.7-76.8	569	51.4	47.2-56.0	474	27.5	25.0-30.3	215	34.0	29.1-39.4	259	24.1	21.2-27.3
1987	962	52.2	48.8-55.7	435	63.1	56.6-70.0	527	46.1	42.2-50.4	479	26.9	24.4-29.6	222	34.5	29.6-39.9	257	23.0	20.2-26.1
1988	985	52.8	49.3-56.3	425	60.8	54.5-67.5	560	48.3	44.3-52.6	476	26.7	24.3-29.4	221	34.2	29.3-39.5	255	22.8	20.0-25.8
1989	1,022	53.3	50.0-56.9	486	69.2	62.5-76.3	536	45.0	41.1-49.0	469	25.2	22.8-27.6	238	36.3	31.3-41.7	231	19.5	17.0-22.2
1990	1,011	52.6	49.3-56.1	435	61.0	54.8-67.5	576	47.8	43.9-52.0	465	25.2	22.9-27.7	212	31.8	27.2-36.9	253	21.8	19.1-24.7
1991	1,082	55.1	51.8-58.7	468	62.4	56.3-68.9	614	50.6	46.6-54.8	495	26.5	24.1-29.0	216	32.9	28.2-38.1	279	23.4	20.6-26.3
1992	1,076	53.1	49.8-56.5	450	60.0	54.0-66.4	626	49.6	45.7-53.7	477	24.4	22.2-26.7	204	29.6	25.3-34.4	273	21.9	19.4-24.7
1993	1,086	52.5	49.3-55.9	492	62.7	56.7-69.1	594	46.6	42.8-50.5	484	24.2	22.0-26.5	211	28.4	24.3-32.9	273	21.8	19.3-24.6
1994	1,174	55.8	52.6-59.2	529	67.7	61.5-74.3	645	49.3	45.5-53.3	535	26.1	23.8-28.5	254	32.8	28.6-37.5	281	22.0	19.5-24.7
1995	1,171	53.5	50.4-56.7	525	63.7	57.9-69.9	646	47.7	44.0-51.6	536	25.8	23.6-28.1	233	30.0	26.0-34.5	303	23.2	20.6-26.0
1996	1,250	56.2	53.0-59.5	560	66.6	60.7-72.8	690	50.1	46.4-54.1	547	25.9	23.7-28.2	263	34.0	29.7-38.6	284	21.3	18.9-24.0
1997	1,291	57.3	54.1-60.6	590	70.6	64.5-77.0	701	50.0	46.3-53.8	570	26.1	23.9-28.4	271	34.3	30.0-39.0	299	21.7	19.3-24.4
1998	1,288	55.6	52.5-58.8	546	62.1	56.6-67.9	742	51.7	48.0-55.6	513	22.7	20.8-24.8	226	26.9	23.2-30.9	287	20.4	18.1-22.9
1999	1,427	59.0	55.9-62.3	608	66.2	60.6-72.0	819	54.9	51.2-58.9	597	25.9	23.8-28.1	255	30.8	26.9-35.1	342	23.4	21.0-26.0
2000	1,287	51.5	48.7-54.5	574	61.7	56.4-67.3	713	46.1	42.8-49.7	548	22.7	20.8-24.8	249	27.9	24.3-31.9	299	19.9	17.6-22.2
2001	1,368	53.6	50.7-56.5	605	63.2	57.9-68.8	763	48.4	45.0-51.9	543	22.0	20.2-24.0	237	27.2	23.6-31.1	306	19.7	17.6-22.1
2002	1,391	53.0	50.1-55.9	602	59.4	54.4-64.7	789	49.1	45.7-52.6	541	21.7	19.9-23.7	248	27.9	24.2-31.8	293	18.6	16.5-20.9
2003	1,460	54.8	52.0-57.7	669	67.1	61.8-72.8	791	48.2	44.9-51.7	547	21.4	19.6-23.3	258	28.1	24.5-32.0	289	17.9	15.9-20.1
2004	1,486	54.2	51.4-57.1	650	62.0	57.0-67.3	836	49.8	46.5-53.3	536	20.6	18.9-22.5	258	26.7	23.3-30.4	278	17.1	15.2-19.3
2005	1,417	50.7	48.1-53.5	650	60.4	55.6-65.5	767	45.1	41.9-48.4	499	18.8	17.2-20.6	220	22.4	19.3-25.7	279	16.7	14.8-18.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,479	52.2	49.5-55.0	636	57.9	53.2-62.8	843	48.9	45.7-52.4	527	19.4	17.8-21.2	233	23.5	20.4-26.9	294	17.3	15.3-19.4
2007	1,456	49.9	47.3-52.6	696	59.8	55.2-64.7	760	43.3	40.3-46.5	547	19.5	17.9-21.2	251	23.3	20.3-26.5	296	17.2	15.3-19.3
2008	1,497	50.4	47.8-53.1	697	59.8	55.1-64.7	800	44.6	41.6-47.9	489	17.2	15.7-18.8	231	21.5	18.6-24.6	258	14.7	12.9-16.6
2009	1,473	48.0	45.5-50.5	710	57.5	53.1-62.1	763	41.4	38.5-44.5	507	17.6	16.0-19.2	246	22.7	19.7-25.8	261	14.5	12.8-16.4
2010	1,408	45.3	42.9-47.8	668	55.5	51.1-60.2	740	39.6	36.7-42.6	481	15.9	14.5-17.4	247	21.7	18.9-24.7	234	12.4	10.8-14.1
2011	1,455	45.1	42.8-47.5	708	54.8	50.6-59.2	747	38.6	35.8-41.5	494	16.1	14.7-17.7	214	18.4	15.9-21.2	280	14.7	13.1-16.6
2012	1,370	41.4	39.2-43.7	667	50.5	46.5-54.7	703	35.6	33.0-38.4	537	16.9	15.5-18.5	260	21.3	18.7-24.3	277	14.3	12.6-16.1
2013	1,417	41.5	39.4-43.8	673	49.2	45.4-53.3	744	36.6	34.0-39.4	481	14.6	13.3-16.0	223	17.0	14.7-19.5	258	12.8	11.3-14.5
2014	1,509	43.4	41.2-45.7	710	50.6	46.8-54.7	799	38.9	36.2-41.7	510	15.1	13.8-16.5	246	18.5	16.1-21.0	264	12.9	11.4-14.6
2015	1,423	39.5	37.4-41.6	714	48.9	45.2-52.8	709	33.3	30.9-35.9	465	13.5	12.3-14.8	205	15.0	12.9-17.3	260	12.4	10.9-14.0
2016	1,507	41.1	39.0-43.3	750	50.9	47.2-54.9	757	34.9	32.4-37.6	502	13.9	12.7-15.2	228	15.7	13.7-18.0	274	12.6	11.1-14.2
2017	1,397	37.7	35.7-39.8	697	46.2	42.7-49.9	700	32.2	29.8-34.7	467	12.8	11.6-14.0	235	16.5	14.4-18.8	232	10.5	9.2-12.0
2018	1,424	37.4	35.5-39.4	665	42.5	39.2-46.0	759	34.0	31.6-36.5	500	13.5	12.3-14.8	237	16.5	14.4-18.8	263	11.7	10.3-13.3
2019	1,463	38.1	36.1-40.1	702	44.6	41.3-48.2	761	33.9	31.5-36.4	490	12.9	11.7-14.1	236	15.9	13.9-18.1	254	10.8	9.5-12.3
2020	1,258	32.3	30.5-34.2	633	39.3	36.2-42.6	625	27.6	25.5-30.0	503	12.7	11.6-13.9	246	15.5	13.5-17.6	257	11.1	9.7-12.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	471	35.6	32.1-39.3	192	37.6	31.7-44.2	279	34.5	30.2-39.1	312	25.2	22.2-28.5	135	28.6	23.2-34.7	177	23.4	19.8-27.4
1977	567	44.7	40.7-48.9	212	44.5	37.9-51.9	355	45.0	40.0-50.3	290	23.1	20.3-26.2	124	26.8	21.8-32.5	166	21.0	17.7-24.7
1978	603	46.2	42.2-50.4	246	50.5	43.4-58.2	357	44.1	39.3-49.3	382	30.1	26.9-33.6	163	34.4	28.5-40.9	219	27.8	24.0-32.0
1979	655	47.2	43.4-51.3	285	53.2	46.4-60.7	370	43.7	39.1-48.7	404	29.9	26.8-33.2	171	34.2	28.6-40.4	233	27.6	24.0-31.6
1980	655	45.2	41.5-49.1	281	51.4	44.7-58.6	374	41.8	37.4-46.5	378	27.1	24.2-30.2	158	29.9	24.8-35.6	220	25.6	22.1-29.4
1981	630	43.2	39.7-47.0	271	50.3	43.7-57.4	359	39.6	35.4-44.1	427	30.2	27.2-33.4	170	33.1	27.7-39.0	257	28.9	25.3-32.8
1982	648	41.2	37.9-44.7	294	49.4	43.2-56.0	354	36.6	32.7-40.7	390	25.6	23.0-28.5	168	29.8	24.9-35.3	222	23.4	20.3-26.9
1983	656	39.9	36.7-43.3	284	48.2	42.1-54.9	372	36.0	32.3-40.0	371	23.0	20.5-25.6	165	26.3	22.0-31.1	206	20.9	18.0-24.1
1984	689	41.8	38.6-45.2	307	47.5	41.8-53.7	382	38.2	34.3-42.3	420	25.6	23.1-28.4	183	30.6	25.8-35.9	237	23.2	20.3-26.5
1985	743	43.8	40.5-47.2	322	52.1	45.9-58.8	421	39.5	35.7-43.6	421	25.6	23.1-28.4	165	27.8	23.2-32.8	256	24.8	21.7-28.1
1986	772	43.1	40.0-46.4	339	50.9	45.0-57.2	433	39.0	35.3-43.0	411	23.9	21.5-26.4	186	29.3	24.8-34.3	225	20.9	18.2-24.0
1987	734	40.0	37.0-43.1	328	47.4	41.9-53.4	406	35.7	32.2-39.4	423	23.7	21.4-26.2	194	29.9	25.4-34.9	229	20.5	17.8-23.4
1988	770	41.5	38.5-44.7	332	47.6	42.1-53.5	438	38.1	34.6-42.0	416	23.1	20.8-25.6	192	29.1	24.6-34.1	224	19.9	17.3-22.8
1989	792	41.7	38.7-44.9	368	53.2	47.3-59.5	424	35.8	32.4-39.5	417	22.3	20.1-24.7	209	32.0	27.3-37.1	208	17.5	15.1-20.1
1990	769	40.2	37.3-43.3	324	46.1	40.8-51.9	445	37.0	33.5-40.6	405	21.9	19.7-24.3	183	27.1	22.9-31.8	222	19.1	16.6-21.9
1991	830	42.7	39.7-45.8	363	49.0	43.6-54.9	467	38.7	35.2-42.5	446	23.9	21.6-26.3	195	29.5	25.1-34.5	251	21.0	18.5-23.9
1992	831	41.3	38.4-44.3	334	45.5	40.3-51.2	497	39.4	36.0-43.1	414	21.2	19.1-23.4	176	26.0	22.0-30.6	238	19.0	16.7-21.7
1993	817	40.2	37.4-43.2	357	47.3	42.0-53.0	460	36.3	33.0-39.8	430	21.3	19.3-23.5	185	24.6	20.8-28.8	245	19.5	17.1-22.1
1994	919	44.2	41.3-47.2	398	51.7	46.3-57.6	521	40.1	36.7-43.8	469	23.0	20.9-25.2	222	29.1	25.1-33.6	247	19.4	17.0-22.0
1995	898	41.7	38.9-44.6	395	49.3	44.1-54.8	503	37.4	34.2-40.9	482	23.3	21.2-25.5	212	27.7	23.8-32.0	270	20.7	18.3-23.4
1996	937	42.3	39.6-45.2	410	49.5	44.4-55.0	527	38.4	35.1-41.8	469	22.3	20.3-24.5	222	29.2	25.2-33.6	247	18.5	16.2-21.0
1997	992	44.0	41.2-46.9	435	52.7	47.4-58.3	557	39.6	36.3-43.0	492	22.6	20.6-24.7	231	29.9	25.8-34.3	261	18.9	16.7-21.4
1998	994	43.3	40.6-46.1	415	48.5	43.5-53.7	579	40.5	37.2-44.0	453	20.0	18.2-22.0	197	23.4	20.0-27.2	256	18.1	15.9-20.5
1999	1,078	45.0	42.3-47.8	438	49.0	44.1-54.1	640	43.0	39.7-46.5	529	23.0	21.0-25.0	217	26.5	22.9-30.6	312	21.3	19.0-23.8
2000	982	39.8	37.3-42.5	426	47.1	42.4-52.1	556	36.2	33.2-39.3	484	20.2	18.4-22.1	218	25.0	21.6-28.9	266	17.7	15.6-19.9
2001	1,027	40.8	38.2-43.4	443	47.6	42.9-52.5	584	37.2	34.2-40.4	487	19.9	18.1-21.8	209	24.2	20.8-27.9	278	18.0	15.9-20.3
2002	1,081	41.6	39.1-44.2	443	44.4	40.1-49.1	638	39.9	36.8-43.1	483	19.5	17.7-21.3	216	24.5	21.1-28.2	267	17.0	15.0-19.1
2003	1,100	41.8	39.3-44.4	487	50.4	45.7-55.4	613	37.5	34.5-40.6	466	18.4	16.7-20.2	209	23.2	20.0-26.8	257	15.9	14.0-18.0
2004	1,120	41.4	38.9-43.9	481	46.7	42.3-51.4	639	38.4	35.5-41.5	466	18.0	16.4-19.8	218	22.5	19.4-25.9	248	15.4	13.5-17.4
2005	1,077	39.0	36.7-41.5	481	45.8	41.6-50.4	596	35.2	32.4-38.2	432	16.4	14.9-18.1	183	18.8	16.1-21.9	249	15.0	13.2-17.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,097	39.2	36.9-41.7	465	43.0	38.9-47.4	632	37.1	34.2-40.1	457	16.9	15.4-18.6	204	20.8	17.8-24.0	253	14.9	13.1-16.9
2007	1,081	37.5	35.3-39.9	506	44.7	40.7-49.0	575	32.9	30.2-35.7	465	16.6	15.1-18.2	208	19.4	16.7-22.4	257	14.9	13.1-16.9
2008	1,137	38.8	36.5-41.2	513	45.1	41.1-49.5	624	35.1	32.3-37.9	421	14.9	13.4-16.4	194	18.2	15.6-21.1	227	12.9	11.2-14.7
2009	1,081	35.7	33.6-37.9	504	41.8	38.1-45.9	577	31.5	28.9-34.2	441	15.4	14.0-17.0	211	19.9	17.2-22.9	230	12.8	11.2-14.6
2010	1,035	33.8	31.8-36.0	473	40.1	36.3-44.1	562	30.4	27.9-33.0	408	13.6	12.3-15.0	209	18.7	16.1-21.6	199	10.6	9.1-12.2
2011	1,079	33.8	31.8-36.0	510	40.4	36.8-44.2	569	29.5	27.1-32.1	418	13.8	12.5-15.2	177	15.7	13.3-18.3	241	12.7	11.1-14.5
2012	1,000	30.7	28.8-32.7	466	36.4	33.0-40.0	534	27.3	25.0-29.7	450	14.3	12.9-15.7	206	17.0	14.6-19.6	244	12.6	11.1-14.3
2013	1,011	30.0	28.1-31.9	474	35.5	32.2-39.0	537	26.5	24.3-28.9	418	12.7	11.5-14.0	186	14.2	12.1-16.5	232	11.6	10.1-13.2
2014	1,105	32.2	30.2-34.1	503	36.5	33.3-40.0	602	29.5	27.2-32.0	420	12.4	11.2-13.6	198	14.8	12.7-17.1	222	10.8	9.4-12.4
2015	1,008	28.2	26.5-30.1	486	33.9	30.9-37.2	522	24.5	22.4-26.7	389	11.3	10.2-12.5	173	12.6	10.7-14.8	216	10.3	8.9-11.8
2016	1,080	29.6	27.8-31.5	513	35.0	31.9-38.3	567	26.3	24.1-28.6	399	11.0	9.9-12.2	173	11.8	10.0-13.7	226	10.4	9.1-11.9
2017	1,012	27.4	25.7-29.2	475	31.9	29.0-35.0	537	24.5	22.5-26.8	385	10.5	9.5-11.7	188	13.2	11.3-15.3	197	8.9	7.7-10.3
2018	1,026	27.0	25.3-28.7	446	28.7	26.0-31.6	580	25.9	23.8-28.2	417	11.3	10.2-12.5	195	13.6	11.7-15.7	222	9.9	8.6-11.3
2019	1,090	28.5	26.8-30.2	495	31.9	29.1-35.0	595	26.3	24.2-28.6	416	11.0	9.9-12.1	194	13.2	11.3-15.2	222	9.5	8.3-10.9
2020	944	24.3	22.7-25.9	456	28.7	26.0-31.5	488	21.5	19.6-23.5	420	10.7	9.7-11.8	200	12.8	11.1-14.8	220	9.4	8.2-10.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	171	12.6	10.6-14.9	81	13.8	10.6-17.6	90	11.6	9.1-14.5	72	6.0	4.5-7.7	33	6.6	4.2-9.6	39	5.5	3.8-7.6
1977	213	15.3	13.1-17.7	109	20.3	16.2-25.0	104	12.1	9.7-14.9	85	6.4	5.0-8.0	50	9.3	6.7-12.5	35	4.4	3.0-6.3
1978	199	14.1	12.0-16.4	89	15.3	11.8-19.3	110	13.3	10.7-16.2	80	6.4	5.0-8.1	39	7.3	5.0-10.2	41	5.7	3.9-7.8
1979	189	13.3	11.3-15.5	85	15.7	12.1-19.9	104	11.9	9.6-14.6	72	5.5	4.2-7.0	35	6.7	4.4-9.6	37	4.7	3.2-6.6
1980	234	15.4	13.4-17.7	111	18.8	15.1-23.1	123	13.4	11.0-16.1	85	5.7	4.5-7.1	39	6.6	4.5-9.2	46	5.1	3.6-6.9
1981	236	14.4	12.5-16.5	104	16.8	13.3-20.7	132	13.1	10.9-15.7	69	4.9	3.7-6.3	31	5.8	3.7-8.4	38	4.4	3.1-6.1
1982	225	13.7	11.9-15.7	104	15.6	12.5-19.2	121	12.3	10.1-14.8	78	4.8	3.7-6.1	40	5.6	3.9-7.9	38	4.2	2.9-5.8
1983	215	13.1	11.3-15.1	99	15.8	12.4-19.6	116	11.5	9.5-13.9	72	4.2	3.2-5.4	38	5.3	3.6-7.5	34	3.4	2.3-4.9
1984	241	13.8	12.0-15.7	123	18.7	15.1-22.7	118	11.0	9.1-13.2	63	3.9	3.0-5.1	28	4.8	3.0-7.0	35	3.5	2.4-4.9
1985	244	13.2	11.5-15.1	125	17.8	14.5-21.6	119	10.6	8.7-12.7	71	4.4	3.4-5.6	36	5.8	3.9-8.3	35	3.6	2.5-5.0
1986	270	14.8	13.0-16.8	134	18.6	15.3-22.4	136	12.4	10.4-14.8	63	3.7	2.8-4.7	29	4.7	2.9-6.9	34	3.2	2.2-4.4
1987	228	12.2	10.6-14.0	107	15.7	12.5-19.3	121	10.5	8.6-12.5	56	3.3	2.4-4.3	28	4.6	2.9-6.8	28	2.5	1.7-3.7
1988	215	11.2	9.7-12.9	93	13.2	10.4-16.5	122	10.2	8.4-12.2	60	3.6	2.7-4.7	29	5.1	3.3-7.4	31	2.9	1.9-4.1
1989	230	11.6	10.1-13.3	118	16.0	12.9-19.4	112	9.1	7.5-11.0	52	2.8	2.1-3.7	29	4.3	2.7-6.4	23	2.0	1.3-3.0
1990	242	12.4	10.8-14.1	111	14.8	12.0-18.1	131	10.9	9.0-12.9	60	3.3	2.5-4.3	29	4.7	2.9-7.0	31	2.7	1.8-3.8
1991	252	12.5	10.9-14.2	105	13.4	10.8-16.5	147	11.8	10.0-13.9	49	2.6	1.9-3.5	21	3.3	1.9-5.2	28	2.3	1.5-3.4
1992	245	11.8	10.3-13.4	116	14.5	11.7-17.7	129	10.2	8.5-12.1	63	3.2	2.4-4.1	28	3.6	2.3-5.3	35	2.9	2.0-4.0
1993	269	12.3	10.8-13.9	135	15.4	12.7-18.5	134	10.2	8.5-12.1	54	2.8	2.1-3.7	26	3.8	2.3-5.7	28	2.4	1.6-3.4
1994	255	11.6	10.2-13.2	131	16.0	13.1-19.2	124	9.2	7.6-10.9	66	3.1	2.4-4.0	32	3.7	2.5-5.3	34	2.6	1.8-3.7
1995	273	11.8	10.4-13.3	130	14.4	11.9-17.4	143	10.3	8.6-12.1	54	2.5	1.9-3.3	21	2.4	1.4-3.6	33	2.5	1.7-3.5
1996	313	13.9	12.3-15.5	150	17.1	14.3-20.3	163	11.8	10.0-13.7	78	3.6	2.8-4.5	41	4.8	3.3-6.6	37	2.8	2.0-3.9
1997	299	13.3	11.8-14.9	155	17.9	15.0-21.1	144	10.4	8.8-12.3	78	3.5	2.8-4.4	40	4.5	3.1-6.2	38	2.8	2.0-3.8
1998	294	12.3	10.9-13.8	131	13.6	11.2-16.3	163	11.2	9.6-13.1	60	2.7	2.1-3.5	29	3.4	2.2-5.0	31	2.3	1.5-3.2
1999	349	14.1	12.6-15.6	170	17.2	14.6-20.2	179	12.0	10.3-13.9	68	2.9	2.2-3.7	38	4.3	2.9-5.9	30	2.1	1.4-3.0
2000	305	11.7	10.4-13.1	148	14.6	12.2-17.3	157	9.9	8.4-11.6	64	2.5	1.9-3.3	31	2.9	1.9-4.1	33	2.2	1.5-3.1
2001	341	12.8	11.5-14.3	162	15.6	13.1-18.4	179	11.1	9.6-12.9	56	2.2	1.6-2.8	28	3.0	1.9-4.5	28	1.7	1.1-2.5
2002	310	11.3	10.1-12.7	159	14.9	12.6-17.6	151	9.2	7.8-10.8	58	2.2	1.7-2.9	32	3.4	2.2-4.9	26	1.7	1.1-2.4
2003	360	13.0	11.7-14.5	182	16.7	14.2-19.5	178	10.7	9.2-12.4	81	3.0	2.4-3.8	49	4.8	3.5-6.5	32	2.0	1.3-2.8
2004	366	12.8	11.5-14.2	169	15.3	12.9-17.9	197	11.4	9.9-13.2	70	2.6	2.0-3.3	40	4.1	2.9-5.7	30	1.8	1.2-2.5
2005	340	11.7	10.5-13.0	169	14.6	12.4-17.1	171	9.8	8.4-11.4	67	2.4	1.9-3.1	37	3.5	2.4-4.9	30	1.8	1.2-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	382	13.0	11.7-14.4	171	14.9	12.6-17.4	211	11.9	10.3-13.6	70	2.5	2.0-3.2	29	2.7	1.8-4.0	41	2.4	1.7-3.2
2007	375	12.4	11.1-13.7	190	15.1	12.9-17.5	185	10.4	9.0-12.0	82	2.9	2.3-3.6	43	3.9	2.7-5.3	39	2.3	1.6-3.1
2008	360	11.6	10.4-12.9	184	14.6	12.5-17.0	176	9.6	8.2-11.1	68	2.4	1.8-3.0	37	3.2	2.2-4.5	31	1.8	1.2-2.5
2009	392	12.3	11.0-13.6	206	15.6	13.5-18.0	186	10.0	8.6-11.5	66	2.1	1.6-2.7	35	2.8	1.9-3.9	31	1.7	1.1-2.4
2010	373	11.5	10.3-12.7	195	15.5	13.2-17.9	178	9.2	7.9-10.7	73	2.3	1.8-2.9	38	3.0	2.0-4.1	35	1.8	1.3-2.6
2011	376	11.3	10.1-12.5	198	14.4	12.3-16.6	178	9.1	7.8-10.5	76	2.3	1.8-3.0	37	2.7	1.9-3.8	39	2.0	1.4-2.8
2012	370	10.7	9.6-11.9	201	14.1	12.1-16.3	169	8.3	7.1-9.7	87	2.7	2.1-3.3	54	4.4	3.2-5.8	33	1.7	1.1-2.4
2013	406	11.6	10.4-12.8	199	13.8	11.8-15.9	207	10.1	8.8-11.6	63	1.8	1.4-2.4	37	2.8	1.9-3.9	26	1.2	0.8-1.8
2014	404	11.3	10.2-12.4	207	14.1	12.1-16.2	197	9.4	8.1-10.8	90	2.7	2.2-3.3	48	3.7	2.7-4.9	42	2.1	1.5-2.8
2015	415	11.3	10.2-12.4	228	14.9	13.0-17.1	187	8.8	7.6-10.2	76	2.2	1.7-2.8	32	2.4	1.6-3.4	44	2.1	1.5-2.8
2016	427	11.5	10.5-12.7	237	15.9	13.9-18.2	190	8.7	7.5-10.0	103	2.9	2.4-3.5	55	4.0	3.0-5.2	48	2.2	1.6-2.9
2017	385	10.3	9.2-11.4	222	14.4	12.5-16.5	163	7.6	6.5-8.9	82	2.3	1.8-2.8	47	3.3	2.4-4.4	35	1.6	1.1-2.2
2018	398	10.4	9.4-11.5	219	13.8	12.0-15.9	179	8.0	6.9-9.3	83	2.2	1.8-2.8	42	2.9	2.1-4.0	41	1.8	1.3-2.5
2019	373	9.6	8.6-10.7	207	12.7	11.0-14.6	166	7.6	6.4-8.8	74	1.9	1.5-2.4	42	2.8	2.0-3.8	32	1.3	0.9-1.9
2020	314	8.0	7.2-9.0	177	10.7	9.1-12.4	137	6.2	5.2-7.3	83	2.1	1.6-2.6	46	2.6	1.9-3.5	37	1.7	1.2-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	79	5.5	4.2-7.0	55	9.5	6.9-12.7	24	2.9	1.8-4.4	82	6.1	4.7-7.7	58	10.5	7.6-13.9	24	3.2	1.9-4.9
1977	74	4.9	3.7-6.2	51	8.0	5.6-10.9	23	2.8	1.7-4.3	92	5.9	4.7-7.4	66	10.9	8.1-14.3	26	2.8	1.8-4.1
1978	101	6.7	5.3-8.2	60	9.7	7.2-12.8	41	4.6	3.2-6.4	89	5.8	4.6-7.3	45	7.0	4.9-9.7	44	5.0	3.5-6.8
1979	75	4.7	3.6-6.1	44	7.0	4.8-9.7	31	3.3	2.2-4.8	93	6.4	5.0-7.9	51	8.9	6.4-12.0	42	4.8	3.4-6.6
1980	88	5.4	4.2-6.7	59	8.7	6.4-11.6	29	3.2	2.1-4.6	85	5.3	4.1-6.6	55	8.5	6.2-11.4	30	3.3	2.2-4.7
1981	91	5.3	4.2-6.6	58	8.5	6.3-11.3	33	3.3	2.2-4.6	93	5.9	4.7-7.2	55	8.1	5.9-10.8	38	4.3	3.0-5.9
1982	101	5.8	4.7-7.1	74	10.6	8.1-13.6	27	2.7	1.8-4.0	95	5.7	4.6-7.0	63	9.0	6.8-11.7	32	3.5	2.3-5.0
1983	96	5.5	4.4-6.7	65	9.5	7.1-12.5	31	2.9	1.9-4.2	121	7.0	5.8-8.5	77	11.3	8.7-14.4	44	4.4	3.1-5.9
1984	101	5.6	4.5-6.9	69	10.4	7.8-13.6	32	2.8	1.9-4.0	113	6.4	5.2-7.7	68	10.9	8.2-14.2	45	3.9	2.8-5.2
1985	100	5.2	4.2-6.4	62	8.5	6.3-11.1	38	3.2	2.2-4.4	102	5.7	4.6-6.9	58	8.2	6.1-10.9	44	4.1	2.9-5.5
1986	98	5.1	4.1-6.3	65	8.9	6.7-11.6	33	2.8	1.9-4.0	105	5.5	4.5-6.7	66	8.9	6.7-11.5	39	3.4	2.4-4.7
1987	102	5.0	4.0-6.1	58	7.1	5.2-9.5	44	3.7	2.7-5.0	112	6.2	5.0-7.5	57	8.5	6.3-11.3	55	4.9	3.7-6.4
1988	100	4.9	3.9-6.0	65	8.1	6.0-10.6	35	2.9	2.0-4.1	104	5.2	4.2-6.4	67	8.8	6.6-11.4	37	3.1	2.2-4.3
1989	110	5.6	4.6-6.8	68	9.1	6.8-11.8	42	3.6	2.6-4.9	111	5.6	4.5-6.8	61	7.7	5.7-10.2	50	4.2	3.1-5.6
1990	131	6.3	5.2-7.5	81	9.6	7.5-12.1	50	4.1	3.1-5.5	118	5.7	4.7-6.9	65	8.2	6.2-10.7	53	4.3	3.2-5.6
1991	136	6.2	5.1-7.3	90	10.5	8.2-13.2	46	3.5	2.5-4.7	129	6.1	5.1-7.3	76	9.4	7.2-12.0	53	4.1	3.1-5.4
1992	132	6.3	5.2-7.5	83	10.2	8.0-12.9	49	3.9	2.8-5.1	124	5.8	4.8-6.9	72	8.8	6.8-11.2	52	4.0	2.9-5.2
1993	130	6.0	5.0-7.2	83	9.5	7.4-12.0	47	3.7	2.7-4.9	132	6.2	5.2-7.4	76	8.7	6.7-11.0	56	4.5	3.4-5.9
1994	140	6.2	5.2-7.3	84	9.2	7.2-11.6	56	4.2	3.1-5.4	120	5.6	4.6-6.8	76	9.3	7.2-11.8	44	3.4	2.5-4.6
1995	155	6.6	5.6-7.8	101	11.0	8.7-13.5	54	3.9	2.9-5.1	130	5.6	4.7-6.7	79	8.9	6.9-11.2	51	3.7	2.7-4.9
1996	170	7.3	6.2-8.6	120	13.3	10.8-16.2	50	3.7	2.7-4.9	146	6.5	5.5-7.7	90	10.4	8.2-12.9	56	4.1	3.1-5.4
1997	174	7.2	6.2-8.4	110	11.5	9.3-14.0	64	4.5	3.5-5.7	129	5.4	4.5-6.5	78	8.3	6.5-10.6	51	3.6	2.7-4.8
1998	190	7.6	6.5-8.7	131	12.4	10.3-14.9	59	4.1	3.1-5.3	159	6.5	5.5-7.6	105	10.6	8.5-12.9	54	3.8	2.9-5.0
1999	181	7.1	6.0-8.2	110	10.3	8.4-12.6	71	4.7	3.6-5.9	112	4.5	3.7-5.4	65	6.2	4.8-8.0	47	3.2	2.3-4.3
2000	188	7.0	6.0-8.1	127	11.1	9.2-13.3	61	3.9	3.0-5.1	157	6.1	5.2-7.1	103	10.1	8.1-12.4	54	3.6	2.7-4.6
2001	190	6.8	5.9-7.9	141	12.3	10.2-14.6	49	3.1	2.3-4.1	143	5.3	4.5-6.3	86	7.8	6.2-9.8	57	3.6	2.7-4.7
2002	206	7.4	6.4-8.5	148	12.8	10.7-15.2	58	3.7	2.8-4.7	136	5.1	4.3-6.1	86	8.0	6.3-10.1	50	3.2	2.4-4.2
2003	230	7.8	6.8-8.9	162	13.0	11.0-15.3	68	4.1	3.2-5.2	154	5.5	4.6-6.4	107	9.4	7.6-11.5	47	2.9	2.1-3.8
2004	279	9.5	8.4-10.7	196	16.1	13.8-18.7	83	4.9	3.9-6.1	195	6.9	6.0-8.0	119	10.0	8.2-12.1	76	4.7	3.7-5.8
2005	285	9.4	8.3-10.5	215	17.1	14.8-19.6	70	4.0	3.1-5.1	176	6.0	5.2-7.0	121	10.3	8.4-12.3	55	3.3	2.4-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	254	8.3	7.3-9.5	189	15.2	13.0-17.6	65	3.7	2.9-4.7	174	5.9	5.0-6.8	127	10.3	8.5-12.4	47	2.8	2.1-3.7
2007	325	10.3	9.2-11.5	237	18.4	16.0-21.0	88	4.9	3.9-6.0	197	6.6	5.7-7.6	131	10.7	8.8-12.8	66	3.8	2.9-4.8
2008	328	10.0	9.0-11.2	245	17.8	15.5-20.3	83	4.5	3.6-5.6	198	6.3	5.4-7.2	144	11.1	9.3-13.2	54	2.9	2.2-3.8
2009	372	11.3	10.2-12.6	268	19.6	17.2-22.3	104	5.5	4.5-6.7	214	6.7	5.8-7.7	163	12.3	10.4-14.5	51	2.8	2.1-3.7
2010	384	11.2	10.1-12.4	272	18.9	16.6-21.4	112	5.8	4.7-7.0	250	7.5	6.6-8.6	172	12.7	10.8-14.9	78	3.9	3.1-4.9
2011	379	10.9	9.8-12.0	268	17.8	15.6-20.1	111	5.8	4.7-7.0	241	7.1	6.3-8.1	159	10.9	9.2-12.8	82	4.3	3.4-5.4
2012	433	12.1	11.0-13.4	312	21.0	18.6-23.6	121	6.0	5.0-7.2	255	7.4	6.5-8.4	179	12.2	10.4-14.2	76	4.0	3.1-5.0
2013	410	11.0	10.0-12.2	276	17.5	15.4-19.7	134	6.4	5.3-7.6	246	6.9	6.1-7.9	173	12.0	10.2-14.0	73	3.5	2.7-4.4
2014	448	11.7	10.6-12.8	313	19.0	16.9-21.3	135	6.3	5.3-7.5	244	6.7	5.9-7.6	156	10.2	8.6-12.0	88	4.2	3.4-5.2
2015	412	10.6	9.6-11.7	292	18.0	15.9-20.2	120	5.4	4.4-6.5	246	6.5	5.7-7.4	160	10.2	8.6-12.0	86	4.0	3.2-5.0
2016	474	11.8	10.7-12.9	349	20.6	18.5-23.0	125	5.4	4.5-6.5	268	7.0	6.2-7.9	199	12.9	11.1-14.9	69	3.1	2.4-4.0
2017	454	11.4	10.3-12.5	325	19.7	17.6-22.1	129	5.5	4.6-6.6	247	6.3	5.5-7.2	165	10.0	8.5-11.7	82	3.6	2.9-4.5
2018	377	9.2	8.3-10.2	253	14.9	13.0-16.9	124	5.2	4.3-6.3	253	6.4	5.6-7.3	162	10.1	8.5-11.8	91	3.9	3.1-4.8
2019	397	9.7	8.8-10.8	264	15.8	13.9-17.9	133	5.6	4.7-6.6	250	6.2	5.4-7.0	156	9.5	8.0-11.2	94	4.0	3.2-4.9
2020	335	8.0	7.2-9.0	206	11.7	10.1-13.5	129	5.3	4.4-6.4	221	5.4	4.7-6.2	138	8.4	7.0-10.0	83	3.4	2.7-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	203	15.3	13.0-17.7	85	14.5	11.3-18.3	118	15.4	12.6-18.7	189	13.4	11.4-15.7	89	15.0	11.7-18.9	100	12.3	9.8-15.1
1977	200	14.1	12.0-16.4	105	17.8	14.0-22.2	95	11.6	9.2-14.3	182	12.8	10.8-15.0	99	16.9	13.3-21.2	83	10.1	7.9-12.7
1978	218	15.3	13.2-17.7	101	16.3	12.9-20.4	117	14.3	11.7-17.4	184	12.5	10.6-14.7	82	13.5	10.4-17.2	102	11.8	9.5-14.5
1979	229	16.1	13.9-18.6	113	19.1	15.3-23.5	116	14.1	11.5-17.0	203	14.3	12.2-16.6	87	13.4	10.5-16.9	116	14.3	11.7-17.4
1980	208	13.7	11.8-15.9	112	18.7	15.0-23.1	96	10.6	8.5-13.0	177	11.9	10.1-13.9	96	16.6	13.0-20.8	81	9.1	7.1-11.4
1981	239	15.5	13.4-17.7	127	20.8	17.0-25.2	112	12.1	9.8-14.7	201	12.9	11.1-15.0	108	17.4	13.8-21.5	93	10.2	8.1-12.5
1982	236	14.9	13.0-17.1	103	17.1	13.6-21.2	133	13.7	11.4-16.3	226	14.6	12.6-16.7	100	17.4	13.8-21.6	126	13.1	10.8-15.7
1983	269	15.7	13.8-17.9	130	19.0	15.5-22.9	139	13.6	11.4-16.2	243	14.5	12.7-16.6	126	19.6	15.9-23.7	117	11.5	9.4-13.9
1984	261	15.6	13.7-17.7	114	18.1	14.5-22.1	147	14.3	12.0-16.9	233	13.9	12.1-15.9	117	18.6	15.0-22.7	116	11.2	9.2-13.5
1985	242	13.9	12.1-15.8	125	17.7	14.5-21.5	117	11.3	9.3-13.6	206	12.2	10.5-14.1	103	15.9	12.7-19.7	103	10.1	8.1-12.2
1986	259	14.3	12.5-16.2	115	15.5	12.6-18.9	144	13.2	11.1-15.6	224	12.7	11.0-14.6	104	14.3	11.4-17.6	120	11.5	9.5-13.7
1987	281	15.1	13.3-17.1	125	16.6	13.5-20.1	156	13.9	11.8-16.3	242	13.1	11.4-14.9	106	15.3	12.1-18.9	136	11.9	9.9-14.1
1988	274	14.9	13.2-16.9	131	18.8	15.5-22.7	143	12.6	10.6-14.9	216	11.8	10.2-13.6	110	15.0	12.1-18.4	106	9.6	7.8-11.7
1989	255	13.4	11.7-15.2	104	14.4	11.5-17.7	151	12.7	10.7-15.0	234	12.4	10.8-14.1	99	13.6	10.8-16.8	135	11.5	9.6-13.6
1990	297	15.9	14.0-17.9	121	17.7	14.3-21.4	176	15.0	12.8-17.4	243	12.8	11.2-14.6	90	13.0	10.1-16.3	153	12.8	10.8-15.0
1991	256	13.0	11.4-14.8	124	16.4	13.4-19.9	132	11.0	9.2-13.1	221	11.1	9.6-12.7	111	14.2	11.4-17.4	110	9.1	7.5-11.0
1992	261	13.1	11.5-14.8	104	14.2	11.3-17.5	157	12.5	10.6-14.7	205	10.8	9.3-12.4	81	12.3	9.5-15.6	124	10.2	8.5-12.2
1993	244	12.0	10.5-13.6	106	13.7	10.9-16.8	138	10.9	9.1-12.9	204	10.1	8.7-11.6	84	11.8	9.1-14.9	120	9.3	7.7-11.2
1994	296	14.2	12.6-16.0	124	15.6	12.8-18.9	172	13.3	11.4-15.5	257	12.6	11.0-14.2	104	13.6	10.9-16.6	153	11.9	10.1-14.0
1995	264	12.5	11.0-14.1	118	14.3	11.7-17.4	146	11.1	9.3-13.1	240	11.2	9.8-12.8	108	13.7	11.0-16.8	132	9.9	8.2-11.7
1996	305	13.7	12.2-15.4	122	14.3	11.7-17.2	183	13.3	11.4-15.4	242	11.1	9.7-12.6	95	11.7	9.3-14.5	147	10.7	9.1-12.6
1997	303	13.5	12.0-15.2	128	14.1	11.7-17.0	175	12.7	10.9-14.8	240	11.0	9.6-12.5	103	11.8	9.5-14.4	137	10.1	8.5-12.0
1998	292	12.8	11.3-14.3	111	12.3	10.0-15.1	181	12.8	11.0-14.8	257	11.5	10.1-13.0	102	11.6	9.3-14.3	155	11.2	9.5-13.1
1999	337	14.3	12.8-16.0	136	15.1	12.5-18.0	201	13.8	11.9-15.8	286	12.2	10.8-13.7	108	11.7	9.5-14.3	178	12.2	10.5-14.1
2000	320	13.5	12.0-15.0	110	12.0	9.7-14.5	210	14.0	12.2-16.1	256	10.7	9.4-12.1	109	12.2	9.9-14.8	147	9.8	8.2-11.5
2001	361	14.4	12.9-16.0	141	14.4	12.0-17.1	220	14.3	12.4-16.3	284	11.7	10.3-13.1	105	11.1	9.0-13.6	179	11.7	10.1-13.6
2002	345	13.6	12.2-15.1	144	14.0	11.7-16.7	201	12.9	11.2-14.9	288	11.4	10.1-12.8	134	13.1	10.9-15.7	154	9.9	8.4-11.6
2003	322	12.4	11.1-13.9	160	16.2	13.7-19.1	162	10.1	8.6-11.8	258	10.2	9.0-11.5	114	11.9	9.6-14.4	144	9.1	7.7-10.8
2004	341	12.8	11.5-14.3	144	13.6	11.3-16.2	197	12.1	10.4-13.9	294	11.1	9.8-12.5	133	13.3	11.0-15.9	161	9.8	8.3-11.4
2005	347	12.6	11.3-14.0	148	13.9	11.7-16.5	199	11.7	10.1-13.5	269	10.0	8.8-11.2	120	11.2	9.1-13.5	149	9.0	7.6-10.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	405	14.4	13.0-15.9	171	15.0	12.7-17.6	234	13.7	12.0-15.6	277	9.9	8.8-11.2	115	10.7	8.7-12.9	162	9.4	8.0-11.0
2007	389	13.8	12.5-15.3	173	15.5	13.2-18.2	216	12.6	11.0-14.4	300	10.8	9.6-12.1	136	12.5	10.3-14.9	164	9.7	8.2-11.3
2008	380	13.2	11.9-14.6	150	12.8	10.8-15.2	230	13.0	11.4-14.9	308	10.8	9.6-12.1	123	11.2	9.2-13.5	185	10.6	9.1-12.2
2009	391	13.2	11.9-14.6	189	16.1	13.7-18.7	202	11.2	9.7-12.9	321	11.0	9.8-12.3	141	12.4	10.3-14.7	180	10.0	8.6-11.6
2010	421	13.8	12.5-15.2	192	15.7	13.4-18.2	229	12.5	10.9-14.2	297	10.1	9.0-11.3	131	11.0	9.1-13.1	166	9.4	8.0-10.9
2011	502	15.9	14.5-17.4	220	16.9	14.6-19.5	282	15.0	13.2-16.8	361	11.8	10.6-13.1	160	13.5	11.4-15.8	201	10.7	9.3-12.3
2012	481	14.6	13.3-16.0	196	14.9	12.8-17.3	285	14.5	12.8-16.3	372	11.3	10.2-12.6	141	10.4	8.7-12.4	231	11.9	10.4-13.6
2013	506	15.1	13.8-16.5	230	16.9	14.7-19.4	276	13.7	12.2-15.5	343	10.4	9.3-11.6	161	12.6	10.6-14.8	182	9.0	7.7-10.5
2014	502	14.4	13.2-15.8	225	15.4	13.4-17.7	277	13.5	12.0-15.2	349	10.3	9.2-11.4	151	10.8	9.1-12.8	198	9.8	8.4-11.2
2015	555	15.7	14.4-17.1	251	17.2	15.0-19.5	304	14.6	12.9-16.3	381	11.1	10.0-12.3	173	12.2	10.4-14.3	208	10.1	8.8-11.6
2016	541	14.9	13.7-16.2	251	16.8	14.7-19.1	290	13.4	11.9-15.1	401	11.1	10.0-12.3	197	13.4	11.5-15.5	204	9.4	8.2-10.9
2017	553	15.0	13.8-16.4	240	15.9	13.9-18.1	313	14.3	12.8-16.1	391	10.7	9.6-11.8	177	12.1	10.3-14.1	214	9.7	8.4-11.1
2018	576	15.2	13.9-16.5	255	16.4	14.4-18.6	321	14.2	12.6-15.8	392	10.4	9.4-11.5	163	10.7	9.1-12.6	229	10.0	8.7-11.4
2019	587	15.2	13.9-16.5	258	16.4	14.4-18.6	329	14.2	12.7-15.9	431	11.1	10.1-12.2	188	11.7	10.0-13.5	243	10.4	9.1-11.8
2020	532	13.2	12.1-14.4	247	14.9	13.0-16.9	285	12.0	10.6-13.5	393	9.9	8.9-11.0	189	11.8	10.1-13.7	204	8.7	7.5-10.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Larynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	115	6.7	5.5-8.2	95	13.3	10.6-16.4	20	2.1	1.2-3.3	66	4.4	3.3-5.7	52	8.8	6.2-11.9	14	1.5	0.8-2.7
1977	112	6.9	5.6-8.4	92	14.3	11.1-18.0	20	2.1	1.3-3.2	54	3.3	2.4-4.4	41	6.8	4.4-9.8	13	1.2	0.7-2.1
1978	124	7.0	5.8-8.4	101	14.2	11.3-17.5	23	2.2	1.4-3.3	83	4.7	3.7-5.8	71	9.9	7.5-12.8	12	1.1	0.6-1.9
1979	147	8.2	6.9-9.7	114	16.0	13.0-19.4	33	3.1	2.1-4.4	68	3.8	2.9-4.8	57	8.0	6.0-10.6	11	1.0	0.5-1.7
1980	138	7.3	6.1-8.7	108	14.2	11.5-17.4	30	2.7	1.8-3.9	66	3.6	2.8-4.6	45	6.4	4.6-8.8	21	1.8	1.1-2.8
1981	132	6.9	5.7-8.2	107	13.5	10.9-16.5	25	2.3	1.5-3.5	74	3.9	3.0-4.9	51	7.0	5.1-9.4	23	1.9	1.2-2.9
1982	122	6.2	5.1-7.4	108	13.4	10.9-16.4	14	1.2	0.7-2.1	58	3.1	2.3-4.1	47	6.5	4.6-8.8	11	1.0	0.5-1.8
1983	138	7.0	5.9-8.3	116	14.4	11.7-17.5	22	2.0	1.3-3.1	64	3.3	2.5-4.2	53	6.7	4.9-8.9	11	1.0	0.5-1.8
1984	146	7.3	6.1-8.6	107	13.1	10.6-16.0	39	3.3	2.3-4.5	73	3.7	2.8-4.6	59	7.3	5.4-9.5	14	1.2	0.7-2.1
1985	188	9.0	7.8-10.4	147	17.9	15.0-21.1	41	3.2	2.3-4.4	67	3.4	2.6-4.4	55	7.1	5.3-9.4	12	1.1	0.5-1.9
1986	179	8.8	7.5-10.2	141	17.8	14.8-21.3	38	3.0	2.1-4.2	81	4.0	3.2-5.1	63	8.1	6.1-10.5	18	1.5	0.9-2.4
1987	152	7.2	6.1-8.4	122	14.4	11.8-17.3	30	2.4	1.6-3.5	78	3.8	3.0-4.8	57	7.2	5.3-9.5	21	1.7	1.1-2.6
1988	153	6.8	5.7-8.0	113	12.6	10.2-15.2	40	3.0	2.1-4.1	88	4.1	3.3-5.1	75	9.1	7.0-11.6	13	0.9	0.5-1.6
1989	180	8.0	6.9-9.3	116	13.2	10.7-16.0	64	4.8	3.7-6.1	71	3.3	2.5-4.1	53	6.5	4.7-8.7	18	1.3	0.8-2.1
1990	180	7.9	6.8-9.2	138	15.5	12.8-18.5	42	3.2	2.3-4.3	62	2.9	2.2-3.7	49	6.0	4.3-8.1	13	1.0	0.5-1.7
1991	151	6.6	5.6-7.8	116	13.2	10.7-16.0	35	2.5	1.8-3.5	72	3.2	2.5-4.1	57	6.5	4.8-8.5	15	1.1	0.6-1.8
1992	191	8.0	6.9-9.2	150	15.9	13.3-18.7	41	2.8	2.0-3.9	82	3.6	2.8-4.4	63	7.2	5.5-9.4	19	1.3	0.8-2.1
1993	154	6.7	5.6-7.8	123	13.9	11.4-16.7	31	2.2	1.5-3.2	83	3.6	2.9-4.5	62	7.1	5.3-9.3	21	1.5	0.9-2.3
1994	149	6.2	5.2-7.3	115	12.2	10.0-14.8	34	2.4	1.6-3.3	78	3.4	2.6-4.2	56	6.2	4.5-8.1	22	1.6	1.0-2.4
1995	157	6.4	5.5-7.5	115	11.9	9.7-14.4	42	2.9	2.1-3.9	90	3.9	3.1-4.8	70	8.2	6.2-10.5	20	1.4	0.8-2.1
1996	153	6.1	5.2-7.2	126	12.9	10.6-15.5	27	1.8	1.2-2.6	80	3.3	2.6-4.2	69	7.8	5.9-10.0	11	0.7	0.4-1.3
1997	142	5.6	4.7-6.6	107	10.6	8.6-12.9	35	2.3	1.6-3.2	68	2.7	2.1-3.4	54	5.4	4.0-7.1	14	0.9	0.5-1.6
1998	161	6.2	5.3-7.3	111	11.2	9.1-13.6	50	3.2	2.4-4.2	67	2.7	2.1-3.4	57	5.7	4.3-7.4	10	0.7	0.3-1.2
1999	138	5.2	4.4-6.2	104	10.1	8.2-12.4	34	2.1	1.5-3.0	65	2.5	1.9-3.2	50	5.0	3.6-6.7	15	1.0	0.5-1.6
2000	129	4.8	4.0-5.7	92	8.8	7.0-10.9	37	2.3	1.6-3.2	67	2.7	2.0-3.4	49	5.0	3.7-6.7	18	1.2	0.7-1.9
2001	125	4.4	3.7-5.3	96	8.2	6.6-10.1	29	1.8	1.2-2.6	45	1.7	1.2-2.3	36	3.6	2.5-5.1	9	0.6	0.3-1.0
2002	125	4.3	3.6-5.2	90	7.6	6.1-9.4	35	2.1	1.4-2.9	70	2.5	1.9-3.2	52	4.5	3.3-6.0	18	1.1	0.7-1.8
2003	146	5.1	4.3-6.0	109	9.7	7.9-11.8	37	2.2	1.5-3.0	57	2.0	1.5-2.6	45	4.1	3.0-5.6	12	0.7	0.4-1.2
2004	132	4.3	3.6-5.1	111	9.0	7.3-10.9	21	1.2	0.7-1.8	61	2.2	1.6-2.8	50	4.5	3.3-6.1	11	0.7	0.3-1.2
2005	124	4.3	3.6-5.1	96	8.6	6.9-10.6	28	1.6	1.0-2.3	51	1.7	1.3-2.3	44	3.8	2.7-5.1	7	0.4	0.2-0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Larynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	125	4.0	3.3-4.7	96	7.6	6.1-9.3	29	1.5	1.0-2.2	50	1.8	1.3-2.3	42	3.8	2.7-5.2	8	0.5	0.2-0.9
2007	140	4.6	3.8-5.4	107	8.8	7.1-10.7	33	1.8	1.2-2.5	54	1.8	1.3-2.3	42	3.6	2.5-4.9	12	0.7	0.3-1.2
2008	153	5.0	4.2-5.9	106	8.9	7.2-10.8	47	2.6	1.9-3.5	61	2.0	1.5-2.6	41	3.5	2.5-4.8	20	1.1	0.6-1.6
2009	121	3.8	3.2-4.6	92	7.5	6.0-9.3	29	1.5	1.0-2.2	57	1.8	1.4-2.4	42	3.6	2.5-4.9	15	0.8	0.4-1.3
2010	128	3.9	3.2-4.7	96	7.4	6.0-9.2	32	1.6	1.1-2.3	47	1.6	1.1-2.1	39	3.4	2.4-4.7	8	0.4	0.2-0.8
2011	130	3.7	3.1-4.4	98	7.0	5.6-8.5	32	1.5	1.0-2.1	43	1.3	0.9-1.8	30	2.4	1.6-3.4	13	0.7	0.4-1.2
2012	103	3.0	2.4-3.6	74	5.4	4.2-6.8	29	1.4	0.9-2.0	52	1.6	1.2-2.1	43	3.5	2.5-4.7	9	0.4	0.2-0.8
2013	117	3.3	2.7-3.9	90	6.5	5.2-8.1	27	1.2	0.8-1.8	47	1.4	1.0-1.8	36	2.8	1.9-3.9	11	0.5	0.3-0.9
2014	134	3.7	3.1-4.4	104	7.1	5.7-8.7	30	1.5	1.0-2.1	54	1.5	1.1-2.0	46	3.4	2.4-4.5	8	0.4	0.2-0.7
2015	152	4.0	3.4-4.8	124	8.4	6.9-10.1	28	1.2	0.8-1.8	35	1.0	0.7-1.4	26	2.0	1.3-3.0	9	0.4	0.2-0.8
2016	114	3.0	2.5-3.7	86	5.8	4.6-7.3	28	1.2	0.8-1.8	49	1.3	1.0-1.8	37	2.6	1.8-3.7	12	0.5	0.3-0.9
2017	108	2.7	2.2-3.3	72	4.5	3.5-5.7	36	1.6	1.1-2.2	35	0.9	0.6-1.3	25	1.7	1.1-2.6	10	0.4	0.2-0.8
2018	107	2.6	2.1-3.2	83	5.1	4.0-6.3	24	0.9	0.6-1.4	36	0.9	0.6-1.3	27	1.8	1.1-2.6	9	0.4	0.2-0.8
2019	111	2.7	2.2-3.3	87	5.3	4.2-6.6	24	1.0	0.7-1.5	30	0.7	0.5-1.0	23	1.3	0.8-2.1	7	0.3	0.1-0.6
2020	97	2.4	1.9-2.9	80	4.8	3.8-6.0	17	0.7	0.4-1.2	21	0.5	0.3-0.8	17	1.1	0.6-1.8	4	0.2	0.0-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	890	57.1	53.1-61.2	667	104.7	96.1-113.9	223	24.9	21.5-28.6	680	43.7	40.2-47.4	517	82.0	74.2-90.3	163	18.0	15.2-21.2
1977	960	60.3	56.3-64.5	708	113.8	104.5-123.6	252	25.7	22.5-29.2	759	47.7	44.1-51.5	573	92.0	83.7-100.8	186	19.0	16.3-22.1
1978	988	61.1	57.1-65.3	730	110.7	102.1-119.9	258	28.2	24.7-32.0	827	51.1	47.5-55.0	613	94.0	86.0-102.5	214	22.9	19.8-26.4
1979	1,065	65.4	61.3-69.7	767	119.4	110.1-129.2	298	30.9	27.3-34.8	855	53.5	49.7-57.4	639	101.4	92.7-110.6	216	23.0	19.9-26.5
1980	1,125	65.4	61.4-69.6	827	122.8	113.7-132.4	298	28.8	25.5-32.4	904	53.6	49.9-57.4	667	100.8	92.5-109.7	237	23.7	20.6-27.0
1981	1,160	65.1	61.3-69.2	824	118.4	109.7-127.6	336	31.1	27.8-34.8	957	54.7	51.1-58.4	685	100.4	92.3-109.1	272	25.7	22.7-29.1
1982	1,177	64.8	60.9-68.7	827	116.3	107.8-125.2	350	32.0	28.6-35.6	945	52.5	49.0-56.1	667	93.3	85.8-101.3	278	26.3	23.2-29.7
1983	1,193	64.2	60.4-68.1	831	114.0	105.8-122.7	362	32.6	29.2-36.3	1,048	57.3	53.7-61.0	753	106.6	98.5-115.2	295	26.5	23.5-29.8
1984	1,243	65.1	61.4-69.0	844	112.4	104.3-120.8	399	35.2	31.7-39.0	978	51.6	48.3-55.1	689	94.5	86.9-102.5	289	25.2	22.3-28.3
1985	1,264	64.9	61.3-68.8	873	116.2	108.0-124.9	391	33.3	30.0-36.8	1,020	53.3	50.0-56.9	714	97.2	89.6-105.3	306	26.5	23.5-29.7
1986	1,305	66.0	62.3-69.8	857	114.0	105.8-122.7	448	37.1	33.6-40.8	1,021	52.0	48.7-55.4	667	89.4	82.1-97.1	354	29.5	26.4-32.8
1987	1,329	65.0	61.4-68.7	899	114.0	106.1-122.2	430	34.8	31.5-38.3	1,097	55.0	51.6-58.4	732	97.1	89.6-105.1	365	30.0	26.9-33.3
1988	1,327	64.7	61.1-68.4	868	112.8	104.8-121.2	459	36.4	33.1-40.0	1,052	52.1	48.8-55.4	698	91.9	84.6-99.6	354	28.6	25.6-31.8
1989	1,403	66.4	62.9-70.1	925	116.5	108.6-124.9	478	36.9	33.6-40.5	1,126	54.5	51.3-57.9	761	99.7	92.1-107.7	365	28.5	25.6-31.6
1990	1,396	65.6	62.0-69.2	917	113.8	105.9-122.1	479	36.8	33.5-40.3	1,111	52.6	49.4-55.8	735	90.8	83.8-98.2	376	29.5	26.5-32.6
1991	1,396	65.1	61.7-68.8	892	110.6	102.8-118.8	504	38.4	35.1-42.0	1,081	50.9	47.9-54.2	707	89.2	82.2-96.6	374	28.6	25.7-31.7
1992	1,437	65.4	61.9-68.9	932	113.2	105.4-121.4	505	37.5	34.2-40.9	1,063	49.9	46.8-53.1	665	84.7	77.8-92.0	398	30.1	27.2-33.3
1993	1,377	62.0	58.7-65.5	854	103.9	96.4-111.8	523	38.3	35.0-41.7	1,085	49.2	46.3-52.4	696	86.8	79.9-94.1	389	28.4	25.6-31.4
1994	1,472	64.2	60.9-67.7	891	103.5	96.3-111.1	581	41.2	37.9-44.8	1,179	52.5	49.5-55.7	769	91.3	84.4-98.6	410	30.0	27.1-33.0
1995	1,412	61.3	58.1-64.7	867	98.1	91.3-105.3	545	39.0	35.8-42.5	1,163	51.0	48.0-54.1	704	82.2	75.8-88.9	459	32.7	29.8-35.9
1996	1,537	65.2	61.9-68.6	908	100.3	93.5-107.6	629	44.0	40.6-47.6	1,187	51.1	48.1-54.1	724	81.1	74.8-87.7	463	33.0	30.0-36.2
1997	1,492	62.6	59.4-65.9	872	96.0	89.3-103.0	620	42.6	39.3-46.1	1,154	49.4	46.5-52.3	696	79.6	73.4-86.2	458	31.8	28.9-34.9
1998	1,501	61.0	57.9-64.2	846	88.5	82.3-95.0	655	44.0	40.7-47.6	1,158	48.3	45.5-51.2	691	75.5	69.7-81.8	467	32.0	29.1-35.0
1999	1,417	56.7	53.8-59.8	811	84.6	78.5-91.0	606	39.9	36.8-43.3	1,082	44.3	41.6-47.1	620	66.7	61.2-72.5	462	31.1	28.3-34.1
2000	1,479	57.3	54.4-60.4	860	86.7	80.7-93.0	619	39.6	36.5-42.8	1,109	43.5	41.0-46.2	654	67.8	62.4-73.5	455	29.2	26.5-32.0
2001	1,472	56.6	53.7-59.7	842	85.7	79.7-92.0	630	39.6	36.6-42.8	1,095	42.7	40.2-45.4	625	65.2	59.9-70.9	470	29.8	27.1-32.6
2002	1,491	55.9	53.0-58.8	827	82.3	76.5-88.5	664	40.7	37.6-43.9	1,123	42.5	40.0-45.1	645	64.8	59.6-70.3	478	29.4	26.8-32.2
2003	1,459	53.5	50.7-56.3	770	73.4	68.0-79.1	689	41.6	38.5-44.8	1,098	40.9	38.4-43.4	613	60.0	55.0-65.2	485	29.5	26.9-32.2
2004	1,556	55.7	52.9-58.6	852	77.3	71.9-83.0	704	41.9	38.8-45.1	1,068	39.1	36.7-41.5	599	57.4	52.7-62.5	469	28.0	25.5-30.7
2005	1,545	54.7	52.0-57.6	857	79.5	74.0-85.4	688	40.2	37.2-43.3	1,112	39.9	37.5-42.4	602	55.7	51.1-60.6	510	30.3	27.7-33.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,518	53.1	50.4-55.8	818	73.1	67.9-78.5	700	40.6	37.6-43.7	1,036	37.0	34.7-39.3	546	50.7	46.3-55.3	490	28.7	26.2-31.4
2007	1,563	53.3	50.6-56.1	835	73.4	68.2-78.8	728	41.1	38.2-44.3	1,081	37.4	35.2-39.8	591	52.7	48.3-57.4	490	28.1	25.7-30.8
2008	1,559	52.7	50.1-55.5	811	70.8	65.8-76.2	748	41.9	38.9-45.1	1,005	34.6	32.5-36.9	526	46.8	42.7-51.2	479	27.1	24.7-29.7
2009	1,632	53.6	51.0-56.4	861	72.4	67.4-77.7	771	42.1	39.1-45.2	1,027	34.4	32.2-36.6	568	49.0	44.8-53.4	459	25.3	23.0-27.8
2010	1,649	52.6	50.0-55.3	841	68.3	63.5-73.3	808	43.1	40.1-46.2	1,120	36.5	34.4-38.8	610	52.6	48.2-57.2	510	27.4	25.0-29.9
2011	1,705	53.1	50.6-55.8	868	69.5	64.7-74.6	837	43.2	40.3-46.2	1,129	35.7	33.6-37.9	599	48.5	44.5-52.8	530	27.7	25.3-30.2
2012	1,707	51.7	49.2-54.3	876	68.0	63.4-72.9	831	41.9	39.1-44.9	1,109	34.3	32.3-36.4	599	48.3	44.3-52.5	510	26.0	23.8-28.4
2013	1,599	47.1	44.8-49.5	819	61.5	57.1-66.1	780	38.2	35.6-41.0	1,081	32.6	30.6-34.6	576	44.7	40.9-48.7	505	25.0	22.9-27.3
2014	1,672	48.0	45.6-50.4	862	62.5	58.2-67.0	810	38.7	36.1-41.5	1,019	29.7	27.9-31.7	543	40.5	37.0-44.2	476	22.9	20.8-25.1
2015	1,731	47.8	45.6-50.2	887	61.9	57.7-66.3	844	39.0	36.4-41.8	1,017	28.9	27.1-30.7	514	37.5	34.2-41.1	503	23.6	21.6-25.8
2016	1,715	46.8	44.6-49.1	858	58.2	54.2-62.4	857	39.2	36.6-42.0	964	26.4	24.7-28.2	530	36.9	33.7-40.3	434	19.7	17.8-21.6
2017	1,717	45.5	43.3-47.7	867	57.6	53.7-61.8	850	37.8	35.2-40.4	885	23.9	22.3-25.5	457	31.7	28.7-34.9	428	19.1	17.3-21.1
2018	1,735	45.1	42.9-47.3	869	57.0	53.2-61.1	866	37.6	35.1-40.2	934	24.6	23.0-26.2	495	33.3	30.3-36.4	439	19.2	17.4-21.1
2019	1,690	42.9	40.8-45.0	827	52.3	48.7-56.1	863	36.9	34.4-39.5	858	21.8	20.3-23.3	453	29.7	27.0-32.7	405	17.0	15.3-18.7
2020	1,447	35.8	33.9-37.7	743	45.5	42.2-49.0	704	29.4	27.2-31.7	746	18.7	17.3-20.1	390	24.7	22.3-27.4	356	14.8	13.3-16.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	10	0.7	0.3-1.4	2	0.3	0.0-1.3	8	1.0	0.4-2.0	9	0.8	0.3-1.6	4	1.1	0.2-3.0	5	0.7	0.2-1.7
1977	14	1.2	0.6-2.1	8	1.9	0.7-4.0	6	0.9	0.3-1.9	7	0.6	0.2-1.3	4	1.3	0.3-3.3	3	0.3	0.1-0.8
1978	13	0.9	0.4-1.6	5	1.2	0.3-2.9	8	0.9	0.3-1.7	6	0.4	0.1-0.9	2	0.2	0.0-1.0	4	0.5	0.1-1.3
1979	18	1.0	0.6-1.6	7	0.9	0.3-1.9	11	1.1	0.5-2.0	4	0.4	0.1-0.9	2	0.4	0.0-1.4	2	0.3	0.0-1.1
1980	14	0.9	0.5-1.6	5	0.6	0.2-1.5	9	1.1	0.5-2.1	14	0.9	0.4-1.5	8	0.9	0.4-1.9	6	0.7	0.3-1.6
1981	25	1.4	0.9-2.2	7	0.8	0.3-1.8	18	1.8	1.0-2.9	15	0.9	0.5-1.6	7	1.0	0.4-2.1	8	0.9	0.3-1.8
1982	19	1.0	0.6-1.5	8	0.9	0.4-1.8	11	1.0	0.5-1.8	10	0.5	0.2-1.0	6	1.1	0.3-2.5	4	0.3	0.1-0.8
1983	12	0.7	0.3-1.2	4	0.5	0.1-1.5	8	0.7	0.3-1.4	20	1.1	0.6-1.7	14	1.7	0.9-2.8	6	0.6	0.2-1.4
1984	26	1.5	0.9-2.2	7	1.0	0.3-2.2	19	1.8	1.0-2.8	13	0.6	0.3-1.0	10	1.1	0.5-2.1	3	0.2	0.0-0.7
1985	23	1.3	0.8-2.0	12	1.4	0.6-2.6	11	1.2	0.6-2.1	14	0.8	0.4-1.4	10	1.7	0.8-3.2	4	0.3	0.1-0.9
1986	20	1.0	0.6-1.6	11	1.7	0.7-3.2	9	0.8	0.3-1.5	21	1.1	0.7-1.7	15	1.8	1.0-3.1	6	0.6	0.2-1.2
1987	17	0.8	0.5-1.3	8	0.9	0.4-1.9	9	0.7	0.3-1.4	9	0.5	0.2-0.9	4	0.5	0.1-1.2	5	0.4	0.1-1.0
1988	20	1.0	0.6-1.5	7	1.1	0.4-2.3	13	1.0	0.5-1.7	9	0.5	0.2-1.0	4	0.7	0.1-1.8	5	0.5	0.1-1.0
1989	17	1.0	0.6-1.6	6	0.8	0.2-1.7	11	1.1	0.5-1.9	17	0.9	0.5-1.4	6	0.6	0.2-1.5	11	0.9	0.5-1.7
1990	21	0.9	0.6-1.5	10	1.2	0.5-2.3	11	0.8	0.4-1.5	6	0.3	0.1-0.7	3	0.5	0.1-1.5	3	0.3	0.1-0.7
1991	27	1.2	0.7-1.7	13	2.0	0.9-3.6	14	0.9	0.5-1.5	11	0.6	0.3-1.0	6	0.8	0.2-1.8	5	0.4	0.1-1.0
1992	13	0.5	0.3-0.9	6	0.5	0.2-1.2	7	0.5	0.2-1.0	7	0.3	0.1-0.6	3	0.3	0.1-0.8	4	0.3	0.1-0.7
1993	25	1.2	0.8-1.8	12	1.5	0.7-2.7	13	1.0	0.5-1.7	8	0.3	0.1-0.6	2	0.2	0.0-0.7	6	0.4	0.1-0.9
1994	18	0.8	0.4-1.2	5	0.6	0.1-1.6	13	0.9	0.5-1.6	6	0.3	0.1-0.7	1	0.2	0.0-0.8	5	0.4	0.1-0.9
1995	25	1.1	0.7-1.6	7	0.6	0.2-1.3	18	1.3	0.8-2.1	6	0.3	0.1-0.6	1	0.1	0.0-0.5	5	0.4	0.1-0.8
1996	23	1.1	0.7-1.6	8	1.0	0.4-2.0	15	1.1	0.6-1.8	13	0.6	0.3-1.1	4	0.5	0.1-1.3	9	0.7	0.3-1.3
1997	38	1.6	1.1-2.2	15	1.6	0.8-2.7	23	1.6	1.0-2.4	9	0.4	0.2-0.8	4	0.3	0.1-0.8	5	0.4	0.1-0.9
1998	28	1.2	0.8-1.7	13	1.5	0.7-2.6	15	1.0	0.6-1.7	11	0.4	0.2-0.8	3	0.2	0.0-0.7	8	0.6	0.2-1.1
1999	13	0.5	0.3-0.9	6	0.6	0.2-1.3	7	0.5	0.2-0.9	8	0.3	0.1-0.6	5	0.6	0.2-1.4	3	0.2	0.0-0.6
2000	13	0.5	0.3-0.9	7	0.7	0.3-1.4	6	0.4	0.1-0.8	13	0.6	0.3-1.0	7	1.1	0.4-2.2	6	0.4	0.2-0.9
2001	29	1.0	0.7-1.5	11	0.9	0.4-1.6	18	1.1	0.7-1.8	6	0.2	0.1-0.5	3	0.3	0.1-0.8	3	0.2	0.0-0.6
2002	26	0.9	0.6-1.4	9	0.8	0.3-1.5	17	1.0	0.6-1.7	7	0.3	0.1-0.6	3	0.4	0.0-1.1	4	0.3	0.1-0.7
2003	20	0.7	0.4-1.1	5	0.6	0.2-1.4	15	0.9	0.5-1.4	10	0.3	0.2-0.6	3	0.3	0.1-0.9	7	0.4	0.2-0.8
2004	25	0.9	0.6-1.3	13	1.3	0.7-2.2	12	0.7	0.3-1.2	4	0.1	0.0-0.4	2	0.1	0.0-0.5	2	0.1	0.0-0.4
2005	28	0.9	0.6-1.4	12	0.9	0.4-1.5	16	0.9	0.5-1.5	6	0.2	0.1-0.4	4	0.3	0.1-0.7	2	0.1	0.0-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	39	1.4	1.0-1.9	20	1.9	1.1-3.0	19	1.1	0.7-1.7	11	0.4	0.2-0.7	6	0.5	0.2-1.1	5	0.3	0.1-0.7
2007	42	1.4	1.0-1.9	12	1.1	0.5-2.0	30	1.7	1.2-2.5	16	0.6	0.3-0.9	6	0.7	0.2-1.5	10	0.6	0.3-1.1
2008	38	1.2	0.9-1.7	15	1.2	0.7-2.1	23	1.3	0.8-1.9	10	0.3	0.2-0.6	3	0.2	0.0-0.6	7	0.4	0.2-0.8
2009	29	0.9	0.6-1.2	11	0.7	0.4-1.3	18	0.9	0.6-1.5	8	0.3	0.1-0.5	5	0.4	0.1-0.9	3	0.2	0.0-0.5
2010	30	0.9	0.6-1.3	11	0.8	0.4-1.5	19	0.9	0.6-1.5	11	0.4	0.2-0.6	3	0.3	0.1-0.9	8	0.4	0.2-0.8
2011	40	1.2	0.9-1.7	19	1.5	0.8-2.3	21	1.1	0.7-1.7	15	0.5	0.3-0.9	6	0.5	0.2-1.1	9	0.5	0.2-1.0
2012	31	0.9	0.6-1.3	14	1.1	0.6-1.8	17	0.8	0.5-1.3	12	0.4	0.2-0.7	3	0.3	0.1-0.8	9	0.4	0.2-0.8
2013	42	1.3	0.9-1.8	14	1.2	0.6-2.0	28	1.4	0.9-2.0	15	0.4	0.2-0.6	8	0.5	0.2-1.0	7	0.3	0.1-0.7
2014	35	1.0	0.7-1.4	15	0.9	0.5-1.5	20	1.0	0.6-1.5	7	0.2	0.1-0.4	2	0.1	0.0-0.4	5	0.3	0.1-0.6
2015	35	1.0	0.7-1.4	18	1.3	0.8-2.1	17	0.8	0.5-1.4	12	0.3	0.2-0.6	7	0.5	0.2-1.1	5	0.2	0.1-0.6
2016	32	0.9	0.6-1.2	14	0.9	0.5-1.5	18	0.8	0.5-1.3	5	0.1	0.0-0.3	2	0.1	0.0-0.5	3	0.1	0.0-0.4
2017	34	1.0	0.7-1.4	16	1.1	0.6-1.9	18	0.9	0.5-1.5	4	0.1	0.0-0.3	3	0.2	0.0-0.6	1	0.0	0.0-0.3
2018	39	1.1	0.7-1.5	16	1.1	0.6-1.7	23	1.1	0.7-1.6	7	0.2	0.1-0.4	3	0.2	0.0-0.6	4	0.2	0.0-0.5
2019	36	1.0	0.7-1.4	9	0.7	0.3-1.3	27	1.2	0.8-1.8	9	0.2	0.1-0.5	5	0.3	0.1-0.7	4	0.2	0.0-0.5
2020	37	1.0	0.7-1.4	14	0.8	0.5-1.4	23	1.0	0.6-1.6	8	0.2	0.1-0.4	3	0.2	0.0-0.5	5	0.2	0.1-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	688	75.3	69.4-81.4	315	36.5	32.3-41.1
1977	716	76.3	70.5-82.4	295	32.2	28.4-36.3
1978	793	82.4	76.5-88.6	313	33.2	29.4-37.3
1979	805	81.6	75.8-87.7	294	30.3	26.7-34.1
1980	870	84.0	78.3-90.0	348	35.0	31.2-39.1
1981	856	82.0	76.3-87.9	357	35.6	31.8-39.7
1982	907	84.4	78.8-90.2	367	34.3	30.7-38.1
1983	933	83.8	78.3-89.6	376	34.3	30.8-38.0
1984	1,046	91.0	85.4-96.8	413	38.2	34.5-42.2
1985	1,059	90.4	84.9-96.2	342	29.6	26.5-33.0
1986	1,057	86.5	81.2-92.0	392	32.6	29.4-36.1
1987	1,142	91.1	85.7-96.6	396	32.2	29.0-35.6
1988	1,234	96.0	90.6-101.7	427	33.5	30.3-36.9
1989	1,162	89.3	84.2-94.7	425	33.6	30.4-37.0
1990	1,241	93.5	88.3-99.0	430	33.3	30.1-36.6
1991	1,383	102.4	96.9-108.0	434	32.8	29.7-36.1
1992	1,339	97.7	92.4-103.1	444	32.8	29.8-36.0
1993	1,356	96.9	91.8-102.3	423	31.4	28.5-34.6
1994	1,440	100.4	95.2-105.7	496	35.0	32.0-38.3
1995	1,462	99.9	94.8-105.2	511	35.9	32.8-39.1
1996	1,606	107.4	102.2-112.8	489	33.5	30.5-36.6
1997	1,649	108.3	103.1-113.7	490	33.3	30.4-36.4
1998	1,806	116.2	110.8-121.7	491	32.4	29.6-35.4
1999	1,652	103.7	98.8-108.9	473	30.5	27.8-33.4
2000	1,627	99.4	94.6-104.4	463	29.2	26.6-32.0
2001	1,629	97.7	93.0-102.6	516	31.7	29.0-34.6
2002	1,732	101.7	96.9-106.6	467	28.2	25.7-30.9
2003	1,784	103.2	98.5-108.2	468	27.7	25.2-30.3
2004	1,840	104.7	99.9-109.6	476	27.8	25.3-30.4
2005	1,857	104.6	99.9-109.5	518	29.5	27.0-32.2
2006	1,973	109.1	104.3-114.1	473	26.8	24.4-29.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	2,017	110.6	105.8-115.5	469	25.9	23.6-28.3
2008	2,100	113.8	109.0-118.9	467	25.5	23.2-27.9
2009	2,131	113.8	108.9-118.8	483	26.1	23.8-28.6
2010	2,100	109.7	105.0-114.5	446	23.3	21.1-25.6
2011	2,406	122.6	117.7-127.7	503	26.1	23.9-28.6
2012	2,273	113.9	109.2-118.7	503	25.5	23.3-27.9
2013	2,438	120.4	115.6-125.3	454	22.4	20.4-24.6
2014	2,488	121.2	116.4-126.1	500	24.3	22.1-26.5
2015	2,545	121.1	116.4-126.0	457	21.6	19.6-23.7
2016	2,568	120.8	116.1-125.7	483	22.6	20.6-24.8
2017	2,655	123.4	118.7-128.3	450	20.6	18.7-22.7
2018	2,656	123.0	118.3-127.9	508	22.5	20.5-24.6
2019	2,947	135.5	130.5-140.6	530	23.7	21.6-25.8
2020	2,396	108.8	104.3-113.3	455	19.8	18.0-21.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	294	30.0	26.5-33.9	130	15.0	12.4-18.1
1977	305	30.8	27.3-34.6	129	14.3	11.8-17.1
1978	290	28.4	25.1-32.1	110	11.7	9.5-14.3
1979	311	29.7	26.3-33.3	105	10.8	8.7-13.2
1980	279	26.6	23.5-30.1	115	11.6	9.5-14.1
1981	236	20.7	18.0-23.6	108	10.1	8.2-12.3
1982	266	23.6	20.8-26.7	97	8.5	6.9-10.5
1983	251	21.4	18.8-24.3	98	8.5	6.9-10.4
1984	260	21.7	19.0-24.6	77	7.2	5.6-9.1
1985	233	19.1	16.7-21.8	94	7.8	6.3-9.7
1986	270	22.0	19.4-24.9	91	7.5	6.0-9.2
1987	247	18.9	16.6-21.5	74	5.9	4.6-7.4
1988	234	17.8	15.5-20.3	108	8.5	6.9-10.2
1989	231	16.7	14.6-19.0	69	5.4	4.2-6.9
1990	299	20.8	18.5-23.3	83	6.4	5.0-7.9
1991	269	18.7	16.5-21.1	82	5.9	4.6-7.3
1992	255	17.1	15.0-19.4	78	5.6	4.4-7.0
1993	253	16.9	14.8-19.2	101	6.9	5.6-8.4
1994	265	17.9	15.8-20.2	79	5.4	4.3-6.8
1995	271	17.7	15.6-19.9	93	6.4	5.1-7.8
1996	293	18.6	16.5-20.9	78	5.1	4.0-6.4
1997	247	15.5	13.6-17.6	91	6.0	4.8-7.4
1998	258	15.4	13.6-17.5	79	5.1	4.0-6.3
1999	248	15.0	13.2-17.0	88	5.4	4.3-6.7
2000	289	17.4	15.4-19.5	91	5.4	4.4-6.7
2001	245	14.4	12.6-16.3	70	4.2	3.3-5.3
2002	245	14.1	12.4-16.0	74	4.4	3.5-5.6
2003	234	13.3	11.6-15.1	85	4.9	3.9-6.0
2004	218	12.1	10.6-13.9	64	3.7	2.8-4.7
2005	234	13.1	11.4-14.9	71	4.1	3.2-5.1
2006	221	12.2	10.7-14.0	72	4.1	3.2-5.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	239	13.2	11.5-15.0	93	5.2	4.2-6.3
2008	218	11.8	10.3-13.5	77	4.1	3.3-5.2
2009	205	10.8	9.4-12.4	90	4.7	3.7-5.8
2010	212	11.3	9.8-13.0	60	3.1	2.4-4.0
2011	210	10.9	9.5-12.5	81	4.2	3.3-5.3
2012	202	10.5	9.0-12.0	67	3.4	2.6-4.3
2013	215	10.8	9.4-12.4	83	4.1	3.2-5.1
2014	198	9.8	8.5-11.3	77	3.7	2.9-4.7
2015	203	10.0	8.6-11.5	60	2.9	2.2-3.7
2016	202	9.8	8.5-11.3	58	2.9	2.2-3.7
2017	209	10.1	8.7-11.6	57	2.7	2.0-3.5
2018	184	8.8	7.5-10.2	65	2.9	2.2-3.7
2019	197	9.4	8.1-10.9	60	2.7	2.1-3.5
2020	158	7.4	6.2-8.7	59	2.7	2.0-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Corpus uteris and uterus, not otherwise specified**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	146	16.4	13.7-19.4	68	8.9	6.8-11.4
1977	143	16.0	13.4-19.0	66	7.4	5.6-9.5
1978	141	15.5	12.9-18.4	69	8.4	6.4-10.8
1979	145	14.5	12.1-17.2	66	7.3	5.6-9.4
1980	185	18.9	16.1-21.9	82	8.4	6.6-10.5
1981	142	13.8	11.5-16.4	72	6.9	5.4-8.8
1982	162	14.7	12.5-17.3	58	5.4	4.0-7.0
1983	164	14.8	12.5-17.3	74	7.1	5.5-8.9
1984	187	17.1	14.7-19.8	75	6.9	5.4-8.7
1985	212	18.2	15.7-20.9	87	8.2	6.5-10.1
1986	149	12.2	10.3-14.4	65	5.7	4.4-7.3
1987	211	17.3	15.0-19.8	55	4.9	3.7-6.4
1988	216	17.0	14.7-19.5	81	6.7	5.2-8.3
1989	182	13.9	11.9-16.1	76	5.9	4.6-7.4
1990	205	15.8	13.7-18.1	76	6.2	4.8-7.7
1991	250	18.8	16.5-21.4	77	6.0	4.7-7.5
1992	242	17.8	15.6-20.2	87	6.8	5.4-8.4
1993	265	19.0	16.7-21.4	85	6.6	5.2-8.1
1994	280	19.5	17.2-21.9	92	6.7	5.4-8.2
1995	259	17.9	15.8-20.3	77	5.6	4.4-7.0
1996	306	20.7	18.5-23.2	110	7.8	6.4-9.5
1997	322	21.7	19.4-24.2	109	7.5	6.2-9.1
1998	330	21.5	19.2-24.0	124	8.5	7.1-10.2
1999	353	22.7	20.4-25.3	114	7.6	6.2-9.1
2000	362	22.8	20.5-25.3	110	7.2	5.9-8.6
2001	384	23.3	21.1-25.8	124	7.9	6.6-9.4
2002	365	21.8	19.6-24.2	110	6.8	5.6-8.2
2003	396	23.1	20.9-25.5	140	8.4	7.0-9.9
2004	452	26.1	23.7-28.6	126	7.5	6.3-9.0
2005	454	25.9	23.6-28.4	144	8.4	7.1-10.0
2006	475	26.5	24.1-29.0	139	8.1	6.8-9.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Corpus uteris and uterus, not otherwise specified**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	467	25.7	23.4-28.2	136	7.9	6.6-9.3
2008	574	30.9	28.4-33.5	161	9.1	7.7-10.6
2009	566	29.9	27.4-32.5	150	8.2	6.9-9.6
2010	581	29.9	27.5-32.5	160	8.6	7.3-10.0
2011	604	30.3	27.9-32.8	181	9.5	8.2-11.0
2012	660	32.2	29.8-34.8	178	9.0	7.7-10.5
2013	645	30.6	28.2-33.1	170	8.2	7.0-9.6
2014	706	32.6	30.2-35.1	231	10.9	9.6-12.5
2015	726	33.0	30.6-35.5	206	9.4	8.2-10.8
2016	774	34.7	32.3-37.3	220	9.9	8.6-11.3
2017	739	32.4	30.1-34.9	211	9.3	8.1-10.7
2018	786	33.6	31.3-36.1	243	10.5	9.2-11.9
2019	819	34.7	32.3-37.2	255	10.9	9.6-12.3
2020	739	30.5	28.3-32.8	222	9.1	7.9-10.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	105	12.1	9.8-14.9	64	7.7	5.8-10.0
1977	120	12.1	9.9-14.6	73	7.9	6.1-10.0
1978	101	10.5	8.4-12.9	72	7.6	5.9-9.7
1979	129	13.2	10.9-15.9	69	6.9	5.3-8.8
1980	130	12.9	10.6-15.4	96	10.1	8.1-12.5
1981	119	10.9	9.0-13.1	62	5.7	4.3-7.4
1982	115	10.5	8.6-12.6	67	6.3	4.9-8.1
1983	151	13.1	11.0-15.4	86	8.0	6.3-9.9
1984	147	12.7	10.7-15.0	90	8.1	6.4-10.0
1985	148	11.8	10.0-14.0	90	7.7	6.2-9.6
1986	151	12.5	10.5-14.7	84	7.0	5.5-8.7
1987	120	9.9	8.2-11.9	82	6.8	5.4-8.5
1988	151	11.9	10.1-14.1	75	6.1	4.8-7.7
1989	133	9.9	8.3-11.8	85	6.6	5.3-8.2
1990	165	12.4	10.5-14.5	107	8.4	6.9-10.2
1991	139	10.5	8.8-12.5	92	7.0	5.6-8.7
1992	184	13.3	11.4-15.4	123	9.2	7.6-11.0
1993	157	10.8	9.1-12.6	114	8.6	7.0-10.3
1994	194	13.6	11.8-15.7	113	8.4	6.9-10.1
1995	180	12.2	10.4-14.1	122	8.8	7.3-10.6
1996	151	10.1	8.5-11.9	77	5.3	4.2-6.6
1997	198	13.0	11.2-14.9	101	7.0	5.7-8.5
1998	145	9.4	8.0-11.1	102	7.0	5.7-8.5
1999	175	11.3	9.6-13.1	100	6.6	5.4-8.1
2000	163	10.3	8.7-12.0	107	7.0	5.7-8.4
2001	170	10.3	8.8-12.0	108	6.8	5.6-8.2
2002	187	11.0	9.5-12.7	108	6.7	5.5-8.1
2003	159	9.2	7.8-10.8	110	6.6	5.4-7.9
2004	185	10.7	9.2-12.3	106	6.4	5.2-7.7
2005	176	10.0	8.6-11.6	114	6.7	5.5-8.0
2006	157	8.7	7.4-10.2	111	6.5	5.3-7.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	163	9.0	7.7-10.5	107	6.1	5.0-7.4
2008	183	10.0	8.6-11.5	103	5.6	4.6-6.8
2009	192	10.2	8.8-11.7	105	5.6	4.6-6.8
2010	181	9.5	8.2-11.0	114	6.1	5.0-7.3
2011	188	9.6	8.3-11.1	116	6.2	5.1-7.4
2012	219	10.9	9.5-12.5	131	6.6	5.5-7.8
2013	186	9.1	7.8-10.6	105	5.1	4.2-6.2
2014	218	10.5	9.1-12.0	121	5.8	4.8-6.9
2015	186	8.9	7.7-10.3	122	5.8	4.8-6.9
2016	200	9.0	7.8-10.4	125	5.7	4.7-6.8
2017	214	10.1	8.8-11.6	111	5.0	4.1-6.1
2018	195	8.9	7.7-10.3	133	5.8	4.8-6.9
2019	212	9.7	8.4-11.1	138	6.0	5.0-7.1
2020	192	8.5	7.3-9.8	113	4.8	3.9-5.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	519	120.0	108.3-132.5	209	55.0	46.6-64.2
1977	551	120.7	109.4-132.6	211	53.8	45.9-62.4
1978	607	132.2	120.4-144.7	226	54.7	46.9-63.3
1979	626	130.9	119.6-142.9	262	62.3	54.1-71.2
1980	763	153.8	141.8-166.5	274	65.1	56.8-74.1
1981	739	147.8	136.1-160.1	314	72.0	63.3-81.3
1982	702	135.1	124.2-146.5	276	60.7	53.1-69.0
1983	711	132.5	121.9-143.6	288	60.9	53.4-68.9
1984	775	138.1	127.6-149.3	286	58.5	51.3-66.3
1985	805	135.8	125.6-146.5	326	65.8	58.2-74.0
1986	845	142.4	132.1-153.3	346	68.3	60.7-76.6
1987	846	136.1	126.2-146.5	348	67.3	59.8-75.4
1988	914	149.3	138.8-160.3	310	60.1	53.1-67.7
1989	953	149.3	139.0-159.9	315	56.3	49.7-63.4
1990	1,102	168.1	157.4-179.2	384	69.3	61.9-77.1
1991	1,258	183.9	172.9-195.3	359	64.9	57.8-72.5
1992	1,563	217.0	205.4-229.0	392	68.3	61.2-76.0
1993	1,745	231.6	219.9-243.7	376	66.6	59.5-74.3
1994	1,869	237.5	225.9-249.3	417	71.4	64.2-79.2
1995	2,046	247.7	236.2-259.5	435	69.9	63.0-77.3
1996	2,092	243.6	232.5-255.0	409	63.5	57.1-70.4
1997	2,201	252.0	240.9-263.5	405	61.6	55.3-68.4
1998	2,288	252.4	241.5-263.7	402	60.8	54.6-67.3
1999	2,611	276.8	265.7-288.3	413	60.3	54.3-66.8
2000	2,899	298.6	287.2-310.4	425	60.3	54.4-66.6
2001	2,775	276.9	266.1-288.1	425	59.5	53.7-65.8
2002	2,817	274.5	263.8-285.5	437	57.4	51.8-63.3
2003	2,809	265.5	255.2-276.1	378	48.7	43.7-54.1
2004	2,833	257.8	247.9-268.0	374	48.1	43.1-53.5
2005	2,853	252.1	242.4-262.0	370	44.6	39.9-49.6
2006	3,034	263.7	253.9-273.8	346	42.4	37.8-47.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	3,300	278.3	268.4-288.5	381	45.5	40.9-50.5
2008	3,216	260.0	250.7-269.6	365	42.5	38.0-47.2
2009	3,224	251.5	242.4-260.7	393	44.7	40.2-49.5
2010	3,194	240.8	232.1-249.8	381	41.4	37.2-45.9
2011	3,482	251.5	242.8-260.4	369	38.5	34.5-42.8
2012	3,152	220.2	212.2-228.3	368	37.5	33.7-41.7
2013	3,233	219.2	211.4-227.2	366	36.1	32.4-40.1
2014	2,958	193.8	186.6-201.2	385	34.5	31.0-38.2
2015	3,027	191.7	184.7-199.0	336	29.5	26.3-32.9
2016	3,181	197.6	190.5-204.8	366	30.5	27.4-33.9
2017	3,313	200.4	193.4-207.6	390	31.1	28.0-34.4
2018	3,293	193.9	187.1-200.9	373	29.0	26.0-32.1
2019	3,452	201.2	194.3-208.2	388	29.1	26.2-32.3
2020	2,813	160.3	154.3-166.6	379	26.9	24.2-29.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Testis**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	8	0.6	0.2-1.4	4	1.0	0.1-2.8
1977	9	1.0	0.4-2.2	3	0.3	0.1-1.1
1978	4	0.5	0.1-1.5	4	0.5	0.1-1.5
1979	6	0.5	0.2-1.3	0	0.0	0.0-0.7
1980	15	1.4	0.8-2.5	3	0.3	0.1-1.0
1981	10	1.0	0.4-1.9	2	0.2	0.0-1.0
1982	7	0.7	0.3-1.5	0	0.0	0.0-0.6
1983	7	0.7	0.2-1.6	7	1.0	0.3-2.3
1984	8	0.6	0.2-1.3	1	0.2	0.0-0.9
1985	18	1.8	0.9-3.1	2	0.1	0.0-0.7
1986	17	1.3	0.7-2.2	2	0.2	0.0-0.7
1987	13	0.8	0.4-1.5	2	0.1	0.0-0.6
1988	14	0.9	0.5-1.7	2	0.4	0.0-1.5
1989	15	1.3	0.7-2.2	1	0.1	0.0-0.6
1990	13	1.1	0.5-1.9	3	0.2	0.0-0.7
1991	10	0.8	0.3-1.7	1	0.1	0.0-0.6
1992	23	1.6	0.9-2.7	2	0.1	0.0-0.6
1993	13	0.9	0.5-1.7	3	0.2	0.0-0.7
1994	18	1.2	0.7-2.0	1	0.1	0.0-0.6
1995	7	0.4	0.2-1.0	0	0.0	0.0-0.4
1996	21	1.3	0.8-2.1	0	0.0	0.0-0.4
1997	13	0.8	0.4-1.4	1	0.1	0.0-0.5
1998	21	1.3	0.8-2.1	1	0.1	0.0-0.5
1999	22	1.4	0.9-2.2	1	0.1	0.0-0.5
2000	30	2.1	1.4-3.1	3	0.3	0.0-0.8
2001	23	1.5	0.9-2.2	1	0.1	0.0-0.6
2002	19	1.2	0.7-1.9	2	0.1	0.0-0.5
2003	27	1.7	1.1-2.5	1	0.1	0.0-0.4
2004	16	1.0	0.5-1.6	1	0.1	0.0-0.4
2005	24	1.6	1.0-2.4	2	0.1	0.0-0.5
2006	30	1.9	1.3-2.7	1	0.1	0.0-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Testis**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	30	1.9	1.3-2.8	1	0.1	0.0-0.4
2008	17	1.1	0.6-1.7	1	0.0	0.0-0.4
2009	18	1.1	0.6-1.7	1	0.1	0.0-0.4
2010	24	1.5	0.9-2.2	2	0.1	0.0-0.5
2011	27	1.6	1.1-2.4	3	0.2	0.0-0.6
2012	24	1.5	0.9-2.2	1	0.1	0.0-0.4
2013	27	1.5	1.0-2.2	1	0.1	0.0-0.4
2014	36	2.2	1.5-3.1	1	0.1	0.0-0.3
2015	33	1.9	1.3-2.8	1	0.1	0.0-0.4
2016	35	2.0	1.4-2.8	2	0.2	0.0-0.6
2017	42	2.5	1.8-3.4	4	0.3	0.1-0.7
2018	28	1.7	1.1-2.4	0	0.0	0.0-0.2
2019	30	1.6	1.1-2.3	1	0.1	0.0-0.4
2020	34	2.1	1.4-2.9	0	0.0	0.0-0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	161	13.8	11.5-16.4	107	24.1	19.0-30.0	54	7.7	5.7-10.3	69	5.9	4.5-7.7	39	7.6	5.1-10.9	30	4.7	3.1-6.9
1977	131	10.5	8.6-12.6	78	16.2	12.4-20.7	53	6.9	5.0-9.2	60	5.5	4.0-7.1	35	9.3	6.0-13.4	25	3.5	2.2-5.2
1978	141	10.6	8.8-12.6	94	19.3	15.1-24.1	47	5.5	3.9-7.4	62	4.8	3.6-6.3	35	7.5	4.9-10.7	27	3.3	2.1-4.9
1979	167	12.6	10.6-14.8	97	19.0	14.9-23.6	70	8.8	6.7-11.2	76	6.6	5.1-8.4	40	9.8	6.7-13.6	36	4.9	3.4-6.9
1980	181	12.6	10.7-14.7	120	21.4	17.4-26.0	61	7.2	5.4-9.4	66	5.2	4.0-6.8	37	8.0	5.3-11.3	29	3.7	2.4-5.4
1981	158	10.7	8.9-12.6	90	15.8	12.3-19.9	68	7.6	5.8-9.8	78	5.5	4.2-6.9	46	8.8	6.2-12.2	32	3.6	2.4-5.1
1982	161	10.6	8.9-12.5	89	16.0	12.5-20.0	72	7.5	5.8-9.6	76	5.2	4.1-6.6	37	6.9	4.6-9.7	39	4.3	3.0-5.9
1983	163	10.0	8.4-11.7	87	13.9	10.8-17.5	76	7.6	5.9-9.6	67	4.5	3.5-5.8	31	5.7	3.7-8.3	36	3.9	2.7-5.4
1984	167	10.0	8.4-11.7	109	17.7	14.2-21.7	58	5.6	4.2-7.3	73	4.8	3.7-6.1	45	7.8	5.5-10.7	28	3.1	2.0-4.4
1985	192	11.7	10.0-13.5	109	17.5	14.0-21.5	83	8.4	6.6-10.4	75	4.9	3.8-6.2	43	8.3	5.9-11.3	32	3.2	2.2-4.6
1986	170	10.2	8.6-11.9	111	19.1	15.4-23.4	59	5.5	4.1-7.1	64	4.1	3.1-5.3	33	6.4	4.2-9.1	31	3.0	2.0-4.3
1987	163	9.5	8.0-11.1	105	16.3	13.0-20.0	58	5.7	4.3-7.4	72	4.5	3.5-5.7	40	7.2	5.0-10.0	32	3.1	2.1-4.4
1988	188	10.8	9.3-12.5	114	17.0	13.8-20.7	74	7.1	5.6-9.0	68	4.1	3.2-5.3	40	7.5	5.1-10.4	28	2.6	1.7-3.7
1989	177	9.6	8.2-11.2	106	15.1	12.1-18.6	71	6.4	4.9-8.1	71	4.5	3.5-5.7	42	7.7	5.4-10.6	29	2.9	1.9-4.1
1990	177	9.8	8.4-11.4	115	18.0	14.6-21.9	62	5.5	4.2-7.1	74	4.3	3.3-5.4	41	6.3	4.4-8.7	33	3.1	2.1-4.4
1991	203	10.3	8.9-11.9	130	18.9	15.5-22.7	73	5.9	4.6-7.4	64	3.6	2.8-4.7	40	6.8	4.7-9.4	24	2.1	1.3-3.1
1992	193	9.6	8.3-11.2	112	15.7	12.7-19.2	81	6.4	5.1-8.0	57	3.1	2.3-4.1	30	4.9	3.2-7.1	27	2.3	1.5-3.3
1993	197	9.8	8.5-11.3	101	14.0	11.2-17.2	96	7.6	6.1-9.3	79	4.0	3.2-5.0	44	6.6	4.6-8.9	35	2.8	1.9-3.8
1994	198	9.7	8.4-11.2	113	15.8	12.8-19.2	85	6.5	5.2-8.1	72	3.9	3.0-4.9	36	5.5	3.8-7.8	36	3.0	2.1-4.1
1995	179	8.6	7.3-9.9	119	15.9	12.9-19.2	60	4.6	3.5-5.9	67	3.5	2.7-4.4	33	5.3	3.4-7.6	34	2.7	1.9-3.8
1996	210	9.8	8.5-11.2	119	15.3	12.4-18.5	91	6.8	5.5-8.3	74	3.7	2.9-4.7	41	6.1	4.2-8.3	33	2.6	1.8-3.6
1997	213	9.9	8.6-11.4	124	16.5	13.5-20.0	89	6.6	5.3-8.2	72	3.5	2.8-4.5	35	5.0	3.3-7.0	37	2.8	2.0-3.9
1998	258	11.6	10.2-13.2	157	19.4	16.3-23.0	101	7.4	6.0-8.9	83	4.0	3.2-5.0	39	5.6	3.9-7.7	44	3.3	2.4-4.4
1999	233	9.9	8.6-11.2	149	17.1	14.3-20.3	84	5.7	4.6-7.1	69	3.1	2.4-3.9	35	4.4	3.0-6.2	34	2.4	1.7-3.3
2000	245	10.2	9.0-11.6	148	16.7	14.0-19.8	97	6.5	5.3-8.0	81	3.6	2.8-4.4	43	5.4	3.8-7.4	38	2.6	1.9-3.6
2001	252	10.3	9.0-11.6	152	16.7	14.0-19.8	100	6.6	5.3-8.0	83	3.6	2.8-4.4	37	4.7	3.2-6.6	46	3.1	2.2-4.1
2002	241	9.8	8.6-11.2	143	15.9	13.2-18.9	98	6.3	5.1-7.7	66	2.8	2.2-3.6	28	3.3	2.1-4.8	38	2.5	1.8-3.4
2003	257	9.8	8.6-11.1	177	18.0	15.3-21.0	80	4.9	3.9-6.2	71	2.9	2.2-3.6	39	4.5	3.1-6.2	32	2.0	1.4-2.9
2004	279	10.7	9.5-12.1	179	18.4	15.6-21.5	100	6.3	5.1-7.6	78	3.1	2.4-3.8	40	4.2	2.9-5.8	38	2.4	1.7-3.3
2005	304	11.2	10.0-12.6	193	18.5	15.8-21.5	111	6.8	5.6-8.2	80	3.4	2.7-4.2	45	5.5	3.9-7.4	35	2.3	1.6-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	300	11.1	9.8-12.4	199	19.9	17.1-23.0	101	5.9	4.8-7.2	61	2.4	1.8-3.1	39	4.3	3.0-6.0	22	1.4	0.9-2.1
2007	261	9.5	8.4-10.8	161	15.9	13.4-18.7	100	5.9	4.8-7.2	78	3.1	2.4-3.8	37	4.2	2.9-5.8	41	2.5	1.8-3.3
2008	298	10.4	9.2-11.6	198	18.1	15.5-21.0	100	5.7	4.6-7.0	67	2.5	1.9-3.2	37	4.2	2.9-5.8	30	1.7	1.1-2.4
2009	320	11.1	9.9-12.5	204	18.8	16.2-21.7	116	6.6	5.5-8.0	73	2.7	2.1-3.5	42	4.7	3.3-6.3	31	1.8	1.2-2.5
2010	305	10.1	9.0-11.4	208	18.4	15.8-21.2	97	5.2	4.2-6.4	88	3.1	2.5-3.9	43	4.6	3.2-6.2	45	2.5	1.8-3.4
2011	361	11.6	10.4-12.9	255	21.7	19.0-24.7	106	5.5	4.5-6.7	88	3.0	2.4-3.7	51	4.9	3.6-6.5	37	2.0	1.4-2.8
2012	335	10.7	9.6-12.0	211	18.3	15.8-21.1	124	6.4	5.3-7.6	88	2.9	2.3-3.5	53	4.6	3.4-6.1	35	1.9	1.3-2.6
2013	388	12.1	10.9-13.4	254	20.7	18.1-23.5	134	6.9	5.7-8.2	88	2.8	2.2-3.4	52	4.6	3.4-6.1	36	1.8	1.2-2.5
2014	360	10.6	9.6-11.8	241	18.3	16.0-20.9	119	5.8	4.8-6.9	91	2.9	2.4-3.6	56	5.2	3.9-6.7	35	1.7	1.2-2.4
2015	391	11.4	10.2-12.6	257	19.3	16.9-21.9	134	6.5	5.4-7.7	84	2.5	2.0-3.1	45	3.6	2.6-4.8	39	1.8	1.3-2.5
2016	373	10.4	9.4-11.6	252	18.1	15.9-20.6	121	5.5	4.6-6.6	89	2.6	2.1-3.3	52	4.3	3.2-5.7	37	1.7	1.2-2.4
2017	450	12.5	11.3-13.7	303	21.9	19.4-24.7	147	6.7	5.7-7.9	86	2.5	2.0-3.1	53	4.3	3.2-5.6	33	1.5	1.1-2.2
2018	401	10.7	9.6-11.8	266	18.0	15.8-20.4	135	6.1	5.1-7.3	91	2.5	2.0-3.1	52	3.9	2.9-5.1	39	1.7	1.2-2.4
2019	411	10.6	9.6-11.7	286	19.1	16.9-21.6	125	5.4	4.5-6.5	86	2.3	1.8-2.9	54	3.9	2.9-5.1	32	1.4	0.9-2.0
2020	358	9.2	8.3-10.2	235	15.4	13.4-17.6	123	5.3	4.4-6.3	78	2.0	1.6-2.5	52	3.7	2.7-4.9	26	1.0	0.7-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	96	6.7	5.3-8.4	58	8.6	6.2-11.5	38	5.3	3.6-7.3	47	3.2	2.2-4.4	28	4.4	2.7-6.7	19	2.4	1.3-3.8
1977	74	4.5	3.4-5.7	47	7.7	5.3-10.8	27	2.5	1.7-3.7	29	1.8	1.1-2.6	20	3.0	1.7-4.8	9	0.9	0.4-1.9
1978	111	6.9	5.6-8.4	72	10.4	7.8-13.4	39	4.4	3.1-6.2	54	3.3	2.4-4.4	36	4.9	3.2-7.2	18	2.1	1.2-3.4
1979	88	5.0	4.0-6.2	65	9.2	6.9-11.9	23	2.2	1.4-3.4	47	2.9	2.1-3.9	37	5.7	3.9-8.1	10	1.0	0.5-1.9
1980	89	5.0	3.9-6.2	47	6.7	4.7-9.2	42	4.0	2.8-5.4	36	2.0	1.4-2.8	28	3.9	2.5-5.9	8	0.8	0.3-1.5
1981	93	5.0	4.0-6.2	65	8.4	6.3-10.9	28	2.7	1.8-3.9	37	1.8	1.2-2.5	25	3.0	1.9-4.5	12	0.9	0.5-1.7
1982	105	5.7	4.6-7.0	65	8.6	6.5-11.2	40	3.8	2.6-5.2	46	2.7	2.0-3.7	25	3.4	2.0-5.1	21	2.2	1.3-3.5
1983	115	6.0	4.9-7.2	63	8.3	6.2-10.9	52	4.6	3.4-6.0	49	2.7	1.9-3.6	30	4.2	2.7-6.2	19	1.7	1.0-2.8
1984	141	7.2	6.0-8.5	90	11.5	9.0-14.5	51	4.5	3.3-5.9	52	2.8	2.1-3.8	30	4.3	2.8-6.3	22	2.0	1.2-3.0
1985	126	6.2	5.1-7.5	70	8.6	6.5-11.1	56	4.7	3.5-6.2	58	3.0	2.3-3.9	33	3.9	2.6-5.7	25	2.3	1.5-3.4
1986	139	6.7	5.6-7.9	79	9.2	7.2-11.6	60	4.9	3.7-6.4	73	3.8	3.0-4.9	43	5.8	4.0-8.0	30	2.6	1.7-3.7
1987	147	7.3	6.1-8.6	90	12.1	9.5-15.1	57	4.6	3.4-6.0	50	2.6	1.9-3.5	25	3.3	2.1-5.0	25	2.2	1.4-3.2
1988	141	6.7	5.6-8.0	75	8.8	6.8-11.2	66	5.3	4.1-6.8	56	2.8	2.1-3.7	34	4.6	3.1-6.6	22	1.8	1.1-2.7
1989	135	6.1	5.1-7.3	78	9.7	7.4-12.4	57	4.2	3.2-5.4	46	2.4	1.8-3.3	20	2.9	1.7-4.6	26	2.2	1.4-3.2
1990	170	7.7	6.6-9.0	98	11.3	8.9-14.0	72	5.5	4.3-7.0	54	2.6	1.9-3.4	29	3.9	2.5-5.8	25	1.9	1.2-2.9
1991	187	8.1	7.0-9.4	119	12.9	10.5-15.7	68	5.0	3.9-6.4	60	2.8	2.1-3.7	35	4.5	3.0-6.4	25	1.9	1.2-2.8
1992	175	7.7	6.6-9.0	114	12.7	10.2-15.5	61	4.6	3.5-5.9	53	2.4	1.8-3.2	32	3.9	2.5-5.6	21	1.6	1.0-2.5
1993	186	7.7	6.6-9.0	121	12.7	10.3-15.4	65	4.6	3.5-5.8	67	3.1	2.4-3.9	39	4.5	3.1-6.3	28	2.2	1.4-3.1
1994	196	7.9	6.8-9.1	98	9.7	7.8-12.0	98	6.7	5.4-8.1	51	2.2	1.6-2.9	30	3.4	2.2-4.9	21	1.5	0.9-2.3
1995	235	9.9	8.6-11.2	135	14.4	11.8-17.2	100	7.1	5.7-8.6	76	3.4	2.7-4.3	46	5.5	3.9-7.5	30	2.2	1.5-3.1
1996	219	8.7	7.5-9.9	142	14.0	11.7-16.8	77	5.2	4.1-6.5	63	2.6	2.0-3.4	36	3.7	2.5-5.3	27	1.9	1.2-2.8
1997	232	9.3	8.1-10.6	131	13.5	11.1-16.1	101	6.8	5.5-8.2	71	3.0	2.3-3.8	42	4.8	3.4-6.7	29	2.0	1.3-2.9
1998	253	9.7	8.5-11.0	153	14.5	12.1-17.1	100	6.6	5.3-8.0	57	2.3	1.7-3.0	34	3.3	2.2-4.7	23	1.6	1.0-2.4
1999	270	10.4	9.1-11.7	138	13.2	10.9-15.7	132	8.5	7.1-10.1	54	2.3	1.7-3.0	26	2.6	1.7-4.0	28	1.9	1.3-2.8
2000	241	9.1	7.9-10.3	125	11.5	9.5-13.8	116	7.4	6.1-8.8	62	2.4	1.8-3.1	34	3.4	2.3-4.8	28	1.8	1.2-2.6
2001	281	10.1	8.9-11.3	160	14.2	11.9-16.7	121	7.3	6.1-8.8	69	2.7	2.1-3.4	37	3.8	2.6-5.3	32	2.0	1.4-2.9
2002	293	10.3	9.1-11.5	168	14.9	12.6-17.4	125	7.4	6.2-8.9	54	2.0	1.5-2.6	37	3.6	2.4-5.0	17	1.1	0.6-1.7
2003	325	11.3	10.1-12.6	190	16.0	13.7-18.6	135	8.0	6.7-9.4	70	2.7	2.1-3.4	44	4.4	3.1-6.0	26	1.6	1.1-2.4
2004	343	11.5	10.3-12.8	200	15.8	13.6-18.3	143	8.2	6.9-9.7	73	2.7	2.1-3.4	43	4.0	2.8-5.5	30	1.8	1.2-2.6
2005	361	12.0	10.8-13.3	190	15.0	12.8-17.4	171	9.8	8.3-11.4	67	2.3	1.8-2.9	44	3.7	2.6-5.1	23	1.3	0.9-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	390	12.9	11.6-14.2	240	19.3	16.8-22.0	150	8.5	7.2-10.0	78	2.7	2.1-3.4	42	3.8	2.7-5.2	36	2.1	1.5-2.9
2007	425	13.7	12.4-15.1	267	21.1	18.5-23.9	158	8.8	7.5-10.3	85	3.0	2.4-3.7	57	5.2	3.9-6.8	28	1.6	1.1-2.3
2008	421	13.5	12.2-14.9	227	17.6	15.2-20.1	194	10.7	9.3-12.4	75	2.4	1.9-3.1	45	3.6	2.6-5.0	30	1.7	1.1-2.4
2009	468	14.4	13.1-15.8	275	20.0	17.6-22.7	193	10.2	8.8-11.8	71	2.3	1.8-2.9	46	3.8	2.7-5.2	25	1.3	0.9-2.0
2010	505	15.4	14.1-16.9	320	23.5	20.9-26.4	185	9.9	8.5-11.4	76	2.6	2.0-3.2	42	3.8	2.7-5.2	34	1.9	1.3-2.6
2011	483	14.6	13.3-16.0	301	22.2	19.6-24.9	182	9.5	8.1-11.0	74	2.3	1.8-2.9	45	3.8	2.7-5.1	29	1.5	1.0-2.2
2012	489	14.3	13.0-15.7	284	19.8	17.5-22.4	205	10.4	9.0-12.0	76	2.3	1.8-2.9	43	3.6	2.6-4.9	33	1.6	1.1-2.3
2013	547	15.3	14.1-16.7	334	22.1	19.7-24.7	213	10.4	9.1-12.0	61	1.8	1.4-2.3	39	2.8	1.9-3.8	22	1.1	0.7-1.7
2014	572	15.7	14.4-17.0	375	24.0	21.5-26.6	197	9.5	8.2-11.0	74	2.2	1.7-2.7	50	3.9	2.8-5.1	24	1.2	0.7-1.7
2015	597	16.0	14.7-17.4	368	23.4	21.0-26.1	229	10.7	9.3-12.2	99	2.8	2.3-3.4	68	5.0	3.8-6.3	31	1.4	1.0-2.0
2016	608	16.2	14.9-17.6	378	24.2	21.8-26.9	230	10.7	9.3-12.2	75	2.1	1.6-2.6	51	3.5	2.5-4.6	24	1.1	0.7-1.7
2017	580	15.1	13.9-16.4	362	22.7	20.3-25.2	218	9.9	8.6-11.3	75	2.1	1.6-2.6	58	4.0	3.0-5.2	17	0.8	0.5-1.3
2018	614	15.7	14.5-17.1	387	23.6	21.2-26.2	227	10.1	8.8-11.5	81	2.1	1.7-2.6	48	3.0	2.2-4.0	33	1.5	1.0-2.1
2019	637	16.2	14.9-17.5	366	21.9	19.6-24.3	271	12.0	10.6-13.5	88	2.2	1.8-2.7	50	3.2	2.3-4.2	38	1.6	1.1-2.2
2020	518	13.0	11.9-14.2	318	18.5	16.4-20.7	200	8.8	7.6-10.2	74	2.0	1.5-2.5	46	3.0	2.2-4.0	28	1.3	0.9-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	66	3.3	2.6-4.3	35	4.1	2.8-5.8	31	2.8	1.9-4.0	52	2.8	2.0-3.8	29	3.5	2.3-5.3	23	2.3	1.4-3.5
1977	84	3.9	3.1-4.9	39	4.5	3.1-6.4	45	3.6	2.6-4.8	42	2.2	1.6-3.1	23	2.9	1.8-4.5	19	1.8	1.1-2.9
1978	52	2.7	2.0-3.6	24	3.1	1.9-4.8	28	2.5	1.6-3.7	40	2.1	1.4-2.8	20	2.2	1.3-3.6	20	1.9	1.1-3.0
1979	77	3.5	2.8-4.5	37	4.1	2.8-5.8	40	3.2	2.3-4.4	51	2.6	1.9-3.5	25	3.2	2.0-4.9	26	2.3	1.5-3.3
1980	76	3.9	3.0-5.0	47	5.6	3.9-7.7	29	2.7	1.8-3.9	47	2.5	1.8-3.3	30	3.8	2.4-5.6	17	1.6	0.9-2.5
1981	55	2.5	1.8-3.3	30	2.8	1.8-4.1	25	2.1	1.3-3.2	35	2.0	1.4-2.9	16	2.0	1.1-3.3	19	2.0	1.1-3.2
1982	79	3.4	2.6-4.3	45	4.3	3.1-6.0	34	2.6	1.8-3.7	65	3.3	2.5-4.3	41	5.1	3.6-7.1	24	2.1	1.3-3.1
1983	75	3.3	2.6-4.2	41	4.1	2.9-5.8	34	2.7	1.8-3.8	61	3.0	2.2-3.8	23	2.6	1.6-4.1	38	3.1	2.2-4.4
1984	99	4.1	3.3-5.1	54	5.1	3.7-6.8	45	3.3	2.4-4.5	70	3.1	2.4-4.0	43	4.5	3.1-6.2	27	2.1	1.4-3.1
1985	101	4.4	3.6-5.4	43	5.0	3.5-6.9	58	4.3	3.2-5.6	56	2.5	1.8-3.2	32	3.6	2.4-5.2	24	1.7	1.1-2.6
1986	94	4.1	3.2-5.0	55	5.4	3.9-7.3	39	3.0	2.1-4.2	63	2.8	2.1-3.7	33	3.4	2.3-4.9	30	2.4	1.6-3.4
1987	101	4.1	3.3-5.1	48	4.6	3.2-6.4	53	3.7	2.8-4.9	55	2.7	2.0-3.5	28	3.5	2.2-5.3	27	2.2	1.4-3.2
1988	88	3.4	2.7-4.3	51	4.7	3.4-6.5	37	2.5	1.8-3.6	48	2.0	1.4-2.6	23	2.4	1.4-3.8	25	1.7	1.1-2.5
1989	94	3.6	2.9-4.4	41	3.6	2.5-5.1	53	3.6	2.7-4.7	40	1.6	1.2-2.2	19	1.8	1.0-3.0	21	1.5	0.9-2.3
1990	102	4.0	3.2-4.9	46	4.4	3.1-6.0	56	3.8	2.9-5.0	46	2.0	1.4-2.7	24	2.8	1.7-4.3	22	1.6	1.0-2.4
1991	130	5.1	4.2-6.1	63	6.0	4.4-7.9	67	4.5	3.4-5.7	53	2.2	1.6-3.0	22	2.2	1.3-3.4	31	2.3	1.5-3.2
1992	131	5.1	4.2-6.1	66	6.1	4.6-8.0	65	4.4	3.3-5.6	55	2.4	1.8-3.2	25	2.9	1.7-4.4	30	2.2	1.5-3.2
1993	93	3.4	2.7-4.2	54	4.6	3.3-6.2	39	2.6	1.8-3.5	62	2.3	1.8-3.0	29	2.8	1.8-4.1	33	2.1	1.5-3.0
1994	108	3.8	3.1-4.6	58	4.7	3.5-6.2	50	3.2	2.3-4.2	59	2.2	1.7-2.9	28	2.6	1.7-3.9	31	2.0	1.4-2.8
1995	116	4.2	3.4-5.1	49	3.8	2.8-5.2	67	4.3	3.4-5.5	50	1.8	1.3-2.4	23	1.9	1.2-3.0	27	1.8	1.1-2.6
1996	137	5.0	4.1-5.9	64	5.5	4.2-7.1	73	4.6	3.6-5.8	51	1.9	1.4-2.5	24	2.3	1.4-3.5	27	1.7	1.1-2.5
1997	111	3.8	3.1-4.6	59	4.8	3.5-6.3	52	3.1	2.3-4.1	49	1.8	1.3-2.4	24	1.9	1.2-2.9	25	1.6	1.0-2.4
1998	118	3.9	3.2-4.7	59	4.1	3.1-5.4	59	3.6	2.7-4.6	45	1.7	1.2-2.2	25	2.2	1.4-3.3	20	1.3	0.8-2.0
1999	125	4.3	3.5-5.1	66	5.4	4.0-7.0	59	3.6	2.7-4.6	42	1.5	1.0-2.0	26	2.2	1.4-3.3	16	1.0	0.5-1.6
2000	124	4.1	3.4-4.9	60	4.7	3.5-6.2	64	3.7	2.9-4.8	54	2.0	1.5-2.6	26	2.2	1.4-3.2	28	1.8	1.2-2.6
2001	147	4.9	4.1-5.8	61	4.4	3.3-5.7	86	5.1	4.1-6.3	63	2.2	1.7-2.9	28	2.5	1.6-3.7	35	2.1	1.4-2.9
2002	127	4.2	3.4-5.0	65	5.2	3.9-6.7	62	3.5	2.7-4.5	62	2.1	1.6-2.6	26	1.9	1.2-2.8	36	2.1	1.5-2.9
2003	147	4.8	4.0-5.7	72	5.2	4.1-6.7	75	4.3	3.4-5.5	55	1.8	1.4-2.4	33	2.6	1.7-3.7	22	1.3	0.8-1.9
2004	132	4.2	3.5-5.0	68	5.0	3.9-6.5	64	3.6	2.8-4.6	62	2.0	1.6-2.6	31	2.4	1.6-3.5	31	1.8	1.2-2.5
2005	131	4.2	3.5-5.0	68	5.5	4.2-7.0	63	3.5	2.7-4.5	49	1.6	1.2-2.1	32	2.5	1.7-3.6	17	0.9	0.5-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	113	3.5	2.9-4.3	55	3.9	2.9-5.2	58	3.2	2.4-4.2	59	1.9	1.4-2.5	33	2.6	1.8-3.8	26	1.4	0.9-2.1
2007	130	4.0	3.3-4.7	67	4.6	3.5-5.8	63	3.5	2.6-4.4	49	1.6	1.2-2.1	21	1.8	1.1-2.7	28	1.5	1.0-2.2
2008	124	3.8	3.2-4.6	58	4.1	3.1-5.4	66	3.6	2.8-4.6	63	2.0	1.5-2.6	31	2.4	1.6-3.4	32	1.7	1.2-2.5
2009	130	4.0	3.3-4.8	67	4.8	3.6-6.1	63	3.4	2.6-4.4	56	1.8	1.4-2.4	30	2.4	1.6-3.5	26	1.4	0.9-2.1
2010	145	4.3	3.6-5.0	77	5.2	4.0-6.5	68	3.6	2.8-4.6	54	1.7	1.2-2.2	28	2.1	1.3-3.0	26	1.4	0.9-2.1
2011	151	4.4	3.7-5.1	76	5.1	4.0-6.5	75	3.8	3.0-4.8	72	2.2	1.7-2.7	37	2.6	1.8-3.6	35	1.8	1.3-2.6
2012	138	4.0	3.4-4.8	81	5.3	4.2-6.7	57	3.0	2.2-3.9	65	1.9	1.4-2.4	33	2.2	1.5-3.2	32	1.6	1.1-2.3
2013	160	4.5	3.8-5.2	75	4.5	3.5-5.7	85	4.4	3.5-5.4	73	2.1	1.6-2.6	38	2.6	1.8-3.6	35	1.7	1.2-2.4
2014	121	3.3	2.8-4.0	72	4.4	3.5-5.6	49	2.4	1.8-3.3	66	1.9	1.5-2.4	38	2.7	1.9-3.8	28	1.3	0.9-1.9
2015	140	3.9	3.3-4.7	73	4.6	3.5-5.8	67	3.5	2.7-4.5	65	1.8	1.4-2.3	32	2.2	1.5-3.1	33	1.6	1.1-2.3
2016	144	4.0	3.4-4.7	88	5.5	4.4-6.8	56	2.8	2.1-3.7	84	2.2	1.8-2.8	44	2.7	1.9-3.6	40	1.9	1.4-2.7
2017	136	3.8	3.2-4.5	66	4.3	3.3-5.5	70	3.4	2.6-4.3	59	1.6	1.2-2.0	35	2.2	1.5-3.1	24	1.0	0.7-1.6
2018	172	4.6	4.0-5.4	99	6.1	5.0-7.5	73	3.5	2.7-4.4	86	2.4	1.9-3.0	38	2.5	1.7-3.4	48	2.4	1.7-3.2
2019	173	4.6	3.9-5.4	87	5.2	4.1-6.4	86	4.1	3.3-5.1	90	2.4	1.9-3.0	55	3.4	2.6-4.5	35	1.6	1.1-2.2
2020	168	4.4	3.7-5.1	83	4.8	3.8-6.0	85	4.0	3.2-5.0	83	2.1	1.7-2.6	41	2.3	1.6-3.1	42	1.9	1.3-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	45	2.4	1.7-3.3	13	1.8	0.9-3.3	32	2.9	2.0-4.2	10	0.6	0.3-1.2	3	0.4	0.1-1.3	7	0.8	0.3-1.6
1977	45	2.3	1.6-3.3	3	0.4	0.1-1.2	42	3.7	2.6-5.2	6	0.5	0.1-1.0	3	0.4	0.1-1.3	3	0.5	0.1-1.3
1978	37	2.1	1.4-2.9	11	1.6	0.7-3.1	26	2.5	1.6-3.7	11	0.8	0.4-1.4	2	0.4	0.0-1.4	9	1.1	0.5-2.0
1979	48	2.9	2.1-4.0	11	1.6	0.7-3.2	37	3.8	2.6-5.4	9	0.6	0.2-1.2	2	0.2	0.0-1.0	7	0.8	0.3-1.8
1980	45	2.1	1.5-2.9	6	0.6	0.2-1.4	39	3.2	2.2-4.5	8	0.5	0.2-1.0	1	0.2	0.0-1.0	7	0.7	0.3-1.5
1981	34	1.9	1.3-2.7	6	0.6	0.2-1.5	28	2.7	1.7-3.9	7	0.5	0.2-1.0	1	0.1	0.0-0.8	6	0.7	0.2-1.5
1982	44	2.3	1.6-3.1	7	0.8	0.3-1.8	37	3.3	2.3-4.5	11	0.7	0.3-1.2	4	0.8	0.2-2.2	7	0.6	0.2-1.3
1983	45	2.0	1.5-2.8	6	0.6	0.2-1.4	39	3.1	2.2-4.3	6	0.4	0.1-0.9	1	0.2	0.0-1.1	5	0.5	0.2-1.2
1984	54	2.5	1.8-3.3	8	0.8	0.3-1.6	46	3.6	2.6-4.9	9	0.5	0.2-1.0	2	0.5	0.0-1.7	7	0.6	0.2-1.3
1985	69	3.1	2.4-4.0	19	2.0	1.2-3.3	50	3.9	2.9-5.2	8	0.4	0.2-0.8	4	0.5	0.1-1.3	4	0.4	0.1-0.9
1986	45	2.2	1.6-3.0	13	1.8	0.9-3.2	32	2.5	1.7-3.6	11	0.6	0.3-1.0	4	0.7	0.1-1.8	7	0.5	0.2-1.1
1987	65	2.9	2.2-3.7	13	1.5	0.8-2.7	52	3.8	2.8-5.1	12	0.7	0.4-1.2	3	0.6	0.1-1.6	9	0.8	0.4-1.5
1988	49	2.2	1.6-2.9	11	1.3	0.6-2.4	38	2.9	2.0-3.9	8	0.5	0.2-0.9	3	0.4	0.1-1.1	5	0.5	0.1-1.1
1989	48	2.1	1.5-2.8	5	0.5	0.2-1.4	43	3.1	2.2-4.3	11	0.7	0.4-1.3	1	0.2	0.0-0.8	10	1.0	0.5-1.8
1990	58	2.3	1.7-3.0	9	0.8	0.4-1.7	49	3.4	2.5-4.5	5	0.3	0.1-0.7	1	0.1	0.0-0.8	4	0.4	0.1-0.9
1991	67	2.8	2.1-3.6	8	1.1	0.4-2.2	59	4.0	3.0-5.2	10	0.6	0.3-1.0	2	0.4	0.1-1.4	8	0.6	0.3-1.3
1992	91	3.3	2.6-4.1	22	1.8	1.1-2.9	69	4.4	3.4-5.6	11	0.5	0.2-0.9	3	0.3	0.1-0.8	8	0.6	0.3-1.2
1993	79	3.0	2.4-3.8	17	1.5	0.8-2.4	62	4.1	3.1-5.3	8	0.4	0.2-0.8	3	0.3	0.1-0.9	5	0.4	0.1-1.0
1994	91	3.5	2.8-4.3	13	1.2	0.6-2.2	78	5.1	4.0-6.4	8	0.3	0.1-0.7	2	0.2	0.0-0.8	6	0.4	0.1-0.9
1995	100	3.7	2.9-4.5	15	1.4	0.8-2.4	85	5.3	4.2-6.6	8	0.4	0.2-0.8	1	0.1	0.0-0.5	7	0.6	0.2-1.1
1996	118	4.1	3.3-4.9	23	2.2	1.3-3.5	95	5.7	4.6-6.9	15	0.7	0.4-1.1	9	1.0	0.4-2.0	6	0.4	0.2-0.9
1997	123	4.2	3.5-5.1	26	2.1	1.3-3.1	97	5.8	4.7-7.1	14	0.5	0.3-0.9	8	0.8	0.3-1.7	6	0.4	0.1-0.8
1998	134	4.6	3.8-5.4	25	2.2	1.4-3.3	109	6.4	5.2-7.7	10	0.4	0.2-0.8	4	0.4	0.1-1.0	6	0.4	0.2-0.9
1999	128	4.4	3.7-5.3	18	1.4	0.8-2.2	110	6.5	5.4-7.9	15	0.6	0.3-1.0	2	0.2	0.0-0.6	13	0.9	0.5-1.5
2000	166	5.5	4.7-6.4	24	1.7	1.1-2.5	142	8.2	6.9-9.7	14	0.6	0.3-1.0	2	0.2	0.0-0.6	12	0.8	0.4-1.4
2001	166	5.3	4.5-6.2	33	2.5	1.7-3.6	133	7.5	6.2-8.8	17	0.6	0.4-1.0	3	0.2	0.0-0.7	14	0.9	0.5-1.5
2002	172	5.5	4.7-6.4	31	2.4	1.6-3.4	141	7.8	6.6-9.2	9	0.4	0.2-0.7	2	0.1	0.0-0.6	7	0.5	0.2-0.9
2003	167	5.2	4.5-6.1	30	2.3	1.5-3.4	137	7.5	6.3-8.9	8	0.3	0.1-0.6	1	0.1	0.0-0.4	7	0.5	0.2-0.9
2004	188	5.9	5.1-6.9	34	3.0	2.0-4.3	154	8.4	7.1-9.8	16	0.6	0.4-1.0	7	0.8	0.3-1.6	9	0.6	0.3-1.1
2005	258	7.9	7.0-9.0	45	3.3	2.4-4.5	213	11.5	10.0-13.2	21	0.7	0.4-1.1	9	0.8	0.3-1.5	12	0.7	0.4-1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	225	7.0	6.1-8.0	36	2.9	2.0-4.1	189	10.2	8.8-11.8	11	0.4	0.2-0.7	3	0.2	0.0-0.7	8	0.5	0.2-0.9
2007	243	7.4	6.5-8.4	50	3.7	2.7-4.9	193	10.4	9.0-12.0	12	0.5	0.2-0.8	3	0.3	0.1-0.9	9	0.5	0.2-1.0
2008	293	8.8	7.8-9.9	60	4.4	3.3-5.8	233	12.3	10.7-14.0	15	0.5	0.3-0.9	5	0.5	0.1-1.2	10	0.6	0.3-1.1
2009	302	9.1	8.1-10.2	60	4.1	3.1-5.3	242	13.0	11.4-14.7	15	0.5	0.3-0.9	5	0.5	0.1-1.1	10	0.5	0.3-1.0
2010	341	9.9	8.9-11.1	52	3.7	2.8-5.0	289	14.8	13.2-16.7	21	0.7	0.4-1.0	11	0.9	0.4-1.7	10	0.5	0.3-1.0
2011	396	11.1	10.0-12.3	79	5.5	4.3-6.9	317	15.7	14.0-17.6	12	0.4	0.2-0.7	1	0.1	0.0-0.6	11	0.6	0.3-1.0
2012	363	10.3	9.3-11.5	58	3.9	2.9-5.1	305	15.4	13.7-17.2	18	0.6	0.3-0.9	6	0.5	0.2-1.2	12	0.6	0.3-1.0
2013	439	12.2	11.0-13.4	57	3.6	2.7-4.7	382	18.8	17.0-20.9	19	0.6	0.4-1.0	5	0.4	0.1-0.9	14	0.7	0.4-1.2
2014	425	11.9	10.8-13.1	65	4.3	3.3-5.5	360	18.0	16.2-20.0	20	0.6	0.4-0.9	3	0.2	0.0-0.7	17	0.8	0.5-1.4
2015	433	12.1	11.0-13.3	81	5.3	4.1-6.6	352	17.7	15.8-19.6	18	0.5	0.3-0.8	6	0.4	0.1-0.9	12	0.6	0.3-1.0
2016	385	10.5	9.5-11.6	73	4.6	3.6-5.8	312	15.2	13.5-17.0	18	0.5	0.3-0.8	5	0.3	0.1-0.7	13	0.6	0.3-1.0
2017	364	9.9	8.9-11.0	68	4.5	3.5-5.8	296	14.4	12.8-16.2	22	0.6	0.4-1.0	8	0.6	0.3-1.2	14	0.6	0.3-1.1
2018	369	10.0	9.0-11.1	71	4.4	3.4-5.6	298	14.5	12.8-16.2	16	0.4	0.2-0.6	9	0.5	0.2-1.0	7	0.3	0.1-0.6
2019	400	10.8	9.8-12.0	79	4.7	3.7-5.9	321	15.9	14.2-17.8	17	0.5	0.3-0.7	1	0.1	0.0-0.4	16	0.7	0.4-1.2
2020	321	8.6	7.7-9.6	61	3.7	2.8-4.7	260	12.8	11.2-14.5	23	0.6	0.4-0.9	6	0.4	0.1-0.9	17	0.7	0.4-1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	46	2.7	1.8-3.8	21	3.3	1.6-5.8	25	2.4	1.5-3.6	18	0.9	0.5-1.5	10	1.1	0.5-2.1	8	0.8	0.3-1.5
1977	38	1.7	1.2-2.4	20	2.1	1.2-3.4	18	1.5	0.9-2.4	19	1.1	0.6-1.9	7	1.3	0.3-3.0	12	1.1	0.6-2.0
1978	41	1.9	1.3-2.6	24	2.4	1.5-3.7	17	1.5	0.8-2.5	14	0.6	0.3-1.1	8	0.9	0.3-1.9	6	0.4	0.2-1.0
1979	48	2.1	1.5-2.8	30	3.0	1.9-4.5	18	1.4	0.8-2.3	21	1.0	0.6-1.6	14	1.5	0.8-2.6	7	0.7	0.2-1.5
1980	60	2.3	1.7-3.0	32	2.7	1.8-3.9	28	1.9	1.2-2.8	13	0.6	0.3-1.0	6	0.7	0.3-1.6	7	0.5	0.2-1.0
1981	54	2.2	1.6-2.9	33	3.0	1.9-4.5	21	1.5	0.9-2.4	22	1.1	0.6-1.7	13	1.6	0.7-2.9	9	0.7	0.3-1.4
1982	59	2.4	1.8-3.1	32	3.0	2.0-4.3	27	1.9	1.3-2.9	18	0.8	0.4-1.2	9	0.9	0.4-1.9	9	0.7	0.3-1.3
1983	67	2.6	2.0-3.4	39	3.7	2.5-5.4	28	1.9	1.2-2.8	14	0.6	0.3-1.1	8	0.7	0.3-1.5	6	0.5	0.2-1.1
1984	58	2.3	1.7-3.0	39	3.6	2.5-5.1	19	1.4	0.8-2.2	21	0.9	0.5-1.4	19	1.7	1.0-2.9	2	0.2	0.0-0.7
1985	51	1.8	1.3-2.4	31	2.6	1.7-3.8	20	1.2	0.7-1.9	27	1.0	0.7-1.5	18	1.6	0.9-2.7	9	0.6	0.3-1.1
1986	66	2.5	1.9-3.2	46	3.9	2.8-5.3	20	1.3	0.8-2.1	19	0.8	0.4-1.2	13	1.2	0.6-2.1	6	0.5	0.2-1.0
1987	64	2.3	1.7-2.9	37	3.3	2.2-4.8	27	1.6	1.0-2.4	17	0.6	0.4-1.1	14	1.4	0.7-2.4	3	0.2	0.0-0.5
1988	70	2.3	1.7-2.9	44	2.9	2.1-4.0	26	1.7	1.1-2.5	17	0.6	0.3-1.0	9	0.8	0.3-1.6	8	0.5	0.2-1.0
1989	71	2.4	1.8-3.1	43	3.3	2.3-4.6	28	1.7	1.1-2.5	19	0.7	0.4-1.2	16	1.2	0.7-2.1	3	0.3	0.1-0.8
1990	69	2.4	1.8-3.0	39	3.0	2.1-4.3	30	1.8	1.2-2.6	20	0.7	0.4-1.1	12	1.0	0.5-1.9	8	0.5	0.2-1.0
1991	94	3.1	2.4-3.8	54	4.2	3.0-5.6	40	2.3	1.6-3.2	18	0.6	0.3-1.0	8	0.6	0.2-1.2	10	0.6	0.3-1.2
1992	66	2.2	1.7-2.8	36	2.5	1.7-3.6	30	1.8	1.2-2.7	22	0.7	0.4-1.1	12	0.8	0.4-1.6	10	0.6	0.3-1.1
1993	73	2.2	1.7-2.8	40	2.8	1.9-3.9	33	1.8	1.3-2.6	17	0.5	0.3-0.9	10	0.7	0.3-1.4	7	0.4	0.2-0.9
1994	60	2.0	1.5-2.6	36	2.8	1.9-4.0	24	1.4	0.9-2.1	20	0.7	0.4-1.1	13	1.1	0.6-2.0	7	0.4	0.2-0.8
1995	80	2.4	1.9-3.1	48	3.2	2.4-4.4	32	1.8	1.2-2.5	17	0.5	0.3-0.8	13	0.9	0.5-1.6	4	0.2	0.1-0.6
1996	69	2.2	1.7-2.8	34	2.4	1.6-3.5	35	2.0	1.4-2.8	13	0.5	0.2-0.8	7	0.6	0.2-1.4	6	0.4	0.1-0.8
1997	80	2.4	1.9-3.0	51	3.3	2.4-4.4	29	1.6	1.1-2.3	16	0.6	0.3-1.0	6	0.4	0.1-0.9	10	0.7	0.3-1.2
1998	82	2.4	1.9-3.0	43	2.9	2.0-3.9	39	2.1	1.5-2.9	12	0.3	0.2-0.6	5	0.3	0.1-0.8	7	0.4	0.1-0.8
1999	79	2.4	1.9-3.1	43	3.0	2.1-4.2	36	2.0	1.4-2.7	14	0.5	0.3-0.8	9	0.7	0.3-1.4	5	0.3	0.1-0.7
2000	92	2.7	2.1-3.3	56	3.6	2.7-4.7	36	1.9	1.3-2.6	5	0.2	0.0-0.4	3	0.2	0.0-0.6	2	0.1	0.0-0.4
2001	85	2.4	1.9-3.0	56	3.5	2.6-4.6	29	1.5	1.0-2.2	12	0.4	0.2-0.7	6	0.4	0.2-1.0	6	0.3	0.1-0.7
2002	89	2.7	2.2-3.3	49	3.3	2.4-4.4	40	2.2	1.6-3.0	10	0.3	0.1-0.5	8	0.5	0.2-1.1	2	0.1	0.0-0.4
2003	84	2.4	1.9-3.0	35	2.3	1.6-3.2	49	2.6	1.9-3.4	14	0.5	0.3-0.8	8	0.8	0.3-1.5	6	0.3	0.1-0.7
2004	94	2.7	2.2-3.3	48	3.0	2.2-4.0	46	2.4	1.8-3.2	9	0.3	0.1-0.5	6	0.4	0.2-1.0	3	0.2	0.0-0.5
2005	99	2.9	2.4-3.6	60	3.8	2.9-5.0	39	2.1	1.5-2.9	11	0.3	0.2-0.5	9	0.6	0.3-1.1	2	0.1	0.0-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	90	2.7	2.1-3.3	49	3.2	2.4-4.3	41	2.2	1.6-3.0	9	0.3	0.1-0.6	5	0.3	0.1-0.8	4	0.2	0.1-0.6
2007	87	2.5	2.0-3.1	51	3.5	2.5-4.6	36	1.9	1.3-2.6	13	0.4	0.2-0.7	4	0.3	0.1-0.8	9	0.5	0.2-0.9
2008	122	3.5	2.9-4.2	71	4.6	3.5-5.8	51	2.7	2.0-3.5	8	0.2	0.1-0.5	5	0.3	0.1-0.8	3	0.2	0.0-0.5
2009	105	3.1	2.5-3.7	47	3.2	2.3-4.3	58	3.0	2.3-3.9	10	0.3	0.1-0.6	6	0.4	0.2-1.0	4	0.2	0.1-0.5
2010	119	3.4	2.8-4.0	69	4.4	3.4-5.6	50	2.6	1.9-3.4	9	0.2	0.1-0.5	5	0.3	0.1-0.7	4	0.2	0.1-0.5
2011	118	3.2	2.6-3.8	65	3.7	2.8-4.7	53	2.7	2.0-3.5	14	0.4	0.2-0.7	8	0.6	0.2-1.2	6	0.3	0.1-0.6
2012	105	2.9	2.4-3.6	49	3.0	2.2-4.0	56	2.9	2.2-3.7	5	0.2	0.1-0.4	4	0.3	0.1-0.7	1	0.1	0.0-0.3
2013	121	3.4	2.8-4.1	64	4.0	3.1-5.2	57	3.0	2.2-3.8	10	0.3	0.1-0.5	5	0.4	0.1-0.8	5	0.2	0.1-0.6
2014	119	3.2	2.6-3.8	61	3.6	2.7-4.6	58	2.9	2.2-3.7	7	0.2	0.1-0.4	4	0.3	0.1-0.7	3	0.1	0.0-0.4
2015	117	3.3	2.7-3.9	59	3.5	2.7-4.6	58	3.1	2.3-4.0	11	0.3	0.2-0.6	6	0.4	0.2-1.0	5	0.3	0.1-0.6
2016	99	2.7	2.2-3.3	43	2.7	1.9-3.7	56	2.8	2.1-3.6	4	0.1	0.0-0.3	4	0.3	0.1-0.7	0	0.0	0.0-0.2
2017	104	2.7	2.2-3.3	52	2.9	2.2-3.9	52	2.6	1.9-3.4	7	0.2	0.1-0.4	2	0.1	0.0-0.4	5	0.2	0.1-0.6
2018	100	2.7	2.2-3.3	60	3.6	2.7-4.6	40	2.0	1.4-2.8	11	0.3	0.1-0.5	7	0.4	0.2-0.9	4	0.2	0.1-0.5
2019	115	3.2	2.6-3.8	58	3.5	2.6-4.5	57	2.9	2.2-3.7	7	0.2	0.1-0.4	3	0.2	0.0-0.5	4	0.2	0.1-0.5
2020	81	2.1	1.7-2.7	41	2.4	1.7-3.3	40	1.9	1.4-2.7	9	0.2	0.1-0.5	6	0.3	0.1-0.7	3	0.1	0.0-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	98	6.1	4.8-7.6	49	7.7	5.4-10.5	49	5.1	3.7-6.8	57	4.0	2.9-5.3	34	6.0	4.0-8.7	23	2.6	1.6-4.0
1977	118	7.1	5.7-8.6	57	8.5	6.2-11.5	61	6.1	4.6-8.0	61	3.8	2.8-5.1	28	4.2	2.5-6.6	33	3.5	2.3-5.1
1978	156	8.9	7.5-10.6	84	10.8	8.4-13.6	72	7.4	5.7-9.5	65	3.7	2.8-4.7	41	5.6	3.9-7.9	24	2.3	1.4-3.5
1979	115	6.8	5.5-8.2	64	8.5	6.3-11.2	51	5.4	3.9-7.2	79	5.2	4.0-6.7	42	6.0	4.1-8.4	37	4.5	3.1-6.4
1980	133	7.5	6.2-9.0	69	9.6	7.1-12.5	64	6.2	4.7-8.0	73	4.3	3.2-5.5	42	5.7	3.9-8.0	31	3.3	2.1-4.7
1981	131	7.5	6.1-9.0	66	7.7	5.7-10.2	65	7.0	5.3-9.0	78	4.4	3.4-5.5	34	4.8	3.2-7.0	44	4.1	2.9-5.6
1982	163	8.5	7.2-10.1	81	9.8	7.6-12.5	82	7.5	5.9-9.4	61	3.6	2.7-4.7	30	4.3	2.7-6.5	31	3.2	2.1-4.5
1983	173	8.7	7.4-10.1	85	9.8	7.6-12.4	88	7.8	6.2-9.7	91	4.8	3.8-5.9	49	5.8	4.1-7.9	42	4.0	2.8-5.4
1984	190	9.3	7.9-10.8	96	11.0	8.7-13.8	94	8.1	6.4-9.9	95	4.8	3.8-5.9	54	6.3	4.5-8.4	41	3.7	2.6-5.0
1985	233	10.9	9.4-12.4	125	13.6	11.0-16.6	108	8.9	7.2-10.8	108	5.3	4.3-6.5	55	6.0	4.3-8.2	53	4.7	3.5-6.2
1986	242	11.5	10.0-13.1	122	14.1	11.4-17.3	120	9.8	8.0-11.8	102	4.9	3.9-6.0	52	6.0	4.2-8.1	50	4.1	3.0-5.5
1987	252	11.2	9.8-12.8	150	16.4	13.5-19.6	102	7.7	6.3-9.4	95	4.4	3.5-5.4	50	5.6	4.0-7.6	45	3.5	2.6-4.8
1988	232	10.2	8.8-11.6	137	14.5	11.9-17.4	95	7.3	5.8-8.9	117	5.7	4.7-6.9	76	9.9	7.5-12.8	41	3.3	2.3-4.5
1989	272	11.2	9.9-12.7	155	14.4	12.0-17.1	117	8.7	7.2-10.5	115	5.1	4.2-6.2	67	7.2	5.3-9.4	48	3.7	2.7-5.0
1990	312	12.4	11.0-14.0	185	16.2	13.8-19.0	127	9.2	7.7-11.1	106	4.7	3.8-5.7	55	5.9	4.2-7.9	51	3.9	2.9-5.2
1991	312	12.7	11.3-14.3	180	16.4	13.8-19.3	132	9.8	8.1-11.6	109	4.8	3.9-5.9	62	6.1	4.5-8.0	47	3.8	2.7-5.0
1992	361	14.4	12.9-16.1	212	19.2	16.5-22.2	149	10.6	9.0-12.5	114	5.0	4.1-6.0	60	6.3	4.6-8.3	54	4.0	3.0-5.3
1993	367	13.5	12.1-15.0	227	19.0	16.5-21.8	140	9.2	7.7-10.9	117	4.9	4.0-5.9	55	5.1	3.7-6.7	62	4.5	3.4-5.8
1994	407	15.7	14.2-17.4	233	20.3	17.6-23.3	174	12.0	10.3-14.0	115	4.9	4.0-6.0	57	5.7	4.2-7.5	58	4.3	3.2-5.5
1995	384	14.6	13.1-16.2	239	21.3	18.4-24.5	145	9.8	8.2-11.6	118	4.9	4.0-5.9	65	5.8	4.4-7.5	53	3.9	2.9-5.1
1996	414	15.4	13.9-17.0	228	18.8	16.3-21.6	186	12.3	10.6-14.2	125	5.3	4.3-6.3	58	6.0	4.4-8.0	67	4.8	3.7-6.0
1997	432	16.1	14.6-17.8	248	21.9	19.0-25.0	184	12.0	10.3-13.9	116	4.9	4.0-5.9	60	6.3	4.7-8.3	56	3.9	2.9-5.1
1998	437	15.8	14.3-17.4	244	21.1	18.3-24.2	193	12.2	10.5-14.0	142	5.8	4.8-6.8	73	7.3	5.6-9.3	69	4.7	3.7-6.0
1999	463	16.4	14.9-18.0	251	21.1	18.4-24.0	212	13.0	11.3-14.9	146	5.5	4.6-6.5	79	7.5	5.8-9.5	67	4.3	3.3-5.5
2000	401	13.6	12.3-15.0	210	16.9	14.5-19.5	191	11.3	9.7-13.0	141	5.2	4.4-6.2	84	7.8	6.1-9.8	57	3.6	2.7-4.6
2001	435	15.0	13.6-16.5	235	19.0	16.5-21.7	200	12.0	10.4-13.8	153	5.7	4.8-6.7	78	7.0	5.4-8.9	75	4.7	3.7-5.9
2002	431	14.2	12.9-15.6	232	17.8	15.5-20.3	199	11.5	9.9-13.2	153	5.4	4.6-6.4	87	7.6	6.0-9.5	66	4.0	3.1-5.0
2003	464	15.4	14.0-16.9	214	17.0	14.7-19.6	250	14.3	12.6-16.2	136	4.7	3.9-5.5	71	6.1	4.7-7.8	65	3.8	2.9-4.8
2004	425	14.2	12.8-15.6	218	17.8	15.4-20.5	207	11.8	10.2-13.5	123	4.3	3.6-5.2	65	5.5	4.2-7.1	58	3.4	2.6-4.5
2005	483	15.6	14.3-17.1	243	18.6	16.2-21.2	240	13.5	11.9-15.4	123	4.4	3.6-5.2	63	5.7	4.3-7.4	60	3.5	2.7-4.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	462	14.9	13.6-16.4	229	17.6	15.3-20.2	233	13.0	11.4-14.8	115	3.8	3.1-4.5	54	4.4	3.2-5.8	61	3.4	2.6-4.4
2007	484	15.3	13.9-16.7	272	20.2	17.8-22.9	212	11.7	10.2-13.4	131	4.4	3.6-5.2	64	5.5	4.2-7.1	67	3.8	2.9-4.8
2008	508	15.8	14.4-17.2	287	21.0	18.6-23.7	221	11.9	10.4-13.6	114	3.7	3.0-4.5	60	5.0	3.7-6.5	54	2.9	2.2-3.8
2009	517	16.2	14.8-17.7	254	18.9	16.5-21.5	263	14.4	12.7-16.2	120	3.8	3.1-4.6	63	5.0	3.8-6.5	57	3.1	2.3-4.0
2010	504	15.4	14.0-16.8	260	18.6	16.2-21.1	244	12.9	11.3-14.6	127	4.1	3.4-4.9	61	4.8	3.6-6.3	66	3.6	2.8-4.6
2011	575	17.1	15.7-18.6	316	23.1	20.5-26.0	259	13.2	11.6-14.9	134	4.3	3.6-5.1	76	6.0	4.6-7.6	58	3.1	2.3-4.0
2012	526	15.4	14.1-16.8	258	18.0	15.7-20.4	268	13.6	12.0-15.4	123	3.7	3.1-4.5	63	5.0	3.8-6.5	60	3.0	2.3-3.9
2013	539	15.4	14.1-16.8	270	18.4	16.2-20.8	269	13.3	11.8-15.0	123	3.7	3.1-4.4	64	4.8	3.7-6.2	59	2.9	2.2-3.8
2014	575	15.9	14.6-17.3	308	20.3	18.0-22.8	267	12.8	11.3-14.5	136	4.0	3.3-4.7	72	5.5	4.3-7.0	64	3.1	2.4-4.0
2015	577	15.9	14.6-17.3	275	18.2	16.0-20.5	302	14.4	12.8-16.1	127	3.7	3.1-4.4	62	4.7	3.6-6.1	65	3.1	2.4-3.9
2016	612	16.8	15.5-18.2	305	20.2	17.9-22.7	307	14.6	13.0-16.3	132	3.7	3.1-4.4	66	4.7	3.6-6.0	66	3.1	2.4-3.9
2017	592	16.0	14.7-17.3	287	18.5	16.4-20.9	305	14.4	12.8-16.1	136	3.7	3.1-4.4	77	5.2	4.1-6.5	59	2.7	2.1-3.5
2018	566	15.1	13.8-16.4	301	19.0	16.8-21.3	265	12.3	10.8-13.9	123	3.3	2.7-3.9	70	4.6	3.5-5.8	53	2.4	1.8-3.2
2019	599	15.7	14.4-17.0	317	19.7	17.5-22.0	282	12.9	11.4-14.6	150	4.0	3.4-4.7	84	5.6	4.4-7.0	66	3.0	2.3-3.8
2020	505	13.2	12.1-14.5	266	16.5	14.5-18.6	239	10.9	9.6-12.5	110	2.8	2.3-3.4	62	3.7	2.8-4.8	48	2.1	1.6-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	122	9.0	7.3-10.9	58	10.9	8.0-14.5	64	7.8	5.9-10.1	72	5.4	4.1-7.0	30	4.9	3.2-7.1	42	5.6	3.9-7.7
1977	122	9.1	7.4-11.0	54	10.4	7.3-14.1	68	8.4	6.4-10.7	81	6.4	4.9-8.1	40	8.9	6.0-12.5	41	5.1	3.5-7.0
1978	121	8.7	7.1-10.5	64	13.1	9.6-17.2	57	6.3	4.7-8.3	61	4.4	3.3-5.7	34	7.3	4.7-10.6	27	2.9	1.9-4.2
1979	141	9.3	7.7-11.0	63	10.3	7.7-13.6	78	8.6	6.7-10.8	72	5.0	3.8-6.3	31	5.4	3.4-7.9	41	4.7	3.3-6.5
1980	153	10.1	8.4-12.0	61	10.7	7.9-14.1	92	9.8	7.8-12.1	99	6.6	5.3-8.2	43	7.5	5.2-10.4	56	6.1	4.5-8.0
1981	152	9.9	8.3-11.7	64	10.7	8.0-13.9	88	9.4	7.5-11.7	82	5.6	4.3-7.0	34	6.5	4.3-9.3	48	5.1	3.7-6.8
1982	166	10.5	8.9-12.3	78	12.2	9.4-15.5	88	9.3	7.3-11.5	113	7.0	5.7-8.5	58	9.1	6.7-12.1	55	5.7	4.2-7.5
1983	146	8.7	7.3-10.3	65	9.5	7.1-12.3	81	8.1	6.4-10.1	102	6.5	5.2-8.0	41	7.1	4.9-9.8	61	6.2	4.7-8.1
1984	181	10.8	9.2-12.6	81	12.6	9.7-16.0	100	9.8	7.9-12.0	116	7.1	5.8-8.6	43	6.8	4.8-9.4	73	7.3	5.6-9.2
1985	172	9.7	8.2-11.3	75	11.6	8.8-14.9	97	8.7	7.0-10.7	109	6.2	5.0-7.5	54	8.2	5.9-10.9	55	5.1	3.8-6.7
1986	186	10.5	9.0-12.2	85	12.8	9.9-16.1	101	9.2	7.5-11.3	114	6.9	5.6-8.3	56	8.7	6.4-11.6	58	5.7	4.3-7.4
1987	164	8.8	7.5-10.3	79	11.5	8.9-14.6	85	7.3	5.8-9.1	99	5.5	4.5-6.8	53	8.2	5.9-10.9	46	4.1	3.0-5.5
1988	154	8.0	6.7-9.4	68	9.1	6.8-11.7	86	7.3	5.8-9.0	83	4.5	3.5-5.6	39	5.5	3.8-7.7	44	3.9	2.8-5.2
1989	191	10.4	8.9-12.0	86	12.3	9.6-15.6	105	9.3	7.6-11.3	90	5.1	4.1-6.3	39	6.5	4.4-9.1	51	4.5	3.3-5.9
1990	193	9.5	8.2-11.0	85	12.1	9.3-15.2	108	8.5	6.9-10.3	98	5.3	4.3-6.5	42	6.5	4.5-8.9	56	4.8	3.6-6.2
1991	204	10.3	8.9-11.9	73	10.0	7.6-12.8	131	10.5	8.8-12.5	110	5.7	4.6-6.9	45	6.7	4.7-9.1	65	5.3	4.1-6.8
1992	238	11.4	9.9-13.0	108	14.0	11.2-17.2	130	10.0	8.3-11.9	132	6.5	5.4-7.7	53	7.8	5.6-10.4	79	6.1	4.8-7.6
1993	188	8.8	7.5-10.2	81	9.8	7.6-12.4	107	8.1	6.6-9.8	114	5.5	4.5-6.7	47	6.5	4.6-8.8	67	5.1	3.9-6.5
1994	209	10.0	8.7-11.5	89	11.3	8.9-14.1	120	9.3	7.7-11.1	119	5.8	4.8-7.0	43	5.4	3.8-7.4	76	5.9	4.6-7.4
1995	215	9.8	8.5-11.3	96	12.2	9.7-15.1	119	8.7	7.2-10.4	138	6.4	5.4-7.6	59	7.5	5.6-9.9	79	5.9	4.6-7.3
1996	222	9.7	8.4-11.0	97	10.8	8.6-13.4	125	8.9	7.4-10.7	109	4.9	4.0-6.0	55	6.9	5.0-9.1	54	3.9	2.9-5.1
1997	249	10.8	9.4-12.2	111	12.3	10.0-15.0	138	9.7	8.2-11.5	134	6.0	5.0-7.1	70	8.7	6.7-11.2	64	4.6	3.5-5.8
1998	248	10.5	9.2-11.9	109	12.1	9.8-14.8	139	9.6	8.1-11.4	112	4.9	4.0-5.9	41	4.9	3.4-6.8	71	5.0	3.9-6.3
1999	253	10.5	9.2-11.9	101	10.3	8.2-12.6	152	10.3	8.8-12.1	121	5.2	4.3-6.2	48	5.3	3.8-7.1	73	5.0	3.9-6.3
2000	280	11.0	9.7-12.3	117	11.6	9.4-14.1	163	10.5	9.0-12.3	116	5.0	4.1-6.0	42	4.8	3.4-6.6	74	5.0	3.9-6.3
2001	300	11.7	10.4-13.1	125	12.4	10.1-14.9	175	11.2	9.6-13.0	136	5.5	4.6-6.6	54	5.9	4.3-7.8	82	5.3	4.2-6.6
2002	290	11.1	9.9-12.5	117	12.0	9.8-14.5	173	10.8	9.2-12.5	147	5.7	4.8-6.8	59	5.6	4.2-7.4	88	5.7	4.6-7.0
2003	315	11.9	10.6-13.3	134	13.3	11.0-15.9	181	11.1	9.5-12.8	124	4.6	3.8-5.5	55	5.1	3.8-6.8	69	4.3	3.3-5.4
2004	306	11.2	9.9-12.5	129	12.1	9.9-14.5	177	10.7	9.1-12.4	136	5.3	4.5-6.3	63	6.7	5.0-8.6	73	4.6	3.6-5.8
2005	367	13.1	11.8-14.6	159	14.3	12.1-16.9	208	12.3	10.6-14.0	125	4.6	3.8-5.5	51	4.7	3.4-6.3	74	4.4	3.5-5.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	359	12.6	11.3-14.0	171	15.3	13.0-17.9	188	10.9	9.4-12.6	125	4.6	3.8-5.4	60	5.6	4.2-7.3	65	3.9	3.0-4.9
2007	315	11.0	9.8-12.3	143	13.0	10.9-15.5	172	9.9	8.4-11.5	153	5.5	4.6-6.4	70	6.4	4.9-8.2	83	4.8	3.8-6.0
2008	412	13.6	12.3-15.0	190	15.9	13.6-18.4	222	12.1	10.6-13.9	118	4.2	3.5-5.0	53	4.9	3.6-6.4	65	3.7	2.9-4.8
2009	440	14.5	13.1-15.9	222	17.9	15.5-20.5	218	12.0	10.4-13.7	129	4.3	3.6-5.2	71	5.9	4.5-7.5	58	3.2	2.5-4.2
2010	440	14.2	12.9-15.6	221	18.7	16.2-21.5	219	11.6	10.1-13.3	151	5.2	4.4-6.1	69	6.3	4.8-8.0	82	4.5	3.6-5.6
2011	468	14.6	13.2-16.0	217	16.8	14.5-19.3	251	13.2	11.6-15.0	141	4.5	3.8-5.3	62	5.2	4.0-6.8	79	4.1	3.2-5.1
2012	491	14.9	13.5-16.3	229	17.7	15.3-20.2	262	13.2	11.6-14.9	144	4.4	3.7-5.2	72	5.4	4.1-6.8	72	3.7	2.9-4.7
2013	497	14.7	13.4-16.1	253	19.0	16.6-21.6	244	12.1	10.6-13.7	144	4.4	3.7-5.2	75	6.0	4.7-7.6	69	3.5	2.7-4.4
2014	514	14.8	13.6-16.2	262	18.6	16.3-21.1	252	12.4	10.9-14.1	148	4.6	3.9-5.4	68	5.6	4.3-7.1	80	4.1	3.2-5.1
2015	489	13.8	12.5-15.1	236	16.4	14.3-18.8	253	12.0	10.5-13.6	181	5.2	4.5-6.0	87	6.5	5.1-8.0	94	4.4	3.5-5.4
2016	548	14.9	13.7-16.2	263	17.9	15.7-20.3	285	13.0	11.6-14.7	170	4.9	4.2-5.7	78	5.7	4.4-7.2	92	4.5	3.6-5.5
2017	538	14.7	13.5-16.0	245	16.7	14.6-19.0	293	13.3	11.8-14.9	162	4.5	3.8-5.3	72	5.4	4.2-6.8	90	4.0	3.2-5.0
2018	543	13.9	12.8-15.2	248	15.3	13.4-17.5	295	12.9	11.5-14.5	153	4.1	3.5-4.9	73	5.3	4.1-6.7	80	3.5	2.7-4.3
2019	538	14.0	12.8-15.2	275	17.5	15.4-19.7	263	11.8	10.4-13.3	151	3.9	3.3-4.6	80	5.2	4.1-6.5	71	3.0	2.4-3.8
2020	425	11.0	9.9-12.1	195	12.5	10.7-14.4	230	10.0	8.7-11.4	126	3.2	2.7-3.8	56	3.7	2.8-4.8	70	3.0	2.4-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	132	7.7	6.3-9.4	70	9.9	7.3-13.1	62	6.3	4.7-8.2	92	5.8	4.5-7.3	53	7.8	5.4-11.0	39	4.4	3.0-6.2
1977	126	8.0	6.5-9.7	67	11.3	8.1-15.2	59	6.1	4.6-8.0	101	6.0	4.8-7.5	54	8.4	6.0-11.5	47	4.6	3.3-6.2
1978	157	9.4	7.8-11.2	76	10.9	8.2-14.2	81	8.4	6.6-10.6	115	7.3	5.9-8.9	62	9.0	6.6-11.9	53	6.0	4.4-8.0
1979	142	8.7	7.2-10.4	72	11.3	8.3-14.9	70	7.2	5.5-9.3	98	5.8	4.6-7.2	56	9.0	6.5-12.1	42	4.0	2.8-5.5
1980	158	9.0	7.5-10.7	74	10.5	7.8-13.7	84	8.1	6.4-10.2	112	6.5	5.3-8.0	49	7.6	5.2-10.6	63	6.1	4.6-7.9
1981	156	9.1	7.6-10.8	83	12.5	9.5-16.1	73	7.1	5.5-9.1	108	6.4	5.1-7.9	46	7.4	5.0-10.5	62	6.0	4.5-7.8
1982	157	8.4	7.0-10.0	79	11.1	8.4-14.3	78	6.9	5.4-8.7	90	5.0	3.9-6.2	44	6.5	4.4-9.2	46	4.2	3.0-5.6
1983	171	9.0	7.6-10.6	89	12.1	9.3-15.3	82	7.2	5.7-9.1	105	5.7	4.6-7.0	50	7.1	5.0-9.7	55	4.9	3.6-6.5
1984	164	8.2	6.9-9.7	86	9.8	7.5-12.5	78	6.9	5.4-8.8	105	5.5	4.4-6.7	54	7.1	5.0-9.6	51	4.5	3.3-6.0
1985	193	9.3	8.0-10.8	111	13.4	10.7-16.5	82	6.8	5.4-8.5	100	5.1	4.1-6.3	61	8.2	6.0-10.9	39	3.2	2.3-4.5
1986	171	8.2	7.0-9.7	96	12.0	9.4-15.1	75	6.0	4.7-7.6	110	5.2	4.2-6.3	61	6.9	5.0-9.1	49	4.0	2.9-5.4
1987	191	9.1	7.7-10.5	94	10.8	8.4-13.6	97	7.9	6.4-9.7	105	5.0	4.0-6.1	51	6.2	4.3-8.5	54	4.4	3.3-5.8
1988	155	7.6	6.4-9.0	77	10.7	8.2-13.6	78	6.1	4.8-7.7	116	6.0	4.9-7.3	55	7.5	5.4-10.0	61	5.2	3.9-6.7
1989	204	9.5	8.2-11.0	105	12.6	9.9-15.7	99	7.7	6.2-9.5	100	4.9	4.0-6.1	61	8.4	6.2-11.0	39	3.1	2.2-4.3
1990	219	9.9	8.5-11.4	107	11.7	9.3-14.6	112	8.7	7.1-10.5	128	6.2	5.1-7.4	59	7.6	5.6-10.1	69	5.4	4.1-6.8
1991	238	10.5	9.1-12.0	115	13.0	10.4-16.0	123	9.1	7.5-10.9	106	4.9	3.9-6.0	55	6.6	4.7-9.0	51	4.0	2.9-5.2
1992	231	10.0	8.6-11.4	106	11.5	9.1-14.4	125	9.1	7.5-10.9	107	5.1	4.1-6.2	48	5.4	3.8-7.4	59	4.7	3.5-6.1
1993	205	8.9	7.7-10.3	103	11.7	9.2-14.5	102	7.4	6.0-9.0	111	4.9	4.0-5.9	55	6.2	4.4-8.3	56	4.1	3.1-5.4
1994	211	9.0	7.7-10.3	109	12.0	9.6-14.7	102	7.2	5.8-8.7	156	7.0	5.9-8.3	74	8.4	6.4-10.9	82	6.1	4.8-7.6
1995	230	9.2	8.0-10.5	121	11.7	9.5-14.3	109	7.5	6.1-9.1	135	5.6	4.7-6.7	66	7.4	5.5-9.6	69	4.7	3.7-6.0
1996	251	9.9	8.7-11.3	140	13.5	11.1-16.2	111	7.5	6.2-9.1	139	5.9	4.9-7.0	66	7.0	5.2-9.1	73	5.2	4.0-6.5
1997	224	9.0	7.8-10.3	100	9.6	7.6-11.9	124	8.4	7.0-10.1	134	5.7	4.7-6.8	72	8.2	6.3-10.6	62	4.3	3.2-5.5
1998	227	8.8	7.7-10.1	103	10.5	8.3-13.0	124	8.1	6.7-9.7	123	4.9	4.1-5.9	59	6.1	4.5-8.1	64	4.3	3.3-5.5
1999	207	7.7	6.6-8.8	105	10.5	8.4-12.9	102	6.3	5.1-7.6	129	5.2	4.3-6.2	65	7.0	5.2-9.1	64	4.2	3.2-5.4
2000	258	9.7	8.5-11.0	131	12.0	9.9-14.4	127	8.1	6.7-9.6	125	4.9	4.1-5.9	53	5.6	4.0-7.5	72	4.6	3.6-5.8
2001	237	8.7	7.6-9.9	109	9.9	8.0-12.1	128	7.9	6.6-9.4	137	5.3	4.4-6.2	65	6.4	4.8-8.2	72	4.6	3.6-5.7
2002	273	9.8	8.7-11.1	147	13.1	10.9-15.5	126	7.7	6.4-9.2	124	4.7	3.9-5.6	61	5.9	4.4-7.8	63	3.9	3.0-5.0
2003	266	9.0	8.0-10.2	130	10.6	8.7-12.7	136	7.9	6.6-9.4	128	4.7	3.9-5.6	57	4.9	3.7-6.5	71	4.4	3.4-5.5
2004	286	9.8	8.7-11.1	163	13.9	11.7-16.4	123	7.2	6.0-8.6	133	5.0	4.2-5.9	70	6.9	5.3-8.9	63	3.9	3.0-5.0
2005	241	8.2	7.2-9.3	123	10.6	8.7-12.8	118	6.8	5.6-8.1	130	4.7	3.9-5.6	62	6.0	4.5-7.8	68	4.0	3.1-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Black Individuals
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	284	9.5	8.4-10.7	165	13.7	11.6-16.2	119	6.8	5.6-8.1	119	4.1	3.4-4.9	66	5.8	4.4-7.5	53	3.1	2.3-4.0
2007	280	9.3	8.2-10.5	133	10.6	8.8-12.7	147	8.3	7.0-9.8	128	4.5	3.8-5.4	68	6.3	4.8-8.1	60	3.5	2.7-4.5
2008	271	8.8	7.8-10.0	134	11.2	9.3-13.4	137	7.5	6.3-8.9	139	4.7	3.9-5.5	74	6.6	5.1-8.4	65	3.6	2.8-4.6
2009	279	8.9	7.9-10.0	152	11.9	10.0-14.1	127	7.0	5.8-8.3	117	3.8	3.1-4.6	65	5.1	3.9-6.6	52	2.9	2.1-3.7
2010	361	11.1	9.9-12.3	200	15.0	12.9-17.4	161	8.5	7.2-10.0	137	4.5	3.7-5.3	74	6.5	5.0-8.2	63	3.4	2.6-4.3
2011	348	10.6	9.5-11.8	187	13.6	11.6-15.8	161	8.4	7.2-9.9	159	5.0	4.2-5.8	85	6.5	5.1-8.1	74	3.9	3.1-4.9
2012	352	10.5	9.4-11.7	181	13.5	11.5-15.7	171	8.7	7.4-10.1	137	4.3	3.6-5.1	77	6.2	4.8-7.8	60	3.1	2.4-4.0
2013	397	11.5	10.4-12.8	192	13.4	11.5-15.6	205	10.3	8.9-11.9	141	4.3	3.6-5.0	85	6.7	5.2-8.3	56	2.7	2.1-3.6
2014	425	12.2	11.0-13.4	233	16.4	14.3-18.7	192	9.5	8.2-11.0	150	4.5	3.8-5.3	76	5.9	4.6-7.4	74	3.7	2.9-4.7
2015	367	10.5	9.4-11.7	188	13.5	11.5-15.6	179	8.6	7.4-10.0	153	4.5	3.8-5.2	75	5.4	4.2-6.9	78	3.9	3.0-4.8
2016	421	11.7	10.6-12.9	216	14.8	12.8-17.0	205	9.8	8.5-11.3	172	4.8	4.1-5.6	86	6.0	4.8-7.5	86	4.0	3.2-5.0
2017	377	10.2	9.2-11.3	180	11.5	9.9-13.4	197	9.3	8.0-10.7	171	4.8	4.1-5.6	86	6.0	4.8-7.5	85	4.0	3.2-5.0
2018	359	9.6	8.6-10.7	196	12.7	10.9-14.7	163	7.6	6.4-8.9	155	4.3	3.6-5.0	79	5.6	4.4-7.0	76	3.6	2.8-4.5
2019	369	9.7	8.7-10.8	187	11.7	10.1-13.6	182	8.3	7.1-9.6	142	3.6	3.0-4.3	73	4.6	3.5-5.8	69	2.9	2.3-3.7
2020	333	8.8	7.9-9.9	178	11.1	9.5-12.9	155	7.4	6.2-8.6	142	3.7	3.1-4.4	71	4.8	3.7-6.1	71	3.1	2.4-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	4,230	374.4	362.1-387.1	2,065	481.4	457.5-506.1	2,165	313.1	299.0-327.5	2,663	245.5	235.2-256.0	1,398	330.6	310.6-351.5	1,265	194.3	182.8-206.2
1977	4,463	386.9	374.5-399.5	2,164	508.3	483.7-533.7	2,299	320.0	306.2-334.3	2,631	231.9	222.2-241.8	1,394	326.4	306.9-346.8	1,237	176.6	166.2-187.4
1978	4,706	402.6	390.1-415.4	2,295	519.1	495.0-544.1	2,411	336.9	322.7-351.6	2,885	250.9	240.9-261.1	1,462	327.9	308.9-347.7	1,423	205.2	193.9-216.9
1979	4,949	411.0	398.6-423.6	2,432	540.8	516.6-565.7	2,517	338.3	324.4-352.6	2,959	254.3	244.4-264.5	1,579	360.1	340.1-381.0	1,380	193.3	182.6-204.5
1980	5,305	424.1	411.9-436.6	2,648	574.3	549.8-599.6	2,657	341.7	328.1-355.7	3,048	252.8	243.1-262.7	1,597	356.6	336.9-377.0	1,451	194.2	183.7-205.0
1981	5,189	401.6	390.0-413.5	2,598	540.2	516.9-564.2	2,591	325.1	312.1-338.5	3,133	249.1	239.8-258.7	1,629	347.9	328.9-367.6	1,504	193.4	183.2-204.0
1982	5,441	410.8	399.2-422.6	2,675	546.6	523.7-570.3	2,766	336.7	323.7-350.0	3,147	243.8	234.8-253.1	1,662	349.8	331.2-369.2	1,485	184.8	175.1-194.9
1983	5,496	395.6	384.5-406.8	2,668	523.3	501.3-546.0	2,828	328.0	315.6-340.8	3,348	248.1	239.2-257.1	1,755	349.1	331.1-367.8	1,593	190.8	181.2-200.8
1984	5,954	417.6	406.4-429.0	2,879	547.0	524.9-569.8	3,075	349.3	336.6-362.3	3,269	238.7	230.1-247.5	1,689	337.2	319.5-355.7	1,580	185.4	176.0-195.1
1985	6,127	419.0	408.0-430.2	3,016	560.0	537.9-582.6	3,111	343.7	331.3-356.5	3,291	235.5	227.1-244.2	1,734	342.4	324.6-360.9	1,557	178.3	169.3-187.7
1986	6,244	415.4	404.6-426.3	3,144	574.3	552.2-597.1	3,100	332.2	320.3-344.5	3,326	230.5	222.3-238.9	1,693	325.9	308.8-343.6	1,633	180.3	171.3-189.5
1987	6,350	409.9	399.4-420.6	3,143	551.3	530.1-573.0	3,207	334.6	322.8-346.7	3,322	225.0	217.1-233.2	1,705	323.1	306.2-340.6	1,617	174.2	165.5-183.1
1988	6,372	407.1	396.7-417.7	3,071	543.2	522.0-565.0	3,301	338.3	326.5-350.3	3,309	220.1	212.3-228.1	1,671	309.6	293.4-326.5	1,638	172.8	164.3-181.7
1989	6,561	410.1	399.7-420.6	3,264	564.9	543.6-586.7	3,297	330.6	319.1-342.3	3,249	211.7	204.2-219.5	1,672	307.1	291.0-323.8	1,577	162.4	154.3-170.8
1990	6,797	419.5	409.2-430.1	3,320	567.4	546.2-589.1	3,477	344.1	332.5-356.0	3,383	220.0	212.3-227.8	1,679	309.0	292.8-325.8	1,704	174.8	166.4-183.5
1991	7,266	442.9	432.4-453.7	3,514	591.6	570.1-613.6	3,752	367.1	355.2-379.3	3,313	211.0	203.6-218.6	1,677	302.4	286.5-318.8	1,636	164.8	156.7-173.1
1992	7,508	447.3	436.8-457.9	3,707	608.2	586.8-630.1	3,801	364.0	352.3-376.0	3,293	207.0	199.7-214.5	1,570	286.3	270.9-302.4	1,723	169.8	161.7-178.1
1993	7,394	432.5	422.4-442.8	3,773	605.5	584.5-627.0	3,621	340.9	329.7-352.4	3,245	200.0	192.9-207.2	1,584	281.0	265.8-296.6	1,661	161.4	153.6-169.5
1994	7,999	460.3	449.9-470.8	4,017	628.3	607.4-649.8	3,982	370.0	358.4-381.8	3,635	220.7	213.4-228.2	1,749	301.7	286.3-317.5	1,886	180.8	172.6-189.2
1995	7,808	439.5	429.6-449.7	4,009	608.4	588.2-629.1	3,799	346.1	335.0-357.4	3,654	217.6	210.4-225.0	1,739	292.2	277.4-307.5	1,915	180.2	172.1-188.6
1996	8,219	454.6	444.5-464.8	4,091	611.6	591.5-632.1	4,128	369.1	357.8-380.7	3,567	209.3	202.3-216.5	1,755	290.7	276.2-305.8	1,812	168.0	160.2-176.0
1997	8,326	456.8	446.8-466.9	4,146	617.5	597.4-638.0	4,180	369.5	358.2-380.9	3,512	203.6	196.7-210.6	1,683	277.5	263.3-292.1	1,829	166.8	159.1-174.7
1998	8,459	452.2	442.4-462.1	4,084	587.7	568.6-607.2	4,375	378.3	367.1-389.8	3,447	194.5	187.9-201.3	1,595	251.1	238.0-264.7	1,852	165.8	158.3-173.6
1999	8,617	450.8	441.1-460.6	4,359	615.1	595.8-634.8	4,258	359.5	348.7-370.5	3,552	195.6	189.1-202.3	1,640	257.5	244.2-271.2	1,912	166.2	158.8-173.9
2000	8,961	456.1	446.5-465.8	4,579	626.0	607.0-645.5	4,382	360.7	350.0-371.6	3,433	184.4	178.2-190.8	1,619	246.0	233.4-259.1	1,814	153.5	146.4-160.7
2001	8,846	441.2	431.9-450.6	4,516	602.6	584.1-621.5	4,330	350.3	339.9-360.9	3,571	188.3	182.1-194.7	1,627	243.8	231.3-256.7	1,944	161.4	154.2-168.8
2002	8,943	437.9	428.7-447.2	4,488	587.4	569.3-605.9	4,455	354.7	344.3-365.3	3,427	177.0	171.0-183.1	1,616	235.2	223.1-247.7	1,811	147.8	141.0-154.8
2003	9,195	442.3	433.1-451.5	4,609	592.0	574.0-610.3	4,586	358.6	348.2-369.2	3,314	167.1	161.4-173.0	1,525	215.0	203.6-226.8	1,789	143.0	136.4-149.8
2004	9,429	446.3	437.2-455.5	4,756	594.8	577.1-612.9	4,673	361.4	351.0-371.9	3,366	168.1	162.4-174.0	1,574	218.4	207.1-230.2	1,792	142.1	135.5-148.9
2005	9,586	446.9	437.8-456.0	4,815	590.2	572.8-608.0	4,771	365.0	354.7-375.6	3,350	164.3	158.7-170.1	1,490	204.3	193.4-215.6	1,860	145.1	138.6-151.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	9,725	449.5	440.5-458.7	4,874	592.9	575.5-610.7	4,851	367.4	357.1-378.0	3,297	160.7	155.2-166.4	1,499	205.9	195.0-217.3	1,798	139.2	132.8-145.8
2007	10,024	456.6	447.5-465.8	5,206	615.9	598.4-633.7	4,818	362.8	352.5-373.2	3,363	160.7	155.2-166.3	1,540	203.8	193.1-214.8	1,823	139.3	132.9-145.9
2008	10,188	455.8	446.8-464.9	5,106	591.6	574.7-608.9	5,082	376.4	366.1-387.0	3,307	156.0	150.7-161.5	1,492	197.1	186.7-208.0	1,815	136.2	130.0-142.7
2009	10,403	456.4	447.5-465.4	5,263	592.0	575.4-609.1	5,140	375.1	364.8-385.6	3,292	152.2	147.0-157.6	1,556	200.0	189.7-210.8	1,736	128.0	122.0-134.3
2010	10,469	449.6	440.8-458.5	5,216	571.5	555.3-588.0	5,253	377.3	367.0-387.7	3,363	152.1	146.9-157.4	1,563	197.2	187.0-207.8	1,800	130.0	124.0-136.2
2011	11,303	471.4	462.6-480.4	5,655	598.5	582.2-615.0	5,648	395.1	384.7-405.7	3,580	157.8	152.6-163.2	1,603	192.8	183.0-203.0	1,977	140.1	133.9-146.4
2012	10,915	446.1	437.6-454.7	5,294	547.3	532.0-563.0	5,621	386.2	376.1-396.6	3,528	150.8	145.8-156.0	1,585	183.8	174.4-193.5	1,943	134.1	128.1-140.2
2013	11,265	448.9	440.5-457.4	5,473	547.2	532.2-562.6	5,792	391.2	381.0-401.6	3,403	141.9	137.1-146.8	1,584	179.7	170.5-189.2	1,819	121.9	116.2-127.7
2014	11,146	436.2	428.0-444.6	5,302	514.0	499.7-528.6	5,844	389.5	379.4-399.8	3,590	146.0	141.2-151.0	1,616	176.1	167.3-185.3	1,974	130.5	124.8-136.5
2015	11,126	425.8	417.7-433.9	5,295	501.8	487.9-516.1	5,831	380.7	370.8-390.8	3,407	135.0	130.4-139.7	1,513	160.4	152.1-169.1	1,894	122.0	116.5-127.7
2016	11,588	435.2	427.2-443.4	5,624	522.4	508.3-536.8	5,964	383.1	373.2-393.1	3,556	137.1	132.6-141.8	1,626	165.1	156.9-173.7	1,930	121.9	116.4-127.6
2017	11,495	426.6	418.7-434.7	5,518	503.6	490.0-517.6	5,977	380.6	370.8-390.6	3,270	124.0	119.7-128.4	1,486	149.7	141.9-157.8	1,784	110.4	105.2-115.7
2018	11,339	414.4	406.7-422.3	5,385	481.1	467.9-494.5	5,954	374.3	364.7-384.2	3,482	129.5	125.1-133.9	1,548	152.4	144.7-160.5	1,934	117.2	111.9-122.6
2019	11,842	429.9	422.0-437.9	5,582	492.8	479.5-506.3	6,260	394.0	384.1-404.2	3,488	127.1	122.8-131.4	1,573	151.0	143.4-158.9	1,915	114.2	109.1-119.6
2020	9,818	352.4	345.3-359.6	4,637	403.9	392.0-416.1	5,181	322.4	313.4-331.6	3,090	110.1	106.2-114.2	1,360	127.2	120.3-134.4	1,730	101.4	96.6-106.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	194	15.2	13.0-17.7	132	26.2	21.3-31.8	62	8.2	6.2-10.6	92	7.3	5.8-9.1	62	12.4	9.3-16.1	30	4.0	2.6-5.8
1977	187	14.3	12.2-16.6	142	27.3	22.6-32.7	45	5.6	4.1-7.6	103	8.1	6.5-9.9	83	15.5	12.2-19.4	20	2.9	1.7-4.7
1978	205	14.7	12.7-17.0	147	26.0	21.7-31.0	58	7.1	5.4-9.3	107	8.0	6.5-9.8	80	15.9	12.1-20.4	27	3.2	2.1-4.7
1979	214	16.0	13.8-18.4	157	29.2	24.5-34.5	57	7.2	5.4-9.4	109	8.1	6.5-9.8	78	14.6	11.2-18.7	31	3.8	2.6-5.5
1980	213	15.2	13.2-17.5	154	27.8	23.3-33.0	59	7.0	5.3-9.1	103	7.5	6.0-9.1	85	15.8	12.2-20.0	18	2.1	1.2-3.3
1981	224	15.2	13.2-17.4	167	27.4	23.2-32.1	57	6.8	5.1-8.9	88	6.1	4.8-7.6	63	11.1	8.4-14.5	25	2.9	1.8-4.3
1982	227	15.2	13.2-17.4	156	26.6	22.3-31.5	71	7.9	6.1-10.0	117	8.1	6.7-9.8	89	15.9	12.5-19.9	28	3.1	2.1-4.6
1983	211	14.2	12.3-16.3	153	25.8	21.6-30.6	58	6.6	5.0-8.6	110	7.7	6.2-9.3	86	15.4	12.1-19.4	24	2.8	1.8-4.3
1984	258	16.2	14.3-18.4	170	26.2	22.3-30.6	88	9.4	7.5-11.6	91	6.0	4.8-7.4	68	11.6	8.8-15.0	23	2.4	1.5-3.7
1985	247	15.5	13.6-17.6	165	25.8	21.8-30.4	82	8.7	6.8-10.8	82	5.4	4.3-6.8	57	9.7	7.2-12.7	25	2.8	1.8-4.1
1986	252	15.2	13.3-17.3	181	27.2	23.2-31.7	71	7.3	5.6-9.2	100	6.6	5.3-8.1	72	12.3	9.4-15.8	28	3.1	2.0-4.6
1987	270	15.9	14.0-18.0	190	28.1	24.1-32.6	80	7.9	6.2-9.8	76	4.6	3.6-5.8	57	8.9	6.6-11.8	19	1.9	1.1-2.9
1988	231	13.4	11.7-15.3	150	22.2	18.5-26.3	81	7.8	6.1-9.7	96	5.6	4.5-6.9	71	10.4	8.0-13.2	25	2.5	1.6-3.7
1989	223	12.5	10.9-14.3	148	21.1	17.7-25.0	75	6.9	5.4-8.7	91	5.2	4.2-6.5	65	9.7	7.3-12.6	26	2.5	1.6-3.7
1990	239	13.9	12.1-15.8	161	24.1	20.2-28.4	78	7.5	5.9-9.4	74	4.6	3.6-5.8	52	9.0	6.5-12.2	22	2.1	1.3-3.2
1991	308	17.6	15.7-19.8	212	32.1	27.6-37.0	96	8.9	7.2-10.9	107	6.3	5.1-7.6	77	11.1	8.7-14.0	30	3.1	2.1-4.4
1992	282	15.7	13.9-17.7	185	26.5	22.6-30.8	97	9.0	7.3-11.0	80	4.5	3.6-5.6	53	7.8	5.7-10.4	27	2.5	1.6-3.6
1993	266	14.6	12.8-16.5	172	24.0	20.4-28.0	94	8.5	6.9-10.4	83	4.7	3.7-5.8	60	8.6	6.5-11.3	23	2.2	1.4-3.3
1994	259	14.1	12.4-16.0	176	24.6	20.9-28.7	83	7.5	5.9-9.3	91	5.1	4.1-6.2	66	9.9	7.4-12.8	25	2.3	1.5-3.3
1995	220	11.6	10.1-13.2	163	21.5	18.2-25.2	57	5.0	3.8-6.5	99	5.5	4.4-6.7	71	10.0	7.7-12.8	28	2.6	1.7-3.7
1996	239	12.4	10.8-14.1	152	19.9	16.8-23.5	87	7.5	6.0-9.2	81	4.4	3.5-5.5	59	8.3	6.2-10.8	22	2.0	1.3-3.1
1997	222	11.5	10.0-13.1	149	19.0	16.0-22.4	73	6.4	5.0-8.1	65	3.4	2.6-4.3	49	6.4	4.6-8.5	16	1.4	0.8-2.3
1998	211	10.6	9.2-12.1	139	17.3	14.4-20.5	72	6.1	4.8-7.7	70	3.7	2.9-4.7	45	6.2	4.4-8.4	25	2.2	1.4-3.2
1999	187	9.2	7.9-10.7	119	14.5	11.9-17.4	68	5.7	4.4-7.2	59	3.1	2.3-4.0	35	4.7	3.2-6.6	24	2.1	1.3-3.1
2000	209	10.3	8.9-11.8	141	17.8	14.8-21.1	68	5.6	4.3-7.1	63	3.2	2.5-4.1	45	6.0	4.3-8.2	18	1.5	0.9-2.4
2001	190	8.9	7.7-10.3	135	16.2	13.5-19.3	55	4.3	3.2-5.6	70	3.4	2.6-4.3	51	6.3	4.6-8.3	19	1.5	0.9-2.4
2002	200	9.4	8.1-10.8	125	14.8	12.2-17.8	75	5.9	4.6-7.4	50	2.3	1.7-3.1	34	3.9	2.6-5.5	16	1.3	0.7-2.1
2003	188	8.6	7.4-9.9	130	14.5	12.1-17.3	58	4.6	3.5-5.9	50	2.4	1.8-3.1	37	4.6	3.2-6.4	13	1.0	0.6-1.8
2004	222	10.0	8.7-11.4	140	15.9	13.3-18.9	82	6.2	4.9-7.7	71	3.3	2.6-4.2	53	6.3	4.6-8.4	18	1.4	0.9-2.3
2005	199	8.9	7.7-10.2	139	16.0	13.3-19.0	60	4.5	3.5-5.9	63	2.9	2.2-3.7	44	5.4	3.8-7.3	19	1.4	0.8-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	191	8.4	7.2-9.7	121	13.3	11.0-16.0	70	5.1	4.0-6.5	42	1.8	1.3-2.5	33	3.7	2.5-5.2	9	0.7	0.3-1.3
2007	223	9.8	8.6-11.2	142	15.6	13.0-18.5	81	6.1	4.9-7.7	59	2.6	2.0-3.4	46	5.3	3.8-7.2	13	1.0	0.5-1.6
2008	192	8.2	7.1-9.5	128	13.5	11.2-16.1	64	4.6	3.5-5.9	57	2.6	1.9-3.4	42	4.9	3.5-6.7	15	1.1	0.6-1.9
2009	191	8.1	6.9-9.3	124	12.8	10.6-15.3	67	4.8	3.7-6.2	54	2.3	1.7-3.0	37	4.0	2.7-5.5	17	1.2	0.7-1.9
2010	205	8.6	7.4-9.8	130	13.5	11.2-16.1	75	5.3	4.1-6.7	50	2.1	1.6-2.8	26	2.9	1.9-4.4	24	1.7	1.1-2.5
2011	209	8.4	7.3-9.6	142	13.9	11.6-16.4	67	4.6	3.6-5.9	72	2.9	2.3-3.7	50	5.1	3.7-6.8	22	1.5	0.9-2.3
2012	211	8.4	7.3-9.6	135	13.1	10.9-15.6	76	5.1	4.0-6.4	52	2.2	1.6-2.9	32	3.3	2.2-4.8	20	1.4	0.9-2.2
2013	208	8.2	7.1-9.4	140	13.3	11.1-15.8	68	4.6	3.6-5.8	66	2.7	2.1-3.5	45	4.7	3.4-6.4	21	1.4	0.9-2.2
2014	196	7.3	6.3-8.5	118	10.6	8.7-12.7	78	5.1	4.0-6.4	46	1.8	1.3-2.4	28	2.8	1.8-4.2	18	1.2	0.7-1.9
2015	196	7.4	6.4-8.5	118	11.0	9.0-13.2	78	5.1	4.0-6.3	50	1.9	1.4-2.5	35	3.2	2.2-4.5	15	1.0	0.6-1.7
2016	215	8.1	7.0-9.3	143	13.3	11.1-15.7	72	4.6	3.6-5.9	39	1.4	1.0-2.0	24	2.3	1.5-3.5	15	0.9	0.5-1.5
2017	231	8.6	7.5-9.9	134	12.3	10.3-14.7	97	6.2	5.0-7.6	48	1.8	1.3-2.4	30	2.7	1.8-3.9	18	1.1	0.7-1.8
2018	214	7.8	6.8-8.9	132	12.2	10.1-14.5	82	5.0	3.9-6.2	53	2.0	1.5-2.7	36	3.5	2.4-4.9	17	1.1	0.7-1.9
2019	185	6.6	5.7-7.7	111	9.6	7.9-11.7	74	4.5	3.5-5.7	52	1.9	1.4-2.5	38	3.4	2.4-4.7	14	0.8	0.4-1.4
2020	190	6.8	5.9-7.9	115	10.1	8.3-12.2	75	4.7	3.7-6.0	47	1.7	1.2-2.2	28	2.6	1.7-3.8	19	1.0	0.6-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	161	12.5	10.6-14.8	116	22.6	18.3-27.6	45	5.9	4.2-8.0	156	12.7	10.7-15.1	118	24.4	19.7-29.8	38	5.2	3.6-7.3
1977	169	13.4	11.3-15.7	118	24.4	19.7-29.9	51	6.6	4.9-8.7	148	11.6	9.7-13.8	107	22.1	17.4-27.4	41	5.2	3.7-7.1
1978	181	14.1	12.0-16.5	128	26.2	21.3-31.7	53	6.6	4.9-8.7	125	9.5	7.8-11.4	83	15.9	12.5-20.0	42	5.3	3.8-7.2
1979	188	13.7	11.7-16.0	127	23.7	19.3-28.6	61	7.5	5.6-9.7	173	13.1	11.1-15.3	125	24.7	20.0-30.1	48	5.9	4.3-7.9
1980	201	14.8	12.7-17.1	150	28.9	24.0-34.5	51	6.1	4.5-8.1	166	12.4	10.4-14.5	108	21.3	17.0-26.2	58	6.9	5.2-9.1
1981	207	14.2	12.3-16.4	144	25.1	20.9-29.8	63	7.2	5.5-9.2	167	11.4	9.7-13.3	123	21.0	17.3-25.3	44	5.1	3.6-6.8
1982	198	13.6	11.7-15.7	143	25.8	21.4-30.7	55	6.0	4.5-7.9	167	11.7	10.0-13.7	117	22.0	17.9-26.8	50	5.6	4.1-7.4
1983	195	12.8	11.0-14.9	134	22.1	18.3-26.5	61	6.9	5.2-8.9	162	10.8	9.1-12.6	113	18.8	15.3-22.8	49	5.6	4.1-7.5
1984	209	13.7	11.8-15.7	139	23.6	19.5-28.3	70	7.5	5.8-9.5	160	10.6	9.0-12.5	112	19.3	15.6-23.6	48	5.3	3.9-7.1
1985	188	12.1	10.4-14.0	126	20.7	17.0-24.9	62	6.7	5.1-8.6	172	11.1	9.4-13.0	126	21.1	17.3-25.4	46	4.9	3.5-6.6
1986	188	12.0	10.3-13.9	141	23.7	19.7-28.3	47	4.9	3.6-6.6	165	10.5	8.9-12.3	114	18.7	15.2-22.7	51	5.5	4.0-7.2
1987	181	11.4	9.8-13.3	115	19.3	15.7-23.4	66	6.7	5.1-8.5	138	8.7	7.3-10.3	99	16.4	13.1-20.2	39	4.1	2.9-5.6
1988	202	12.6	10.9-14.6	144	24.1	20.0-28.7	58	5.8	4.4-7.6	148	9.4	7.9-11.1	112	19.0	15.4-23.2	36	3.7	2.5-5.1
1989	192	11.6	10.0-13.4	132	21.0	17.3-25.2	60	5.9	4.5-7.7	142	8.8	7.4-10.4	99	16.0	12.8-19.7	43	4.5	3.2-6.0
1990	173	10.7	9.1-12.4	119	19.3	15.8-23.4	54	5.6	4.2-7.3	135	8.2	6.8-9.7	86	14.0	11.0-17.5	49	4.8	3.5-6.4
1991	154	9.1	7.7-10.7	102	16.5	13.2-20.3	52	4.9	3.6-6.4	128	7.7	6.4-9.1	85	13.7	10.7-17.2	43	4.1	3.0-5.6
1992	171	10.2	8.7-11.8	111	17.3	14.0-21.1	60	5.8	4.4-7.5	110	6.4	5.2-7.7	73	11.0	8.5-14.1	37	3.6	2.5-4.9
1993	183	10.5	9.0-12.1	121	17.7	14.5-21.4	62	5.9	4.5-7.6	138	8.1	6.8-9.6	95	15.2	12.0-18.9	43	4.1	2.9-5.5
1994	176	10.2	8.7-11.9	115	18.2	14.7-22.2	61	5.8	4.4-7.4	133	7.7	6.4-9.2	88	13.7	10.8-17.1	45	4.3	3.1-5.7
1995	142	8.1	6.8-9.6	94	13.8	11.0-17.1	48	4.6	3.4-6.1	124	7.2	6.0-8.7	84	13.0	10.2-16.3	40	3.9	2.8-5.3
1996	138	8.0	6.7-9.5	78	11.5	9.0-14.5	60	5.7	4.3-7.3	123	7.1	5.8-8.4	80	12.1	9.4-15.2	43	4.0	2.9-5.4
1997	144	8.0	6.7-9.4	100	14.0	11.3-17.2	44	4.1	3.0-5.5	112	6.5	5.3-7.8	75	11.4	8.8-14.4	37	3.5	2.5-4.8
1998	128	6.8	5.7-8.1	88	12.2	9.7-15.1	40	3.5	2.5-4.8	95	5.2	4.2-6.3	59	8.4	6.3-11.0	36	3.3	2.3-4.5
1999	130	6.7	5.6-8.0	88	12.2	9.7-15.2	42	3.5	2.5-4.7	100	5.1	4.2-6.3	70	9.4	7.2-11.9	30	2.5	1.7-3.6
2000	134	7.2	6.0-8.5	67	9.6	7.3-12.3	67	5.8	4.5-7.4	112	5.9	4.8-7.1	65	9.0	6.8-11.5	47	4.0	2.9-5.3
2001	121	6.0	4.9-7.1	83	10.6	8.4-13.3	38	3.1	2.2-4.2	91	4.6	3.7-5.7	67	9.2	7.0-11.8	24	2.0	1.3-3.0
2002	117	5.8	4.8-7.0	69	9.1	7.0-11.6	48	3.9	2.9-5.2	81	4.2	3.3-5.2	47	6.7	4.8-8.9	34	2.8	1.9-3.9
2003	129	6.4	5.3-7.6	81	10.9	8.5-13.7	48	3.8	2.8-5.1	91	4.3	3.5-5.3	54	6.4	4.8-8.4	37	2.9	2.1-4.0
2004	124	5.9	4.9-7.0	78	9.3	7.3-11.8	46	3.7	2.7-4.9	96	4.7	3.8-5.7	54	6.6	4.9-8.7	42	3.4	2.4-4.5
2005	107	5.0	4.0-6.0	56	6.5	4.9-8.5	51	3.9	2.9-5.1	72	3.4	2.7-4.3	43	5.1	3.7-7.0	29	2.3	1.5-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	92	4.5	3.6-5.5	64	8.2	6.2-10.6	28	2.3	1.5-3.3	89	4.3	3.4-5.3	52	6.9	5.1-9.2	37	2.8	2.0-3.9
2007	91	4.1	3.3-5.1	64	7.5	5.7-9.7	27	2.0	1.3-3.0	65	2.9	2.2-3.7	44	4.9	3.5-6.6	21	1.6	1.0-2.4
2008	111	5.0	4.1-6.1	75	9.2	7.2-11.7	36	2.6	1.8-3.6	84	3.9	3.1-4.8	52	6.5	4.8-8.6	32	2.4	1.6-3.4
2009	105	4.7	3.8-5.7	68	8.0	6.1-10.2	37	2.7	1.9-3.8	74	3.3	2.5-4.1	58	6.6	4.9-8.7	16	1.2	0.7-1.9
2010	85	3.7	2.9-4.5	49	5.4	3.9-7.2	36	2.6	1.8-3.6	67	3.0	2.3-3.9	42	5.4	3.8-7.3	25	1.8	1.1-2.7
2011	108	4.6	3.8-5.6	73	8.1	6.3-10.3	35	2.5	1.7-3.5	66	2.8	2.1-3.6	45	4.8	3.4-6.4	21	1.5	0.9-2.3
2012	86	3.6	2.8-4.4	52	5.9	4.4-7.9	34	2.2	1.5-3.1	56	2.2	1.7-2.9	37	3.7	2.6-5.2	19	1.2	0.7-1.9
2013	80	3.3	2.6-4.2	55	6.2	4.6-8.1	25	1.7	1.1-2.5	49	2.1	1.5-2.7	29	3.1	2.0-4.5	20	1.4	0.9-2.2
2014	92	3.7	3.0-4.5	68	7.1	5.4-9.1	24	1.6	1.0-2.4	61	2.5	1.9-3.3	39	4.4	3.1-6.1	22	1.5	0.9-2.3
2015	86	3.3	2.6-4.1	51	5.2	3.8-6.9	35	2.2	1.5-3.1	49	2.0	1.5-2.6	29	3.1	2.0-4.5	20	1.3	0.8-2.0
2016	106	3.9	3.2-4.7	70	6.7	5.2-8.5	36	2.2	1.5-3.1	54	2.0	1.5-2.6	34	3.3	2.3-4.7	20	1.2	0.7-1.9
2017	83	3.1	2.4-3.8	53	4.7	3.5-6.2	30	1.9	1.3-2.7	58	2.2	1.6-2.8	37	3.7	2.6-5.2	21	1.3	0.8-2.0
2018	77	2.8	2.2-3.6	47	4.5	3.2-6.0	30	1.8	1.2-2.6	47	1.7	1.3-2.3	33	3.1	2.1-4.4	14	0.9	0.5-1.5
2019	81	2.8	2.2-3.5	56	4.9	3.6-6.4	25	1.5	1.0-2.3	40	1.4	1.0-1.9	26	2.4	1.6-3.6	14	0.7	0.4-1.3
2020	78	2.7	2.2-3.4	47	4.2	3.0-5.6	31	1.9	1.3-2.7	49	1.6	1.2-2.1	27	2.2	1.4-3.2	22	1.2	0.8-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	178	17.4	14.7-20.5	100	24.4	19.1-30.5	78	13.1	10.2-16.6	148	14.3	11.9-17.0	88	22.6	17.4-28.6	60	9.5	7.1-12.3
1977	213	20.3	17.4-23.4	115	27.7	22.1-34.0	98	15.6	12.4-19.2	161	15.8	13.2-18.7	93	23.3	18.1-29.4	68	11.2	8.5-14.5
1978	176	15.9	13.4-18.7	91	18.8	14.8-23.5	85	13.7	10.7-17.1	159	14.9	12.5-17.7	81	18.6	14.2-23.7	78	12.6	9.8-15.9
1979	186	17.5	14.8-20.4	112	26.6	21.2-32.7	74	12.0	9.2-15.2	138	12.6	10.4-15.0	76	17.1	13.0-21.9	62	9.7	7.3-12.6
1980	238	20.8	18.0-23.9	132	29.5	24.0-35.6	106	15.6	12.6-19.0	161	14.9	12.5-17.6	92	22.3	17.5-28.0	69	10.5	8.1-13.5
1981	222	18.6	16.0-21.5	134	28.7	23.4-34.7	88	12.6	10.0-15.7	178	15.1	12.8-17.7	111	24.6	19.7-30.3	67	9.6	7.3-12.3
1982	231	19.3	16.7-22.1	130	29.9	24.3-36.1	101	13.5	10.9-16.5	167	15.1	12.7-17.7	106	27.8	22.1-34.3	61	8.4	6.3-10.9
1983	222	17.3	15.0-19.9	112	22.9	18.4-27.9	110	14.0	11.4-17.0	173	13.5	11.5-15.8	96	19.8	15.7-24.6	77	9.8	7.7-12.4
1984	201	15.3	13.2-17.8	113	24.5	19.7-30.0	88	10.5	8.3-13.0	134	10.3	8.6-12.3	73	16.4	12.4-21.0	61	7.3	5.5-9.4
1985	225	17.0	14.8-19.5	115	22.4	18.1-27.3	110	13.7	11.2-16.6	138	10.4	8.6-12.3	72	15.0	11.3-19.3	66	7.8	6.0-10.0
1986	258	18.5	16.2-21.0	153	30.9	25.6-36.8	105	11.9	9.7-14.5	165	12.4	10.5-14.6	96	19.5	15.3-24.3	69	8.5	6.6-10.8
1987	243	17.4	15.2-19.8	139	24.9	20.6-29.8	104	12.6	10.2-15.3	154	11.5	9.7-13.6	80	15.8	12.1-20.2	74	9.1	7.1-11.4
1988	235	16.3	14.2-18.7	132	24.4	20.0-29.3	103	11.8	9.5-14.3	169	11.9	10.1-14.0	98	18.4	14.6-22.8	71	8.2	6.4-10.4
1989	236	16.7	14.5-19.0	119	21.3	17.3-25.9	117	13.6	11.2-16.4	159	11.2	9.4-13.1	82	15.3	11.9-19.3	77	8.7	6.8-10.9
1990	248	16.6	14.5-18.9	128	23.1	18.9-27.8	120	13.0	10.7-15.5	163	11.6	9.8-13.7	82	15.4	11.9-19.5	81	9.3	7.4-11.6
1991	227	14.8	12.9-17.0	118	20.3	16.5-24.6	109	11.6	9.5-14.0	151	10.0	8.4-11.8	86	15.1	11.8-19.0	65	7.1	5.4-9.1
1992	253	16.4	14.3-18.6	127	21.0	17.2-25.4	126	13.3	11.0-15.9	168	11.5	9.8-13.5	78	13.9	10.7-17.7	90	9.9	7.9-12.1
1993	248	15.6	13.7-17.7	122	20.8	17.0-25.2	126	12.7	10.5-15.1	149	9.5	8.0-11.2	69	12.1	9.2-15.5	80	8.1	6.4-10.1
1994	287	18.3	16.2-20.6	143	24.2	20.1-28.9	144	14.7	12.4-17.3	189	12.2	10.5-14.1	102	17.6	14.1-21.7	87	9.0	7.2-11.1
1995	241	14.9	13.0-17.0	125	20.3	16.7-24.5	116	11.5	9.5-13.9	160	10.0	8.5-11.8	71	11.3	8.7-14.4	89	8.9	7.2-11.0
1996	260	15.6	13.7-17.6	143	23.1	19.2-27.4	117	11.2	9.2-13.4	180	11.2	9.6-12.9	96	16.5	13.1-20.4	84	8.2	6.5-10.2
1997	258	15.7	13.8-17.8	139	24.1	19.9-28.7	119	11.4	9.4-13.6	157	9.2	7.8-10.7	86	13.9	10.9-17.3	71	6.6	5.2-8.4
1998	253	14.6	12.8-16.5	128	20.2	16.6-24.2	125	11.5	9.6-13.7	149	8.8	7.4-10.3	76	12.3	9.5-15.5	73	6.8	5.3-8.5
1999	238	13.4	11.7-15.2	121	18.8	15.4-22.7	117	10.4	8.6-12.4	145	8.3	7.0-9.8	72	11.8	9.1-15.1	73	6.5	5.1-8.2
2000	262	14.6	12.8-16.5	133	20.9	17.3-25.0	129	11.2	9.4-13.3	148	8.1	6.8-9.6	85	13.2	10.4-16.6	63	5.4	4.2-6.9
2001	245	13.5	11.9-15.3	117	17.4	14.2-21.0	128	11.2	9.3-13.3	131	7.4	6.2-8.8	69	10.6	8.1-13.5	62	5.5	4.2-7.0
2002	246	12.8	11.2-14.6	117	17.0	13.8-20.6	129	10.6	8.8-12.6	138	7.3	6.1-8.7	70	11.0	8.4-14.0	68	5.6	4.3-7.1
2003	256	13.2	11.6-15.0	123	17.2	14.1-20.7	133	10.9	9.1-13.0	136	7.0	5.9-8.4	65	8.6	6.6-11.1	71	5.9	4.6-7.4
2004	244	12.5	11.0-14.2	120	17.3	14.2-20.8	124	10.0	8.3-11.9	135	7.0	5.9-8.3	64	9.3	7.0-11.9	71	5.7	4.5-7.2
2005	308	15.7	14.0-17.6	166	23.5	19.9-27.6	142	11.4	9.6-13.4	148	7.6	6.4-8.9	83	12.5	9.7-15.6	65	5.2	4.0-6.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	268	13.2	11.7-14.9	142	19.0	15.9-22.6	126	9.8	8.2-11.7	143	7.4	6.2-8.7	79	11.6	9.1-14.6	64	5.1	3.9-6.5
2007	239	11.5	10.1-13.1	121	16.1	13.2-19.4	118	9.2	7.6-11.0	127	6.3	5.2-7.5	57	7.1	5.3-9.3	70	5.5	4.3-7.0
2008	228	11.1	9.7-12.6	112	15.4	12.6-18.7	116	8.9	7.3-10.6	123	5.9	4.9-7.1	60	8.0	6.0-10.5	63	4.9	3.7-6.2
2009	268	12.7	11.2-14.3	151	18.6	15.6-22.0	117	9.0	7.4-10.8	116	5.7	4.7-6.8	62	8.6	6.5-11.1	54	4.1	3.1-5.4
2010	263	12.1	10.7-13.7	133	16.2	13.4-19.3	130	9.5	8.0-11.3	113	5.4	4.4-6.5	60	8.1	6.1-10.5	53	3.9	2.9-5.2
2011	265	11.7	10.3-13.2	127	15.1	12.5-18.1	138	9.7	8.2-11.5	128	5.9	4.9-7.0	73	9.3	7.2-11.7	55	3.9	2.9-5.1
2012	282	12.0	10.6-13.5	154	17.1	14.4-20.2	128	8.9	7.4-10.6	115	5.1	4.2-6.1	56	6.3	4.7-8.2	59	4.2	3.2-5.5
2013	280	11.7	10.4-13.2	152	16.8	14.1-19.8	128	8.7	7.3-10.4	120	5.1	4.2-6.2	60	6.9	5.2-9.0	60	4.1	3.1-5.3
2014	268	10.9	9.6-12.3	138	14.7	12.3-17.5	130	8.6	7.2-10.2	134	5.7	4.7-6.7	73	7.8	6.1-9.9	61	4.3	3.3-5.6
2015	281	11.0	9.8-12.4	145	14.6	12.2-17.3	136	8.8	7.4-10.5	96	3.9	3.1-4.7	52	5.6	4.2-7.5	44	2.7	2.0-3.7
2016	286	11.1	9.9-12.5	159	15.9	13.5-18.7	127	8.3	6.9-9.9	151	5.9	5.0-6.9	93	9.3	7.5-11.5	58	3.7	2.8-4.8
2017	278	10.8	9.6-12.2	145	15.0	12.5-17.7	133	8.5	7.1-10.1	84	3.4	2.7-4.2	42	4.4	3.2-6.0	42	2.7	1.9-3.7
2018	244	9.2	8.1-10.5	130	12.7	10.5-15.1	114	7.2	5.9-8.7	101	3.8	3.1-4.6	59	5.8	4.4-7.5	42	2.5	1.8-3.5
2019	293	10.9	9.6-12.2	149	13.8	11.6-16.3	144	9.1	7.6-10.7	130	4.9	4.1-5.8	80	8.0	6.3-10.1	50	3.0	2.2-4.0
2020	234	8.5	7.4-9.6	125	11.4	9.4-13.7	109	6.4	5.3-7.8	94	3.5	2.8-4.3	52	4.8	3.6-6.4	42	2.5	1.8-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	481	46.5	42.0-51.2	209	52.5	44.6-61.1	272	42.8	37.4-48.6	302	31.5	27.6-35.6	133	37.1	30.2-45.1	169	28.3	23.9-33.3
1977	591	59.2	54.0-64.7	239	64.5	55.4-74.5	352	56.1	49.9-62.9	279	28.4	24.8-32.2	123	34.7	28.1-42.1	156	24.8	20.8-29.3
1978	609	60.3	55.1-65.8	248	65.4	56.3-75.4	361	57.6	51.3-64.3	364	37.3	33.2-41.8	148	39.8	32.8-47.7	216	35.8	30.8-41.3
1979	640	59.8	54.9-65.1	266	67.6	58.6-77.3	374	55.8	49.9-62.1	355	33.9	30.1-37.9	161	42.8	35.6-50.8	194	29.2	25.0-33.9
1980	668	58.1	53.4-63.0	294	67.3	58.8-76.5	374	52.9	47.4-58.9	356	32.8	29.2-36.7	146	36.7	30.2-44.0	210	30.9	26.6-35.6
1981	659	56.0	51.5-60.8	278	64.3	56.0-73.4	381	51.7	46.4-57.4	370	33.7	30.1-37.6	140	35.8	29.4-43.1	230	32.9	28.6-37.7
1982	680	55.3	50.9-59.9	312	67.2	59.2-76.0	368	48.3	43.3-53.7	355	29.4	26.2-32.9	160	34.3	28.6-40.7	195	26.4	22.6-30.5
1983	671	51.6	47.5-55.9	297	64.0	56.1-72.7	374	45.6	40.9-50.6	344	26.8	23.9-30.0	157	31.6	26.3-37.5	187	23.9	20.4-27.7
1984	700	53.4	49.3-57.7	317	62.7	55.3-70.8	383	47.7	42.9-52.9	380	30.0	26.9-33.4	168	36.9	30.9-43.6	212	26.6	23.0-30.6
1985	771	57.0	52.8-61.4	346	71.0	62.9-79.8	425	49.8	45.0-54.9	376	29.4	26.3-32.7	152	33.4	27.7-39.8	224	27.5	23.9-31.5
1986	803	57.1	53.0-61.4	357	69.8	62.0-78.2	446	50.7	45.9-55.7	355	26.2	23.4-29.2	159	32.4	27.1-38.4	196	22.9	19.7-26.4
1987	715	49.7	46.0-53.7	309	58.6	51.5-66.2	406	45.0	40.6-49.7	354	25.6	22.9-28.6	151	29.8	24.7-35.6	203	23.3	20.1-26.9
1988	739	51.2	47.4-55.2	314	59.7	52.6-67.4	425	46.6	42.2-51.4	351	25.1	22.4-28.0	157	31.4	26.1-37.2	194	21.8	18.8-25.2
1989	760	51.4	47.6-55.3	361	67.9	60.4-76.0	399	42.7	38.5-47.2	342	23.2	20.7-25.9	163	30.9	25.9-36.6	179	19.1	16.3-22.2
1990	770	52.1	48.4-56.1	324	61.1	54.1-68.8	446	47.3	42.9-52.0	333	23.6	21.1-26.4	148	30.3	25.1-36.2	185	20.4	17.5-23.6
1991	839	56.4	52.5-60.5	349	64.2	57.0-71.9	490	51.9	47.4-56.8	369	25.9	23.2-28.8	152	31.5	26.2-37.5	217	23.4	20.3-26.7
1992	796	51.5	47.8-55.3	313	56.5	49.8-63.8	483	49.3	44.9-53.9	361	23.9	21.4-26.6	151	29.8	24.8-35.4	210	21.4	18.6-24.6
1993	821	52.4	48.7-56.2	368	64.5	57.5-72.1	453	45.8	41.6-50.2	345	22.6	20.2-25.2	141	25.7	21.2-30.7	204	21.0	18.2-24.1
1994	871	54.6	50.9-58.4	367	63.6	56.7-71.0	504	49.9	45.6-54.5	381	24.6	22.1-27.3	176	30.9	26.2-36.2	205	20.9	18.1-24.0
1995	852	51.4	47.9-55.0	378	62.0	55.4-69.1	474	45.4	41.4-49.7	409	25.8	23.3-28.5	170	28.8	24.3-33.7	239	23.8	20.8-27.0
1996	921	54.9	51.3-58.6	392	63.9	57.2-71.1	529	50.1	45.9-54.6	421	26.2	23.7-28.8	203	36.0	30.9-41.7	218	21.1	18.3-24.1
1997	971	57.4	53.8-61.2	432	70.6	63.6-78.2	539	50.3	46.1-54.8	422	25.7	23.2-28.3	193	34.1	29.0-39.6	229	21.6	18.9-24.6
1998	950	54.7	51.2-58.4	388	60.7	54.3-67.4	562	51.3	47.1-55.7	384	22.7	20.4-25.1	161	26.4	22.2-31.2	223	20.7	18.1-23.6
1999	1,035	57.0	53.5-60.7	414	62.0	55.8-68.6	621	54.2	50.0-58.7	444	25.2	22.9-27.7	182	29.3	24.9-34.1	262	23.2	20.5-26.2
2000	932	49.6	46.4-52.9	401	59.3	53.3-65.7	531	44.7	41.0-48.7	388	21.5	19.4-23.8	165	25.1	21.1-29.5	223	19.4	16.9-22.1
2001	997	52.2	48.9-55.6	413	59.6	53.6-66.0	584	48.6	44.7-52.7	417	22.6	20.4-24.9	175	27.7	23.5-32.4	242	20.4	17.9-23.1
2002	977	49.8	46.6-53.0	414	55.8	50.3-61.8	563	46.1	42.3-50.1	388	20.8	18.7-23.0	179	27.5	23.3-32.1	209	17.4	15.1-19.9
2003	1,061	53.2	50.0-56.6	463	63.7	57.7-70.1	598	47.8	44.0-51.8	409	21.4	19.4-23.6	191	28.6	24.5-33.2	218	17.8	15.5-20.3
2004	1,063	52.1	48.9-55.4	432	57.2	51.7-63.2	631	49.6	45.7-53.6	381	19.6	17.7-21.7	179	25.6	21.8-29.8	202	16.3	14.1-18.7
2005	1,020	49.3	46.2-52.4	451	57.9	52.4-63.8	569	44.4	40.8-48.2	358	18.2	16.4-20.2	153	21.7	18.2-25.6	205	16.3	14.1-18.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,082	51.9	48.8-55.1	441	57.3	51.7-63.2	641	49.3	45.6-53.4	381	19.0	17.1-21.0	159	22.7	19.1-26.6	222	17.2	15.0-19.7
2007	1,069	50.1	47.1-53.3	500	60.8	55.3-66.7	569	43.3	39.8-47.1	370	17.9	16.1-19.9	162	21.2	17.9-24.9	208	16.1	14.0-18.5
2008	1,059	49.1	46.1-52.2	475	57.6	52.3-63.3	584	43.9	40.4-47.6	367	17.7	15.9-19.6	170	22.1	18.7-25.9	197	15.1	13.0-17.3
2009	1,058	47.6	44.7-50.6	498	57.5	52.3-63.0	560	41.1	37.8-44.7	362	17.3	15.5-19.2	170	22.3	18.9-26.1	192	14.5	12.5-16.7
2010	1,020	45.4	42.6-48.3	460	54.6	49.4-60.1	560	40.7	37.4-44.3	355	16.2	14.5-18.0	183	23.0	19.6-26.7	172	12.3	10.5-14.3
2011	1,044	44.7	42.0-47.5	493	54.5	49.6-59.7	551	38.4	35.2-41.8	357	16.1	14.4-17.9	143	17.4	14.6-20.7	214	15.2	13.2-17.4
2012	1,006	42.2	39.6-44.9	477	52.0	47.2-57.1	529	36.4	33.3-39.7	413	17.8	16.1-19.6	201	22.7	19.5-26.2	212	14.6	12.7-16.7
2013	1,028	41.8	39.2-44.5	479	50.1	45.5-55.0	549	36.7	33.7-40.0	344	14.5	13.0-16.1	152	16.6	13.9-19.6	192	13.0	11.2-15.0
2014	1,084	43.4	40.8-46.1	496	50.6	46.0-55.4	588	39.2	36.1-42.6	362	14.8	13.3-16.4	167	18.1	15.3-21.1	195	12.7	11.0-14.7
2015	987	38.4	36.0-40.9	487	48.3	43.9-53.0	500	32.2	29.4-35.2	309	12.4	11.0-13.9	133	13.9	11.6-16.6	176	11.3	9.7-13.2
2016	1,104	42.1	39.6-44.7	535	51.9	47.4-56.7	569	36.0	33.1-39.2	379	14.7	13.2-16.2	165	16.6	14.1-19.5	214	13.4	11.6-15.3
2017	958	36.3	34.0-38.8	463	44.4	40.3-48.7	495	31.3	28.5-34.2	326	12.5	11.1-13.9	150	15.2	12.8-17.9	176	10.9	9.3-12.7
2018	945	35.2	33.0-37.6	434	40.1	36.3-44.2	511	31.9	29.1-34.9	348	13.2	11.8-14.7	162	16.3	13.8-19.1	186	11.3	9.7-13.1
2019	1,012	37.3	35.0-39.7	486	44.9	40.9-49.2	526	32.5	29.7-35.5	346	12.7	11.4-14.2	162	15.5	13.1-18.2	184	10.8	9.3-12.6
2020	848	31.0	28.9-33.2	406	36.6	33.0-40.5	442	27.3	24.7-30.0	328	11.8	10.5-13.2	155	14.1	11.9-16.6	173	10.3	8.8-12.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	361	34.9	31.1-39.1	151	39.3	32.5-47.0	210	32.5	27.9-37.6	244	25.5	22.0-29.2	105	29.9	23.6-37.2	139	23.1	19.2-27.6
1977	428	44.1	39.5-48.9	156	44.0	36.3-52.5	272	44.3	38.7-50.3	217	22.5	19.4-26.0	85	25.2	19.5-31.8	132	21.2	17.4-25.4
1978	462	47.0	42.4-52.0	182	51.3	43.0-60.4	280	45.1	39.6-51.1	304	31.0	27.3-35.1	120	33.0	26.5-40.3	184	30.1	25.6-35.1
1979	501	47.1	42.7-51.8	209	53.0	45.1-61.8	292	43.9	38.7-49.6	296	28.1	24.7-31.7	130	34.9	28.4-42.3	166	24.6	20.8-28.9
1980	496	43.9	39.8-48.3	213	49.8	42.4-57.8	283	40.6	35.7-45.9	295	27.5	24.2-31.2	117	29.9	24.0-36.6	178	26.5	22.5-30.9
1981	477	41.8	37.9-46.1	200	48.1	40.8-56.2	277	38.7	34.0-43.8	323	29.6	26.2-33.2	119	30.4	24.6-37.1	204	29.4	25.3-33.9
1982	502	41.4	37.6-45.4	225	50.0	42.9-57.7	277	36.6	32.2-41.3	295	24.6	21.7-27.8	130	28.4	23.2-34.3	165	22.3	18.9-26.1
1983	502	38.5	35.0-42.3	224	49.0	42.1-56.7	278	33.6	29.6-37.9	284	22.3	19.6-25.2	126	25.8	21.0-31.2	158	20.1	16.9-23.6
1984	512	39.7	36.1-43.5	224	44.4	38.2-51.2	288	36.5	32.3-41.1	331	26.0	23.1-29.1	147	32.0	26.4-38.2	184	23.0	19.7-26.7
1985	572	43.2	39.5-47.1	245	52.2	45.2-59.9	327	38.7	34.5-43.3	316	24.5	21.7-27.5	123	27.1	21.9-32.9	193	23.5	20.2-27.1
1986	599	43.0	39.4-46.7	262	52.4	45.6-59.9	337	38.3	34.2-42.7	312	23.0	20.4-25.8	141	28.9	23.9-34.7	171	19.9	16.9-23.2
1987	546	38.1	34.8-41.5	231	43.2	37.3-49.8	315	35.0	31.1-39.2	308	22.2	19.7-25.0	128	25.2	20.5-30.4	180	20.6	17.6-24.0
1988	573	40.1	36.7-43.7	246	47.1	40.8-54.0	327	36.3	32.4-40.6	311	22.0	19.5-24.7	137	26.9	22.1-32.4	174	19.5	16.6-22.6
1989	597	40.7	37.4-44.3	278	52.9	46.3-60.1	319	34.5	30.7-38.6	305	20.8	18.4-23.3	143	27.4	22.7-32.8	162	17.3	14.7-20.2
1990	590	40.3	37.0-43.8	242	46.5	40.3-53.3	348	37.0	33.1-41.2	293	20.8	18.4-23.5	128	26.2	21.4-31.6	165	18.2	15.5-21.3
1991	641	43.5	40.0-47.1	270	50.1	43.7-57.0	371	39.7	35.7-44.0	335	23.5	20.9-26.2	138	28.4	23.4-34.1	197	21.2	18.3-24.4
1992	612	39.8	36.6-43.2	230	42.7	36.8-49.2	382	38.9	35.0-43.0	318	21.2	18.9-23.8	132	26.9	22.1-32.3	186	19.0	16.3-21.9
1993	618	40.1	36.9-43.5	267	48.3	42.2-55.0	351	35.8	32.1-39.8	306	20.0	17.7-22.4	124	22.4	18.3-27.2	182	18.7	16.0-21.6
1994	693	43.8	40.5-47.2	277	48.8	42.7-55.4	416	41.3	37.4-45.5	334	21.6	19.3-24.2	155	27.5	23.0-32.6	179	18.3	15.6-21.2
1995	655	40.1	37.0-43.3	286	48.2	42.3-54.5	369	35.6	32.0-39.5	375	23.7	21.3-26.3	158	26.8	22.5-31.7	217	21.6	18.8-24.7
1996	702	42.0	38.8-45.3	304	50.3	44.3-56.8	398	37.7	34.1-41.6	370	23.1	20.8-25.7	178	32.2	27.3-37.6	192	18.6	16.0-21.4
1997	750	44.2	41.1-47.6	322	53.2	47.1-59.8	428	39.8	36.1-43.8	364	22.2	19.9-24.6	164	29.5	24.8-34.7	200	18.9	16.3-21.7
1998	730	42.4	39.3-45.6	291	46.6	41.1-52.7	439	40.1	36.4-44.0	341	20.1	17.9-22.4	142	23.4	19.4-27.8	199	18.4	15.9-21.1
1999	786	43.7	40.6-46.9	296	45.4	40.1-51.2	490	42.9	39.1-46.9	394	22.5	20.3-24.8	156	25.6	21.5-30.2	238	21.1	18.5-24.0
2000	706	38.1	35.2-41.0	292	44.3	39.0-50.0	414	35.1	31.7-38.6	349	19.3	17.3-21.5	146	22.5	18.8-26.7	203	17.6	15.3-20.2
2001	740	39.3	36.4-42.2	305	45.1	39.9-50.8	435	36.4	33.1-40.0	377	20.4	18.4-22.6	156	24.7	20.7-29.1	221	18.7	16.3-21.3
2002	747	38.3	35.6-41.2	296	39.8	35.2-44.9	451	37.1	33.7-40.7	350	18.8	16.8-20.9	159	24.5	20.6-28.8	191	15.9	13.7-18.3
2003	797	40.4	37.6-43.3	332	47.1	41.8-52.7	465	37.3	33.9-40.8	353	18.6	16.7-20.7	160	24.5	20.6-28.8	193	15.8	13.6-18.2
2004	791	39.2	36.5-42.1	315	42.4	37.5-47.6	476	37.6	34.3-41.2	328	17.0	15.2-19.0	150	21.5	18.0-25.4	178	14.4	12.4-16.7
2005	777	37.9	35.2-40.7	329	43.0	38.2-48.1	448	35.0	31.8-38.4	312	16.0	14.3-17.9	129	18.4	15.2-22.1	183	14.6	12.5-16.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	801	38.9	36.2-41.7	323	42.6	37.9-47.8	478	37.1	33.8-40.6	332	16.6	14.8-18.5	139	20.1	16.7-23.9	193	15.0	13.0-17.3
2007	785	37.2	34.6-40.0	356	44.7	39.9-49.9	429	32.7	29.7-36.0	324	15.7	14.0-17.6	138	18.1	15.0-21.6	186	14.4	12.4-16.7
2008	804	37.8	35.2-40.6	355	44.2	39.5-49.3	449	34.0	30.9-37.3	315	15.2	13.5-17.0	142	18.6	15.5-22.1	173	13.2	11.3-15.3
2009	771	35.3	32.8-37.9	345	41.2	36.7-46.0	426	31.4	28.4-34.6	317	15.3	13.6-17.1	149	19.9	16.7-23.5	168	12.8	10.9-14.9
2010	752	33.9	31.5-36.5	323	39.0	34.7-43.8	429	31.5	28.5-34.6	305	14.0	12.4-15.7	157	20.0	16.9-23.6	148	10.6	8.9-12.5
2011	772	33.4	31.0-35.9	359	40.5	36.2-45.1	413	28.9	26.1-31.9	300	13.6	12.1-15.3	116	14.5	11.9-17.5	184	13.1	11.3-15.2
2012	740	31.5	29.2-33.9	332	37.3	33.2-41.8	408	28.3	25.6-31.2	339	14.7	13.1-16.3	156	17.8	15.0-21.0	183	12.6	10.8-14.6
2013	740	30.4	28.2-32.7	341	36.3	32.4-40.6	399	26.8	24.2-29.6	297	12.6	11.2-14.2	123	13.6	11.2-16.3	174	11.8	10.1-13.7
2014	800	32.4	30.1-34.7	353	36.6	32.7-40.8	447	30.0	27.2-33.0	296	12.0	10.6-13.4	136	14.6	12.1-17.4	160	10.4	8.8-12.1
2015	696	27.3	25.3-29.4	326	33.2	29.5-37.1	370	23.7	21.3-26.3	263	10.6	9.3-11.9	114	11.9	9.7-14.4	149	9.6	8.1-11.3
2016	789	30.2	28.1-32.4	366	35.6	31.9-39.6	423	26.9	24.3-29.6	292	11.2	10.0-12.6	119	11.9	9.8-14.3	173	10.8	9.2-12.6
2017	699	26.5	24.5-28.6	318	30.7	27.3-34.3	381	23.8	21.4-26.3	268	10.3	9.0-11.6	114	11.6	9.5-14.0	154	9.5	8.0-11.2
2018	672	25.0	23.1-27.0	282	25.9	22.8-29.2	390	24.4	21.9-27.0	294	11.1	9.9-12.5	135	13.5	11.3-16.1	159	9.7	8.2-11.4
2019	768	28.2	26.2-30.3	348	32.5	29.1-36.2	420	25.6	23.2-28.3	295	10.9	9.6-12.2	133	12.8	10.6-15.2	162	9.6	8.1-11.2
2020	648	23.6	21.8-25.5	301	27.3	24.2-30.7	347	21.3	19.0-23.7	277	9.9	8.8-11.2	129	11.9	9.9-14.2	148	8.7	7.3-10.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	120	11.5	9.3-14.1	58	13.2	9.6-17.6	62	10.3	7.6-13.4	58	6.0	4.4-8.0	28	7.2	4.4-11.0	30	5.2	3.3-7.6
1977	163	15.1	12.7-17.8	83	20.6	15.9-26.1	80	11.9	9.2-15.0	62	5.8	4.4-7.6	38	9.4	6.5-13.2	24	3.6	2.2-5.5
1978	147	13.2	11.0-15.8	66	14.1	10.5-18.5	81	12.5	9.7-15.7	60	6.3	4.6-8.3	28	6.8	4.3-10.1	32	5.7	3.7-8.2
1979	139	12.7	10.5-15.2	57	14.5	10.6-19.3	82	11.9	9.3-14.9	59	5.8	4.3-7.6	31	7.9	5.1-11.5	28	4.6	2.9-6.7
1980	172	14.2	12.0-16.6	81	17.5	13.5-22.3	91	12.3	9.8-15.2	61	5.2	3.9-6.8	29	6.8	4.3-10.0	32	4.4	2.9-6.2
1981	182	14.1	12.0-16.5	78	16.2	12.4-20.7	104	13.0	10.5-15.9	47	4.2	3.0-5.6	21	5.4	3.1-8.6	26	3.6	2.3-5.2
1982	178	13.9	11.8-16.2	87	17.3	13.5-21.6	91	11.7	9.3-14.5	60	4.8	3.6-6.3	30	5.9	3.8-8.7	30	4.1	2.7-5.9
1983	169	13.1	11.1-15.4	73	15.0	11.3-19.3	96	12.0	9.7-14.8	60	4.6	3.4-6.0	31	5.8	3.7-8.5	29	3.8	2.5-5.5
1984	188	13.7	11.7-15.9	93	18.3	14.4-22.9	95	11.2	9.0-13.7	49	4.0	2.9-5.4	21	5.0	2.9-7.7	28	3.6	2.3-5.2
1985	199	13.8	11.9-16.0	101	18.9	15.0-23.4	98	11.1	8.9-13.6	60	4.9	3.6-6.3	29	6.4	4.0-9.4	31	4.0	2.7-5.7
1986	204	14.2	12.2-16.4	95	17.3	13.7-21.5	109	12.4	10.1-15.0	43	3.2	2.3-4.3	18	3.5	1.9-5.7	25	3.0	1.9-4.4
1987	169	11.7	9.9-13.7	78	15.3	11.7-19.5	91	9.9	7.9-12.3	46	3.4	2.4-4.6	23	4.7	2.8-7.2	23	2.7	1.7-4.1
1988	166	11.1	9.4-13.0	68	12.6	9.5-16.3	98	10.3	8.3-12.6	40	3.1	2.2-4.3	20	4.4	2.6-6.9	20	2.4	1.4-3.7
1989	163	10.6	9.0-12.5	83	15.0	11.7-19.0	80	8.2	6.5-10.3	37	2.4	1.7-3.4	20	3.5	2.0-5.6	17	1.8	1.0-2.9
1990	180	11.8	10.1-13.8	82	14.6	11.4-18.4	98	10.3	8.3-12.5	40	2.8	2.0-3.8	20	4.1	2.3-6.6	20	2.2	1.3-3.4
1991	198	12.9	11.1-14.9	79	14.1	10.9-17.8	119	12.2	10.1-14.7	34	2.4	1.6-3.4	14	3.1	1.6-5.3	20	2.1	1.3-3.3
1992	184	11.7	10.0-13.5	83	13.8	10.7-17.4	101	10.4	8.5-12.7	43	2.7	1.9-3.6	19	2.9	1.7-4.6	24	2.5	1.6-3.7
1993	203	12.3	10.6-14.1	101	16.2	12.9-20.0	102	10.0	8.1-12.1	39	2.6	1.8-3.6	17	3.2	1.7-5.3	22	2.3	1.4-3.5
1994	178	10.8	9.2-12.6	90	14.8	11.7-18.5	88	8.6	6.8-10.6	47	3.0	2.2-4.0	21	3.4	2.0-5.3	26	2.6	1.7-3.8
1995	197	11.3	9.7-13.0	92	13.8	10.9-17.3	105	9.8	8.0-11.9	34	2.2	1.5-3.0	12	1.9	1.0-3.4	22	2.2	1.4-3.3
1996	219	12.9	11.2-14.8	88	13.6	10.7-16.9	131	12.3	10.3-14.7	51	3.0	2.2-4.0	25	3.8	2.4-5.8	26	2.5	1.6-3.6
1997	221	13.2	11.4-15.0	110	17.4	14.1-21.3	111	10.5	8.7-12.7	58	3.5	2.6-4.5	29	4.6	3.0-6.8	29	2.7	1.8-3.9
1998	220	12.4	10.8-14.2	97	14.0	11.2-17.3	123	11.2	9.3-13.3	43	2.6	1.9-3.5	19	3.1	1.8-4.9	24	2.3	1.5-3.4
1999	249	13.3	11.7-15.1	118	16.5	13.5-20.0	131	11.3	9.5-13.5	50	2.7	2.0-3.6	26	3.7	2.3-5.5	24	2.1	1.4-3.2
2000	226	11.5	10.1-13.2	109	15.0	12.2-18.2	117	9.6	8.0-11.6	39	2.1	1.5-2.9	19	2.6	1.5-4.0	20	1.8	1.1-2.7
2001	257	12.9	11.4-14.6	108	14.5	11.7-17.7	149	12.1	10.3-14.3	40	2.1	1.5-2.9	19	3.0	1.7-4.7	21	1.7	1.1-2.6
2002	230	11.4	10.0-13.0	118	16.0	13.1-19.3	112	9.0	7.4-10.8	38	2.0	1.4-2.7	20	3.0	1.7-4.7	18	1.5	0.9-2.4
2003	264	12.8	11.3-14.5	131	16.6	13.7-19.9	133	10.6	8.8-12.5	56	2.8	2.1-3.6	31	4.1	2.8-5.9	25	2.0	1.3-2.9
2004	272	12.9	11.4-14.5	117	14.9	12.2-18.0	155	11.9	10.1-14.0	53	2.6	1.9-3.4	29	4.1	2.6-5.9	24	1.9	1.2-2.8
2005	243	11.4	10.0-13.0	122	14.9	12.2-17.9	121	9.3	7.7-11.2	46	2.2	1.6-3.0	24	3.3	2.0-4.9	22	1.7	1.1-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	281	13.0	11.5-14.7	118	14.6	12.0-17.6	163	12.2	10.4-14.3	49	2.4	1.8-3.2	20	2.6	1.5-4.0	29	2.2	1.5-3.2
2007	284	12.9	11.4-14.5	144	16.1	13.5-19.1	140	10.6	8.9-12.6	46	2.2	1.6-2.9	24	3.1	2.0-4.7	22	1.7	1.0-2.5
2008	255	11.3	9.9-12.8	120	13.4	11.0-16.2	135	9.8	8.2-11.7	52	2.5	1.9-3.3	28	3.5	2.3-5.2	24	1.9	1.2-2.8
2009	287	12.3	10.9-13.9	153	16.3	13.7-19.2	134	9.8	8.2-11.6	45	2.0	1.4-2.7	21	2.4	1.4-3.7	24	1.7	1.1-2.5
2010	268	11.5	10.1-13.0	137	15.5	12.9-18.5	131	9.3	7.7-11.0	50	2.2	1.6-2.9	26	2.9	1.8-4.3	24	1.7	1.1-2.5
2011	272	11.3	10.0-12.7	134	14.0	11.6-16.7	138	9.5	8.0-11.3	57	2.5	1.8-3.2	27	2.9	1.9-4.3	30	2.1	1.4-3.0
2012	266	10.7	9.4-12.1	145	14.7	12.3-17.4	121	8.1	6.7-9.8	74	3.1	2.4-3.9	45	4.9	3.5-6.6	29	2.0	1.3-2.9
2013	288	11.4	10.1-12.9	138	13.8	11.5-16.4	150	9.9	8.4-11.7	47	1.9	1.4-2.5	29	3.0	2.0-4.4	18	1.2	0.7-1.9
2014	284	11.1	9.8-12.5	143	13.9	11.7-16.5	141	9.2	7.7-10.9	66	2.8	2.2-3.6	31	3.5	2.3-5.0	35	2.4	1.6-3.3
2015	291	11.1	9.8-12.5	161	15.1	12.8-17.7	130	8.5	7.1-10.2	46	1.8	1.3-2.5	19	2.1	1.2-3.3	27	1.7	1.1-2.6
2016	315	11.9	10.6-13.3	169	16.3	13.9-19.1	146	9.1	7.7-10.8	87	3.4	2.7-4.2	46	4.7	3.4-6.4	41	2.6	1.8-3.6
2017	259	9.8	8.7-11.2	145	13.7	11.5-16.2	114	7.5	6.1-9.0	58	2.2	1.7-2.9	36	3.6	2.5-5.1	22	1.4	0.8-2.1
2018	273	10.2	9.0-11.5	152	14.2	12.0-16.7	121	7.5	6.2-9.1	54	2.0	1.5-2.7	27	2.8	1.8-4.1	27	1.6	1.0-2.4
2019	244	9.0	7.9-10.3	138	12.4	10.4-14.8	106	6.8	5.6-8.3	51	1.9	1.4-2.5	29	2.7	1.8-4.0	22	1.3	0.8-2.0
2020	200	7.4	6.4-8.5	105	9.3	7.6-11.3	95	6.0	4.8-7.4	51	1.9	1.4-2.5	26	2.2	1.4-3.3	25	1.5	1.0-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	63	5.9	4.4-7.7	41	9.9	6.7-13.8	22	3.4	2.0-5.3	64	6.3	4.7-8.2	43	11.1	7.7-15.4	21	3.5	2.0-5.4
1977	61	5.1	3.8-6.8	42	9.0	6.1-12.7	19	2.8	1.6-4.5	80	6.8	5.3-8.6	54	12.4	8.9-16.7	26	3.5	2.3-5.3
1978	81	7.0	5.4-8.8	46	10.0	7.1-13.6	35	5.0	3.4-7.1	78	6.6	5.1-8.3	40	8.0	5.5-11.3	38	5.5	3.8-7.7
1979	62	5.1	3.8-6.6	36	7.8	5.1-11.1	26	3.4	2.2-5.1	83	7.3	5.7-9.2	44	10.6	7.4-14.6	39	5.5	3.8-7.6
1980	67	5.2	4.0-6.7	45	8.9	6.3-12.3	22	3.0	1.8-4.6	69	5.5	4.2-7.0	43	8.8	6.1-12.3	26	3.5	2.2-5.1
1981	67	5.1	3.9-6.5	44	8.4	5.9-11.6	23	3.0	1.9-4.5	68	5.4	4.2-7.0	41	7.8	5.4-10.9	27	3.8	2.5-5.6
1982	77	5.6	4.4-7.0	53	9.6	7.0-12.7	24	3.1	1.9-4.6	75	5.7	4.4-7.2	48	8.6	6.2-11.6	27	3.7	2.4-5.4
1983	77	5.7	4.4-7.2	48	9.4	6.6-12.8	29	3.5	2.3-5.0	97	7.3	5.8-9.0	58	11.1	8.2-14.7	39	4.9	3.5-6.8
1984	86	6.2	4.9-7.8	58	12.0	8.7-16.0	28	3.1	2.0-4.6	94	6.8	5.4-8.5	54	11.7	8.4-15.6	40	4.3	3.1-6.0
1985	85	5.7	4.5-7.1	57	10.4	7.7-13.8	28	3.0	2.0-4.4	93	6.6	5.2-8.1	55	10.3	7.5-13.7	38	4.4	3.1-6.1
1986	79	5.5	4.3-6.9	53	9.9	7.2-13.2	26	3.0	1.9-4.4	90	6.3	5.0-7.8	57	10.5	7.7-13.8	33	3.8	2.6-5.4
1987	85	5.4	4.3-6.8	49	8.2	5.8-11.2	36	3.8	2.6-5.3	92	6.6	5.3-8.2	45	9.4	6.6-12.8	47	5.3	3.9-7.1
1988	76	4.9	3.8-6.1	49	8.2	5.8-11.1	27	2.9	1.9-4.2	79	5.2	4.1-6.5	46	7.8	5.5-10.6	33	3.6	2.5-5.1
1989	82	5.5	4.3-6.9	45	8.3	5.8-11.4	37	4.0	2.8-5.5	90	5.8	4.6-7.2	49	8.0	5.7-10.7	41	4.4	3.1-6.0
1990	100	6.3	5.1-7.7	60	9.4	7.0-12.4	40	4.2	3.0-5.8	93	6.1	4.8-7.5	46	8.1	5.8-11.1	47	4.9	3.5-6.5
1991	103	6.3	5.1-7.6	66	10.9	8.1-14.1	37	3.6	2.5-5.0	102	6.6	5.3-8.0	57	10.1	7.4-13.4	45	4.6	3.3-6.2
1992	109	6.8	5.6-8.3	68	11.6	8.8-14.9	41	4.1	3.0-5.6	95	6.0	4.8-7.3	54	9.3	6.9-12.3	41	4.1	2.9-5.5
1993	104	6.2	5.1-7.6	72	11.3	8.6-14.5	32	3.2	2.2-4.5	110	6.8	5.5-8.2	68	10.7	8.1-13.7	42	4.4	3.1-5.9
1994	115	6.7	5.5-8.1	65	9.4	7.1-12.2	50	4.8	3.6-6.4	92	5.7	4.6-7.1	57	9.3	6.9-12.2	35	3.6	2.5-4.9
1995	113	6.3	5.2-7.6	70	10.1	7.7-13.0	43	4.0	2.9-5.4	100	5.7	4.6-7.0	56	8.2	6.0-10.9	44	4.1	3.0-5.6
1996	130	7.4	6.1-8.8	95	14.2	11.3-17.6	35	3.3	2.3-4.6	107	6.3	5.1-7.6	65	10.2	7.7-13.2	42	4.0	2.9-5.4
1997	123	6.7	5.6-8.0	77	10.6	8.2-13.4	46	4.3	3.1-5.7	96	5.4	4.3-6.6	58	8.3	6.2-11.0	38	3.5	2.5-4.8
1998	136	7.3	6.1-8.6	94	12.6	10.1-15.6	42	3.8	2.7-5.1	112	6.1	5.0-7.3	75	10.3	7.9-13.0	37	3.4	2.4-4.6
1999	151	7.9	6.6-9.2	89	11.7	9.3-14.4	62	5.3	4.1-6.8	87	4.6	3.7-5.7	51	6.7	4.9-8.8	36	3.2	2.2-4.4
2000	154	7.6	6.5-9.0	102	12.1	9.8-14.7	52	4.4	3.3-5.8	118	6.1	5.0-7.3	77	10.4	8.1-13.2	41	3.5	2.5-4.7
2001	146	7.1	6.0-8.4	110	13.5	10.9-16.4	36	3.0	2.1-4.2	106	5.3	4.4-6.5	65	8.3	6.3-10.6	41	3.4	2.5-4.7
2002	144	7.0	5.9-8.3	98	12.1	9.6-14.8	46	3.8	2.8-5.1	103	5.2	4.3-6.4	64	8.3	6.3-10.8	39	3.3	2.3-4.5
2003	168	7.7	6.6-9.0	120	13.7	11.2-16.5	48	3.8	2.8-5.0	112	5.4	4.5-6.6	72	9.2	7.1-11.8	40	3.2	2.3-4.3
2004	210	9.8	8.5-11.2	147	16.8	14.1-19.9	63	5.0	3.8-6.4	147	7.1	6.0-8.3	89	10.4	8.3-12.9	58	4.7	3.6-6.1
2005	203	8.9	7.7-10.2	154	16.8	14.1-19.8	49	3.7	2.7-4.9	123	5.9	4.9-7.0	81	10.0	7.9-12.6	42	3.3	2.4-4.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	188	8.5	7.3-9.8	137	15.7	13.0-18.6	51	3.9	2.9-5.1	136	6.1	5.1-7.3	105	11.9	9.6-14.6	31	2.4	1.6-3.4
2007	226	9.9	8.7-11.3	165	18.4	15.6-21.7	61	4.6	3.5-5.9	144	6.5	5.4-7.7	96	11.0	8.8-13.6	48	3.6	2.7-4.8
2008	230	9.9	8.6-11.2	171	17.8	15.1-20.8	59	4.4	3.4-5.7	139	6.1	5.1-7.2	98	10.8	8.7-13.3	41	3.0	2.1-4.0
2009	260	11.1	9.8-12.6	177	19.0	16.1-22.1	83	6.0	4.7-7.4	141	6.2	5.2-7.3	99	10.8	8.7-13.2	42	3.1	2.2-4.2
2010	283	11.5	10.2-13.0	189	19.0	16.3-22.1	94	6.6	5.3-8.1	171	7.2	6.2-8.4	113	12.2	9.9-14.8	58	4.0	3.0-5.2
2011	281	11.4	10.1-12.8	187	18.2	15.6-21.2	94	6.6	5.3-8.1	164	6.9	5.9-8.0	99	9.9	8.0-12.1	65	4.7	3.6-6.0
2012	331	13.1	11.7-14.6	234	23.1	20.1-26.4	97	6.5	5.3-8.0	166	6.8	5.8-7.9	114	11.4	9.3-13.8	52	3.6	2.7-4.8
2013	304	11.4	10.2-12.8	207	18.9	16.3-21.7	97	6.3	5.1-7.7	171	6.7	5.7-7.8	112	11.1	9.1-13.5	59	3.8	2.9-4.9
2014	303	11.0	9.8-12.4	197	16.9	14.5-19.5	106	6.8	5.6-8.3	172	6.6	5.6-7.6	106	9.8	7.9-11.9	66	4.3	3.3-5.5
2015	299	10.8	9.6-12.2	214	19.0	16.5-21.9	85	5.2	4.2-6.5	160	5.8	5.0-6.8	99	8.9	7.1-10.9	61	3.8	2.9-4.9
2016	312	11.0	9.8-12.4	225	19.5	17.0-22.3	87	5.1	4.1-6.4	181	6.6	5.7-7.7	132	12.2	10.1-14.5	49	3.0	2.2-4.0
2017	314	11.2	9.9-12.5	228	20.1	17.5-23.0	86	5.1	4.1-6.4	159	5.7	4.8-6.7	108	9.6	7.8-11.6	51	3.0	2.2-4.0
2018	270	9.3	8.2-10.5	177	15.2	13.0-17.7	93	5.3	4.3-6.6	172	6.2	5.3-7.2	110	10.1	8.3-12.3	62	3.6	2.8-4.7
2019	280	9.8	8.6-11.0	178	15.4	13.2-18.0	102	5.9	4.8-7.3	168	5.9	5.0-6.9	100	9.1	7.3-11.1	68	4.0	3.1-5.1
2020	221	7.5	6.5-8.6	132	11.0	9.1-13.1	89	5.1	4.1-6.3	150	5.2	4.3-6.1	91	8.0	6.4-9.9	59	3.3	2.5-4.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	149	14.6	12.2-17.4	64	14.7	10.9-19.2	85	14.3	11.2-18.0	151	13.9	11.6-16.5	70	15.9	11.9-20.6	81	12.6	9.8-15.9
1977	145	13.4	11.1-16.1	72	17.1	12.7-22.4	73	11.3	8.7-14.4	125	11.6	9.5-14.1	65	15.5	11.4-20.6	60	9.3	6.9-12.2
1978	148	13.5	11.2-16.1	61	12.8	9.4-17.0	87	13.6	10.7-17.0	134	11.7	9.7-14.1	55	11.8	8.5-15.8	79	11.6	9.0-14.6
1979	159	14.1	11.8-16.7	72	14.8	11.3-19.0	87	13.3	10.5-16.6	146	13.3	11.0-15.8	56	11.1	8.2-14.6	90	14.2	11.2-17.6
1980	158	13.5	11.3-16.0	85	18.3	14.2-23.3	73	10.5	8.1-13.3	130	11.4	9.4-13.7	69	15.9	11.9-20.8	61	8.8	6.6-11.4
1981	180	14.8	12.5-17.3	99	20.7	16.4-25.7	81	11.1	8.6-13.9	147	12.0	10.0-14.3	82	17.1	13.1-21.9	65	8.9	6.8-11.5
1982	172	14.3	12.1-16.8	70	16.2	12.2-20.9	102	13.4	10.8-16.4	160	13.5	11.4-15.9	67	15.6	11.7-20.2	93	12.4	9.9-15.3
1983	196	14.7	12.6-17.0	89	16.7	13.1-20.9	107	13.3	10.8-16.1	178	13.6	11.6-15.9	90	18.3	14.3-22.9	88	10.9	8.7-13.5
1984	188	14.3	12.2-16.6	84	16.4	12.7-20.8	104	13.0	10.5-15.8	165	12.9	10.9-15.1	81	17.0	13.1-21.5	84	10.5	8.3-13.1
1985	175	12.8	10.9-14.9	92	16.7	13.2-20.8	83	10.2	8.1-12.7	149	11.3	9.4-13.3	75	14.4	11.0-18.5	74	9.3	7.2-11.7
1986	190	13.7	11.8-15.9	83	15.1	11.8-19.0	107	12.6	10.3-15.3	158	11.7	9.9-13.7	70	13.0	9.9-16.7	88	10.7	8.5-13.2
1987	214	14.8	12.8-17.0	104	17.4	13.9-21.4	110	12.7	10.4-15.4	175	12.2	10.3-14.2	80	14.9	11.5-19.0	95	10.6	8.5-13.0
1988	209	14.2	12.3-16.4	97	17.8	14.1-22.1	112	12.2	10.0-14.7	149	10.2	8.5-12.0	72	12.2	9.3-15.6	77	8.7	6.8-10.9
1989	195	13.2	11.3-15.2	79	14.0	10.9-17.7	116	12.5	10.3-15.0	180	12.0	10.3-14.0	75	13.1	10.1-16.7	105	11.2	9.1-13.6
1990	205	14.1	12.1-16.2	86	16.1	12.5-20.2	119	12.9	10.6-15.5	171	11.6	9.8-13.5	70	12.8	9.7-16.5	101	10.8	8.8-13.2
1991	180	12.0	10.2-13.9	82	14.6	11.3-18.4	98	10.5	8.5-12.8	165	10.5	8.9-12.3	82	13.6	10.6-17.1	83	8.6	6.8-10.7
1992	191	12.4	10.6-14.3	71	12.7	9.6-16.4	120	12.2	10.1-14.6	143	9.7	8.1-11.4	53	10.6	7.7-14.1	90	9.4	7.5-11.5
1993	193	12.2	10.5-14.1	84	14.4	11.2-18.2	109	11.0	9.0-13.3	148	9.5	8.0-11.2	58	11.0	8.1-14.6	90	9.0	7.2-11.1
1994	217	13.5	11.8-15.5	88	14.6	11.5-18.3	129	12.8	10.7-15.2	191	12.0	10.3-13.9	71	11.9	9.1-15.3	120	11.9	9.8-14.2
1995	187	11.7	10.1-13.6	82	13.9	10.8-17.4	105	10.4	8.5-12.6	173	10.7	9.1-12.4	75	12.8	9.8-16.3	98	9.5	7.7-11.6
1996	218	13.1	11.3-14.9	79	12.4	9.6-15.6	139	13.2	11.1-15.6	167	10.1	8.6-11.8	62	9.7	7.3-12.6	105	10.1	8.3-12.3
1997	224	13.2	11.5-15.0	96	14.0	11.2-17.2	128	12.2	10.1-14.5	177	10.7	9.1-12.4	72	11.0	8.4-13.9	105	10.1	8.2-12.2
1998	210	12.2	10.6-14.0	81	12.0	9.4-15.1	129	12.0	10.0-14.2	171	10.0	8.5-11.7	69	10.3	7.8-13.1	102	9.6	7.8-11.6
1999	250	14.2	12.5-16.1	97	15.3	12.2-18.9	153	13.6	11.5-16.0	196	11.3	9.7-13.0	71	11.0	8.4-14.0	125	11.2	9.3-13.4
2000	237	13.2	11.5-15.0	78	11.8	9.2-14.9	159	13.8	11.7-16.1	184	10.2	8.7-11.8	78	12.0	9.3-15.0	106	9.1	7.5-11.0
2001	244	12.9	11.3-14.7	91	12.8	10.2-15.9	153	12.9	10.9-15.1	200	10.8	9.4-12.5	67	9.7	7.4-12.4	133	11.3	9.5-13.4
2002	245	12.9	11.3-14.7	94	12.5	10.0-15.5	151	12.8	10.8-15.0	207	10.8	9.4-12.4	92	12.4	9.8-15.3	115	9.6	8.0-11.6
2003	229	11.8	10.3-13.5	111	15.4	12.5-18.7	118	9.6	8.0-11.5	171	9.0	7.7-10.5	74	10.3	7.9-13.1	97	8.1	6.6-9.9
2004	241	12.1	10.6-13.7	102	13.2	10.6-16.2	139	11.2	9.4-13.2	210	10.5	9.1-12.0	91	12.4	9.8-15.3	119	9.4	7.8-11.3
2005	221	10.7	9.3-12.3	89	11.9	9.4-14.8	132	10.2	8.6-12.2	185	9.2	7.9-10.7	76	9.7	7.6-12.3	109	8.7	7.2-10.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	291	14.0	12.4-15.7	118	14.1	11.6-17.1	173	13.6	11.6-15.8	194	9.5	8.2-10.9	83	10.8	8.5-13.5	111	8.7	7.1-10.5
2007	283	13.6	12.0-15.3	131	16.4	13.5-19.6	152	11.8	10.0-13.9	224	10.9	9.5-12.5	102	13.1	10.5-16.1	122	9.6	8.0-11.5
2008	255	12.1	10.6-13.7	92	11.1	8.9-13.8	163	12.3	10.5-14.4	218	10.4	9.1-11.9	88	11.1	8.8-13.9	130	9.9	8.3-11.8
2009	267	12.3	10.9-13.9	120	14.9	12.2-17.9	147	10.8	9.1-12.7	209	9.8	8.5-11.3	91	11.6	9.2-14.3	118	8.7	7.2-10.5
2010	309	13.9	12.3-15.5	137	15.9	13.2-18.9	172	12.5	10.7-14.6	208	9.8	8.4-11.2	88	10.6	8.4-13.1	120	9.1	7.6-10.9
2011	355	15.5	13.9-17.2	151	16.6	14.0-19.7	204	14.6	12.7-16.8	262	11.7	10.3-13.3	117	13.9	11.4-16.8	145	10.4	8.7-12.2
2012	339	14.3	12.8-15.9	130	14.2	11.7-16.9	209	14.5	12.6-16.6	257	10.9	9.6-12.3	86	9.3	7.3-11.6	171	11.9	10.2-13.9
2013	378	15.5	14.0-17.2	167	17.4	14.7-20.4	211	14.2	12.3-16.3	246	10.2	8.9-11.6	115	12.6	10.3-15.2	131	8.8	7.3-10.5
2014	338	13.4	12.0-14.9	148	14.2	12.0-16.8	190	12.6	10.8-14.6	246	9.9	8.7-11.2	103	10.2	8.3-12.5	143	9.6	8.0-11.3
2015	388	15.3	13.8-16.9	183	18.1	15.4-21.0	205	13.4	11.6-15.4	269	10.8	9.5-12.2	123	12.5	10.3-15.0	146	9.6	8.1-11.3
2016	377	14.3	12.9-15.9	176	16.7	14.3-19.5	201	12.6	10.9-14.5	269	10.2	9.0-11.6	134	12.8	10.7-15.3	135	8.4	7.1-10.0
2017	376	14.3	12.8-15.8	159	15.3	12.9-18.0	217	13.6	11.8-15.6	265	10.0	8.9-11.4	116	11.5	9.4-13.9	149	9.2	7.8-10.8
2018	396	14.6	13.1-16.1	169	15.5	13.2-18.1	227	13.7	12.0-15.7	263	9.6	8.4-10.8	103	9.4	7.6-11.5	160	9.5	8.0-11.1
2019	382	13.8	12.4-15.3	166	14.9	12.7-17.5	216	12.9	11.2-14.8	283	10.2	9.0-11.5	116	10.4	8.5-12.6	167	9.8	8.4-11.4
2020	354	12.2	11.0-13.6	161	13.8	11.7-16.2	193	11.2	9.7-13.0	268	9.4	8.3-10.6	130	11.4	9.5-13.6	138	8.1	6.8-9.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Larynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	79	6.0	4.7-7.6	63	11.7	8.8-15.1	16	2.1	1.1-3.5	50	4.2	3.0-5.7	39	8.3	5.6-11.8	11	1.6	0.7-2.9
1977	90	7.1	5.6-8.8	72	14.8	11.1-19.1	18	2.3	1.4-3.7	43	3.5	2.4-4.9	32	7.4	4.5-11.2	11	1.3	0.7-2.4
1978	99	7.3	5.9-9.0	81	15.1	11.7-19.1	18	2.2	1.3-3.4	64	4.5	3.5-5.8	55	9.7	7.2-12.9	9	1.0	0.5-2.0
1979	106	7.8	6.3-9.4	82	15.4	12.0-19.3	24	2.9	1.8-4.3	58	4.1	3.1-5.4	47	8.7	6.2-11.7	11	1.2	0.6-2.2
1980	100	6.9	5.6-8.5	74	12.8	9.9-16.4	26	3.0	1.9-4.5	52	3.7	2.8-4.9	34	6.6	4.4-9.4	18	2.0	1.2-3.2
1981	93	6.3	5.1-7.8	75	12.7	9.8-16.2	18	2.1	1.2-3.4	55	3.6	2.7-4.8	35	6.1	4.2-8.5	20	2.1	1.3-3.3
1982	86	5.7	4.5-7.0	74	12.2	9.5-15.4	12	1.3	0.7-2.4	39	2.7	1.9-3.7	31	5.6	3.7-8.1	8	0.9	0.4-1.8
1983	114	7.5	6.1-9.1	94	15.7	12.4-19.5	20	2.3	1.4-3.5	46	3.1	2.2-4.1	38	6.3	4.3-8.9	8	0.9	0.4-1.9
1984	110	7.0	5.8-8.5	80	12.8	10.1-16.1	30	3.2	2.2-4.6	62	3.9	3.0-5.0	49	7.6	5.6-10.2	13	1.4	0.7-2.4
1985	146	9.1	7.6-10.7	117	19.1	15.6-23.0	29	2.8	1.9-4.0	52	3.4	2.5-4.5	42	6.9	4.9-9.4	10	1.1	0.5-2.1
1986	140	8.9	7.4-10.5	109	18.3	14.8-22.4	31	3.2	2.1-4.5	65	4.2	3.3-5.4	52	8.8	6.5-11.8	13	1.4	0.8-2.4
1987	123	7.6	6.2-9.0	98	15.4	12.3-19.0	25	2.6	1.7-3.8	66	4.2	3.2-5.4	50	8.3	6.0-11.2	16	1.7	0.9-2.7
1988	129	7.4	6.1-8.8	96	14.0	11.3-17.3	33	3.1	2.1-4.3	71	4.3	3.3-5.4	58	9.4	7.0-12.3	13	1.2	0.6-2.0
1989	142	8.3	6.9-9.8	91	13.8	10.9-17.1	51	4.9	3.6-6.5	60	3.6	2.7-4.7	43	7.1	5.0-9.8	17	1.6	0.9-2.6
1990	147	8.5	7.1-10.0	110	16.6	13.5-20.3	37	3.6	2.5-4.9	46	2.8	2.0-3.7	35	5.7	3.9-8.2	11	1.1	0.5-1.9
1991	123	7.0	5.8-8.4	92	14.1	11.2-17.4	31	2.9	1.9-4.1	55	3.2	2.4-4.2	40	6.0	4.2-8.3	15	1.4	0.8-2.4
1992	152	8.3	7.0-9.8	122	17.2	14.2-20.7	30	2.6	1.8-3.8	65	3.7	2.9-4.8	51	7.9	5.7-10.5	14	1.3	0.7-2.2
1993	116	6.7	5.5-8.0	89	13.8	10.9-17.1	27	2.5	1.7-3.7	67	3.9	3.0-4.9	48	7.7	5.5-10.4	19	1.8	1.1-2.8
1994	108	6.0	4.9-7.2	77	11.1	8.7-14.1	31	2.8	1.9-4.0	60	3.4	2.5-4.3	42	6.0	4.2-8.2	18	1.7	1.0-2.6
1995	120	6.5	5.4-7.8	85	11.8	9.3-14.7	35	3.2	2.2-4.4	72	4.1	3.2-5.1	55	8.5	6.2-11.2	17	1.5	0.9-2.5
1996	113	6.0	4.9-7.2	92	12.6	10.0-15.6	21	1.8	1.1-2.8	63	3.5	2.7-4.5	54	8.2	6.0-10.8	9	0.8	0.4-1.5
1997	100	5.2	4.2-6.4	74	9.9	7.7-12.5	26	2.3	1.5-3.3	56	3.0	2.2-3.9	45	6.2	4.4-8.4	11	1.0	0.5-1.7
1998	112	5.8	4.8-7.0	76	10.6	8.2-13.4	36	3.0	2.1-4.2	52	2.7	2.0-3.6	43	5.8	4.2-7.9	9	0.8	0.4-1.5
1999	101	5.2	4.2-6.3	75	10.2	7.9-13.0	26	2.2	1.4-3.2	45	2.3	1.7-3.1	31	4.2	2.8-6.1	14	1.2	0.6-2.0
2000	96	4.9	3.9-5.9	69	9.3	7.1-11.9	27	2.2	1.5-3.3	54	2.8	2.1-3.7	42	5.7	4.0-7.8	12	1.1	0.6-1.9
2001	87	4.2	3.3-5.2	64	7.6	5.8-9.8	23	1.9	1.2-2.8	35	1.8	1.2-2.5	27	3.7	2.4-5.4	8	0.6	0.3-1.3
2002	88	4.2	3.3-5.1	62	7.4	5.6-9.5	26	2.1	1.3-3.0	46	2.2	1.6-3.0	34	4.2	2.9-5.9	12	1.0	0.5-1.7
2003	104	4.9	4.0-5.9	77	9.5	7.4-12.0	27	2.1	1.4-3.1	37	1.8	1.3-2.5	31	4.0	2.7-5.7	6	0.5	0.2-1.0
2004	92	4.1	3.3-5.0	75	8.4	6.6-10.7	17	1.2	0.7-2.0	47	2.3	1.6-3.0	38	4.9	3.4-6.8	9	0.7	0.3-1.4
2005	91	4.2	3.4-5.2	68	8.4	6.4-10.7	23	1.7	1.1-2.6	37	1.7	1.2-2.4	31	3.7	2.5-5.3	6	0.5	0.2-1.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Larynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	77	3.3	2.6-4.1	57	6.2	4.6-8.1	20	1.4	0.8-2.1	39	1.9	1.3-2.5	32	4.0	2.7-5.7	7	0.5	0.2-1.1
2007	94	4.1	3.3-5.1	76	8.5	6.7-10.7	18	1.3	0.8-2.1	43	1.9	1.4-2.6	35	4.2	2.9-5.9	8	0.6	0.2-1.2
2008	106	4.8	3.9-5.8	69	8.2	6.3-10.4	37	2.7	1.9-3.8	41	1.8	1.3-2.5	26	3.3	2.1-4.8	15	1.1	0.6-1.8
2009	87	3.7	3.0-4.6	68	7.8	6.0-10.0	19	1.3	0.8-2.1	39	1.7	1.2-2.3	29	3.4	2.2-5.0	10	0.7	0.3-1.3
2010	85	3.5	2.8-4.4	63	6.8	5.1-8.7	22	1.5	1.0-2.3	31	1.4	0.9-2.0	25	3.1	1.9-4.6	6	0.4	0.2-0.9
2011	88	3.5	2.8-4.3	62	6.3	4.8-8.2	26	1.7	1.1-2.4	32	1.4	0.9-2.0	21	2.5	1.5-3.9	11	0.8	0.4-1.4
2012	72	2.8	2.2-3.6	53	5.4	4.0-7.1	19	1.2	0.7-2.0	31	1.3	0.9-1.8	24	2.7	1.7-4.1	7	0.4	0.2-0.9
2013	84	3.3	2.6-4.1	67	6.8	5.2-8.7	17	1.1	0.6-1.7	32	1.3	0.9-1.8	22	2.2	1.4-3.4	10	0.6	0.3-1.2
2014	89	3.5	2.8-4.3	66	6.3	4.9-8.1	23	1.6	1.0-2.4	37	1.4	1.0-2.0	31	3.1	2.1-4.5	6	0.4	0.1-0.8
2015	107	4.0	3.2-4.8	85	8.3	6.6-10.3	22	1.3	0.8-2.0	22	0.9	0.6-1.4	14	1.5	0.8-2.6	8	0.5	0.2-1.1
2016	69	2.6	2.0-3.3	51	5.0	3.7-6.6	18	1.1	0.6-1.7	34	1.3	0.9-1.8	26	2.6	1.7-3.9	8	0.5	0.2-0.9
2017	76	2.7	2.1-3.4	48	4.2	3.1-5.7	28	1.7	1.1-2.5	26	0.9	0.6-1.3	18	1.7	1.0-2.7	8	0.4	0.2-0.9
2018	76	2.6	2.1-3.3	60	5.3	4.0-6.9	16	0.9	0.5-1.5	26	0.9	0.6-1.4	18	1.8	1.0-2.9	8	0.5	0.2-1.0
2019	79	2.8	2.2-3.5	60	5.4	4.1-7.1	19	1.1	0.7-1.8	17	0.6	0.3-0.9	14	1.2	0.7-2.1	3	0.2	0.0-0.5
2020	55	1.9	1.4-2.5	46	4.0	2.9-5.4	9	0.5	0.2-1.0	13	0.5	0.2-0.8	9	0.9	0.4-1.7	4	0.2	0.1-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	666	55.5	51.0-60.2	493	102.5	92.6-113.1	173	25.0	21.2-29.3	524	43.5	39.6-47.8	395	82.5	73.5-92.2	129	18.3	15.1-22.0
1977	674	54.1	49.8-58.7	488	102.6	92.6-113.4	186	23.9	20.5-27.8	554	44.5	40.6-48.6	408	85.5	76.3-95.3	146	18.9	15.8-22.4
1978	718	57.3	52.9-62.0	527	105.2	95.5-115.5	191	26.7	22.8-31.0	611	48.4	44.4-52.6	444	88.5	79.7-97.9	167	22.7	19.2-26.7
1979	787	62.4	57.8-67.2	567	114.8	104.5-125.8	220	29.7	25.7-34.1	638	51.3	47.1-55.8	473	98.1	88.4-108.6	165	22.5	19.0-26.4
1980	791	59.2	54.9-63.7	573	110.9	101.0-121.3	218	27.1	23.5-31.1	642	48.6	44.7-52.7	474	91.9	83.0-101.4	168	21.6	18.3-25.3
1981	828	59.3	55.2-63.7	574	107.1	97.8-117.1	254	29.6	26.0-33.6	699	50.9	47.0-55.0	495	93.3	84.5-102.7	204	24.6	21.2-28.3
1982	851	60.4	56.2-64.8	583	107.3	98.1-117.1	268	31.3	27.5-35.4	680	48.7	45.0-52.7	470	86.3	78.0-95.1	210	25.5	22.0-29.3
1983	856	59.5	55.4-63.8	590	106.5	97.3-116.2	266	30.7	27.0-34.7	755	52.6	48.8-56.7	528	96.9	88.1-106.3	227	25.7	22.4-29.4
1984	918	61.9	57.8-66.2	616	108.6	99.5-118.2	302	33.6	29.8-37.7	705	48.2	44.5-52.0	492	89.5	81.1-98.6	213	23.6	20.5-27.1
1985	911	60.7	56.7-64.9	623	110.8	101.5-120.7	288	31.2	27.6-35.1	746	50.6	46.8-54.5	516	94.0	85.4-103.3	230	25.3	22.0-28.9
1986	934	60.6	56.6-64.8	595	103.0	94.2-112.4	339	35.7	31.9-39.8	728	47.8	44.2-51.5	468	82.2	74.3-90.8	260	27.6	24.3-31.3
1987	992	62.8	58.8-66.9	665	112.1	103.1-121.6	327	33.6	30.0-37.6	772	50.1	46.5-53.9	498	87.8	79.5-96.5	274	28.7	25.4-32.4
1988	933	58.6	54.7-62.6	595	101.5	92.9-110.7	338	34.1	30.4-38.0	760	48.7	45.2-52.4	494	86.3	78.2-95.0	266	27.4	24.1-31.0
1989	993	61.0	57.2-65.1	639	107.4	98.6-116.7	354	34.9	31.3-38.8	748	47.3	43.8-50.9	503	88.6	80.4-97.4	245	24.4	21.4-27.8
1990	1,012	61.2	57.4-65.2	644	105.4	96.7-114.5	368	35.9	32.2-39.8	780	47.6	44.2-51.1	511	83.0	75.4-91.1	269	26.7	23.6-30.2
1991	984	59.6	55.9-63.6	627	102.6	94.1-111.6	357	35.0	31.4-38.9	735	44.9	41.6-48.4	486	81.1	73.5-89.2	249	24.5	21.5-27.8
1992	1,017	60.4	56.6-64.3	637	102.6	94.2-111.6	380	36.4	32.8-40.3	737	45.2	41.9-48.6	438	74.1	66.8-82.0	299	29.3	26.0-32.8
1993	971	57.3	53.6-61.1	586	95.3	87.1-104.1	385	36.4	32.8-40.2	723	42.8	39.6-46.1	444	74.2	66.8-82.0	279	26.1	23.1-29.4
1994	1,032	59.3	55.7-63.1	635	98.5	90.4-107.0	397	36.7	33.2-40.6	810	47.7	44.4-51.2	509	81.8	74.3-89.9	301	28.7	25.5-32.1
1995	970	55.5	52.0-59.2	586	89.9	82.4-98.0	384	35.5	32.0-39.3	811	46.8	43.5-50.2	492	77.0	69.9-84.5	319	29.5	26.4-33.0
1996	1,081	60.5	56.9-64.3	631	94.0	86.3-102.1	450	41.0	37.3-45.0	828	47.1	43.8-50.5	509	77.6	70.5-85.1	319	29.5	26.3-33.0
1997	1,028	57.3	53.8-61.0	579	87.0	79.7-94.9	449	40.3	36.6-44.2	783	44.6	41.5-47.9	460	71.7	64.9-79.0	323	29.4	26.3-32.8
1998	1,034	55.8	52.4-59.4	573	81.5	74.7-88.8	461	40.5	36.9-44.4	783	43.4	40.3-46.6	452	67.1	60.7-73.9	331	29.7	26.6-33.1
1999	957	50.8	47.6-54.2	546	77.4	70.7-84.5	411	35.3	32.0-38.9	746	40.5	37.6-43.5	416	60.9	54.9-67.4	330	28.9	25.8-32.2
2000	996	51.6	48.4-54.9	553	76.9	70.3-83.9	443	37.0	33.6-40.6	735	38.5	35.7-41.4	423	60.0	54.1-66.3	312	26.2	23.4-29.3
2001	979	50.6	47.4-53.9	541	76.1	69.5-83.2	438	36.3	32.9-39.8	742	38.5	35.8-41.5	414	59.2	53.3-65.5	328	27.2	24.3-30.3
2002	1,026	51.3	48.2-54.6	554	75.6	69.1-82.5	472	38.0	34.6-41.6	752	38.1	35.4-40.9	412	57.1	51.4-63.1	340	27.5	24.6-30.6
2003	983	48.6	45.5-51.8	498	66.1	60.1-72.4	485	38.7	35.3-42.3	725	36.1	33.5-38.9	387	52.3	47.0-58.1	338	27.0	24.2-30.1
2004	1,048	51.0	47.9-54.2	551	69.7	63.8-76.1	497	39.3	35.9-42.9	696	34.0	31.5-36.7	375	48.7	43.6-54.1	321	25.3	22.6-28.2
2005	1,044	50.0	46.9-53.1	560	71.9	65.8-78.5	484	37.6	34.3-41.1	728	35.4	32.9-38.1	383	49.2	44.1-54.6	345	27.3	24.4-30.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	1,022	48.1	45.2-51.2	529	65.2	59.5-71.3	493	37.7	34.4-41.2	712	34.4	31.9-37.1	360	46.7	41.8-52.0	352	27.4	24.6-30.5
2007	1,039	48.3	45.3-51.4	532	65.3	59.6-71.4	507	38.5	35.2-42.0	712	33.7	31.2-36.3	382	47.9	43.0-53.2	330	25.4	22.7-28.4
2008	1,056	49.0	46.0-52.1	540	66.5	60.7-72.6	516	38.9	35.6-42.5	679	32.0	29.6-34.5	344	43.0	38.3-48.0	335	25.5	22.8-28.4
2009	1,076	48.7	45.8-51.7	554	65.9	60.2-71.9	522	38.4	35.2-41.9	671	30.8	28.5-33.3	365	44.4	39.7-49.4	306	22.7	20.2-25.5
2010	1,085	47.6	44.8-50.6	543	62.2	56.8-67.9	542	39.0	35.8-42.5	742	33.0	30.6-35.5	387	46.5	41.7-51.6	355	25.6	23.0-28.5
2011	1,161	50.1	47.2-53.1	569	65.1	59.6-71.0	592	41.4	38.1-44.9	728	31.6	29.3-34.1	371	42.4	38.0-47.1	357	25.1	22.6-27.9
2012	1,129	47.3	44.5-50.2	538	59.7	54.5-65.2	591	40.4	37.2-43.8	730	31.2	28.9-33.6	384	43.8	39.4-48.7	346	23.9	21.4-26.6
2013	1,041	42.7	40.1-45.5	522	55.8	50.9-61.0	519	35.0	32.0-38.2	705	29.3	27.1-31.5	365	40.1	35.9-44.6	340	22.8	20.4-25.4
2014	1,108	44.1	41.5-46.9	554	57.2	52.4-62.4	554	36.1	33.1-39.3	658	26.7	24.7-28.9	338	36.2	32.3-40.4	320	20.9	18.7-23.4
2015	1,097	42.0	39.5-44.7	536	52.5	48.0-57.3	561	35.7	32.7-38.8	656	25.6	23.6-27.6	320	32.8	29.2-36.8	336	21.4	19.1-23.8
2016	1,097	41.6	39.1-44.2	525	50.4	46.0-55.1	572	35.8	32.9-38.9	616	23.5	21.6-25.4	328	32.0	28.5-35.8	288	18.0	15.9-20.2
2017	1,130	41.7	39.2-44.2	558	52.4	48.0-57.2	572	34.8	32.0-37.8	559	21.0	19.2-22.8	280	27.5	24.3-31.1	279	17.1	15.1-19.2
2018	1,157	42.1	39.7-44.7	578	54.1	49.6-58.8	579	34.6	31.8-37.6	608	22.2	20.5-24.1	319	30.0	26.7-33.6	289	17.4	15.4-19.6
2019	1,073	38.4	36.1-40.8	507	46.1	42.0-50.4	566	33.5	30.8-36.5	537	19.2	17.6-20.9	274	25.8	22.8-29.2	263	15.3	13.5-17.3
2020	897	31.5	29.4-33.6	458	41.1	37.3-45.2	439	25.3	23.0-27.8	464	16.3	14.9-17.9	230	21.1	18.4-24.1	234	13.4	11.7-15.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	8	0.8	0.3-1.6	1	0.2	0.0-1.2	7	1.1	0.4-2.4	5	0.6	0.1-1.4	3	1.3	0.2-3.8	2	0.3	0.0-1.0
1977	8	0.7	0.3-1.5	3	0.7	0.1-2.1	5	0.8	0.2-1.8	4	0.4	0.1-1.0	2	0.7	0.1-2.4	2	0.2	0.0-0.9
1978	9	0.8	0.3-1.5	3	0.6	0.1-2.0	6	0.9	0.3-1.9	5	0.4	0.1-1.0	2	0.3	0.0-1.3	3	0.5	0.1-1.4
1979	17	1.2	0.7-2.0	6	1.0	0.4-2.3	11	1.4	0.6-2.5	2	0.2	0.0-0.7	1	0.3	0.0-1.6	1	0.1	0.0-0.7
1980	10	0.8	0.3-1.5	3	0.5	0.1-1.5	7	1.0	0.4-2.0	14	1.1	0.6-1.9	8	1.2	0.5-2.5	6	0.9	0.3-2.1
1981	18	1.4	0.8-2.3	5	0.8	0.3-2.0	13	1.8	0.9-3.1	14	1.1	0.5-1.9	6	0.9	0.3-2.2	8	1.1	0.4-2.3
1982	10	0.7	0.4-1.4	2	0.3	0.0-1.3	8	1.0	0.4-1.9	7	0.5	0.2-1.1	4	1.0	0.2-2.8	3	0.3	0.1-0.8
1983	10	0.7	0.3-1.3	2	0.3	0.0-1.2	8	0.9	0.4-1.8	18	1.2	0.7-2.0	13	2.1	1.1-3.6	5	0.6	0.2-1.5
1984	21	1.5	0.9-2.3	5	0.8	0.2-2.0	16	1.9	1.1-3.1	12	0.7	0.4-1.3	9	1.4	0.6-2.7	3	0.3	0.1-0.8
1985	17	1.2	0.7-2.0	9	1.4	0.5-2.9	8	1.1	0.4-2.1	12	0.9	0.4-1.5	9	1.9	0.8-3.6	3	0.3	0.1-0.9
1986	14	1.0	0.5-1.7	8	1.8	0.6-3.8	6	0.6	0.2-1.4	14	1.0	0.5-1.7	10	1.7	0.8-3.3	4	0.5	0.1-1.3
1987	12	0.7	0.4-1.3	5	0.7	0.2-1.7	7	0.7	0.3-1.5	8	0.5	0.2-1.1	3	0.5	0.1-1.4	5	0.5	0.2-1.3
1988	15	1.0	0.5-1.6	5	1.1	0.3-2.6	10	1.0	0.5-1.8	7	0.5	0.2-1.1	2	0.6	0.0-2.0	5	0.6	0.2-1.3
1989	13	1.0	0.5-1.7	4	0.8	0.2-2.1	9	1.1	0.5-2.1	12	0.8	0.4-1.4	4	0.5	0.1-1.4	8	0.9	0.4-1.8
1990	17	1.0	0.5-1.6	9	1.4	0.5-2.9	8	0.8	0.3-1.5	6	0.4	0.1-0.9	3	0.6	0.1-2.0	3	0.3	0.1-0.9
1991	22	1.2	0.7-1.9	12	2.2	1.0-4.1	10	0.8	0.4-1.5	9	0.6	0.3-1.1	6	1.0	0.3-2.4	3	0.3	0.1-0.9
1992	10	0.5	0.2-1.0	4	0.5	0.1-1.4	6	0.5	0.2-1.2	6	0.3	0.1-0.7	2	0.2	0.0-1.0	4	0.4	0.1-0.9
1993	17	1.1	0.6-1.7	11	1.6	0.7-2.9	6	0.7	0.2-1.4	5	0.3	0.1-0.7	2	0.2	0.0-1.0	3	0.3	0.1-0.8
1994	13	0.7	0.4-1.3	3	0.7	0.1-1.9	10	0.9	0.4-1.6	3	0.3	0.1-0.7	1	0.2	0.0-1.1	2	0.2	0.0-0.8
1995	19	1.1	0.7-1.8	4	0.5	0.1-1.4	15	1.4	0.8-2.4	2	0.2	0.0-0.6	0	0.0	0.0-0.5	2	0.2	0.0-0.8
1996	18	1.2	0.7-1.8	5	0.8	0.2-1.8	13	1.3	0.7-2.2	9	0.6	0.3-1.2	3	0.5	0.1-1.6	6	0.6	0.2-1.3
1997	31	1.7	1.2-2.5	14	2.0	1.1-3.5	17	1.5	0.9-2.5	7	0.4	0.2-0.9	3	0.3	0.1-1.1	4	0.4	0.1-1.0
1998	20	1.1	0.7-1.8	10	1.6	0.7-3.1	10	0.9	0.4-1.6	8	0.4	0.2-0.9	2	0.2	0.0-0.9	6	0.5	0.2-1.2
1999	9	0.5	0.2-0.9	4	0.6	0.2-1.5	5	0.4	0.1-1.0	7	0.4	0.1-0.8	4	0.7	0.1-1.8	3	0.3	0.1-0.7
2000	13	0.7	0.4-1.2	7	1.0	0.4-2.0	6	0.5	0.2-1.1	10	0.6	0.3-1.1	5	1.1	0.3-2.5	5	0.4	0.1-1.0
2001	23	1.1	0.7-1.7	9	1.0	0.4-2.0	14	1.2	0.6-1.9	2	0.1	0.0-0.4	1	0.1	0.0-0.6	1	0.1	0.0-0.5
2002	17	0.8	0.5-1.3	5	0.5	0.2-1.2	12	1.0	0.5-1.7	5	0.3	0.1-0.6	2	0.2	0.0-0.8	3	0.3	0.1-0.7
2003	12	0.6	0.3-1.0	3	0.5	0.1-1.4	9	0.7	0.3-1.3	5	0.2	0.1-0.5	1	0.1	0.0-0.6	4	0.3	0.1-0.8
2004	16	0.8	0.4-1.3	8	1.2	0.5-2.4	8	0.6	0.3-1.2	3	0.1	0.0-0.4	1	0.1	0.0-0.6	2	0.2	0.0-0.6
2005	20	0.9	0.6-1.5	7	0.8	0.3-1.6	13	1.0	0.5-1.7	4	0.2	0.0-0.5	3	0.3	0.1-0.9	1	0.1	0.0-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	28	1.3	0.9-2.0	13	1.9	0.9-3.3	15	1.1	0.6-1.9	7	0.3	0.1-0.6	4	0.4	0.1-1.1	3	0.2	0.0-0.7
2007	32	1.5	1.0-2.1	11	1.4	0.7-2.6	21	1.6	1.0-2.5	15	0.7	0.4-1.2	6	0.9	0.3-2.0	9	0.7	0.3-1.3
2008	26	1.2	0.8-1.7	11	1.4	0.6-2.5	15	1.1	0.6-1.9	8	0.4	0.1-0.7	3	0.3	0.1-0.9	5	0.4	0.1-0.9
2009	21	0.9	0.5-1.4	9	0.9	0.4-1.7	12	0.9	0.4-1.5	4	0.2	0.0-0.4	3	0.3	0.1-0.9	1	0.1	0.0-0.4
2010	20	0.8	0.5-1.3	8	0.9	0.4-1.8	12	0.8	0.4-1.5	6	0.3	0.1-0.6	2	0.3	0.0-1.1	4	0.3	0.1-0.7
2011	31	1.3	0.9-1.9	15	1.7	0.9-2.8	16	1.2	0.7-1.9	11	0.5	0.3-0.9	5	0.6	0.2-1.5	6	0.5	0.2-1.0
2012	24	1.0	0.6-1.5	10	1.1	0.5-2.0	14	0.9	0.5-1.6	8	0.3	0.1-0.7	3	0.4	0.1-1.1	5	0.3	0.1-0.8
2013	27	1.2	0.8-1.7	9	1.1	0.5-2.1	18	1.2	0.7-1.9	9	0.3	0.1-0.6	5	0.4	0.1-1.0	4	0.3	0.1-0.7
2014	22	0.9	0.5-1.3	9	0.8	0.4-1.6	13	0.9	0.5-1.6	6	0.3	0.1-0.6	2	0.2	0.0-0.6	4	0.3	0.1-0.7
2015	22	0.8	0.5-1.3	11	1.2	0.6-2.2	11	0.7	0.3-1.3	9	0.3	0.1-0.6	5	0.5	0.1-1.2	4	0.3	0.1-0.7
2016	21	0.8	0.5-1.2	8	0.7	0.3-1.5	13	0.8	0.4-1.4	4	0.2	0.0-0.4	1	0.1	0.0-0.5	3	0.2	0.0-0.6
2017	22	0.9	0.6-1.4	11	1.1	0.5-2.0	11	0.8	0.4-1.4	1	0.0	0.0-0.2	1	0.1	0.0-0.5	0	0.0	0.0-0.3
2018	27	1.0	0.7-1.5	9	0.8	0.4-1.6	18	1.2	0.7-1.9	4	0.1	0.0-0.4	0	0.0	0.0-0.4	4	0.2	0.1-0.6
2019	28	1.1	0.7-1.6	8	0.9	0.4-1.8	20	1.3	0.8-2.0	7	0.3	0.1-0.6	5	0.4	0.1-1.0	2	0.1	0.0-0.5
2020	24	0.9	0.6-1.4	10	0.9	0.4-1.6	14	0.9	0.5-1.6	5	0.2	0.1-0.5	2	0.2	0.0-0.6	3	0.2	0.0-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	537	74.1	67.6-81.1	248	36.3	31.6-41.5
1977	546	72.9	66.6-79.7	230	31.5	27.3-36.1
1978	608	80.2	73.7-87.2	258	35.1	30.7-39.9
1979	630	80.8	74.3-87.7	229	30.1	26.2-34.5
1980	666	82.1	75.7-88.8	278	35.7	31.4-40.3
1981	651	79.6	73.3-86.2	273	34.5	30.3-39.1
1982	703	83.9	77.5-90.5	277	33.1	29.2-37.4
1983	710	81.3	75.2-87.7	299	35.1	31.0-39.4
1984	823	91.4	85.1-98.0	316	36.8	32.7-41.3
1985	814	88.7	82.5-95.2	251	27.7	24.3-31.4
1986	799	83.6	77.7-89.7	303	32.1	28.5-36.0
1987	890	90.0	84.1-96.3	312	32.1	28.6-36.0
1988	955	95.3	89.2-101.7	321	32.2	28.7-36.0
1989	913	89.7	83.9-95.9	319	32.1	28.6-35.9
1990	938	91.0	85.1-97.1	340	33.9	30.4-37.8
1991	1,066	102.1	96.0-108.6	321	31.3	27.9-35.0
1992	1,009	95.6	89.7-101.8	330	31.7	28.3-35.3
1993	1,017	94.3	88.5-100.3	314	30.4	27.1-34.0
1994	1,089	99.3	93.4-105.4	380	34.7	31.2-38.4
1995	1,113	99.4	93.6-105.5	403	36.7	33.2-40.5
1996	1,209	106.2	100.3-112.5	381	34.1	30.7-37.7
1997	1,220	105.1	99.3-111.3	375	33.1	29.8-36.7
1998	1,366	116.1	110.0-122.5	369	32.0	28.8-35.4
1999	1,206	100.0	94.4-105.8	375	31.8	28.7-35.2
2000	1,226	99.0	93.5-104.7	336	27.7	24.8-30.9
2001	1,211	96.1	90.7-101.7	395	32.1	29.0-35.4
2002	1,267	99.0	93.6-104.7	344	27.6	24.7-30.7
2003	1,315	101.4	96.0-107.1	349	27.4	24.6-30.5
2004	1,329	101.1	95.7-106.7	348	27.0	24.2-30.0
2005	1,342	101.2	95.8-106.8	392	29.7	26.8-32.8
2006	1,420	105.9	100.4-111.6	352	26.7	24.0-29.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1,460	108.2	102.7-113.9	334	24.7	22.1-27.5
2008	1,515	111.6	106.0-117.4	334	24.8	22.2-27.6
2009	1,536	111.6	106.0-117.4	332	24.3	21.8-27.1
2010	1,503	107.3	101.9-112.9	320	22.8	20.3-25.4
2011	1,719	119.9	114.2-125.8	355	25.2	22.6-28.0
2012	1,655	113.2	107.8-118.9	382	26.3	23.7-29.1
2013	1,776	120.2	114.6-126.0	314	21.0	18.7-23.5
2014	1,766	118.7	113.2-124.5	362	24.1	21.6-26.8
2015	1,829	120.1	114.5-125.8	333	21.9	19.5-24.4
2016	1,786	116.4	111.0-122.1	351	22.6	20.2-25.1
2017	1,819	117.6	112.1-123.3	327	20.6	18.4-23.0
2018	1,906	123.3	117.7-129.2	360	21.8	19.6-24.3
2019	2,085	134.4	128.5-140.5	377	23.3	21.0-25.9
2020	1,652	106.8	101.5-112.3	301	18.1	16.1-20.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	224	28.5	24.7-32.7	101	14.6	11.8-18.0
1977	247	31.7	27.7-36.1	102	14.1	11.4-17.4
1978	233	29.3	25.5-33.5	89	12.1	9.5-15.0
1979	250	30.7	26.9-34.9	81	10.9	8.5-13.7
1980	221	26.9	23.3-30.9	90	11.5	9.1-14.2
1981	199	22.2	19.1-25.6	81	9.3	7.3-11.7
1982	223	25.7	22.3-29.4	81	9.0	7.1-11.2
1983	202	22.2	19.1-25.6	76	8.5	6.6-10.7
1984	208	22.3	19.3-25.6	64	7.3	5.6-9.4
1985	192	20.0	17.2-23.1	71	7.6	5.9-9.6
1986	217	22.8	19.8-26.1	71	7.3	5.7-9.3
1987	198	19.3	16.7-22.3	58	5.8	4.4-7.6
1988	187	18.3	15.7-21.1	90	9.0	7.2-11.0
1989	180	16.5	14.1-19.2	50	5.0	3.7-6.7
1990	235	21.2	18.5-24.1	65	6.4	4.9-8.1
1991	216	19.5	17.0-22.4	63	5.8	4.4-7.5
1992	214	18.8	16.4-21.6	64	6.0	4.6-7.6
1993	201	17.7	15.3-20.3	76	6.8	5.4-8.5
1994	217	19.4	16.8-22.2	64	5.8	4.4-7.4
1995	218	18.7	16.3-21.4	77	6.9	5.4-8.6
1996	220	18.4	16.0-21.0	59	5.1	3.8-6.5
1997	213	17.6	15.3-20.2	67	5.9	4.5-7.5
1998	207	16.5	14.3-18.9	62	5.2	4.0-6.7
1999	197	15.7	13.6-18.1	75	6.1	4.8-7.6
2000	247	19.7	17.3-22.3	72	5.7	4.5-7.2
2001	191	14.9	12.8-17.1	60	4.8	3.7-6.2
2002	199	15.3	13.2-17.6	58	4.6	3.5-5.9
2003	177	13.4	11.5-15.6	66	5.1	3.9-6.4
2004	167	12.5	10.6-14.5	50	3.8	2.8-5.1
2005	184	13.8	11.9-16.0	54	4.1	3.1-5.4
2006	172	12.9	11.0-15.0	57	4.4	3.3-5.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	191	14.3	12.3-16.5	72	5.4	4.2-6.8
2008	166	12.3	10.5-14.3	62	4.5	3.4-5.8
2009	162	11.6	9.9-13.6	64	4.5	3.4-5.7
2010	167	12.2	10.4-14.3	47	3.3	2.4-4.4
2011	165	11.8	10.0-13.7	67	4.8	3.7-6.1
2012	158	11.3	9.6-13.3	48	3.3	2.4-4.4
2013	153	10.7	9.1-12.6	60	4.0	3.1-5.2
2014	147	10.3	8.7-12.2	62	4.2	3.2-5.5
2015	150	10.2	8.6-12.0	43	2.8	2.0-3.8
2016	155	10.6	9.0-12.5	52	3.5	2.6-4.6
2017	158	10.7	9.1-12.6	39	2.5	1.8-3.5
2018	141	9.5	8.0-11.3	49	3.0	2.2-4.1
2019	157	10.8	9.1-12.7	41	2.7	1.9-3.7
2020	117	7.7	6.3-9.3	47	2.9	2.1-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals**

Corpus uterus and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	115	16.0	13.1-19.4	55	8.7	6.4-11.4
1977	107	15.2	12.3-18.5	51	7.2	5.3-9.5
1978	116	15.9	13.0-19.2	60	9.5	7.1-12.3
1979	107	13.9	11.3-16.9	52	7.3	5.3-9.6
1980	135	17.5	14.6-20.9	61	8.1	6.1-10.5
1981	113	14.1	11.5-17.1	51	6.2	4.5-8.2
1982	131	15.1	12.5-18.0	47	5.7	4.1-7.6
1983	132	15.2	12.7-18.2	61	7.3	5.6-9.5
1984	154	17.9	15.1-21.1	59	7.0	5.2-9.0
1985	163	17.7	15.0-20.7	68	7.8	6.0-10.0
1986	111	11.3	9.3-13.7	53	5.9	4.4-7.8
1987	161	16.8	14.2-19.6	44	5.0	3.6-6.7
1988	170	16.7	14.2-19.5	63	6.5	4.9-8.3
1989	147	14.2	12.0-16.8	52	4.9	3.7-6.5
1990	154	15.3	12.9-17.9	66	6.7	5.2-8.6
1991	208	20.1	17.4-23.0	60	6.0	4.6-7.8
1992	179	16.8	14.4-19.4	64	6.3	4.8-8.0
1993	205	18.9	16.4-21.7	69	6.8	5.2-8.6
1994	213	19.1	16.6-21.9	79	7.5	5.9-9.3
1995	207	18.6	16.1-21.4	59	5.5	4.2-7.1
1996	227	20.0	17.4-22.8	86	8.0	6.4-9.9
1997	252	22.0	19.4-24.9	97	8.7	7.0-10.6
1998	250	21.4	18.8-24.2	96	8.5	6.9-10.4
1999	269	22.7	20.1-25.6	90	7.8	6.2-9.6
2000	281	23.2	20.5-26.0	80	6.8	5.4-8.4
2001	304	24.2	21.6-27.1	99	8.2	6.7-10.0
2002	290	22.9	20.3-25.7	81	6.5	5.2-8.1
2003	312	24.0	21.4-26.9	104	8.2	6.7-9.9
2004	342	26.2	23.4-29.1	106	8.5	6.9-10.2
2005	352	26.7	23.9-29.6	111	8.5	7.0-10.3
2006	374	28.0	25.2-31.0	107	8.3	6.8-10.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals**

Corpus uterus and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	341	25.3	22.7-28.2	95	7.3	5.9-8.9
2008	460	33.1	30.2-36.3	115	8.7	7.1-10.4
2009	429	30.4	27.5-33.4	120	8.8	7.2-10.5
2010	450	31.4	28.5-34.4	114	8.2	6.7-9.8
2011	468	32.0	29.1-35.1	145	10.2	8.6-12.1
2012	501	33.2	30.3-36.3	131	8.9	7.4-10.6
2013	509	33.0	30.2-36.1	125	8.2	6.8-9.8
2014	533	33.5	30.7-36.6	174	11.1	9.5-12.9
2015	558	34.7	31.8-37.7	166	10.2	8.7-12.0
2016	580	35.5	32.6-38.6	173	10.6	9.0-12.3
2017	574	34.7	31.9-37.7	155	9.4	8.0-11.0
2018	589	34.7	31.9-37.7	185	10.9	9.4-12.7
2019	604	35.6	32.8-38.7	198	11.6	10.0-13.4
2020	519	29.7	27.1-32.4	172	9.7	8.3-11.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	76	10.8	8.4-13.7	47	6.7	4.8-9.0
1977	91	11.8	9.3-14.6	56	7.7	5.7-10.2
1978	77	10.4	8.1-13.2	59	7.8	5.8-10.2
1979	95	12.4	9.9-15.4	47	6.0	4.4-8.1
1980	104	13.1	10.6-16.1	76	10.3	8.0-13.1
1981	91	10.5	8.4-12.9	48	5.6	4.1-7.5
1982	79	9.4	7.4-11.7	51	6.3	4.6-8.3
1983	122	13.3	10.9-16.0	72	8.5	6.6-10.8
1984	119	13.2	10.9-15.9	66	7.4	5.7-9.5
1985	113	11.2	9.2-13.6	77	8.4	6.6-10.6
1986	114	12.0	9.8-14.5	63	6.6	5.0-8.5
1987	92	9.6	7.7-11.8	55	5.7	4.3-7.5
1988	112	11.5	9.4-13.8	55	5.7	4.3-7.5
1989	95	9.0	7.3-11.1	62	6.2	4.7-7.9
1990	128	12.5	10.4-14.9	79	8.0	6.3-10.0
1991	116	11.1	9.1-13.3	72	7.0	5.4-8.8
1992	153	14.4	12.1-16.9	89	8.5	6.8-10.4
1993	116	10.4	8.6-12.5	95	9.2	7.4-11.3
1994	162	14.8	12.6-17.3	90	8.7	6.9-10.7
1995	138	12.3	10.3-14.5	99	9.2	7.4-11.2
1996	111	9.6	7.9-11.6	58	5.2	3.9-6.7
1997	147	12.6	10.6-14.9	75	6.8	5.3-8.5
1998	108	9.3	7.6-11.2	79	7.0	5.5-8.7
1999	128	10.9	9.0-12.9	74	6.4	5.0-8.1
2000	121	10.0	8.3-12.0	76	6.5	5.1-8.1
2001	129	10.4	8.7-12.4	84	6.9	5.5-8.5
2002	142	11.1	9.4-13.1	79	6.5	5.1-8.1
2003	131	10.1	8.4-12.0	75	6.0	4.7-7.5
2004	138	10.5	8.9-12.5	78	6.2	4.9-7.8
2005	136	10.4	8.7-12.3	88	6.8	5.5-8.4
2006	117	8.7	7.2-10.4	80	6.2	4.9-7.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	128	9.5	7.9-11.4	81	6.1	4.8-7.6
2008	134	9.8	8.2-11.6	82	5.9	4.7-7.3
2009	140	10.1	8.5-11.9	80	5.8	4.6-7.3
2010	139	10.1	8.5-11.9	88	6.3	5.1-7.8
2011	138	9.6	8.1-11.4	86	6.2	4.9-7.6
2012	155	10.6	9.0-12.5	97	6.7	5.4-8.2
2013	146	10.0	8.4-11.8	71	4.7	3.6-5.9
2014	156	10.3	8.8-12.1	84	5.5	4.4-6.9
2015	139	9.1	7.6-10.7	92	5.9	4.8-7.3
2016	146	9.3	7.8-11.0	97	6.0	4.8-7.4
2017	159	10.3	8.7-12.1	81	5.0	4.0-6.2
2018	153	9.9	8.3-11.6	104	6.2	5.1-7.6
2019	159	10.3	8.7-12.1	107	6.4	5.2-7.7
2020	151	9.4	7.9-11.1	90	5.1	4.1-6.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	396	118.4	105.2-132.6	144	49.8	40.7-60.0
1977	422	122.4	109.3-136.4	147	45.9	38.0-54.7
1978	464	133.4	119.8-147.9	167	54.9	45.8-64.9
1979	482	132.4	119.4-146.2	189	60.6	51.2-70.9
1980	587	157.3	143.3-172.1	218	67.4	57.8-77.9
1981	569	149.2	135.8-163.4	232	70.7	60.9-81.4
1982	542	138.8	126.1-152.3	202	59.6	50.9-69.1
1983	526	129.1	117.1-141.9	215	59.5	51.1-68.7
1984	604	142.8	130.4-155.9	219	60.5	52.1-69.7
1985	623	138.0	126.2-150.4	254	68.3	59.4-78.0
1986	639	141.1	129.3-153.5	254	66.4	57.8-75.8
1987	659	138.2	126.8-150.3	265	64.9	56.6-73.8
1988	702	152.4	140.2-165.2	210	53.0	45.6-61.2
1989	746	154.7	142.8-167.3	233	55.4	47.9-63.6
1990	844	169.9	157.6-182.8	288	69.7	61.2-78.9
1991	899	175.0	162.7-188.0	255	59.5	51.8-67.9
1992	1,129	209.8	196.7-223.4	305	70.7	62.3-79.8
1993	1,228	218.2	205.2-231.8	274	64.7	56.7-73.4
1994	1,353	231.2	218.1-244.8	296	67.6	59.5-76.3
1995	1,482	240.6	227.6-254.0	322	69.7	61.8-78.3
1996	1,463	234.1	221.5-247.2	293	61.2	53.9-69.0
1997	1,581	246.5	233.7-259.7	297	61.0	53.8-68.8
1998	1,616	244.5	232.0-257.5	288	57.5	50.7-64.9
1999	1,905	277.1	264.0-290.5	319	62.3	55.2-69.9
2000	2,121	296.1	283.0-309.7	296	55.7	49.2-62.6
2001	1,983	270.7	258.3-283.5	315	58.4	51.8-65.5
2002	1,986	265.0	252.8-277.5	323	56.2	49.9-62.9
2003	1,981	258.4	246.6-270.6	266	46.1	40.5-52.3
2004	2,065	261.7	250.0-273.8	277	48.4	42.6-54.7
2005	2,044	251.5	240.2-263.1	263	44.1	38.7-50.0
2006	2,140	260.7	249.2-272.5	263	44.8	39.4-50.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	2,329	275.0	263.5-287.0	282	46.4	40.9-52.3
2008	2,280	259.3	248.3-270.7	262	43.2	37.9-48.9
2009	2,266	250.6	240.0-261.7	282	44.7	39.4-50.3
2010	2,202	234.8	224.7-245.3	268	40.3	35.4-45.5
2011	2,450	250.8	240.5-261.4	281	41.3	36.5-46.6
2012	2,243	223.5	214.0-233.3	271	38.4	33.8-43.3
2013	2,267	218.8	209.6-228.4	269	36.9	32.5-41.7
2014	2,024	189.4	180.9-198.2	283	35.2	31.1-39.7
2015	2,064	187.1	178.8-195.6	259	31.5	27.7-35.7
2016	2,202	195.7	187.4-204.4	266	31.2	27.4-35.2
2017	2,237	193.3	185.1-201.7	264	29.2	25.7-33.0
2018	2,200	186.3	178.3-194.5	279	30.5	26.9-34.4
2019	2,316	194.9	186.8-203.2	293	31.1	27.6-35.0
2020	1,806	149.6	142.5-156.9	260	26.6	23.4-30.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Testis**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	5	0.5	0.2-1.5	3	1.0	0.1-3.4
1977	4	0.8	0.2-2.3	2	0.2	0.0-1.2
1978	2	0.5	0.0-1.8	3	0.6	0.1-1.9
1979	6	0.7	0.2-1.8	0	0.0	0.0-0.9
1980	8	1.2	0.5-2.5	1	0.1	0.0-1.0
1981	6	0.7	0.3-1.8	2	0.3	0.0-1.3
1982	4	0.5	0.1-1.5	0	0.0	0.0-0.8
1983	5	0.8	0.2-2.0	6	1.2	0.3-2.9
1984	3	0.2	0.0-1.0	1	0.2	0.0-1.1
1985	12	1.8	0.8-3.4	1	0.1	0.0-0.8
1986	11	1.1	0.6-2.2	2	0.2	0.0-1.0
1987	8	0.7	0.3-1.5	1	0.1	0.0-0.7
1988	9	0.8	0.4-1.7	2	0.5	0.0-1.9
1989	13	1.5	0.8-2.7	1	0.1	0.0-0.8
1990	10	1.2	0.5-2.2	3	0.3	0.1-1.0
1991	7	0.9	0.3-2.0	0	0.0	0.0-0.6
1992	15	1.3	0.7-2.2	1	0.1	0.0-0.7
1993	10	1.0	0.5-2.0	2	0.2	0.0-0.8
1994	14	1.3	0.7-2.4	1	0.1	0.0-0.8
1995	5	0.4	0.1-1.1	0	0.0	0.0-0.5
1996	17	1.5	0.9-2.5	0	0.0	0.0-0.5
1997	7	0.6	0.3-1.4	1	0.1	0.0-0.7
1998	15	1.4	0.8-2.4	1	0.1	0.0-0.6
1999	14	1.3	0.7-2.3	0	0.0	0.0-0.5
2000	24	2.3	1.5-3.5	3	0.4	0.1-1.1
2001	15	1.3	0.8-2.3	1	0.2	0.0-0.9
2002	13	1.2	0.6-2.1	2	0.2	0.0-0.8
2003	18	1.7	1.0-2.7	1	0.1	0.0-0.6
2004	11	1.0	0.5-1.8	1	0.1	0.0-0.6
2005	21	2.0	1.3-3.2	2	0.2	0.0-0.7
2006	24	2.3	1.4-3.4	1	0.1	0.0-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Testis**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	21	2.0	1.2-3.1	1	0.1	0.0-0.6
2008	11	1.0	0.5-1.8	0	0.0	0.0-0.4
2009	14	1.3	0.7-2.2	1	0.1	0.0-0.6
2010	19	1.7	1.0-2.8	1	0.1	0.0-0.5
2011	20	1.8	1.1-2.8	3	0.3	0.1-0.8
2012	18	1.7	1.0-2.6	1	0.1	0.0-0.5
2013	17	1.4	0.8-2.3	0	0.0	0.0-0.4
2014	28	2.6	1.7-3.8	1	0.1	0.0-0.5
2015	25	2.1	1.4-3.2	1	0.1	0.0-0.6
2016	26	2.3	1.5-3.4	2	0.3	0.0-0.9
2017	32	2.8	1.9-4.0	4	0.4	0.1-1.1
2018	18	1.6	0.9-2.6	0	0.0	0.0-0.4
2019	20	1.7	1.0-2.7	1	0.1	0.0-0.6
2020	22	2.2	1.4-3.4	0	0.0	0.0-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	117	12.5	10.1-15.2	76	20.9	15.8-26.9	41	7.5	5.2-10.3	54	6.0	4.4-8.1	32	8.2	5.3-12.0	22	4.6	2.7-7.0
1977	102	10.3	8.2-12.7	65	18.0	13.3-23.5	37	5.9	4.0-8.2	46	5.1	3.6-6.9	27	8.7	5.2-13.3	19	3.2	1.9-5.1
1978	99	9.4	7.5-11.5	64	16.5	12.3-21.6	35	5.3	3.6-7.4	44	4.7	3.3-6.4	27	8.3	5.2-12.4	17	2.8	1.6-4.5
1979	112	11.0	8.9-13.4	67	17.7	13.2-23.1	45	7.2	5.2-9.7	53	5.7	4.2-7.6	26	7.5	4.7-11.2	27	4.7	3.0-6.8
1980	141	12.6	10.4-15.0	91	21.5	16.9-26.9	50	7.3	5.3-9.8	46	4.9	3.5-6.7	25	7.3	4.4-11.1	21	3.6	2.2-5.6
1981	116	10.1	8.2-12.3	67	15.7	11.6-20.5	49	7.0	5.1-9.3	62	5.6	4.2-7.3	33	7.9	5.2-11.5	29	4.2	2.8-6.2
1982	128	10.7	8.8-12.9	68	16.1	12.1-20.7	60	7.8	5.9-10.2	57	5.0	3.7-6.5	25	6.0	3.7-9.1	32	4.4	2.9-6.3
1983	127	10.0	8.2-12.0	69	14.7	11.1-19.1	58	7.2	5.4-9.4	49	4.3	3.1-5.8	24	5.9	3.6-8.9	25	3.5	2.2-5.1
1984	112	8.4	6.8-10.1	71	14.4	11.0-18.5	41	5.0	3.5-6.8	53	4.2	3.1-5.5	31	6.1	4.0-8.9	22	3.0	1.8-4.5
1985	147	11.6	9.7-13.7	82	17.3	13.3-22.0	65	8.4	6.4-10.8	59	4.9	3.7-6.4	33	8.3	5.5-11.8	26	3.3	2.1-4.9
1986	129	10.1	8.3-12.0	85	19.9	15.5-25.0	44	5.2	3.7-7.0	48	3.9	2.9-5.3	25	6.6	4.0-9.8	23	2.8	1.7-4.2
1987	129	9.6	8.0-11.5	85	17.4	13.6-21.9	44	5.4	3.9-7.3	59	4.7	3.5-6.0	34	8.0	5.3-11.3	25	3.0	1.9-4.5
1988	135	10.0	8.3-11.9	81	16.3	12.6-20.7	54	6.5	4.9-8.5	52	3.9	2.9-5.1	31	7.5	4.8-10.9	21	2.3	1.4-3.5
1989	123	8.9	7.3-10.6	72	13.8	10.5-17.7	51	6.0	4.4-7.9	49	3.8	2.8-5.1	27	6.1	3.8-9.0	22	2.7	1.7-4.1
1990	136	9.7	8.0-11.5	85	17.7	13.8-22.1	51	5.7	4.2-7.4	50	3.7	2.7-4.9	25	5.3	3.3-7.9	25	2.8	1.8-4.2
1991	139	9.3	7.7-11.0	83	16.0	12.5-20.1	56	5.9	4.4-7.6	40	2.9	2.1-4.0	28	6.5	4.2-9.5	12	1.3	0.7-2.2
1992	154	10.0	8.5-11.8	82	15.6	12.1-19.6	72	7.3	5.7-9.2	45	3.3	2.4-4.4	21	4.9	2.9-7.6	24	2.6	1.7-3.9
1993	148	9.8	8.2-11.5	77	15.2	11.7-19.2	71	7.2	5.6-9.0	58	3.8	2.8-4.9	34	6.5	4.3-9.2	24	2.4	1.5-3.6
1994	153	9.8	8.2-11.5	81	14.8	11.5-18.6	72	7.1	5.5-9.0	57	3.9	2.9-5.1	27	5.4	3.4-8.0	30	3.1	2.1-4.5
1995	131	8.3	6.9-9.8	85	15.4	12.0-19.3	46	4.6	3.3-6.1	53	3.5	2.6-4.6	25	4.8	2.9-7.3	28	2.9	1.9-4.1
1996	150	9.2	7.8-10.9	89	15.5	12.2-19.3	61	5.9	4.5-7.6	51	3.4	2.5-4.4	28	5.5	3.5-8.0	23	2.3	1.5-3.5
1997	148	9.1	7.6-10.7	76	13.1	10.1-16.6	72	6.9	5.4-8.7	56	3.7	2.8-4.8	26	5.1	3.2-7.6	30	3.0	2.0-4.3
1998	171	10.0	8.5-11.6	99	16.0	12.8-19.6	72	6.7	5.2-8.4	63	4.0	3.1-5.1	29	5.6	3.6-8.1	34	3.3	2.3-4.6
1999	169	9.4	8.0-11.0	109	16.7	13.5-20.3	60	5.3	4.1-6.9	49	2.9	2.1-3.8	23	3.9	2.4-6.0	26	2.3	1.5-3.4
2000	166	9.3	8.0-10.9	91	14.6	11.6-18.1	75	6.6	5.2-8.2	66	3.7	2.9-4.8	36	5.8	3.9-8.1	30	2.7	1.8-3.8
2001	175	9.4	8.1-11.0	102	15.1	12.2-18.5	73	6.2	4.9-7.8	64	3.6	2.8-4.6	27	4.5	2.8-6.6	37	3.2	2.2-4.4
2002	175	9.5	8.1-11.0	100	15.5	12.4-19.0	75	6.3	4.9-7.9	47	2.6	1.9-3.5	18	2.7	1.5-4.3	29	2.5	1.7-3.6
2003	173	8.9	7.6-10.3	121	17.1	14.0-20.6	52	4.3	3.2-5.6	47	2.5	1.8-3.3	26	3.9	2.5-5.8	21	1.7	1.1-2.6
2004	191	9.8	8.5-11.3	119	17.1	14.0-20.7	72	5.9	4.6-7.4	57	3.0	2.2-3.8	29	4.2	2.7-6.2	28	2.2	1.5-3.3
2005	219	10.8	9.4-12.4	131	16.9	14.0-20.2	88	7.1	5.7-8.7	57	3.1	2.4-4.0	32	5.0	3.3-7.1	25	2.1	1.4-3.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	219	11.0	9.5-12.5	145	20.1	16.8-23.9	74	5.9	4.6-7.4	49	2.6	1.9-3.4	31	4.8	3.2-6.9	18	1.5	0.9-2.3
2007	176	8.8	7.5-10.2	103	14.6	11.8-17.9	73	5.8	4.5-7.3	47	2.5	1.8-3.3	22	3.6	2.2-5.5	25	2.0	1.3-3.0
2008	195	9.2	8.0-10.6	128	16.2	13.4-19.5	67	5.1	4.0-6.5	49	2.4	1.8-3.2	25	3.9	2.5-5.8	24	1.8	1.1-2.6
2009	220	10.4	9.1-11.9	137	17.7	14.7-21.1	83	6.4	5.0-7.9	46	2.4	1.7-3.1	22	3.5	2.2-5.4	24	1.8	1.1-2.7
2010	212	9.7	8.4-11.1	139	17.0	14.2-20.3	73	5.4	4.2-6.8	65	3.2	2.4-4.0	34	4.9	3.3-6.9	31	2.4	1.6-3.4
2011	229	10.2	8.9-11.6	153	18.6	15.6-21.9	76	5.4	4.2-6.7	61	2.8	2.2-3.6	32	4.3	2.9-6.1	29	2.1	1.4-3.0
2012	215	9.3	8.1-10.7	131	15.8	13.1-18.9	84	5.7	4.5-7.0	65	2.9	2.2-3.7	38	4.6	3.2-6.4	27	1.9	1.2-2.8
2013	265	11.3	10.0-12.8	171	19.6	16.7-23.0	94	6.5	5.3-8.0	58	2.5	1.9-3.3	33	4.2	2.8-5.9	25	1.6	1.1-2.5
2014	246	10.0	8.8-11.3	165	17.7	15.0-20.7	81	5.4	4.2-6.7	67	3.0	2.3-3.8	39	5.1	3.6-6.9	28	1.9	1.2-2.7
2015	263	10.5	9.2-11.9	170	18.0	15.2-21.0	93	6.0	4.9-7.4	58	2.4	1.8-3.1	31	3.6	2.4-5.1	27	1.7	1.1-2.5
2016	254	9.9	8.7-11.2	176	18.2	15.5-21.2	78	4.8	3.8-6.1	67	2.7	2.1-3.5	42	4.8	3.4-6.5	25	1.5	1.0-2.3
2017	308	11.9	10.6-13.4	203	21.1	18.2-24.3	105	6.6	5.4-8.0	62	2.5	1.9-3.3	37	4.3	3.0-5.9	25	1.6	1.0-2.4
2018	257	9.6	8.4-10.8	172	16.8	14.3-19.6	85	5.2	4.2-6.5	63	2.4	1.8-3.1	33	3.4	2.3-4.8	30	1.8	1.2-2.6
2019	269	9.8	8.6-11.1	184	17.8	15.3-20.7	85	5.1	4.0-6.3	62	2.2	1.7-2.9	40	4.0	2.8-5.4	22	1.2	0.8-1.9
2020	244	8.7	7.7-9.9	159	14.8	12.5-17.3	85	5.0	4.0-6.2	50	1.8	1.3-2.4	34	3.6	2.5-5.0	16	0.9	0.5-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	65	5.9	4.4-7.6	38	7.3	4.8-10.5	27	4.8	3.0-7.0	37	3.3	2.2-4.7	22	4.7	2.6-7.6	15	2.4	1.3-4.1
1977	47	3.5	2.5-4.7	33	6.5	4.2-9.6	14	1.6	0.9-2.8	22	1.7	1.0-2.7	16	2.9	1.6-4.9	6	0.9	0.3-2.0
1978	81	6.2	4.8-7.8	55	10.3	7.5-13.7	26	3.5	2.2-5.2	39	3.0	2.0-4.2	28	5.4	3.3-8.2	11	1.4	0.7-2.6
1979	58	4.2	3.2-5.6	44	8.6	6.0-11.8	14	1.5	0.8-2.6	32	2.4	1.6-3.4	24	4.4	2.7-6.6	8	1.1	0.5-2.2
1980	68	4.9	3.7-6.2	36	6.6	4.5-9.3	32	3.8	2.6-5.5	26	1.8	1.2-2.7	21	3.7	2.2-5.7	5	0.6	0.2-1.4
1981	63	4.4	3.3-5.7	43	7.3	5.2-9.9	20	2.4	1.4-3.8	25	1.6	1.0-2.3	17	2.8	1.6-4.5	8	0.8	0.3-1.6
1982	74	5.1	4.0-6.5	47	8.2	5.9-11.0	27	3.1	2.0-4.6	27	1.9	1.2-2.9	16	3.0	1.6-5.1	11	1.3	0.6-2.3
1983	88	5.9	4.7-7.3	43	7.7	5.4-10.6	45	4.9	3.6-6.6	39	2.7	1.9-3.8	23	4.1	2.5-6.4	16	1.9	1.0-3.1
1984	97	6.4	5.1-7.8	62	10.4	7.8-13.6	35	3.9	2.7-5.4	40	2.8	2.0-3.9	23	4.2	2.6-6.5	17	2.0	1.1-3.2
1985	80	5.1	3.9-6.4	45	6.9	4.9-9.4	35	3.8	2.6-5.3	42	2.8	2.0-3.8	22	3.6	2.1-5.7	20	2.3	1.4-3.5
1986	100	6.1	4.9-7.5	59	8.9	6.6-11.6	41	4.3	3.1-5.9	53	3.6	2.7-4.8	32	5.7	3.7-8.2	21	2.4	1.4-3.7
1987	111	7.0	5.7-8.5	67	11.4	8.6-14.7	44	4.5	3.2-6.1	33	2.1	1.4-3.0	20	3.4	2.0-5.4	13	1.4	0.7-2.4
1988	101	6.2	5.0-7.6	57	8.5	6.3-11.2	44	4.6	3.3-6.2	39	2.5	1.8-3.4	25	4.1	2.6-6.1	14	1.5	0.8-2.5
1989	102	6.0	4.8-7.3	59	9.8	7.2-12.9	43	4.0	2.9-5.4	34	2.3	1.5-3.2	14	2.6	1.3-4.4	20	2.1	1.3-3.2
1990	108	6.6	5.4-8.0	61	9.8	7.2-12.9	47	4.7	3.5-6.3	38	2.4	1.7-3.3	20	3.8	2.2-6.1	18	1.8	1.0-2.8
1991	128	7.4	6.1-8.9	81	12.0	9.3-15.2	47	4.6	3.3-6.1	40	2.3	1.6-3.2	23	3.7	2.2-5.6	17	1.6	0.9-2.6
1992	115	6.8	5.6-8.2	74	11.2	8.6-14.4	41	4.1	2.9-5.6	34	2.1	1.5-3.0	20	3.3	1.9-5.3	14	1.5	0.8-2.5
1993	121	6.7	5.5-8.0	79	11.3	8.7-14.3	42	3.9	2.8-5.3	45	2.7	2.0-3.7	29	4.4	2.8-6.5	16	1.7	0.9-2.7
1994	121	6.5	5.4-7.8	61	8.6	6.4-11.2	60	5.3	4.0-6.8	31	1.7	1.2-2.5	17	2.5	1.4-4.1	14	1.3	0.7-2.2
1995	159	8.7	7.4-10.2	97	13.4	10.7-16.5	62	5.7	4.4-7.3	53	3.1	2.3-4.0	32	4.9	3.3-6.9	21	2.0	1.2-3.0
1996	148	7.8	6.5-9.2	94	12.5	9.9-15.4	54	4.8	3.6-6.2	41	2.2	1.6-3.0	26	3.3	2.1-4.9	15	1.4	0.8-2.4
1997	159	8.5	7.2-9.9	86	12.0	9.4-14.9	73	6.4	5.0-8.0	44	2.4	1.8-3.3	28	4.5	2.9-6.7	16	1.4	0.8-2.2
1998	150	7.8	6.5-9.1	90	11.5	9.1-14.3	60	5.3	4.0-6.8	43	2.3	1.6-3.1	27	3.5	2.2-5.2	16	1.5	0.8-2.4
1999	200	10.2	8.8-11.8	100	13.1	10.5-16.2	100	8.5	6.9-10.3	34	1.9	1.3-2.6	16	2.2	1.2-3.7	18	1.6	0.9-2.5
2000	168	8.6	7.3-10.0	87	11.1	8.8-13.8	81	6.9	5.4-8.5	40	2.1	1.5-2.9	22	3.1	1.8-4.8	18	1.5	0.9-2.4
2001	198	9.5	8.2-11.0	110	13.7	11.2-16.7	88	7.0	5.6-8.6	46	2.4	1.7-3.2	24	3.6	2.3-5.5	22	1.8	1.1-2.7
2002	189	9.0	7.7-10.4	108	13.2	10.7-16.1	81	6.4	5.1-8.0	44	2.2	1.6-3.0	30	4.0	2.6-5.8	14	1.2	0.7-2.0
2003	232	10.8	9.4-12.2	137	16.1	13.4-19.2	95	7.3	5.9-8.9	44	2.2	1.6-3.0	24	3.1	2.0-4.7	20	1.6	1.0-2.5
2004	228	10.5	9.1-12.0	132	15.0	12.5-17.9	96	7.4	6.0-9.0	52	2.6	1.9-3.4	32	4.1	2.8-5.8	20	1.6	1.0-2.5
2005	253	11.5	10.1-13.1	131	14.4	12.0-17.3	122	9.4	7.8-11.2	39	1.8	1.3-2.5	23	2.8	1.7-4.3	16	1.3	0.7-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	259	11.8	10.4-13.3	151	17.1	14.4-20.2	108	8.2	6.7-9.9	57	2.8	2.1-3.6	30	3.8	2.5-5.5	27	2.2	1.4-3.2
2007	278	12.3	10.9-13.9	184	20.6	17.6-23.9	94	7.1	5.8-8.8	53	2.6	1.9-3.4	34	4.4	3.0-6.2	19	1.4	0.9-2.3
2008	273	12.2	10.7-13.7	142	16.1	13.4-19.1	131	9.8	8.2-11.6	47	2.1	1.6-2.9	26	3.2	2.0-4.7	21	1.6	1.0-2.4
2009	291	12.3	10.9-13.8	171	17.9	15.2-20.9	120	8.4	7.0-10.1	38	1.7	1.2-2.4	26	3.3	2.1-4.9	12	0.9	0.4-1.5
2010	333	14.1	12.6-15.7	212	22.4	19.4-25.8	121	8.8	7.2-10.5	51	2.4	1.8-3.1	31	4.0	2.7-5.8	20	1.5	0.9-2.3
2011	317	13.4	11.9-14.9	189	19.6	16.8-22.7	128	9.2	7.6-10.9	50	2.2	1.6-2.9	31	3.8	2.5-5.4	19	1.3	0.8-2.1
2012	326	13.3	11.9-14.9	181	17.8	15.2-20.7	145	10.2	8.6-12.0	47	1.9	1.4-2.6	24	2.6	1.6-4.0	23	1.5	1.0-2.3
2013	372	14.6	13.1-16.2	217	20.7	18.0-23.8	155	10.4	8.8-12.2	44	1.8	1.3-2.4	28	2.8	1.8-4.1	16	1.1	0.6-1.8
2014	377	14.5	13.1-16.1	249	23.2	20.3-26.4	128	8.4	7.0-10.0	46	1.9	1.3-2.5	27	2.9	1.9-4.3	19	1.2	0.7-2.0
2015	368	13.8	12.4-15.3	220	20.1	17.4-23.0	148	9.5	8.0-11.2	59	2.4	1.8-3.0	43	4.6	3.3-6.3	16	0.9	0.5-1.6
2016	403	15.0	13.6-16.6	241	22.3	19.5-25.5	162	10.3	8.8-12.1	50	2.0	1.5-2.6	31	3.1	2.1-4.5	19	1.2	0.7-2.0
2017	383	14.1	12.7-15.6	249	22.7	19.9-25.8	134	8.2	6.9-9.8	52	2.0	1.5-2.6	39	3.9	2.7-5.3	13	0.8	0.4-1.4
2018	364	13.1	11.8-14.5	229	20.4	17.8-23.3	135	8.1	6.8-9.6	56	2.0	1.5-2.6	34	3.1	2.1-4.4	22	1.2	0.8-1.9
2019	428	15.3	13.9-16.9	240	20.9	18.2-23.8	188	11.4	9.8-13.3	60	2.1	1.6-2.7	35	3.1	2.1-4.3	25	1.4	0.9-2.1
2020	328	11.8	10.5-13.2	188	15.9	13.6-18.4	140	8.7	7.3-10.3	50	1.9	1.4-2.5	34	3.2	2.2-4.6	16	1.0	0.6-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	46	3.0	2.2-4.1	25	3.8	2.4-5.9	21	2.4	1.5-3.8	36	2.6	1.8-3.7	21	3.7	2.2-5.9	15	1.9	1.0-3.3
1977	58	3.5	2.6-4.6	29	4.3	2.8-6.4	29	3.0	2.0-4.4	32	2.3	1.5-3.3	19	3.2	1.8-5.2	13	1.6	0.8-2.9
1978	33	2.3	1.6-3.3	18	3.0	1.7-5.0	15	1.9	1.0-3.1	30	2.0	1.3-2.9	15	2.2	1.2-3.9	15	1.8	1.0-3.1
1979	53	3.1	2.3-4.2	21	2.9	1.7-4.7	32	3.4	2.3-4.8	39	2.7	1.9-3.7	19	3.4	2.0-5.5	20	2.2	1.4-3.5
1980	57	3.9	2.9-5.2	31	5.1	3.3-7.6	26	3.1	2.0-4.6	38	2.6	1.8-3.7	22	3.9	2.3-6.2	16	1.9	1.0-3.0
1981	41	2.5	1.8-3.5	23	2.9	1.8-4.5	18	2.1	1.2-3.4	29	2.3	1.5-3.4	11	1.8	0.9-3.4	18	2.5	1.4-4.0
1982	65	3.7	2.8-4.7	35	4.6	3.1-6.5	30	3.0	2.0-4.3	56	3.7	2.8-4.9	36	5.8	3.9-8.1	20	2.3	1.4-3.6
1983	60	3.6	2.7-4.7	34	4.7	3.2-6.8	26	2.8	1.8-4.1	50	3.2	2.3-4.2	18	2.7	1.6-4.4	32	3.4	2.3-4.9
1984	77	4.3	3.4-5.5	40	5.2	3.5-7.4	37	3.6	2.5-5.1	52	3.2	2.3-4.2	32	4.7	3.0-6.8	20	2.1	1.3-3.3
1985	80	4.6	3.6-5.8	34	5.1	3.4-7.4	46	4.4	3.2-6.0	45	2.6	1.9-3.6	27	4.2	2.7-6.2	18	1.7	1.0-2.7
1986	75	4.1	3.2-5.2	47	5.8	4.2-8.0	28	2.8	1.8-4.1	54	3.1	2.3-4.0	30	4.1	2.7-5.9	24	2.4	1.5-3.6
1987	83	4.3	3.4-5.5	41	5.4	3.7-7.6	42	3.7	2.6-5.1	46	2.8	2.0-3.8	25	4.3	2.6-6.5	21	2.0	1.2-3.1
1988	64	3.3	2.5-4.3	37	4.8	3.2-6.9	27	2.4	1.6-3.5	37	2.0	1.4-2.8	16	2.3	1.2-4.0	21	1.9	1.2-2.9
1989	76	3.9	3.0-4.9	36	4.5	3.0-6.4	40	3.5	2.5-4.8	32	1.7	1.2-2.5	15	2.1	1.1-3.5	17	1.6	0.9-2.5
1990	81	4.3	3.3-5.3	36	4.3	3.0-6.1	45	4.1	3.0-5.6	37	2.1	1.5-3.0	21	3.3	1.9-5.3	16	1.5	0.9-2.5
1991	93	4.9	3.9-6.1	44	6.1	4.2-8.4	49	4.3	3.2-5.7	42	2.3	1.6-3.2	19	2.5	1.4-4.0	23	2.2	1.4-3.3
1992	93	4.8	3.8-5.9	47	6.1	4.4-8.4	46	4.0	2.9-5.4	39	2.4	1.7-3.3	20	3.2	1.8-5.1	19	1.9	1.1-3.0
1993	69	3.4	2.6-4.4	43	5.2	3.6-7.2	26	2.3	1.5-3.3	38	1.9	1.3-2.6	18	2.4	1.4-4.0	20	1.7	1.0-2.6
1994	88	4.1	3.3-5.1	49	5.5	4.0-7.4	39	3.2	2.3-4.4	51	2.6	1.9-3.4	24	3.2	1.9-4.9	27	2.3	1.5-3.3
1995	83	4.0	3.1-5.0	33	3.4	2.3-4.9	50	4.3	3.1-5.6	34	1.6	1.1-2.3	14	1.5	0.8-2.7	20	1.7	1.0-2.6
1996	98	4.8	3.9-5.9	42	5.0	3.6-7.0	56	4.7	3.5-6.1	33	1.6	1.1-2.3	12	1.6	0.7-2.9	21	1.7	1.1-2.6
1997	79	3.7	2.9-4.6	37	4.1	2.8-5.8	42	3.4	2.4-4.6	38	1.9	1.3-2.6	17	1.9	1.1-3.2	21	1.8	1.1-2.7
1998	97	4.4	3.5-5.4	45	4.5	3.2-6.1	52	4.2	3.1-5.5	32	1.6	1.1-2.3	15	1.8	1.0-3.1	17	1.4	0.8-2.3
1999	90	4.3	3.4-5.3	43	5.2	3.6-7.2	47	3.8	2.8-5.0	29	1.4	0.9-2.0	18	2.1	1.2-3.4	11	0.9	0.5-1.6
2000	97	4.5	3.6-5.5	49	5.6	4.0-7.5	48	3.8	2.8-5.0	41	2.0	1.4-2.8	18	2.1	1.2-3.4	23	1.9	1.2-2.9
2001	95	4.3	3.4-5.2	39	3.8	2.7-5.3	56	4.4	3.3-5.7	41	1.9	1.3-2.6	19	2.1	1.2-3.5	22	1.7	1.1-2.6
2002	85	3.9	3.1-4.8	46	5.2	3.7-7.1	39	3.1	2.2-4.2	42	1.9	1.4-2.6	16	1.7	0.9-2.9	26	2.0	1.3-3.0
2003	109	4.9	4.0-6.0	54	5.5	4.1-7.3	55	4.3	3.2-5.6	34	1.6	1.1-2.2	21	2.4	1.4-3.7	13	1.0	0.5-1.7
2004	89	4.0	3.2-4.9	49	5.3	3.9-7.1	40	3.1	2.2-4.2	49	2.2	1.6-2.9	27	2.9	1.9-4.2	22	1.7	1.1-2.6
2005	97	4.3	3.5-5.3	44	5.0	3.6-6.9	53	4.0	3.0-5.3	36	1.6	1.1-2.2	24	2.8	1.7-4.2	12	0.9	0.5-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	83	3.7	2.9-4.6	41	4.3	3.0-5.9	42	3.2	2.3-4.3	45	2.0	1.5-2.7	22	2.6	1.6-4.0	23	1.7	1.1-2.6
2007	102	4.3	3.5-5.3	53	5.2	3.9-6.9	49	3.7	2.7-4.9	33	1.5	1.0-2.1	14	1.5	0.8-2.6	19	1.4	0.8-2.2
2008	85	3.7	2.9-4.6	42	4.4	3.1-6.0	43	3.2	2.3-4.3	44	2.0	1.4-2.7	22	2.6	1.6-3.9	22	1.6	1.0-2.5
2009	91	3.9	3.2-4.9	45	4.6	3.3-6.2	46	3.5	2.5-4.6	39	1.7	1.2-2.3	22	2.4	1.5-3.7	17	1.2	0.7-2.0
2010	91	3.8	3.1-4.7	46	4.5	3.3-6.1	45	3.3	2.4-4.4	33	1.4	1.0-2.0	16	1.7	0.9-2.9	17	1.3	0.7-2.0
2011	105	4.4	3.6-5.3	52	5.4	4.0-7.1	53	3.7	2.8-4.9	49	2.1	1.6-2.8	22	2.3	1.4-3.5	27	2.0	1.3-2.9
2012	91	3.8	3.1-4.7	56	5.5	4.1-7.2	35	2.6	1.8-3.6	40	1.6	1.1-2.2	20	2.0	1.2-3.2	20	1.4	0.8-2.1
2013	117	4.7	3.9-5.7	56	5.0	3.8-6.6	61	4.5	3.4-5.8	47	1.9	1.4-2.6	23	2.3	1.4-3.5	24	1.7	1.1-2.5
2014	82	3.3	2.6-4.1	51	4.6	3.4-6.1	31	2.3	1.5-3.2	47	1.9	1.4-2.5	26	2.7	1.7-3.9	21	1.3	0.8-2.1
2015	95	3.9	3.1-4.8	48	4.5	3.3-6.0	47	3.5	2.6-4.7	37	1.4	1.0-2.0	21	2.0	1.2-3.1	16	1.1	0.6-1.8
2016	107	4.3	3.5-5.2	67	6.2	4.8-7.9	40	2.8	2.0-3.9	58	2.2	1.7-2.9	32	2.9	1.9-4.1	26	1.8	1.1-2.6
2017	91	3.7	3.0-4.6	45	4.3	3.1-5.9	46	3.2	2.3-4.3	38	1.5	1.0-2.0	25	2.3	1.5-3.5	13	0.8	0.4-1.4
2018	119	4.6	3.8-5.5	66	6.0	4.6-7.6	53	3.5	2.6-4.6	57	2.4	1.8-3.2	24	2.5	1.6-3.7	33	2.4	1.6-3.4
2019	127	4.9	4.0-5.8	67	6.0	4.6-7.6	60	4.0	3.0-5.2	67	2.5	2.0-3.2	42	3.9	2.8-5.3	25	1.6	1.0-2.4
2020	115	4.4	3.6-5.3	58	5.0	3.8-6.5	57	3.8	2.9-5.0	49	1.7	1.3-2.3	25	2.0	1.3-2.9	24	1.4	0.9-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	29	2.1	1.4-3.0	7	1.3	0.4-2.8	22	2.7	1.6-4.1	8	0.6	0.2-1.2	2	0.4	0.0-1.5	6	0.8	0.3-1.7
1977	33	2.1	1.4-3.1	1	0.2	0.0-1.2	32	3.5	2.3-5.1	4	0.3	0.1-0.9	2	0.3	0.0-1.4	2	0.3	0.0-1.1
1978	25	1.8	1.1-2.7	8	1.7	0.6-3.6	17	1.9	1.1-3.1	7	0.6	0.2-1.2	0	0.0	0.0-0.9	7	0.9	0.4-1.9
1979	37	3.1	2.1-4.3	9	2.0	0.7-4.1	28	3.8	2.4-5.6	6	0.4	0.1-0.9	2	0.3	0.0-1.4	4	0.4	0.1-1.1
1980	34	2.1	1.4-3.0	4	0.6	0.1-1.6	30	3.1	2.1-4.6	7	0.6	0.2-1.2	1	0.2	0.0-1.3	6	0.8	0.3-1.7
1981	27	2.1	1.3-3.1	4	0.6	0.2-1.7	23	3.0	1.9-4.6	5	0.5	0.1-1.2	0	0.0	0.0-0.8	5	0.8	0.2-1.8
1982	32	2.3	1.5-3.2	6	1.0	0.3-2.2	26	3.1	2.0-4.6	9	0.7	0.3-1.4	3	0.9	0.1-2.7	6	0.7	0.2-1.5
1983	29	1.8	1.2-2.6	4	0.5	0.1-1.5	25	2.6	1.7-3.9	6	0.5	0.2-1.1	1	0.3	0.0-1.4	5	0.6	0.2-1.5
1984	44	2.7	1.9-3.7	6	0.8	0.3-1.8	38	4.0	2.8-5.6	9	0.7	0.3-1.3	2	0.6	0.0-2.2	7	0.8	0.3-1.6
1985	49	2.9	2.1-3.9	16	2.3	1.3-3.8	33	3.3	2.2-4.7	7	0.5	0.2-1.0	3	0.5	0.1-1.5	4	0.5	0.1-1.2
1986	35	2.3	1.5-3.2	10	1.9	0.8-3.7	25	2.5	1.6-3.8	8	0.5	0.2-1.1	3	0.7	0.1-2.2	5	0.5	0.2-1.2
1987	53	3.1	2.3-4.1	10	1.5	0.7-3.0	43	4.1	2.9-5.6	8	0.7	0.3-1.3	2	0.6	0.1-2.0	6	0.7	0.3-1.5
1988	41	2.4	1.7-3.3	10	1.6	0.7-3.0	31	3.0	2.0-4.3	5	0.4	0.1-0.9	1	0.1	0.0-0.8	4	0.5	0.1-1.3
1989	37	2.2	1.5-3.0	4	0.7	0.2-1.7	33	3.1	2.1-4.5	8	0.7	0.3-1.4	1	0.2	0.0-1.0	7	0.9	0.4-1.9
1990	35	1.9	1.3-2.6	5	0.7	0.2-1.7	30	2.7	1.8-3.9	3	0.2	0.0-0.6	1	0.2	0.0-1.0	2	0.2	0.0-0.8
1991	41	2.2	1.6-3.1	4	0.7	0.2-1.9	37	3.2	2.3-4.5	7	0.5	0.2-1.1	1	0.3	0.0-1.4	6	0.7	0.2-1.4
1992	71	3.4	2.6-4.3	16	1.6	0.9-2.8	55	4.6	3.5-6.0	9	0.5	0.2-1.0	2	0.2	0.0-0.9	7	0.7	0.3-1.4
1993	53	2.8	2.1-3.7	10	1.3	0.6-2.5	43	3.8	2.7-5.1	5	0.4	0.1-0.8	1	0.1	0.0-0.8	4	0.4	0.1-1.1
1994	62	3.2	2.4-4.1	11	1.5	0.7-2.7	51	4.4	3.2-5.7	7	0.4	0.2-0.8	2	0.3	0.0-1.1	5	0.4	0.1-1.0
1995	68	3.3	2.5-4.2	11	1.4	0.7-2.6	57	4.6	3.5-6.0	5	0.3	0.1-0.8	0	0.0	0.0-0.5	5	0.5	0.2-1.2
1996	88	4.1	3.2-5.0	17	2.3	1.2-3.8	71	5.5	4.3-7.0	12	0.7	0.4-1.2	7	1.1	0.4-2.3	5	0.5	0.2-1.1
1997	89	4.1	3.3-5.1	18	2.0	1.2-3.2	71	5.6	4.3-7.1	9	0.5	0.2-0.9	4	0.6	0.2-1.7	5	0.4	0.1-1.0
1998	105	4.8	3.9-5.9	19	2.4	1.4-3.7	86	6.7	5.3-8.3	8	0.4	0.2-0.9	3	0.3	0.1-1.0	5	0.5	0.2-1.1
1999	101	4.7	3.8-5.8	15	1.6	0.9-2.7	86	6.8	5.4-8.4	11	0.6	0.3-1.1	2	0.2	0.0-0.8	9	0.8	0.4-1.5
2000	115	5.2	4.2-6.2	14	1.4	0.7-2.3	101	7.8	6.3-9.5	12	0.6	0.3-1.1	2	0.2	0.0-0.8	10	0.9	0.4-1.6
2001	115	5.1	4.2-6.1	21	2.2	1.4-3.5	94	7.1	5.7-8.6	15	0.7	0.4-1.2	3	0.3	0.1-1.0	12	1.0	0.5-1.7
2002	124	5.4	4.5-6.5	21	2.3	1.4-3.5	103	7.6	6.2-9.3	6	0.3	0.1-0.7	1	0.1	0.0-0.6	5	0.4	0.1-1.0
2003	113	4.9	4.0-5.9	15	1.7	0.9-2.8	98	7.2	5.8-8.8	4	0.2	0.1-0.6	0	0.0	0.0-0.5	4	0.3	0.1-0.9
2004	127	5.5	4.5-6.5	26	3.1	2.0-4.6	101	7.4	6.0-9.0	10	0.5	0.2-0.9	5	0.8	0.2-2.0	5	0.4	0.1-0.9
2005	188	8.1	7.0-9.3	33	3.5	2.4-5.0	155	11.4	9.7-13.4	16	0.7	0.4-1.2	5	0.6	0.2-1.4	11	0.8	0.4-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	157	6.8	5.8-8.0	20	2.3	1.4-3.5	137	10.1	8.5-12.0	9	0.4	0.2-0.8	2	0.2	0.0-0.8	7	0.5	0.2-1.1
2007	162	6.8	5.8-8.0	32	3.3	2.2-4.7	130	9.5	7.9-11.3	9	0.5	0.2-0.9	1	0.2	0.0-0.8	8	0.6	0.3-1.2
2008	215	9.0	7.8-10.3	42	4.4	3.1-6.1	173	12.4	10.6-14.4	11	0.6	0.3-1.0	3	0.5	0.1-1.5	8	0.6	0.3-1.2
2009	208	8.8	7.6-10.1	38	3.6	2.5-5.0	170	12.5	10.7-14.6	10	0.5	0.2-0.9	4	0.5	0.1-1.4	6	0.4	0.2-1.0
2010	245	10.2	8.9-11.5	34	3.7	2.5-5.2	211	15.0	13.0-17.2	14	0.6	0.3-1.1	7	0.9	0.3-1.8	7	0.5	0.2-1.1
2011	280	11.1	9.8-12.4	54	5.4	4.0-7.1	226	15.4	13.5-17.6	8	0.3	0.1-0.7	1	0.2	0.0-0.8	7	0.5	0.2-1.0
2012	251	10.1	8.9-11.4	32	3.2	2.2-4.6	219	15.2	13.2-17.4	9	0.4	0.2-0.8	3	0.4	0.1-1.2	6	0.4	0.1-0.9
2013	318	12.4	11.1-13.9	40	3.6	2.5-4.9	278	19.1	16.8-21.5	12	0.5	0.3-0.9	5	0.5	0.2-1.2	7	0.5	0.2-1.0
2014	298	11.9	10.6-13.4	44	4.3	3.1-5.8	254	17.7	15.6-20.1	17	0.7	0.4-1.1	2	0.2	0.0-0.7	15	1.0	0.6-1.7
2015	287	11.6	10.2-13.0	47	4.5	3.3-6.0	240	17.0	14.9-19.3	12	0.5	0.3-0.9	3	0.3	0.1-0.8	9	0.6	0.3-1.2
2016	271	10.6	9.3-11.9	49	4.5	3.3-6.0	222	15.1	13.1-17.3	10	0.4	0.2-0.7	1	0.1	0.0-0.5	9	0.5	0.2-1.0
2017	250	9.8	8.6-11.1	41	4.0	2.8-5.4	209	14.3	12.4-16.5	17	0.7	0.4-1.1	5	0.5	0.2-1.2	12	0.8	0.4-1.4
2018	239	9.3	8.2-10.6	42	3.9	2.8-5.3	197	13.5	11.6-15.6	12	0.4	0.2-0.7	6	0.5	0.2-1.2	6	0.3	0.1-0.8
2019	267	10.5	9.2-11.9	55	4.8	3.6-6.3	212	15.0	13.0-17.2	11	0.4	0.2-0.7	0	0.0	0.0-0.4	11	0.7	0.3-1.2
2020	215	8.5	7.4-9.8	42	4.0	2.8-5.4	173	12.2	10.4-14.3	15	0.5	0.3-0.9	4	0.3	0.1-0.9	11	0.6	0.3-1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	33	2.5	1.6-3.8	16	3.8	1.6-7.0	17	2.0	1.1-3.3	15	0.9	0.5-1.6	9	1.2	0.5-2.6	6	0.7	0.3-1.6
1977	30	1.8	1.2-2.7	16	2.3	1.3-4.0	14	1.5	0.8-2.6	15	1.2	0.6-2.1	4	1.2	0.2-3.5	11	1.4	0.7-2.5
1978	29	1.8	1.1-2.7	14	1.9	0.9-3.5	15	1.7	0.9-2.9	12	0.7	0.3-1.2	6	0.8	0.3-2.1	6	0.6	0.2-1.4
1979	40	2.4	1.7-3.4	25	3.6	2.2-5.5	15	1.6	0.9-2.7	17	1.0	0.6-1.7	11	1.6	0.8-3.1	6	0.6	0.2-1.4
1980	46	2.4	1.7-3.2	22	2.5	1.6-4.0	24	2.2	1.4-3.3	9	0.5	0.2-1.0	4	0.6	0.2-1.8	5	0.4	0.1-1.1
1981	41	2.2	1.6-3.1	25	3.3	2.0-5.2	16	1.5	0.8-2.5	17	1.1	0.6-1.8	9	1.6	0.6-3.3	8	0.8	0.3-1.5
1982	41	2.1	1.5-2.9	25	2.9	1.8-4.5	16	1.5	0.8-2.4	15	0.8	0.4-1.4	7	0.9	0.3-2.0	8	0.7	0.3-1.5
1983	55	2.9	2.2-3.9	32	4.3	2.7-6.4	23	2.1	1.3-3.1	12	0.7	0.4-1.3	6	0.8	0.3-1.9	6	0.6	0.2-1.4
1984	43	2.2	1.6-3.0	28	3.2	2.0-4.8	15	1.5	0.8-2.5	18	1.0	0.6-1.6	16	1.9	1.0-3.2	2	0.3	0.0-0.9
1985	34	1.6	1.1-2.2	22	2.4	1.4-3.8	12	0.9	0.5-1.7	21	1.1	0.6-1.7	16	1.9	1.0-3.2	5	0.4	0.1-1.1
1986	44	2.3	1.6-3.1	30	3.6	2.4-5.3	14	1.3	0.7-2.2	11	0.6	0.3-1.1	6	0.8	0.3-1.8	5	0.5	0.1-1.2
1987	39	1.9	1.3-2.7	27	3.4	2.1-5.3	12	1.0	0.5-1.7	13	0.6	0.3-1.1	11	1.4	0.7-2.7	2	0.1	0.0-0.6
1988	56	2.5	1.9-3.3	33	3.1	2.1-4.5	23	2.0	1.2-3.0	10	0.5	0.2-0.9	4	0.4	0.1-1.2	6	0.5	0.2-1.1
1989	52	2.4	1.8-3.2	33	3.7	2.5-5.4	19	1.5	0.9-2.4	16	0.7	0.4-1.2	15	1.6	0.9-2.7	1	0.1	0.0-0.6
1990	44	2.0	1.4-2.7	26	3.0	1.9-4.6	18	1.4	0.8-2.2	15	0.7	0.4-1.2	10	1.2	0.5-2.4	5	0.4	0.1-1.0
1991	69	3.1	2.4-4.0	41	4.6	3.1-6.4	28	2.1	1.4-3.1	11	0.5	0.2-1.0	5	0.5	0.2-1.3	6	0.5	0.2-1.1
1992	53	2.4	1.8-3.1	25	2.3	1.5-3.6	28	2.3	1.5-3.3	18	0.8	0.5-1.3	9	0.8	0.4-1.7	9	0.7	0.3-1.4
1993	47	2.0	1.4-2.7	23	2.4	1.4-3.7	24	1.8	1.1-2.6	14	0.6	0.3-1.0	8	0.7	0.3-1.6	6	0.5	0.2-1.1
1994	47	2.1	1.5-2.9	30	3.2	2.1-4.8	17	1.3	0.7-2.1	15	0.7	0.4-1.2	9	1.1	0.5-2.1	6	0.4	0.2-1.0
1995	54	2.3	1.7-3.0	30	3.0	2.0-4.4	24	1.7	1.1-2.6	9	0.4	0.2-0.7	5	0.5	0.2-1.3	4	0.3	0.1-0.7
1996	45	2.0	1.4-2.7	23	2.3	1.4-3.6	22	1.7	1.0-2.6	12	0.6	0.3-1.0	6	0.8	0.3-1.7	6	0.5	0.2-1.0
1997	62	2.6	2.0-3.3	37	3.4	2.4-4.8	25	1.9	1.2-2.8	12	0.6	0.3-1.1	4	0.4	0.1-1.1	8	0.7	0.3-1.4
1998	56	2.3	1.7-3.0	27	2.4	1.6-3.6	29	2.1	1.4-3.1	10	0.4	0.2-0.7	5	0.4	0.1-1.1	5	0.4	0.1-0.8
1999	50	2.1	1.6-2.8	27	2.8	1.8-4.2	23	1.7	1.1-2.6	10	0.4	0.2-0.8	7	0.8	0.3-1.6	3	0.2	0.0-0.7
2000	59	2.4	1.8-3.1	36	3.3	2.3-4.6	23	1.7	1.1-2.5	5	0.2	0.1-0.5	3	0.3	0.1-0.9	2	0.2	0.0-0.6
2001	53	2.1	1.5-2.7	36	3.2	2.2-4.4	17	1.2	0.7-1.9	8	0.3	0.1-0.7	4	0.4	0.1-1.1	4	0.3	0.1-0.8
2002	63	2.7	2.0-3.4	32	3.3	2.2-4.7	31	2.3	1.5-3.2	7	0.3	0.1-0.6	5	0.5	0.2-1.2	2	0.1	0.0-0.5
2003	61	2.5	1.9-3.2	25	2.4	1.5-3.6	36	2.6	1.8-3.6	13	0.6	0.3-1.0	7	1.0	0.4-2.0	6	0.4	0.2-0.9
2004	65	2.6	2.0-3.4	35	3.2	2.2-4.5	30	2.2	1.5-3.1	8	0.3	0.1-0.7	5	0.5	0.2-1.3	3	0.2	0.0-0.6
2005	74	3.1	2.4-3.9	45	4.1	3.0-5.5	29	2.2	1.5-3.1	5	0.2	0.1-0.5	4	0.4	0.1-1.0	1	0.1	0.0-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	57	2.3	1.8-3.0	30	2.8	1.9-4.1	27	1.9	1.3-2.8	5	0.2	0.1-0.6	1	0.1	0.0-0.6	4	0.3	0.1-0.8
2007	55	2.3	1.7-3.0	30	3.0	2.0-4.4	25	1.8	1.2-2.7	8	0.4	0.2-0.7	2	0.2	0.0-0.7	6	0.5	0.2-1.0
2008	92	3.8	3.0-4.6	52	4.8	3.6-6.4	40	3.0	2.1-4.1	5	0.2	0.1-0.5	3	0.3	0.1-0.8	2	0.2	0.0-0.6
2009	71	3.0	2.3-3.8	27	2.7	1.7-4.0	44	3.2	2.4-4.4	7	0.3	0.1-0.6	4	0.4	0.1-1.1	3	0.2	0.0-0.6
2010	81	3.3	2.6-4.2	48	4.6	3.3-6.2	33	2.4	1.6-3.4	8	0.3	0.1-0.6	4	0.3	0.1-0.9	4	0.3	0.1-0.8
2011	84	3.3	2.6-4.1	45	3.8	2.7-5.1	39	2.8	2.0-3.8	7	0.3	0.1-0.6	4	0.5	0.1-1.3	3	0.2	0.0-0.6
2012	66	2.6	2.0-3.3	32	2.8	1.9-4.0	34	2.5	1.7-3.5	4	0.2	0.0-0.5	3	0.3	0.1-0.9	1	0.1	0.0-0.4
2013	87	3.5	2.8-4.4	45	4.2	3.0-5.7	42	3.1	2.2-4.2	8	0.3	0.1-0.7	4	0.4	0.1-1.1	4	0.3	0.1-0.7
2014	89	3.4	2.7-4.2	46	4.0	2.9-5.3	43	3.0	2.2-4.1	6	0.2	0.1-0.5	3	0.3	0.1-1.0	3	0.2	0.0-0.6
2015	73	3.0	2.3-3.8	36	3.3	2.3-4.6	37	2.8	1.9-3.9	6	0.2	0.1-0.5	2	0.2	0.0-0.8	4	0.3	0.1-0.7
2016	68	2.7	2.1-3.4	31	2.8	1.9-4.0	37	2.6	1.8-3.6	2	0.1	0.0-0.3	2	0.2	0.0-0.8	0	0.0	0.0-0.3
2017	76	2.8	2.2-3.6	35	2.8	1.9-4.0	41	2.9	2.1-4.0	5	0.2	0.1-0.5	1	0.1	0.0-0.6	4	0.3	0.1-0.8
2018	60	2.4	1.8-3.1	37	3.3	2.3-4.6	23	1.7	1.1-2.6	8	0.3	0.1-0.6	4	0.3	0.1-0.9	4	0.3	0.1-0.8
2019	85	3.6	2.8-4.4	44	4.1	3.0-5.5	41	3.1	2.2-4.3	6	0.2	0.1-0.5	2	0.1	0.0-0.6	4	0.3	0.1-0.8
2020	52	2.1	1.5-2.8	26	2.4	1.6-3.6	26	1.9	1.2-2.8	6	0.3	0.1-0.6	4	0.4	0.1-1.0	2	0.2	0.0-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	69	5.6	4.3-7.3	32	7.1	4.5-10.5	37	4.8	3.3-6.8	45	3.8	2.7-5.1	28	6.2	3.9-9.3	17	2.2	1.3-3.7
1977	72	5.6	4.3-7.2	29	5.7	3.5-8.6	43	5.6	4.0-7.7	37	3.0	1.9-4.3	14	2.2	1.1-3.9	23	3.2	1.9-5.1
1978	109	8.0	6.4-9.7	60	10.4	7.7-13.7	49	6.2	4.5-8.4	52	3.7	2.8-5.0	33	6.1	4.1-8.9	19	2.2	1.3-3.5
1979	85	6.8	5.3-8.6	48	9.2	6.5-12.5	37	5.2	3.5-7.3	50	4.5	3.2-6.1	24	5.2	3.0-8.1	26	4.0	2.5-6.0
1980	107	7.7	6.2-9.4	53	9.1	6.6-12.2	54	6.7	4.9-8.9	55	4.1	3.0-5.4	34	5.6	3.8-8.1	21	2.9	1.7-4.6
1981	99	7.4	5.9-9.2	46	7.1	4.9-10.0	53	7.2	5.3-9.6	60	4.4	3.3-5.8	26	4.8	3.0-7.3	34	4.2	2.8-5.9
1982	128	8.8	7.3-10.6	57	9.1	6.7-12.0	71	8.4	6.5-10.7	53	4.2	3.1-5.6	24	4.8	2.8-7.4	29	3.8	2.5-5.6
1983	132	8.7	7.2-10.4	62	9.9	7.4-13.0	70	7.9	6.1-10.0	73	5.1	3.9-6.4	39	6.3	4.3-9.0	34	4.2	2.8-5.9
1984	149	9.5	7.9-11.3	72	11.4	8.6-14.8	77	8.3	6.5-10.4	60	3.9	2.9-5.1	34	5.3	3.5-7.8	26	2.9	1.9-4.3
1985	186	11.3	9.6-13.1	97	13.7	10.8-17.1	89	9.5	7.6-11.8	83	5.3	4.1-6.7	41	5.9	4.0-8.4	42	4.8	3.4-6.5
1986	191	11.9	10.1-13.8	97	15.3	12.0-19.2	94	9.8	7.9-12.1	82	5.1	4.0-6.4	40	6.2	4.1-8.8	42	4.5	3.2-6.1
1987	202	11.8	10.1-13.6	121	17.8	14.4-21.7	81	7.9	6.2-9.9	77	4.5	3.5-5.6	38	5.5	3.7-7.9	39	3.8	2.7-5.3
1988	175	9.9	8.4-11.6	102	14.1	11.3-17.4	73	7.1	5.6-9.0	85	5.3	4.2-6.6	55	8.9	6.5-12.0	30	3.1	2.1-4.5
1989	203	11.3	9.7-13.0	118	15.1	12.2-18.4	85	8.3	6.6-10.3	87	5.1	4.1-6.4	51	7.5	5.3-10.1	36	3.6	2.5-5.0
1990	233	12.0	10.5-13.7	139	16.4	13.6-19.6	94	8.6	6.9-10.6	75	4.4	3.4-5.6	39	5.6	3.7-8.0	36	3.6	2.5-5.0
1991	238	12.9	11.2-14.7	135	16.9	13.9-20.4	103	9.8	8.0-12.0	75	4.4	3.4-5.6	43	5.8	4.0-8.1	32	3.3	2.2-4.7
1992	284	15.1	13.3-17.0	157	19.3	16.2-22.9	127	11.7	9.7-14.0	85	4.8	3.8-6.0	41	5.6	3.8-7.8	44	4.2	3.0-5.7
1993	272	13.2	11.7-14.9	170	19.1	16.2-22.4	102	8.8	7.1-10.7	78	4.2	3.3-5.3	39	5.0	3.5-7.0	39	3.6	2.5-5.0
1994	321	16.6	14.8-18.6	186	22.3	19.1-26.0	135	12.2	10.2-14.5	87	4.8	3.8-6.0	38	4.8	3.3-6.7	49	4.6	3.4-6.1
1995	283	14.6	12.9-16.4	168	20.5	17.3-24.1	115	10.3	8.5-12.4	85	4.6	3.6-5.7	49	6.1	4.4-8.2	36	3.4	2.4-4.7
1996	298	14.9	13.2-16.7	158	18.3	15.4-21.5	140	12.0	10.1-14.2	93	5.2	4.2-6.4	39	5.4	3.6-7.6	54	5.0	3.7-6.5
1997	330	16.6	14.8-18.5	187	22.6	19.2-26.3	143	12.4	10.4-14.6	95	5.3	4.2-6.5	49	6.8	4.9-9.1	46	4.2	3.1-5.6
1998	319	15.7	13.9-17.5	171	20.6	17.4-24.1	148	12.4	10.4-14.6	108	5.9	4.8-7.1	53	7.2	5.3-9.5	55	4.9	3.7-6.4
1999	333	15.9	14.2-17.7	176	20.5	17.4-24.0	157	12.7	10.8-14.9	114	5.7	4.7-6.9	64	8.5	6.4-11.0	50	4.2	3.1-5.5
2000	300	13.9	12.3-15.5	155	17.3	14.6-20.4	145	11.4	9.6-13.4	99	4.9	4.0-6.0	53	6.8	5.0-9.0	46	3.8	2.7-5.0
2001	311	14.6	13.0-16.3	161	18.4	15.6-21.7	150	11.9	10.1-14.0	111	5.4	4.5-6.6	57	6.7	5.0-8.7	54	4.4	3.3-5.8
2002	302	13.5	12.0-15.1	158	16.9	14.3-19.9	144	11.0	9.3-13.0	105	5.0	4.1-6.1	61	7.5	5.6-9.7	44	3.5	2.5-4.6
2003	342	15.4	13.8-17.1	158	17.5	14.7-20.6	184	14.0	12.0-16.2	109	5.0	4.1-6.1	57	6.8	5.1-9.0	52	4.0	3.0-5.2
2004	313	14.3	12.7-16.0	155	17.5	14.8-20.7	158	12.1	10.2-14.1	87	4.1	3.2-5.0	46	5.4	3.9-7.3	41	3.2	2.3-4.3
2005	365	16.1	14.5-17.9	175	18.7	15.9-21.8	190	14.3	12.3-16.5	91	4.3	3.4-5.3	45	5.5	3.9-7.4	46	3.6	2.6-4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	322	14.4	12.8-16.0	154	16.8	14.2-19.9	168	12.8	10.9-14.9	81	3.7	2.9-4.6	35	4.2	2.8-5.9	46	3.5	2.5-4.6
2007	356	15.6	14.0-17.3	196	20.3	17.5-23.5	160	12.0	10.2-14.1	100	4.5	3.7-5.5	45	5.3	3.8-7.2	55	4.1	3.1-5.4
2008	347	15.0	13.4-16.7	191	19.9	17.1-23.0	156	11.5	9.8-13.5	82	3.7	2.9-4.6	39	4.5	3.2-6.3	43	3.2	2.3-4.3
2009	381	16.8	15.1-18.6	185	19.7	16.9-22.9	196	14.7	12.7-16.9	86	3.8	3.0-4.7	40	4.5	3.2-6.2	46	3.4	2.5-4.5
2010	360	15.3	13.7-17.0	182	18.8	16.0-21.9	178	12.8	11.0-14.8	85	3.8	3.0-4.7	39	4.4	3.1-6.1	46	3.3	2.4-4.5
2011	413	17.2	15.5-19.0	213	22.4	19.4-25.8	200	13.8	12.0-15.9	92	4.0	3.2-4.9	49	5.4	3.9-7.2	43	3.1	2.2-4.2
2012	384	15.8	14.2-17.5	184	18.4	15.7-21.4	200	14.1	12.2-16.2	90	3.8	3.0-4.6	47	5.1	3.7-6.9	43	2.9	2.1-4.0
2013	392	15.6	14.0-17.2	192	18.5	15.9-21.5	200	13.5	11.7-15.6	95	3.9	3.1-4.8	50	5.4	4.0-7.2	45	3.0	2.2-4.0
2014	421	16.3	14.8-18.0	219	20.6	17.9-23.6	202	13.3	11.5-15.4	93	3.7	2.9-4.5	47	4.9	3.5-6.5	46	3.0	2.2-4.0
2015	397	15.5	14.0-17.1	189	18.2	15.6-21.1	208	13.6	11.8-15.6	88	3.6	2.9-4.4	41	4.6	3.2-6.3	47	3.1	2.3-4.2
2016	438	17.0	15.4-18.7	213	20.4	17.6-23.4	225	14.8	12.9-16.9	96	3.8	3.1-4.7	45	4.5	3.2-6.0	51	3.3	2.4-4.4
2017	419	16.1	14.5-17.7	193	17.9	15.4-20.8	226	15.0	13.0-17.1	87	3.4	2.7-4.2	43	4.2	3.0-5.8	44	2.8	2.0-3.8
2018	400	15.3	13.8-16.9	208	19.3	16.7-22.2	192	12.4	10.7-14.3	86	3.2	2.6-4.0	49	4.8	3.5-6.4	37	2.3	1.6-3.2
2019	408	15.3	13.8-16.9	204	18.6	16.1-21.4	204	13.3	11.5-15.4	103	3.8	3.1-4.7	56	5.2	3.9-6.8	47	2.9	2.1-3.9
2020	352	13.2	11.9-14.7	187	17.1	14.6-19.8	165	10.6	9.0-12.4	69	2.5	2.0-3.2	42	3.7	2.7-5.1	27	1.7	1.1-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	91	9.0	7.1-11.2	45	11.7	8.2-16.0	46	7.3	5.2-9.9	50	5.2	3.7-6.9	19	4.2	2.4-6.7	31	5.5	3.6-8.0
1977	92	8.7	6.9-10.9	41	10.4	7.0-14.9	51	7.8	5.7-10.4	62	6.1	4.5-8.0	28	7.9	4.9-11.9	34	5.2	3.5-7.3
1978	92	8.7	6.9-10.8	52	14.0	10.0-19.0	40	5.8	4.0-7.9	45	4.1	2.9-5.6	25	6.8	4.1-10.5	20	2.7	1.6-4.2
1979	106	8.8	7.1-10.7	48	9.2	6.6-12.4	58	8.3	6.2-10.8	56	5.0	3.7-6.6	23	5.1	3.1-7.9	33	4.9	3.2-6.9
1980	120	10.2	8.3-12.4	48	11.4	8.1-15.6	72	9.7	7.5-12.4	77	6.7	5.2-8.5	34	8.1	5.3-11.6	43	5.9	4.2-8.0
1981	112	9.1	7.4-11.1	49	10.2	7.3-13.7	63	8.5	6.4-11.0	56	4.9	3.6-6.5	21	5.1	2.9-8.1	35	4.8	3.3-6.8
1982	119	9.7	7.9-11.7	52	10.7	7.7-14.3	67	9.0	6.8-11.5	85	7.0	5.5-8.7	42	8.9	6.1-12.4	43	5.8	4.2-7.9
1983	120	9.2	7.6-11.2	49	9.2	6.6-12.3	71	9.1	7.0-11.5	76	6.2	4.8-7.8	28	6.0	3.8-8.9	48	6.2	4.5-8.3
1984	132	10.4	8.6-12.4	54	11.8	8.5-15.9	78	9.7	7.6-12.1	89	7.0	5.5-8.7	30	5.9	3.8-8.6	59	7.5	5.6-9.8
1985	134	9.7	8.1-11.6	54	11.6	8.3-15.5	80	9.0	7.1-11.3	80	5.9	4.6-7.4	38	8.0	5.4-11.3	42	4.8	3.4-6.6
1986	134	9.5	7.9-11.3	60	11.7	8.6-15.3	74	8.3	6.5-10.5	79	6.0	4.7-7.5	34	6.9	4.6-9.8	45	5.4	3.9-7.3
1987	125	8.5	7.0-10.2	60	10.7	8.0-14.0	65	7.1	5.5-9.1	77	5.4	4.2-6.8	40	7.6	5.2-10.7	37	4.1	2.9-5.7
1988	109	7.2	5.9-8.8	48	8.3	5.9-11.2	61	6.6	5.0-8.5	63	4.4	3.4-5.7	29	5.6	3.5-8.2	34	3.8	2.6-5.3
1989	134	9.5	7.9-11.3	62	12.0	8.9-15.8	72	8.1	6.3-10.2	61	4.5	3.4-5.8	28	6.2	3.9-9.1	33	3.7	2.5-5.2
1990	139	8.8	7.3-10.4	56	10.6	7.7-14.1	83	8.2	6.5-10.2	65	4.6	3.5-5.9	26	5.7	3.6-8.5	39	4.2	3.0-5.7
1991	145	9.7	8.1-11.4	47	8.7	6.2-11.8	98	10.2	8.2-12.5	84	5.7	4.5-7.1	33	6.6	4.3-9.5	51	5.4	4.0-7.1
1992	180	11.2	9.6-13.0	81	14.1	10.9-17.8	99	9.8	7.9-11.9	82	5.1	4.0-6.3	30	5.7	3.6-8.4	52	5.0	3.7-6.6
1993	131	8.0	6.6-9.5	55	8.4	6.2-11.0	76	7.5	5.9-9.4	75	4.8	3.8-6.1	32	6.0	3.9-8.6	43	4.3	3.1-5.8
1994	148	9.2	7.7-10.8	62	10.6	8.0-13.8	86	8.5	6.7-10.5	78	4.9	3.9-6.2	27	4.4	2.8-6.6	51	5.1	3.8-6.7
1995	163	9.8	8.3-11.5	68	11.5	8.7-14.8	95	9.0	7.3-11.0	108	6.6	5.4-8.0	45	7.6	5.4-10.4	63	6.0	4.6-7.7
1996	161	9.3	7.9-10.8	67	10.1	7.7-13.0	94	8.7	7.0-10.7	81	4.8	3.8-6.0	38	6.4	4.4-8.9	43	4.0	2.9-5.4
1997	177	10.1	8.6-11.7	75	11.1	8.6-14.1	102	9.4	7.6-11.4	87	5.1	4.1-6.3	39	6.5	4.5-9.0	48	4.4	3.3-5.9
1998	181	10.2	8.7-11.8	78	11.7	9.1-14.8	103	9.3	7.6-11.3	76	4.4	3.4-5.5	26	4.1	2.6-6.1	50	4.5	3.3-6.0
1999	186	10.3	8.8-11.9	66	9.1	6.9-11.7	120	10.6	8.8-12.7	84	4.6	3.7-5.8	31	4.4	2.9-6.3	53	4.7	3.5-6.1
2000	193	10.1	8.7-11.6	76	10.3	7.9-13.0	117	9.8	8.1-11.8	71	4.1	3.2-5.1	17	2.8	1.6-4.6	54	4.8	3.6-6.2
2001	213	11.0	9.5-12.6	84	11.6	9.1-14.5	129	10.8	9.0-12.8	87	4.7	3.7-5.8	36	5.4	3.7-7.6	51	4.3	3.2-5.6
2002	218	11.1	9.6-12.7	87	12.1	9.5-15.0	131	10.7	8.9-12.7	107	5.6	4.6-6.8	43	5.7	4.0-7.8	64	5.4	4.2-6.9
2003	228	11.6	10.1-13.3	93	13.0	10.3-16.1	135	10.9	9.2-13.0	89	4.5	3.6-5.5	38	4.8	3.3-6.8	51	4.2	3.1-5.5
2004	212	10.2	8.9-11.7	87	10.7	8.5-13.3	125	9.8	8.2-11.7	94	4.8	3.9-5.9	39	5.3	3.7-7.3	55	4.6	3.4-6.0
2005	252	12.1	10.6-13.7	109	13.5	11.0-16.4	143	11.1	9.3-13.1	87	4.3	3.4-5.3	35	4.6	3.1-6.5	52	4.1	3.0-5.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	255	12.3	10.8-13.9	119	14.8	12.2-17.8	136	10.6	8.9-12.5	80	3.9	3.1-4.9	34	4.4	3.0-6.2	46	3.6	2.6-4.8
2007	222	10.7	9.3-12.2	100	13.0	10.4-15.9	122	9.4	7.8-11.2	109	5.2	4.3-6.3	48	6.0	4.3-8.0	61	4.7	3.6-6.0
2008	293	13.4	11.9-15.0	136	16.5	13.7-19.6	157	11.6	9.8-13.5	76	3.7	2.9-4.6	31	4.0	2.7-5.7	45	3.4	2.5-4.6
2009	314	14.2	12.7-15.9	156	17.8	15.0-20.9	158	11.8	10.0-13.8	93	4.3	3.5-5.3	52	6.2	4.6-8.2	41	3.1	2.2-4.2
2010	307	13.6	12.1-15.2	159	18.7	15.7-22.0	148	10.7	9.0-12.6	101	4.7	3.8-5.7	45	5.7	4.1-7.7	56	4.1	3.1-5.4
2011	352	15.0	13.5-16.7	173	18.9	16.1-22.1	179	12.8	11.0-14.8	98	4.3	3.5-5.2	47	5.5	4.0-7.4	51	3.5	2.6-4.6
2012	352	14.9	13.3-16.5	151	16.8	14.1-19.8	201	13.8	11.9-15.8	95	4.0	3.2-4.9	47	5.0	3.6-6.7	48	3.3	2.4-4.4
2013	336	13.6	12.1-15.1	165	17.6	14.9-20.6	171	11.3	9.6-13.1	96	3.9	3.2-4.8	52	5.6	4.1-7.5	44	3.0	2.2-4.0
2014	365	14.7	13.2-16.4	171	17.5	14.9-20.4	194	13.0	11.2-15.0	100	4.3	3.5-5.2	49	5.6	4.1-7.4	51	3.5	2.6-4.6
2015	339	13.3	11.9-14.8	162	15.9	13.5-18.7	177	11.6	9.9-13.4	127	5.0	4.2-6.0	59	6.4	4.8-8.3	68	4.3	3.3-5.4
2016	381	14.5	13.0-16.1	173	16.8	14.3-19.6	208	13.1	11.3-15.0	125	5.0	4.1-5.9	57	5.8	4.4-7.6	68	4.5	3.5-5.7
2017	385	14.7	13.3-16.3	171	16.7	14.2-19.5	214	13.4	11.6-15.3	117	4.4	3.7-5.3	52	5.3	3.9-7.1	65	4.0	3.0-5.1
2018	363	13.1	11.8-14.6	150	13.3	11.1-15.6	213	12.9	11.2-14.9	97	3.5	2.9-4.3	44	4.3	3.1-5.8	53	3.1	2.3-4.1
2019	359	13.2	11.8-14.6	185	17.1	14.6-19.8	174	10.8	9.2-12.6	104	3.7	3.0-4.5	52	4.8	3.5-6.3	52	2.9	2.2-3.9
2020	307	11.1	9.9-12.5	139	12.7	10.6-15.0	168	10.2	8.7-11.9	72	2.6	2.0-3.3	28	2.5	1.7-3.7	44	2.7	2.0-3.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	101	7.9	6.2-9.8	51	9.6	6.8-13.1	50	6.7	4.8-9.1	74	6.1	4.6-7.9	41	8.1	5.2-11.9	33	4.8	3.2-6.9
1977	91	7.2	5.6-9.0	46	9.5	6.3-13.5	45	6.0	4.3-8.1	81	6.3	4.9-8.0	40	8.5	5.7-12.0	41	5.0	3.5-6.9
1978	118	9.3	7.5-11.3	60	11.9	8.5-16.0	58	7.7	5.7-10.1	83	6.6	5.1-8.3	47	8.6	6.1-11.8	36	5.1	3.5-7.3
1979	106	8.6	6.9-10.6	57	12.6	8.9-17.2	49	6.4	4.7-8.6	76	6.0	4.6-7.6	44	9.6	6.5-13.4	32	4.0	2.6-5.7
1980	115	8.8	7.1-10.7	53	10.1	7.1-13.9	62	7.9	6.0-10.3	81	6.5	5.0-8.2	36	7.9	5.0-11.6	45	5.8	4.1-7.9
1981	116	8.7	7.1-10.6	61	12.1	8.8-16.1	55	6.9	5.1-9.0	81	6.1	4.7-7.7	33	6.4	4.0-9.6	48	6.0	4.3-8.0
1982	125	8.9	7.3-10.8	64	12.3	9.0-16.3	61	7.1	5.3-9.2	72	5.4	4.1-6.9	36	7.5	4.9-11.0	36	4.3	3.0-6.1
1983	129	8.8	7.2-10.5	65	11.6	8.6-15.3	64	7.2	5.5-9.3	79	5.5	4.2-6.9	38	6.5	4.3-9.2	41	4.8	3.4-6.6
1984	131	8.7	7.1-10.4	70	10.7	8.0-13.9	61	7.1	5.4-9.2	83	5.5	4.3-7.0	41	7.2	4.8-10.1	42	4.7	3.3-6.4
1985	135	8.6	7.1-10.3	82	13.4	10.3-17.0	53	5.7	4.2-7.6	78	5.2	4.1-6.6	49	8.9	6.3-12.2	29	3.1	2.0-4.5
1986	129	8.2	6.7-9.8	68	11.7	8.7-15.3	61	6.3	4.8-8.1	75	4.6	3.6-5.9	40	6.6	4.5-9.3	35	3.5	2.4-5.0
1987	141	8.8	7.3-10.5	67	10.4	7.7-13.6	74	7.8	6.1-9.8	74	4.4	3.4-5.6	36	5.5	3.5-8.0	38	3.9	2.7-5.3
1988	110	7.1	5.8-8.7	58	10.6	7.7-14.0	52	5.3	3.9-7.0	76	5.2	4.0-6.6	35	6.2	4.1-9.0	41	4.6	3.3-6.3
1989	156	9.4	7.9-11.1	79	12.5	9.5-16.1	77	7.6	6.0-9.6	82	5.2	4.1-6.5	47	8.4	5.9-11.5	35	3.6	2.4-5.0
1990	155	9.2	7.7-10.9	74	10.9	8.2-14.1	81	8.1	6.4-10.1	95	5.9	4.7-7.3	45	7.5	5.2-10.3	50	5.0	3.6-6.6
1991	180	10.4	8.9-12.2	87	13.2	10.3-16.8	93	8.8	7.1-10.9	71	4.3	3.3-5.5	32	5.3	3.4-7.8	39	3.9	2.7-5.3
1992	174	10.0	8.5-11.7	76	11.4	8.6-14.8	98	9.3	7.5-11.4	86	5.4	4.2-6.7	36	5.4	3.6-7.8	50	5.1	3.8-6.8
1993	151	8.6	7.3-10.2	74	11.2	8.5-14.4	77	7.3	5.7-9.1	84	5.0	3.9-6.2	39	5.9	4.0-8.4	45	4.4	3.2-5.9
1994	155	8.7	7.4-10.3	79	11.6	9.0-14.8	76	7.1	5.5-8.9	113	6.6	5.4-8.0	51	7.4	5.3-10.1	62	6.0	4.6-7.7
1995	163	8.7	7.4-10.2	80	10.2	7.9-13.0	83	7.5	6.0-9.4	93	5.2	4.1-6.4	43	6.4	4.5-8.9	50	4.6	3.4-6.0
1996	177	9.5	8.1-11.0	96	12.7	10.0-15.8	81	7.3	5.8-9.1	97	5.5	4.4-6.8	45	6.5	4.5-9.0	52	4.9	3.6-6.4
1997	160	8.6	7.3-10.1	64	8.8	6.6-11.5	96	8.5	6.9-10.4	95	5.4	4.3-6.6	49	7.8	5.6-10.5	46	4.1	3.0-5.5
1998	154	8.0	6.8-9.5	67	9.3	7.0-12.1	87	7.4	5.9-9.2	76	4.0	3.1-5.0	34	4.6	3.0-6.6	42	3.7	2.6-4.9
1999	146	7.3	6.1-8.6	74	10.0	7.6-12.8	72	5.9	4.6-7.4	92	5.0	4.0-6.1	47	7.0	5.0-9.6	45	3.9	2.8-5.2
2000	192	9.8	8.4-11.3	92	11.7	9.3-14.5	100	8.4	6.8-10.2	93	5.0	4.0-6.2	35	5.4	3.6-7.7	58	4.9	3.7-6.4
2001	164	8.2	7.0-9.6	70	8.9	6.8-11.5	94	7.7	6.2-9.5	90	4.7	3.8-5.8	41	5.6	3.9-7.7	49	4.1	3.0-5.5
2002	188	9.1	7.8-10.5	101	12.4	9.9-15.3	87	7.0	5.6-8.7	93	4.6	3.7-5.7	47	6.4	4.6-8.7	46	3.7	2.7-4.9
2003	187	8.7	7.5-10.0	91	10.2	8.1-12.6	96	7.6	6.1-9.2	90	4.4	3.5-5.4	40	4.8	3.3-6.6	50	4.1	3.0-5.4
2004	188	8.9	7.6-10.2	103	12.2	9.8-14.9	85	6.7	5.3-8.3	81	4.1	3.2-5.1	42	5.5	3.9-7.5	39	3.2	2.3-4.3
2005	172	8.0	6.8-9.3	87	10.4	8.2-13.0	85	6.6	5.2-8.1	89	4.4	3.5-5.5	41	5.6	3.9-7.7	48	3.8	2.8-5.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Black Individuals
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	184	8.6	7.3-9.9	108	13.0	10.6-15.9	76	5.8	4.6-7.3	80	3.8	3.0-4.7	47	5.9	4.2-7.9	33	2.5	1.7-3.6
2007	189	8.7	7.5-10.1	84	9.4	7.4-11.7	105	8.0	6.6-9.7	85	4.2	3.3-5.2	45	6.0	4.3-8.1	40	3.2	2.3-4.4
2008	179	8.2	7.1-9.6	82	10.5	8.2-13.2	97	7.3	5.9-8.9	95	4.4	3.6-5.4	51	6.5	4.8-8.7	44	3.3	2.4-4.4
2009	203	9.0	7.8-10.3	105	12.0	9.7-14.6	98	7.2	5.9-8.9	83	3.8	3.0-4.7	43	4.9	3.5-6.6	40	3.0	2.1-4.1
2010	256	11.1	9.7-12.5	134	14.8	12.2-17.6	122	8.9	7.4-10.7	102	4.6	3.7-5.6	51	6.3	4.6-8.4	51	3.8	2.8-5.0
2011	237	10.2	8.9-11.6	121	12.4	10.2-14.9	116	8.5	7.0-10.3	112	4.8	3.9-5.7	57	5.9	4.4-7.7	55	4.0	3.0-5.2
2012	226	9.4	8.2-10.7	120	12.7	10.4-15.2	106	7.5	6.1-9.1	90	4.0	3.2-4.9	46	5.4	3.9-7.2	44	3.2	2.3-4.3
2013	270	10.9	9.7-12.3	126	12.5	10.4-15.0	144	10.0	8.4-11.8	92	3.8	3.1-4.7	57	6.3	4.7-8.3	35	2.3	1.6-3.3
2014	306	12.3	11.0-13.8	159	15.8	13.4-18.6	147	10.1	8.5-11.9	101	4.3	3.5-5.2	52	5.7	4.2-7.6	49	3.4	2.5-4.6
2015	260	10.5	9.2-11.9	126	13.0	10.7-15.5	134	9.0	7.5-10.7	113	4.6	3.8-5.6	55	5.6	4.2-7.4	58	4.0	3.0-5.2
2016	284	11.2	9.9-12.6	138	13.5	11.2-16.0	146	9.8	8.2-11.5	111	4.3	3.5-5.2	54	5.4	4.0-7.0	57	3.7	2.7-4.8
2017	263	10.2	9.0-11.5	123	11.9	9.8-14.2	140	9.1	7.6-10.8	116	4.6	3.8-5.5	59	5.9	4.5-7.7	57	3.7	2.8-4.8
2018	251	9.6	8.4-10.9	129	12.3	10.2-14.7	122	7.9	6.6-9.5	108	4.2	3.4-5.1	51	5.1	3.7-6.7	57	3.8	2.9-5.0
2019	249	9.6	8.4-10.9	131	12.2	10.1-14.5	118	7.7	6.4-9.3	96	3.5	2.8-4.3	50	4.5	3.3-6.0	46	2.7	2.0-3.7
2020	227	8.8	7.7-10.1	123	11.4	9.4-13.6	104	7.1	5.7-8.6	92	3.4	2.7-4.2	46	4.6	3.4-6.2	46	2.8	2.0-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
All Invasive Malignant Tumors

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,389	427.4	402.9-452.9	715	523.4	479.5-569.8	674	362.6	333.7-393.1	772	251.2	231.8-271.6	435	330.5	294.7-369.0	337	195.2	173.3-219.0
1977	1,506	451.9	427.1-477.7	787	561.0	516.8-607.7	719	375.9	346.9-406.6	867	275.8	255.9-296.7	517	397.0	358.2-438.5	350	192.6	171.5-215.4
1978	1,583	463.8	439.1-489.4	837	597.3	551.0-646.0	746	376.7	348.5-406.5	884	266.5	247.6-286.3	517	371.4	335.3-410.0	367	194.0	173.4-216.2
1979	1,579	452.8	428.7-477.9	828	577.9	533.1-625.0	751	371.1	343.4-400.3	945	285.6	266.0-306.2	531	386.8	349.8-426.2	414	217.0	195.2-240.4
1980	1,723	470.7	447.0-495.3	923	617.3	572.6-664.2	800	375.3	348.4-403.6	965	273.8	255.4-293.0	545	379.6	344.0-417.5	420	204.4	184.3-225.9
1981	1,673	448.5	425.6-472.3	908	606.0	561.6-652.6	765	349.1	323.6-376.0	1,034	291.9	272.9-311.7	590	416.5	378.8-456.5	444	213.0	192.6-234.7
1982	1,655	420.7	399.3-442.8	893	562.3	521.8-604.9	762	329.6	305.6-354.8	1,009	267.7	250.3-285.9	579	383.6	349.1-420.2	430	193.6	175.0-213.5
1983	1,720	427.8	406.4-449.9	925	570.8	529.9-613.7	795	338.5	314.4-363.8	1,030	266.2	249.2-284.0	603	390.0	355.8-426.3	427	187.3	169.3-206.5
1984	1,815	439.1	417.8-461.1	980	577.5	537.6-619.3	835	348.8	324.6-374.3	1,061	269.9	252.8-287.7	593	369.7	336.8-404.6	468	205.1	186.3-225.3
1985	1,897	447.1	426.0-469.0	979	563.9	525.0-604.6	918	373.7	349.1-399.6	1,052	258.3	242.0-275.3	582	347.4	316.7-380.0	470	199.3	181.0-218.7
1986	1,968	455.7	434.5-477.6	1,040	599.5	559.0-641.8	928	367.9	343.8-393.2	1,153	279.2	262.3-296.7	655	397.4	363.6-433.2	498	206.6	188.3-226.0
1987	1,901	428.7	408.5-449.5	973	551.0	512.8-591.1	928	356.7	333.4-381.1	1,108	263.8	247.6-280.7	629	387.5	353.9-423.1	479	191.3	174.0-209.7
1988	2,048	449.9	429.6-471.0	1,043	576.3	537.7-616.7	1,005	376.8	353.2-401.4	1,172	274.5	258.2-291.6	660	404.5	370.5-440.4	512	199.3	181.9-217.8
1989	2,147	451.4	431.5-472.0	1,135	594.2	555.8-634.2	1,012	365.8	343.0-389.7	1,228	275.4	259.4-292.0	691	395.3	362.9-429.4	537	203.1	185.9-221.5
1990	2,254	466.3	446.2-487.1	1,175	618.5	579.2-659.6	1,079	378.6	355.7-402.5	1,209	263.3	247.9-279.4	661	362.1	331.6-394.4	548	203.6	186.5-221.8
1991	2,395	473.4	453.6-493.8	1,301	651.0	611.5-692.0	1,094	367.9	345.9-391.0	1,242	263.2	248.0-279.1	683	377.2	346.0-410.3	559	197.1	180.7-214.5
1992	2,520	483.9	464.2-504.1	1,411	689.1	649.0-730.9	1,109	362.5	340.9-385.0	1,247	255.5	240.8-270.8	651	341.9	312.8-372.6	596	205.5	189.0-223.1
1993	2,655	494.3	474.7-514.4	1,485	693.2	654.3-733.7	1,170	371.5	350.0-393.9	1,274	253.5	239.1-268.5	707	364.3	334.7-395.6	567	189.1	173.5-205.6
1994	2,703	489.1	470.0-508.7	1,465	672.1	634.3-711.3	1,238	379.7	358.4-401.9	1,339	260.9	246.5-275.9	772	391.4	361.1-423.4	567	184.4	169.2-200.4
1995	2,871	505.0	485.9-524.7	1,595	696.7	659.2-735.7	1,276	384.8	363.5-407.0	1,319	249.7	235.8-264.1	708	346.1	318.2-375.6	611	194.0	178.6-210.2
1996	3,105	523.1	504.1-542.6	1,729	713.7	677.1-751.6	1,376	399.7	378.4-421.8	1,289	235.6	222.4-249.3	679	317.2	291.2-344.6	610	187.7	172.8-203.4
1997	3,074	511.3	492.7-530.5	1,698	700.2	663.9-737.9	1,376	392.6	371.8-414.3	1,323	235.5	222.5-249.0	727	332.5	306.2-360.2	596	178.4	164.2-193.6
1998	3,258	520.7	502.2-539.5	1,772	693.1	658.1-729.3	1,486	409.9	388.9-431.6	1,393	243.9	230.8-257.5	747	333.8	307.9-361.1	646	190.5	175.9-206.0
1999	3,266	511.2	493.2-529.7	1,779	672.8	639.1-707.7	1,487	404.3	383.6-425.7	1,332	227.1	214.6-240.1	695	300.3	276.0-326.0	637	184.0	169.8-199.1
2000	3,245	495.3	477.8-513.3	1,821	679.5	645.8-714.3	1,424	375.1	355.5-395.5	1,446	236.6	224.1-249.5	760	321.0	296.3-347.1	686	188.9	174.8-203.8
2001	3,444	506.3	488.9-524.1	1,949	696.0	662.6-730.5	1,495	381.5	362.1-401.7	1,352	216.9	205.0-229.1	684	287.8	264.4-312.5	668	178.9	165.3-193.1
2002	3,560	506.0	488.9-523.6	1,966	687.6	654.8-721.6	1,594	390.3	371.1-410.3	1,426	218.5	206.9-230.6	752	294.6	271.6-318.8	674	175.4	162.2-189.4
2003	3,600	497.6	480.9-514.7	1,991	667.2	635.7-699.8	1,609	386.9	367.9-406.5	1,413	211.4	200.1-223.1	740	278.2	256.5-301.0	673	170.7	157.8-184.2
2004	3,735	498.8	482.4-515.7	1,985	633.9	604.0-664.9	1,750	409.5	390.2-429.4	1,418	209.2	198.0-220.7	745	280.0	258.3-302.8	673	167.3	154.7-180.6
2005	3,772	496.4	480.1-513.1	2,028	640.6	610.7-671.4	1,744	400.8	381.9-420.4	1,431	202.0	191.3-213.2	747	261.4	241.3-282.6	684	164.5	152.2-177.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
All Invasive Malignant Tumors

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	3,935	499.2	483.2-515.5	2,175	658.5	629.0-689.0	1,760	392.1	373.8-411.2	1,309	179.3	169.4-189.6	653	220.5	202.5-239.6	656	153.5	141.8-165.8
2007	4,092	503.4	487.6-519.6	2,266	662.2	633.2-692.0	1,826	394.7	376.5-413.5	1,454	194.2	184.0-204.7	719	240.4	221.7-260.1	735	167.3	155.3-180.0
2008	4,212	500.0	484.5-515.8	2,288	636.5	608.8-665.1	1,924	405.6	387.4-424.4	1,391	177.4	167.8-187.2	693	217.3	200.2-235.5	698	152.8	141.5-164.7
2009	4,332	499.5	484.3-515.1	2,343	621.3	594.6-648.8	1,989	410.7	392.6-429.5	1,471	183.6	174.0-193.5	759	235.3	217.5-254.1	712	151.9	140.8-163.6
2010	4,352	485.8	470.9-500.9	2,406	629.1	602.3-656.7	1,946	387.0	369.7-404.9	1,485	180.6	171.2-190.4	782	240.3	222.4-259.1	703	145.0	134.3-156.3
2011	4,641	501.2	486.5-516.4	2,537	635.6	609.4-662.6	2,104	406.6	389.2-424.7	1,531	179.1	170.0-188.6	777	220.9	204.5-238.2	754	152.7	141.8-164.1
2012	4,430	463.4	449.5-477.7	2,345	567.0	542.7-592.0	2,085	392.4	375.4-409.9	1,540	175.2	166.3-184.5	791	220.9	204.6-238.1	749	147.5	137.0-158.6
2013	4,582	469.4	455.4-483.6	2,386	562.2	538.5-586.7	2,196	403.1	386.2-420.6	1,527	168.9	160.2-177.8	765	208.7	193.2-225.1	762	143.9	133.7-154.7
2014	4,802	472.8	459.2-486.8	2,529	573.3	549.8-597.5	2,273	404.0	387.3-421.3	1,529	163.1	154.8-171.8	774	204.9	189.8-220.8	755	137.5	127.7-147.9
2015	4,996	481.1	467.5-495.0	2,608	577.7	554.4-601.6	2,388	415.6	398.8-432.9	1,506	157.5	149.4-165.9	748	189.6	175.4-204.5	758	136.5	126.8-146.7
2016	5,089	478.1	464.6-491.8	2,627	565.7	543.0-589.0	2,462	418.2	401.5-435.4	1,526	153.3	145.4-161.4	777	188.3	174.5-202.9	749	130.1	120.8-139.9
2017	5,122	470.8	457.6-484.2	2,644	556.4	534.3-579.3	2,478	412.8	396.4-429.7	1,533	149.3	141.7-157.2	787	187.7	174.1-202.1	746	125.0	116.0-134.5
2018	5,234	465.5	452.7-478.7	2,741	555.6	533.9-577.9	2,493	403.9	387.9-420.3	1,573	148.8	141.3-156.5	755	176.0	163.0-189.7	818	133.3	124.1-142.9
2019	5,412	470.8	458.1-483.9	2,730	537.1	516.1-558.8	2,682	427.3	411.0-444.0	1,619	148.1	140.8-155.6	782	173.0	160.4-186.2	837	132.3	123.3-141.7
2020	4,809	404.1	392.5-416.0	2,490	474.9	455.6-494.9	2,319	355.6	341.0-370.7	1,586	140.5	133.5-147.8	773	165.3	153.3-178.0	813	126.1	117.4-135.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Oral cavity and pharynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	57	14.2	10.6-18.6	39	22.1	15.5-30.9	18	7.7	4.5-12.5	19	6.6	3.7-10.6	13	13.2	5.9-24.0	6	2.6	1.0-5.9
1977	50	12.4	9.1-16.5	39	22.2	15.4-30.9	11	4.8	2.3-8.7	24	6.1	3.8-9.1	17	9.0	5.2-14.8	7	3.5	1.4-7.4
1978	63	14.9	11.4-19.2	38	20.6	14.4-28.7	25	10.4	6.7-15.5	26	6.8	4.4-10.0	20	12.4	7.4-19.5	6	2.5	0.9-5.6
1979	58	14.1	10.6-18.3	47	26.0	18.8-35.1	11	4.7	2.3-8.5	39	9.5	6.7-13.2	31	17.9	11.7-26.0	8	3.2	1.4-6.5
1980	64	14.2	10.9-18.3	42	21.4	15.2-29.3	22	8.5	5.3-13.0	37	9.1	6.3-12.6	29	16.9	11.1-24.6	8	3.3	1.4-6.5
1981	79	17.8	13.9-22.3	52	26.1	18.9-35.1	27	11.2	7.3-16.4	32	7.5	5.1-10.7	22	11.7	7.2-18.0	10	4.3	2.0-7.9
1982	65	14.3	11.0-18.3	44	21.5	15.5-29.2	21	8.4	5.2-13.0	35	7.5	5.2-10.5	27	13.4	8.6-19.8	8	2.9	1.2-5.8
1983	74	16.4	12.7-20.8	55	28.2	20.6-37.5	19	7.5	4.5-11.9	35	7.7	5.3-10.9	24	12.3	7.4-19.1	11	4.4	2.2-7.9
1984	76	16.0	12.6-20.2	56	27.7	20.6-36.4	20	7.3	4.4-11.3	30	6.7	4.5-9.6	24	12.5	7.7-18.9	6	2.4	0.9-5.2
1985	78	17.1	13.4-21.5	57	29.0	21.5-38.1	21	8.2	5.0-12.5	34	7.1	4.8-10.0	25	12.0	7.5-18.0	9	3.3	1.5-6.4
1986	70	14.4	11.1-18.3	49	23.5	16.9-31.7	21	7.5	4.6-11.6	35	7.8	5.3-11.0	25	12.8	7.9-19.4	10	4.0	1.9-7.3
1987	80	16.5	13.0-20.7	50	23.0	16.8-30.7	30	11.3	7.6-16.2	29	6.2	4.1-9.1	20	10.4	6.0-16.6	9	3.4	1.5-6.4
1988	75	14.5	11.3-18.2	53	24.0	17.5-32.0	22	7.4	4.6-11.2	37	7.3	5.1-10.1	25	12.0	7.3-18.4	12	4.2	2.1-7.2
1989	109	20.2	16.5-24.5	78	32.8	25.6-41.5	31	10.3	6.9-14.7	43	8.0	5.8-10.9	29	12.0	7.9-17.4	14	4.8	2.6-8.1
1990	68	12.9	9.9-16.5	53	23.5	17.1-31.4	15	5.1	2.8-8.4	27	4.8	3.2-7.0	18	6.9	4.1-11.1	9	3.0	1.4-5.7
1991	91	16.6	13.3-20.5	71	30.2	23.3-38.4	20	6.4	3.9-9.9	35	6.7	4.6-9.3	24	10.5	6.5-16.0	11	3.8	1.9-6.7
1992	87	15.1	11.9-18.7	57	23.7	17.2-31.6	30	9.2	6.1-13.1	38	6.6	4.6-9.1	31	12.2	8.1-17.5	7	2.2	0.9-4.5
1993	83	14.1	11.2-17.6	57	21.8	16.2-28.7	26	8.1	5.2-11.8	20	3.5	2.1-5.4	13	5.8	2.9-10.2	7	2.0	0.8-4.2
1994	77	12.1	9.5-15.3	56	20.0	14.9-26.4	21	5.9	3.6-9.0	28	4.6	3.0-6.7	23	8.9	5.5-13.6	5	1.4	0.5-3.3
1995	84	13.8	10.9-17.2	62	22.3	16.8-29.0	22	6.9	4.3-10.4	40	6.4	4.5-8.8	30	10.5	7.0-15.2	10	3.0	1.4-5.5
1996	94	15.7	12.5-19.3	67	27.2	20.6-35.2	27	7.7	5.0-11.2	27	4.6	3.0-6.7	17	6.7	3.7-10.9	10	3.0	1.4-5.4
1997	106	15.7	12.8-19.0	71	23.9	18.4-30.5	35	9.3	6.5-13.0	24	4.8	3.0-7.1	17	9.4	5.1-15.4	7	2.3	0.9-4.6
1998	85	12.4	9.8-15.4	56	18.4	13.6-24.3	29	7.7	5.1-11.0	27	4.1	2.7-6.0	25	8.6	5.4-13.0	2	0.6	0.1-2.0
1999	82	12.1	9.5-15.0	62	20.8	15.5-27.1	20	5.5	3.3-8.5	32	4.8	3.2-6.8	22	8.3	4.9-12.9	10	2.5	1.2-4.6
2000	71	10.2	7.9-13.0	45	15.4	11.0-20.9	26	6.5	4.2-9.5	38	5.8	4.1-8.1	26	9.0	5.7-13.5	12	3.4	1.7-5.8
2001	63	8.5	6.5-10.9	50	14.3	10.5-19.0	13	3.5	1.8-5.9	16	2.7	1.5-4.4	9	3.7	1.5-7.4	7	2.1	0.8-4.1
2002	93	13.2	10.5-16.2	63	19.9	14.9-25.9	30	7.8	5.2-11.2	26	3.8	2.4-5.6	20	6.7	3.9-10.7	6	1.7	0.6-3.6
2003	81	10.4	8.2-13.0	50	14.0	10.1-18.8	31	7.4	5.0-10.5	29	4.1	2.7-5.9	23	6.9	4.3-10.5	6	1.7	0.6-3.5
2004	75	9.2	7.2-11.6	54	14.7	10.9-19.4	21	4.8	2.9-7.3	31	3.9	2.6-5.6	25	7.1	4.5-10.6	6	1.4	0.5-3.1
2005	67	8.4	6.4-10.7	45	13.0	9.3-17.6	22	4.9	3.0-7.4	20	2.8	1.7-4.3	16	5.2	2.9-8.6	4	1.0	0.3-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Oral cavity and pharynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	69	8.1	6.2-10.3	51	13.7	10.0-18.3	18	3.8	2.2-6.0	22	2.8	1.7-4.2	18	5.2	3.0-8.4	4	0.9	0.2-2.2
2007	83	9.6	7.6-12.0	61	16.0	12.0-20.9	22	4.7	2.9-7.2	23	2.9	1.8-4.4	15	4.8	2.4-8.1	8	1.8	0.7-3.5
2008	87	9.9	7.9-12.3	53	12.9	9.5-17.2	34	7.1	4.9-10.0	19	2.2	1.3-3.5	12	3.0	1.5-5.4	7	1.5	0.6-3.1
2009	88	9.5	7.5-11.8	55	13.3	9.8-17.6	33	6.4	4.4-9.1	17	1.9	1.1-3.1	12	3.3	1.6-5.9	5	1.0	0.3-2.4
2010	92	9.3	7.4-11.4	62	14.2	10.7-18.4	30	5.4	3.6-7.8	17	1.8	1.0-3.0	12	2.7	1.3-4.8	5	1.0	0.3-2.4
2011	78	7.9	6.2-10.0	61	13.4	10.1-17.4	17	3.4	2.0-5.5	27	2.9	1.9-4.2	18	4.1	2.4-6.6	9	1.8	0.8-3.4
2012	76	7.5	5.8-9.4	41	9.5	6.6-13.1	35	6.2	4.3-8.7	22	2.3	1.4-3.5	18	4.3	2.4-6.9	4	0.8	0.2-1.9
2013	67	6.4	4.9-8.2	44	9.3	6.7-12.7	23	4.1	2.6-6.1	22	2.2	1.3-3.3	16	3.6	2.0-5.9	6	1.1	0.4-2.3
2014	85	8.1	6.4-10.1	61	12.9	9.7-16.8	24	4.3	2.7-6.5	19	1.8	1.1-2.8	12	2.5	1.3-4.4	7	1.2	0.5-2.6
2015	81	7.6	6.0-9.5	60	13.0	9.8-17.0	21	3.5	2.1-5.4	20	2.0	1.2-3.2	12	2.7	1.4-4.8	8	1.4	0.6-2.9
2016	97	8.9	7.2-11.0	60	12.3	9.2-16.0	37	6.4	4.4-8.8	20	1.9	1.1-2.9	14	3.1	1.6-5.3	6	1.0	0.3-2.1
2017	79	7.0	5.5-8.7	57	11.5	8.6-15.0	22	3.5	2.2-5.3	24	2.3	1.5-3.5	20	4.6	2.7-7.2	4	0.7	0.2-1.8
2018	104	9.1	7.4-11.1	70	13.2	10.2-16.8	34	5.7	4.0-8.1	19	1.6	0.9-2.5	14	2.4	1.3-4.2	5	0.8	0.2-1.8
2019	90	7.7	6.1-9.5	59	11.3	8.5-14.8	31	5.1	3.4-7.2	24	2.1	1.3-3.1	20	4.1	2.4-6.4	4	0.6	0.2-1.6
2020	91	7.2	5.8-8.9	62	11.1	8.4-14.3	29	4.1	2.8-6.0	20	1.7	1.0-2.6	13	2.4	1.2-4.2	7	1.1	0.4-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Esophagus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	41	12.4	8.7-17.1	28	19.1	12.3-28.1	13	7.0	3.5-12.3	48	14.1	10.0-19.1	37	25.5	16.8-36.8	11	5.6	2.6-10.4
1977	51	13.6	9.9-18.1	37	22.3	15.4-31.2	14	6.7	3.5-11.5	48	13.0	9.5-17.4	38	24.7	17.0-34.5	10	4.3	2.1-8.1
1978	66	18.5	14.0-23.9	50	30.2	21.7-40.9	16	9.1	5.0-15.1	54	14.2	10.5-18.9	41	24.3	16.7-34.1	13	6.4	3.2-11.1
1979	49	14.4	10.4-19.3	41	29.0	20.0-40.4	8	4.0	1.7-8.0	42	11.9	8.4-16.3	36	25.3	16.8-36.3	6	2.6	0.9-5.7
1980	57	14.7	11.0-19.2	44	26.2	18.6-35.8	13	6.0	3.1-10.4	47	12.8	9.2-17.3	40	25.4	17.4-35.5	7	3.7	1.4-7.6
1981	55	13.4	10.0-17.6	42	23.8	16.6-33.0	13	5.7	3.0-9.8	45	10.8	7.8-14.6	34	19.6	13.2-28.0	11	4.4	2.2-7.9
1982	41	10.3	7.3-14.2	33	20.5	13.4-29.6	8	3.4	1.4-6.6	46	11.9	8.6-16.0	38	23.2	15.9-32.6	8	3.8	1.6-7.5
1983	52	12.0	8.8-15.9	41	23.3	16.0-32.5	11	4.2	2.1-7.5	49	12.2	8.8-16.3	38	23.1	15.5-32.8	11	4.6	2.3-8.3
1984	54	12.3	9.1-16.1	40	20.7	14.5-28.7	14	5.8	3.1-9.8	47	11.6	8.4-15.6	33	20.8	13.6-30.1	14	5.7	3.1-9.6
1985	52	12.3	9.0-16.2	42	24.3	16.8-33.6	10	4.0	1.9-7.4	45	9.9	7.1-13.3	37	18.4	12.6-25.9	8	3.2	1.4-6.4
1986	66	15.0	11.4-19.2	49	27.3	19.5-37.0	17	6.4	3.7-10.4	53	12.7	9.3-16.7	38	23.8	15.9-33.7	15	5.8	3.2-9.6
1987	44	9.6	6.9-13.0	31	15.8	10.4-22.9	13	5.0	2.6-8.6	41	9.1	6.4-12.4	32	16.4	10.7-23.7	9	3.7	1.7-7.1
1988	52	10.8	8.0-14.2	37	18.4	12.6-25.7	15	5.5	3.0-9.0	54	12.0	8.9-15.8	37	20.7	14.0-29.2	17	6.4	3.7-10.2
1989	65	13.7	10.4-17.5	48	25.0	17.8-34.0	17	6.1	3.6-9.8	42	8.3	5.9-11.2	35	16.0	10.8-22.7	7	2.5	1.0-5.1
1990	57	10.8	8.1-14.1	39	17.6	12.1-24.7	18	6.0	3.5-9.5	56	11.6	8.6-15.1	37	19.3	13.0-27.2	19	6.7	4.0-10.5
1991	57	11.0	8.2-14.3	42	19.8	13.8-27.2	15	4.8	2.7-8.0	49	9.1	6.7-12.1	35	15.2	10.3-21.4	14	4.6	2.5-7.7
1992	52	10.1	7.5-13.3	42	19.8	13.7-27.4	10	3.5	1.6-6.3	38	7.6	5.3-10.5	27	13.3	8.3-19.9	11	3.8	1.9-6.7
1993	53	9.1	6.7-11.9	38	15.1	10.4-21.2	15	4.5	2.5-7.5	44	7.9	5.7-10.7	39	16.8	11.5-23.5	5	1.6	0.5-3.7
1994	57	10.0	7.5-13.0	37	14.8	10.2-20.6	20	6.3	3.8-9.8	44	8.1	5.8-10.9	34	15.2	10.3-21.5	10	3.2	1.5-5.8
1995	44	7.6	5.5-10.2	29	11.3	7.3-16.5	15	4.7	2.6-7.7	48	8.3	6.1-11.1	35	13.8	9.4-19.4	13	4.1	2.2-6.9
1996	57	9.7	7.3-12.7	44	17.4	12.3-23.8	13	4.1	2.2-6.9	43	7.3	5.2-9.9	29	11.4	7.4-16.7	14	4.2	2.3-7.1
1997	45	7.7	5.5-10.3	32	13.9	9.1-20.1	13	3.7	2.0-6.3	38	6.3	4.4-8.7	24	9.8	6.0-15.0	14	3.9	2.1-6.6
1998	54	9.1	6.8-11.9	38	14.7	10.2-20.5	16	4.9	2.8-7.8	52	8.7	6.5-11.5	36	13.7	9.3-19.3	16	5.0	2.8-8.1
1999	64	9.9	7.5-12.6	47	17.5	12.6-23.6	17	4.5	2.6-7.2	31	5.2	3.4-7.4	21	8.6	5.0-13.4	10	2.9	1.4-5.3
2000	53	8.5	6.3-11.2	35	13.2	8.9-18.6	18	5.1	3.0-8.0	54	8.7	6.5-11.4	38	15.4	10.6-21.4	16	4.3	2.4-6.9
2001	53	8.0	5.9-10.5	35	13.7	9.2-19.5	18	4.5	2.6-7.1	36	5.4	3.7-7.5	25	9.1	5.6-13.7	11	2.9	1.4-5.1
2002	40	5.9	4.1-8.0	29	10.3	6.6-15.2	11	2.8	1.4-5.0	51	7.0	5.2-9.3	39	12.7	8.7-17.7	12	2.9	1.5-5.1
2003	56	8.2	6.1-10.7	43	14.1	10.0-19.3	13	3.6	1.9-6.2	37	5.6	3.9-7.8	26	9.6	5.9-14.6	11	3.0	1.5-5.4
2004	50	7.3	5.4-9.7	31	10.0	6.7-14.3	19	5.1	3.0-7.9	43	6.3	4.5-8.6	32	11.9	7.8-17.1	11	2.8	1.4-5.0
2005	48	6.1	4.4-8.1	36	10.6	7.2-14.9	12	2.7	1.4-4.7	36	5.3	3.7-7.3	24	8.0	5.0-12.0	12	3.1	1.6-5.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Esophagus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	38	5.3	3.7-7.3	27	11.0	6.9-16.3	11	2.4	1.2-4.3	25	3.5	2.2-5.2	19	7.4	4.2-11.7	6	1.3	0.5-2.8
2007	48	5.9	4.3-7.9	36	9.9	6.8-13.8	12	2.7	1.4-4.8	33	4.5	3.0-6.3	23	7.7	4.7-11.7	10	2.4	1.1-4.3
2008	37	4.4	3.0-6.0	27	7.9	5.0-11.8	10	2.1	1.0-3.8	36	4.2	2.9-5.9	22	5.8	3.5-9.0	14	3.1	1.7-5.2
2009	43	5.0	3.5-6.7	34	9.5	6.3-13.6	9	2.0	0.9-3.7	28	3.4	2.2-5.0	23	7.4	4.5-11.4	5	1.0	0.3-2.4
2010	42	4.9	3.5-6.7	32	9.2	6.1-13.2	10	2.1	1.0-3.8	43	4.9	3.5-6.6	32	8.4	5.5-12.1	11	2.4	1.2-4.3
2011	48	5.4	3.9-7.2	32	8.6	5.7-12.4	16	3.2	1.8-5.2	26	3.0	1.9-4.4	19	5.3	3.0-8.5	7	1.5	0.6-3.0
2012	42	4.2	3.0-5.7	31	6.7	4.5-9.7	11	2.0	1.0-3.6	32	3.5	2.4-5.0	22	5.3	3.2-8.2	10	2.0	1.0-3.7
2013	48	5.2	3.8-6.9	34	8.8	5.9-12.4	14	2.7	1.4-4.5	39	4.2	3.0-5.8	26	6.5	4.1-9.6	13	2.5	1.3-4.4
2014	36	3.9	2.7-5.4	23	6.2	3.8-9.4	13	2.3	1.2-4.0	38	4.1	2.9-5.7	25	6.6	4.1-9.9	13	2.4	1.3-4.2
2015	38	3.8	2.6-5.2	28	6.7	4.4-9.9	10	1.7	0.8-3.2	23	2.3	1.4-3.5	18	4.5	2.6-7.2	5	0.8	0.2-1.9
2016	43	4.6	3.3-6.3	26	6.6	4.2-9.8	17	3.2	1.8-5.1	40	4.3	3.1-5.9	30	7.9	5.2-11.4	10	1.9	0.9-3.5
2017	45	4.4	3.2-5.9	31	7.4	4.9-10.6	14	2.4	1.3-4.0	22	2.0	1.2-3.1	15	3.6	1.9-6.0	7	1.0	0.4-2.2
2018	46	4.2	3.0-5.6	26	5.9	3.8-8.7	20	3.1	1.9-4.9	27	2.4	1.6-3.6	21	4.4	2.6-6.9	6	1.0	0.4-2.2
2019	41	3.5	2.5-4.8	26	5.5	3.5-8.1	15	2.2	1.2-3.6	35	3.1	2.1-4.4	24	5.5	3.5-8.3	11	1.5	0.7-2.7
2020	24	2.3	1.4-3.4	20	4.6	2.7-7.2	4	0.6	0.2-1.6	30	2.7	1.8-3.9	21	4.6	2.8-7.2	9	1.4	0.6-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Stomach

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	65	22.9	17.1-29.7	38	32.3	21.4-46.2	27	16.3	10.3-24.2	41	13.9	9.6-19.4	29	24.3	15.1-36.4	12	6.8	3.3-12.1
1977	58	18.3	13.6-24.1	34	25.1	16.6-36.1	24	13.3	8.2-20.2	51	17.5	12.6-23.4	31	25.5	16.3-37.4	20	12.1	7.0-18.9
1978	68	23.8	18.0-30.6	47	39.8	27.9-54.6	21	13.0	7.7-20.0	50	17.2	12.4-23.2	25	21.5	12.8-33.3	25	14.6	9.1-21.8
1979	67	21.5	16.2-27.8	31	24.0	14.9-35.9	36	20.3	13.9-28.3	42	13.9	9.7-19.1	23	18.9	11.0-29.7	19	10.8	6.3-17.0
1980	78	25.5	19.8-32.2	41	31.8	21.7-44.4	37	21.3	14.7-29.5	47	16.0	11.5-21.5	22	19.5	11.6-30.1	25	13.9	8.8-20.7
1981	61	19.1	14.3-24.8	40	29.8	20.1-42.1	21	12.0	7.3-18.4	62	20.0	15.1-26.0	38	27.0	18.2-38.0	24	14.8	9.3-22.0
1982	51	14.8	10.8-19.7	34	24.8	16.5-35.3	17	8.3	4.7-13.3	41	12.8	9.0-17.5	24	18.9	11.4-28.8	17	9.1	5.2-14.5
1983	61	16.0	12.0-20.7	33	21.4	14.2-30.8	28	12.5	8.1-18.3	41	11.2	7.8-15.4	23	15.7	9.4-24.4	18	8.4	4.8-13.4
1984	63	16.2	12.3-21.0	41	25.1	17.3-35.0	22	10.1	6.2-15.3	45	12.8	9.1-17.2	29	23.7	15.2-34.8	16	6.6	3.7-10.8
1985	63	16.5	12.4-21.4	38	21.6	14.4-30.8	25	12.3	7.8-18.1	46	11.3	8.2-15.2	25	13.1	8.2-19.7	21	9.5	5.8-14.6
1986	50	12.8	9.3-17.1	31	18.7	11.9-27.6	19	8.8	5.2-13.7	44	11.5	8.1-15.6	28	18.0	11.2-27.0	16	7.3	4.1-11.8
1987	76	19.7	15.3-24.8	46	27.9	19.6-38.2	30	13.9	9.3-19.8	46	11.5	8.2-15.5	27	14.9	9.5-22.0	19	8.6	5.1-13.5
1988	71	18.6	14.4-23.7	44	29.6	20.6-40.6	27	12.1	7.9-17.5	46	13.0	9.3-17.4	23	17.4	10.3-26.8	23	10.6	6.7-15.9
1989	74	17.3	13.4-21.9	43	22.7	15.6-31.5	31	13.1	8.8-18.6	58	13.8	10.3-17.9	39	22.2	15.2-31.1	19	8.0	4.7-12.4
1990	75	17.3	13.4-21.9	52	31.3	22.3-42.4	23	9.0	5.6-13.5	48	11.1	8.0-14.9	31	16.7	10.5-24.8	17	7.3	4.2-11.6
1991	69	14.8	11.3-18.8	38	20.8	13.9-29.6	31	11.3	7.6-16.1	54	11.4	8.4-15.1	36	19.5	12.9-28.0	18	6.5	3.8-10.2
1992	72	14.7	11.3-18.7	36	18.4	12.2-26.4	36	12.6	8.7-17.4	61	12.3	9.3-15.9	24	11.0	6.8-16.6	37	12.9	9.0-17.8
1993	85	18.4	14.6-23.0	54	27.3	19.8-36.6	31	12.0	8.1-17.0	44	9.6	6.8-12.9	29	15.0	9.4-22.4	15	5.9	3.3-9.7
1994	82	15.7	12.4-19.7	56	27.8	20.1-37.2	26	8.5	5.5-12.5	56	11.1	8.2-14.5	39	21.0	14.1-29.8	17	5.7	3.3-9.1
1995	85	16.7	13.2-20.8	41	20.8	14.2-29.0	44	14.3	10.3-19.2	53	10.7	7.9-14.1	31	16.6	10.6-24.3	22	7.3	4.5-11.0
1996	79	15.0	11.7-18.8	50	24.9	17.8-33.5	29	9.1	6.0-13.0	36	7.2	4.9-10.0	26	13.6	8.3-20.6	10	3.4	1.6-6.2
1997	74	12.7	9.9-16.1	46	19.8	13.9-27.0	28	8.4	5.5-12.1	38	6.6	4.6-9.1	27	11.7	7.5-17.2	11	3.2	1.6-5.7
1998	89	15.2	12.1-18.8	56	22.5	16.4-30.0	33	9.9	6.7-13.9	56	10.3	7.7-13.4	37	16.4	11.1-23.2	19	6.0	3.6-9.4
1999	80	13.9	10.9-17.4	50	22.5	15.9-30.5	30	9.1	6.1-13.0	56	10.0	7.4-13.0	35	16.3	10.7-23.5	21	6.4	3.9-9.7
2000	76	12.7	9.9-16.0	41	16.0	11.1-22.0	35	10.0	6.9-13.9	47	8.3	6.1-11.1	23	10.6	6.5-16.1	24	7.0	4.5-10.4
2001	81	13.5	10.6-16.8	42	18.1	12.5-25.1	39	11.0	7.8-15.0	36	5.7	3.9-7.9	24	8.4	5.1-13.0	12	3.5	1.8-6.1
2002	78	12.0	9.4-15.0	45	16.7	11.7-22.9	33	8.8	6.0-12.4	51	8.6	6.3-11.3	31	13.6	8.7-19.9	20	5.7	3.5-8.8
2003	71	10.0	7.7-12.7	45	14.9	10.5-20.3	26	6.6	4.3-9.7	38	5.4	3.8-7.4	28	9.3	5.9-13.9	10	2.7	1.3-4.9
2004	72	10.9	8.4-13.8	33	13.2	8.8-18.8	39	9.7	6.8-13.2	39	5.8	4.1-8.0	20	7.9	4.6-12.5	19	4.7	2.8-7.4
2005	101	14.3	11.6-17.5	45	15.3	10.8-20.9	56	13.6	10.2-17.7	64	10.1	7.7-12.9	39	15.0	10.3-20.9	25	6.7	4.3-9.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Stomach

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	100	14.1	11.4-17.2	60	20.9	15.6-27.3	40	9.6	6.9-13.1	45	6.5	4.7-8.7	28	9.4	6.0-13.8	17	4.4	2.5-6.9
2007	97	13.2	10.6-16.2	52	18.0	13.1-24.0	45	10.3	7.5-13.8	51	7.4	5.4-9.7	29	10.8	6.9-15.9	22	5.4	3.4-8.1
2008	79	10.2	8.0-12.8	51	16.1	11.7-21.4	28	6.0	4.0-8.7	57	7.6	5.7-9.9	33	11.2	7.5-16.0	24	5.3	3.4-7.9
2009	94	12.0	9.6-14.7	52	15.7	11.4-20.8	42	9.4	6.7-12.7	43	5.9	4.2-7.9	23	8.2	5.0-12.5	20	4.5	2.7-6.9
2010	85	10.5	8.4-13.1	51	14.8	10.7-19.7	34	7.6	5.2-10.6	46	6.0	4.3-8.0	32	10.5	7.0-15.0	14	3.1	1.7-5.1
2011	111	13.1	10.7-15.8	63	17.0	12.8-22.0	48	10.1	7.4-13.5	43	5.2	3.7-7.0	25	7.0	4.4-10.5	18	3.9	2.3-6.1
2012	103	12.1	9.8-14.7	58	15.3	11.4-20.0	45	9.2	6.7-12.3	50	6.3	4.7-8.4	29	8.4	5.5-12.3	21	4.6	2.9-7.1
2013	101	11.0	8.9-13.5	52	13.7	10.0-18.3	49	9.3	6.8-12.3	52	5.9	4.4-7.8	31	8.3	5.5-11.9	21	4.1	2.5-6.3
2014	114	11.7	9.6-14.1	71	17.0	13.0-21.7	43	7.8	5.6-10.6	55	6.0	4.5-7.9	38	10.1	7.0-14.1	17	3.3	1.9-5.2
2015	126	13.0	10.8-15.6	74	17.7	13.7-22.6	52	9.6	7.2-12.7	53	5.7	4.2-7.4	38	10.5	7.2-14.5	15	2.7	1.5-4.4
2016	119	11.9	9.8-14.3	66	15.0	11.4-19.4	53	9.5	7.1-12.4	45	4.4	3.1-5.9	26	5.5	3.5-8.3	19	3.3	2.0-5.2
2017	108	10.9	8.9-13.2	54	12.6	9.3-16.6	54	9.7	7.3-12.7	40	3.8	2.7-5.2	23	4.7	2.9-7.1	17	2.9	1.7-4.7
2018	136	13.0	10.8-15.4	74	16.8	13.0-21.3	62	10.3	7.9-13.3	46	4.5	3.2-6.0	28	6.9	4.4-10.0	18	3.0	1.8-4.7
2019	115	10.4	8.5-12.5	54	10.9	8.0-14.4	61	9.9	7.5-12.8	54	5.1	3.8-6.7	31	7.1	4.7-10.1	23	3.6	2.3-5.5
2020	100	8.6	6.9-10.5	50	9.9	7.2-13.2	50	7.6	5.6-10.0	39	3.6	2.5-4.9	18	3.9	2.3-6.3	21	3.2	2.0-5.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Colorectal

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	161	54.3	45.4-64.4	64	48.3	35.4-63.9	97	58.5	46.6-72.1	82	30.2	23.2-38.3	35	28.6	18.5-41.8	47	30.7	21.9-41.6
1977	189	62.8	53.3-73.3	82	66.1	50.8-84.2	107	60.7	49.0-74.2	96	33.5	26.6-41.5	51	41.0	29.5-55.0	45	27.8	19.7-37.6
1978	193	60.5	51.5-70.4	87	66.5	51.1-84.6	106	57.2	46.3-69.8	98	33.8	26.9-41.7	54	47.9	34.1-64.7	44	25.4	18.2-34.4
1979	204	63.0	53.9-73.0	104	72.9	57.8-90.5	100	55.4	44.5-68.0	121	40.8	33.2-49.4	45	34.9	24.3-48.1	76	44.1	34.2-55.8
1980	221	69.4	59.8-79.9	98	79.7	62.6-99.4	123	64.0	52.6-76.9	107	32.8	26.5-40.1	51	35.8	25.7-48.1	56	30.2	22.5-39.6
1981	207	63.5	54.6-73.3	97	75.7	59.7-94.1	110	56.4	45.9-68.4	126	39.8	32.7-47.8	61	48.3	35.9-63.2	65	34.3	26.1-44.1
1982	193	53.6	45.9-62.3	86	57.9	44.8-73.1	107	51.3	41.6-62.3	113	34.1	27.7-41.4	48	39.2	27.4-53.5	65	32.2	24.5-41.2
1983	200	57.9	49.7-67.1	86	63.9	49.5-80.8	114	55.1	45.1-66.4	99	28.5	22.8-35.0	46	31.8	22.1-43.8	53	26.3	19.5-34.6
1984	230	63.1	54.7-72.2	113	77.3	62.2-94.6	117	54.4	44.7-65.5	103	28.2	22.7-34.5	43	30.3	20.9-41.9	60	27.5	20.8-35.5
1985	216	57.2	49.4-65.9	101	66.1	52.3-82.0	115	51.7	42.4-62.3	116	32.6	26.6-39.4	49	34.2	24.4-46.0	67	31.6	24.3-40.3
1986	239	60.7	52.8-69.3	116	68.5	55.3-83.6	123	54.2	44.7-64.9	119	32.5	26.6-39.2	56	39.1	28.3-52.2	63	28.7	21.9-36.9
1987	247	60.6	52.9-69.1	126	77.4	62.8-94.1	121	50.6	41.8-60.7	125	31.6	26.0-38.0	71	49.7	37.5-64.0	54	21.9	16.3-28.7
1988	246	57.9	50.6-66.0	111	64.0	51.3-78.6	135	54.3	45.3-64.4	125	32.5	26.7-38.9	64	43.2	32.0-56.6	61	26.5	20.1-34.1
1989	262	60.0	52.6-68.1	125	72.9	59.0-88.6	137	53.2	44.4-63.0	127	32.1	26.5-38.4	75	53.8	41.0-68.7	52	20.9	15.5-27.5
1990	241	54.0	47.0-61.6	111	60.0	48.2-73.7	130	49.5	41.2-59.0	132	30.7	25.4-36.6	64	36.6	27.2-47.9	68	26.8	20.7-34.1
1991	243	50.6	44.2-57.7	119	56.5	45.8-68.9	124	45.5	37.7-54.4	126	28.3	23.3-33.9	64	36.8	27.3-48.3	62	23.3	17.8-29.9
1992	280	58.1	51.2-65.6	137	70.0	57.4-84.2	143	50.8	42.6-59.9	116	25.8	21.1-31.2	53	28.8	20.8-38.6	63	23.6	18.0-30.2
1993	265	53.1	46.6-60.1	124	56.9	46.2-69.1	141	49.3	41.4-58.3	139	29.3	24.4-34.8	70	36.2	27.3-46.7	69	24.8	19.2-31.5
1994	303	59.7	52.9-67.1	162	79.6	66.4-94.3	141	47.2	39.6-55.7	154	30.6	25.8-36.1	78	38.2	29.3-48.8	76	25.7	20.1-32.2
1995	319	60.1	53.4-67.3	147	68.5	56.7-81.9	172	55.4	47.3-64.4	127	25.7	21.2-30.7	63	34.1	25.2-44.7	64	21.4	16.4-27.3
1996	329	59.9	53.4-67.0	168	74.1	62.3-87.3	161	50.2	42.6-58.7	126	25.1	20.7-30.0	60	27.8	20.6-36.4	66	22.2	17.1-28.3
1997	320	56.6	50.4-63.4	158	70.2	58.6-83.2	162	48.5	41.2-56.6	148	27.2	22.8-32.1	78	34.6	26.5-44.2	70	22.2	17.3-28.1
1998	338	57.7	51.5-64.4	158	65.9	55.0-78.1	180	53.0	45.4-61.4	129	22.7	18.8-27.0	65	27.9	20.9-36.2	64	19.2	14.7-24.5
1999	392	65.2	58.7-72.3	194	77.9	66.2-90.9	198	57.3	49.5-65.9	153	28.3	23.8-33.2	73	35.5	27.0-45.5	80	24.3	19.2-30.2
2000	354	57.3	51.3-63.8	173	68.0	57.2-80.0	181	50.8	43.5-58.8	160	26.4	22.3-31.0	84	36.0	27.9-45.5	76	21.1	16.6-26.5
2001	371	57.3	51.4-63.6	192	72.6	61.8-84.5	179	47.3	40.5-54.8	126	20.4	16.9-24.4	62	25.7	19.2-33.4	64	17.6	13.5-22.4
2002	414	62.5	56.4-69.0	188	69.3	58.8-81.0	226	58.4	50.9-66.6	153	24.7	20.8-29.0	69	29.1	21.9-37.6	84	22.6	18.0-28.0
2003	399	59.8	53.8-66.1	206	76.8	65.7-89.1	193	49.7	42.8-57.3	138	21.4	17.9-25.4	67	26.4	19.9-34.1	71	18.4	14.3-23.2
2004	423	60.5	54.7-66.8	218	74.8	64.2-86.5	205	51.2	44.3-58.7	155	23.8	20.1-28.0	79	29.8	23.0-37.8	76	19.9	15.6-24.9
2005	397	55.0	49.5-60.8	199	67.3	57.5-78.1	198	47.0	40.6-54.1	141	20.6	17.2-24.4	67	23.9	18.1-30.8	74	18.1	14.2-22.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Colorectal

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	397	53.0	47.7-58.7	195	58.8	50.1-68.5	202	48.0	41.5-55.2	146	20.9	17.5-24.7	74	25.7	19.7-32.8	72	17.6	13.7-22.2
2007	387	49.1	44.2-54.4	196	56.9	48.6-66.1	191	43.0	37.1-49.7	177	23.8	20.3-27.7	89	28.3	22.3-35.4	88	20.5	16.4-25.3
2008	437	53.8	48.7-59.2	221	64.9	55.9-74.8	216	46.6	40.5-53.4	122	16.0	13.2-19.2	61	19.8	14.8-25.8	61	13.6	10.3-17.4
2009	415	48.5	43.8-53.5	212	57.3	49.3-66.1	203	41.9	36.2-48.2	145	18.0	15.1-21.2	76	23.1	17.9-29.3	69	14.4	11.1-18.2
2010	388	45.3	40.8-50.2	208	57.8	49.7-66.9	180	36.5	31.3-42.4	126	15.0	12.4-18.0	64	18.3	13.7-23.7	62	12.7	9.7-16.3
2011	411	46.2	41.7-51.1	215	55.5	47.9-64.0	196	39.0	33.6-44.9	137	16.2	13.5-19.3	71	20.8	15.9-26.6	66	13.3	10.2-16.9
2012	364	38.9	34.8-43.2	190	46.4	39.5-54.0	174	33.3	28.5-38.7	124	14.9	12.3-17.8	59	18.5	13.8-24.2	65	13.2	10.1-16.8
2013	389	40.8	36.8-45.2	194	47.2	40.3-54.8	195	36.3	31.3-41.9	137	14.7	12.2-17.4	71	18.0	13.8-23.0	66	12.2	9.4-15.5
2014	425	43.3	39.2-47.8	214	50.5	43.6-58.2	211	38.0	32.9-43.5	148	15.8	13.3-18.6	79	19.5	15.2-24.6	69	13.1	10.2-16.7
2015	436	42.3	38.3-46.6	227	50.1	43.4-57.4	209	36.2	31.4-41.5	156	16.5	13.9-19.3	72	17.8	13.7-22.6	84	15.3	12.2-19.0
2016	403	38.7	34.9-42.8	215	48.9	42.2-56.3	188	31.7	27.2-36.6	123	12.0	9.9-14.3	63	13.6	10.3-17.6	60	10.4	7.9-13.4
2017	439	41.4	37.5-45.5	234	51.0	44.3-58.3	205	34.7	30.0-39.9	141	13.5	11.3-16.0	85	19.6	15.4-24.4	56	9.3	7.0-12.1
2018	479	43.1	39.2-47.3	231	48.8	42.3-55.9	248	39.5	34.7-44.8	152	14.7	12.4-17.3	75	17.4	13.5-22.1	77	12.9	10.2-16.2
2019	451	39.8	36.1-43.8	216	43.8	37.8-50.4	235	37.3	32.6-42.5	144	13.3	11.2-15.8	74	17.0	13.2-21.5	70	10.8	8.4-13.7
2020	409	35.6	32.1-39.3	226	45.2	39.3-51.8	183	28.5	24.5-33.0	175	15.1	12.9-17.5	91	18.8	14.9-23.4	84	12.9	10.3-16.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Colon excluding rectum

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	110	37.6	30.2-46.2	41	32.0	21.2-45.7	69	41.7	31.7-53.4	68	24.3	18.1-31.6	30	24.1	14.8-36.6	38	24.1	16.4-33.8
1977	139	46.9	38.7-56.2	56	46.5	33.6-62.3	83	47.7	37.4-59.8	73	25.3	19.4-32.3	39	32.0	21.9-44.7	34	20.5	13.8-29.0
1978	141	43.5	36.1-51.9	64	47.7	35.1-62.9	77	41.1	32.1-51.8	78	26.7	20.6-33.9	43	39.0	26.4-54.7	35	19.7	13.4-27.6
1979	154	47.8	39.9-56.6	76	53.6	40.9-68.8	78	43.1	33.5-54.3	108	36.4	29.3-44.6	41	31.9	21.8-44.5	67	38.9	29.6-49.9
1980	159	49.6	41.6-58.5	68	56.5	42.1-73.7	91	46.5	37.0-57.5	83	25.6	20.0-32.2	41	30.0	20.6-41.7	42	22.5	15.9-30.6
1981	153	48.0	40.2-56.7	71	57.2	43.2-73.7	82	42.8	33.7-53.4	104	32.3	26.0-39.5	51	41.4	29.9-55.5	53	27.0	19.9-35.6
1982	146	40.7	34.0-48.3	69	47.5	35.8-61.5	77	36.7	28.6-46.1	95	29.2	23.3-36.1	38	34.4	23.1-48.4	57	27.7	20.8-36.1
1983	154	44.8	37.5-52.9	60	45.6	33.3-60.2	94	45.3	36.3-55.7	87	25.6	20.2-31.9	39	28.1	18.8-39.7	48	24.1	17.5-32.1
1984	177	49.2	41.8-57.4	83	57.5	44.6-72.6	94	44.1	35.3-54.1	89	24.5	19.4-30.4	36	26.2	17.4-37.3	53	24.2	18.0-31.7
1985	171	46.1	39.0-53.9	77	51.8	39.5-66.2	94	42.8	34.3-52.5	105	29.7	24.0-36.2	42	30.1	20.9-41.5	63	29.6	22.6-38.0
1986	173	43.6	36.9-50.9	77	45.7	35.0-58.3	96	41.5	33.4-51.0	99	27.2	21.8-33.4	45	30.6	21.4-42.0	54	24.9	18.5-32.6
1987	188	46.7	39.9-54.2	97	60.7	47.8-75.7	91	38.3	30.6-47.1	115	28.9	23.6-35.0	66	45.3	33.9-59.0	49	19.9	14.6-26.4
1988	197	46.1	39.6-53.4	86	48.8	37.8-61.6	111	44.7	36.5-53.9	105	27.0	21.8-32.9	55	36.1	26.0-48.3	50	21.7	16.0-28.7
1989	195	45.1	38.7-52.2	90	54.1	42.1-68.0	105	40.7	33.1-49.4	112	27.8	22.7-33.7	66	46.7	34.9-60.7	46	18.2	13.2-24.3
1990	179	39.7	33.8-46.2	82	44.6	34.4-56.7	97	36.6	29.5-44.8	112	25.8	21.0-31.2	55	30.1	21.9-40.0	57	22.5	16.9-29.3
1991	189	39.7	33.9-46.0	93	45.5	35.7-56.9	96	35.1	28.3-43.0	111	24.9	20.3-30.2	57	32.7	23.7-43.6	54	20.3	15.2-26.6
1992	219	46.1	40.0-52.9	104	53.5	42.6-66.0	115	41.4	34.0-49.7	96	21.0	16.8-25.8	44	23.1	16.2-31.8	52	19.2	14.3-25.2
1993	199	40.7	35.0-47.0	90	44.0	34.3-55.2	109	38.3	31.3-46.2	124	25.7	21.2-30.8	61	30.7	22.8-40.3	63	22.3	17.1-28.6
1994	226	45.5	39.5-52.1	121	60.3	48.8-73.4	105	36.1	29.4-43.7	135	27.1	22.5-32.3	67	33.7	25.2-43.8	68	23.0	17.8-29.2
1995	243	46.6	40.7-53.1	109	52.4	41.9-64.5	134	43.7	36.5-51.9	107	22.2	18.1-27.0	54	30.6	22.1-41.0	53	18.0	13.4-23.6
1996	235	43.2	37.7-49.3	106	47.0	37.6-57.7	129	40.4	33.6-48.1	99	19.5	15.7-23.9	44	20.2	14.3-27.6	55	18.2	13.7-23.7
1997	242	43.0	37.6-49.0	113	51.1	41.1-62.5	129	38.5	32.1-45.9	128	23.6	19.5-28.2	67	30.7	22.9-39.8	61	19.3	14.7-24.7
1998	264	46.0	40.4-52.1	124	53.4	43.6-64.6	140	41.7	35.0-49.3	112	19.7	16.1-23.8	55	23.4	17.1-31.1	57	17.1	12.9-22.2
1999	292	48.8	43.2-55.0	142	58.9	48.6-70.6	150	43.2	36.4-50.7	135	24.7	20.6-29.4	61	29.5	21.8-38.7	74	22.3	17.5-28.0
2000	275	45.1	39.8-51.0	134	54.6	44.9-65.7	141	39.8	33.4-46.9	135	22.7	18.9-27.0	72	32.5	24.6-41.7	63	17.6	13.5-22.6
2001	287	45.1	39.8-50.8	138	54.2	44.9-64.7	149	39.4	33.2-46.3	110	18.2	14.8-22.0	53	22.5	16.4-29.9	57	15.9	12.0-20.6
2002	334	51.7	46.1-57.7	147	57.8	48.0-68.9	187	48.6	41.8-56.2	133	21.6	18.0-25.7	57	24.6	18.0-32.6	76	20.5	16.1-25.6
2003	303	46.1	40.9-51.8	155	59.8	49.9-70.9	148	38.4	32.3-45.1	113	17.5	14.4-21.2	49	19.6	14.1-26.4	64	16.4	12.6-21.0
2004	329	48.0	42.8-53.6	166	58.5	49.0-69.2	163	41.3	35.1-48.1	138	21.2	17.7-25.1	68	25.3	19.1-32.7	70	18.3	14.2-23.2
2005	300	42.7	37.8-48.0	152	53.7	44.8-63.6	148	35.9	30.2-42.2	120	17.6	14.5-21.1	54	19.8	14.5-26.3	66	16.1	12.4-20.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Colon excluding rectum

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	296	40.1	35.5-45.1	142	43.4	35.9-51.9	154	37.1	31.4-43.5	125	18.0	14.9-21.5	65	22.4	16.9-29.0	60	14.7	11.2-19.0
2007	296	38.1	33.7-42.8	150	44.3	36.9-52.5	146	33.3	28.0-39.2	141	19.0	15.9-22.5	70	22.7	17.3-29.1	71	16.4	12.8-20.7
2008	332	41.4	37.0-46.3	157	47.5	39.8-56.1	175	37.9	32.4-44.0	106	14.1	11.5-17.1	52	17.4	12.6-23.1	54	12.1	9.0-15.8
2009	310	36.4	32.3-40.8	159	43.1	36.2-50.9	151	31.3	26.5-36.8	124	15.6	12.8-18.6	62	19.5	14.7-25.4	62	12.9	9.8-16.5
2010	283	33.6	29.7-37.9	150	42.3	35.3-50.1	133	27.3	22.8-32.4	103	12.5	10.1-15.2	52	15.1	11.0-20.2	51	10.5	7.8-13.8
2011	307	34.9	31.0-39.1	151	40.2	33.6-47.6	156	31.0	26.3-36.3	118	14.1	11.6-17.0	61	18.5	13.9-24.1	57	11.4	8.6-14.9
2012	260	28.4	24.9-32.1	134	33.7	27.8-40.5	126	24.5	20.3-29.2	111	13.2	10.8-16.0	50	15.2	11.0-20.3	61	12.4	9.5-16.0
2013	271	28.8	25.4-32.5	133	33.2	27.5-39.8	138	25.6	21.5-30.3	121	12.9	10.6-15.4	63	15.6	11.7-20.2	58	10.7	8.1-13.9
2014	305	31.6	28.1-35.5	150	36.2	30.3-42.8	155	28.3	24.0-33.2	124	13.3	11.0-15.9	62	15.3	11.5-19.8	62	11.9	9.1-15.3
2015	312	30.6	27.2-34.3	160	35.5	29.9-41.8	152	26.6	22.4-31.2	126	13.4	11.1-16.0	59	14.7	11.0-19.1	67	12.2	9.4-15.5
2016	291	28.0	24.8-31.5	147	33.9	28.4-40.2	144	24.2	20.3-28.6	107	10.3	8.4-12.5	54	11.4	8.4-15.1	53	9.2	6.8-12.1
2017	313	30.0	26.7-33.6	157	35.1	29.5-41.3	156	26.5	22.5-31.1	117	11.1	9.1-13.3	74	17.1	13.3-21.7	43	7.0	5.0-9.5
2018	354	32.2	28.9-35.9	164	35.6	30.1-41.8	190	30.3	26.0-35.0	123	12.0	9.9-14.3	60	14.2	10.6-18.4	63	10.6	8.1-13.5
2019	322	29.0	25.8-32.4	147	30.8	25.8-36.5	175	28.0	24.0-32.6	121	11.3	9.3-13.5	61	14.2	10.7-18.4	60	9.3	7.1-12.0
2020	296	26.1	23.2-29.3	155	31.9	26.9-37.5	141	22.0	18.5-26.0	143	12.5	10.5-14.8	71	15.2	11.7-19.3	72	11.0	8.6-13.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Rectum & rectosigmoid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	51	16.7	12.1-22.3	23	16.3	9.9-25.1	28	16.8	10.8-24.6	14	5.9	3.1-9.9	5	4.5	1.4-10.4	9	6.6	2.9-12.5
1977	50	15.9	11.4-21.4	26	19.6	12.1-29.6	24	13.0	8.0-19.8	23	8.2	4.9-12.7	12	9.0	4.2-16.3	11	7.3	3.4-13.2
1978	52	17.0	12.2-22.8	23	18.8	10.7-29.9	29	16.1	10.4-23.4	20	7.1	4.2-11.0	11	8.9	4.1-16.4	9	5.8	2.6-10.7
1979	50	15.2	11.0-20.4	28	19.3	11.9-29.2	22	12.4	7.5-18.9	13	4.4	2.1-7.7	4	3.1	0.6-8.3	9	5.2	2.2-10.0
1980	62	19.8	14.8-25.8	30	23.1	14.7-34.1	32	17.5	11.7-24.9	24	7.2	4.4-10.9	10	5.8	2.7-11.0	14	7.8	4.1-13.1
1981	54	15.5	11.4-20.5	26	18.5	11.4-27.8	28	13.7	8.9-19.9	22	7.4	4.4-11.5	10	6.9	3.0-13.0	12	7.3	3.6-12.8
1982	47	12.9	9.3-17.5	17	10.3	5.4-17.6	30	14.6	9.7-21.0	18	4.9	2.7-7.9	10	4.8	2.3-9.1	8	4.4	1.8-8.7
1983	46	13.1	9.4-17.7	26	18.4	11.3-27.7	20	9.8	5.9-15.1	12	2.9	1.5-5.1	7	3.7	1.4-7.9	5	2.2	0.7-5.1
1984	53	13.9	10.2-18.3	30	19.8	12.5-29.2	23	10.4	6.5-15.5	14	3.7	1.9-6.2	7	4.1	1.4-8.9	7	3.3	1.3-6.8
1985	45	11.2	8.0-15.1	24	14.3	8.7-21.9	21	9.0	5.4-13.8	11	2.9	1.4-5.3	7	4.1	1.5-8.6	4	2.0	0.5-4.9
1986	66	17.1	13.0-22.0	39	22.8	15.6-31.9	27	12.6	8.2-18.4	20	5.3	3.1-8.3	11	8.5	3.6-15.9	9	3.8	1.7-7.2
1987	59	14.0	10.4-18.2	29	16.7	10.5-25.0	30	12.4	8.2-17.7	10	2.7	1.2-5.0	5	4.3	1.2-10.1	5	2.0	0.6-4.6
1988	49	11.8	8.6-15.8	25	15.2	9.3-23.1	24	9.6	6.1-14.4	20	5.5	3.3-8.5	9	7.1	3.0-13.6	11	4.8	2.3-8.4
1989	67	14.9	11.4-19.0	35	18.8	12.3-27.0	32	12.4	8.4-17.5	15	4.3	2.3-7.0	9	7.1	3.0-13.4	6	2.7	1.0-5.7
1990	62	14.3	10.8-18.5	29	15.4	9.9-22.5	33	12.9	8.8-18.2	20	4.9	2.9-7.6	9	6.5	2.6-12.8	11	4.3	2.1-7.6
1991	54	11.0	8.1-14.4	26	11.1	7.0-16.6	28	10.4	6.8-15.0	15	3.4	1.8-5.6	7	4.1	1.4-8.7	8	3.0	1.3-5.8
1992	61	12.0	9.0-15.5	33	16.5	10.7-24.1	28	9.4	6.2-13.6	20	4.8	2.9-7.5	9	5.7	2.2-11.1	11	4.4	2.1-7.7
1993	66	12.4	9.4-15.8	34	12.9	8.8-18.3	32	11.1	7.5-15.6	15	3.6	2.0-6.0	9	5.4	2.1-10.7	6	2.5	0.9-5.3
1994	77	14.2	11.1-17.8	41	19.3	13.3-26.8	36	11.1	7.8-15.4	19	3.5	2.1-5.5	11	4.5	2.1-8.3	8	2.6	1.1-5.1
1995	76	13.4	10.5-16.9	38	16.1	11.0-22.6	38	11.6	8.2-16.0	20	3.5	2.1-5.4	9	3.5	1.5-6.8	11	3.4	1.7-6.0
1996	94	16.7	13.4-20.5	62	27.1	20.2-35.4	32	9.8	6.7-13.8	27	5.5	3.6-8.1	16	7.5	4.0-12.6	11	4.0	2.0-7.0
1997	78	13.6	10.7-17.1	45	19.1	13.5-26.0	33	9.9	6.8-13.9	20	3.6	2.1-5.6	11	4.0	1.9-7.4	9	3.0	1.3-5.5
1998	74	11.7	9.1-14.8	34	12.5	8.3-17.9	40	11.3	8.0-15.4	17	3.0	1.7-4.8	10	4.5	2.0-8.3	7	2.0	0.8-4.1
1999	100	16.4	13.2-20.0	52	19.0	13.8-25.3	48	14.1	10.4-18.7	18	3.6	2.1-5.6	12	6.0	2.9-10.7	6	2.0	0.7-4.1
2000	79	12.2	9.6-15.3	39	13.4	9.2-18.7	40	11.0	7.8-15.0	25	3.7	2.4-5.5	12	3.6	1.8-6.3	13	3.5	1.9-6.0
2001	84	12.3	9.7-15.2	54	18.3	13.4-24.5	30	7.9	5.3-11.3	16	2.2	1.3-3.6	9	3.2	1.3-6.2	7	1.7	0.7-3.4
2002	80	10.8	8.5-13.5	41	11.5	8.0-15.9	39	9.8	6.9-13.4	20	3.1	1.8-4.7	12	4.4	2.1-8.1	8	2.1	0.9-4.2
2003	96	13.7	11.0-16.8	51	17.0	12.2-22.9	45	11.4	8.3-15.2	25	3.9	2.4-5.7	18	6.7	3.7-11.1	7	2.0	0.8-4.0
2004	94	12.5	10.1-15.4	52	16.3	11.9-21.7	42	9.9	7.1-13.4	17	2.6	1.5-4.2	11	4.5	2.0-8.2	6	1.5	0.6-3.3
2005	97	12.3	9.9-15.0	47	13.6	9.8-18.4	50	11.1	8.2-14.7	21	3.0	1.8-4.7	13	4.1	2.1-7.1	8	2.1	0.9-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Rectum & rectosigmoid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	101	12.9	10.4-15.7	53	15.4	11.2-20.5	48	10.9	8.0-14.5	21	2.9	1.8-4.5	9	3.3	1.3-6.5	12	2.9	1.5-5.0
2007	91	11.0	8.8-13.6	46	12.7	9.0-17.2	45	9.8	7.1-13.1	36	4.8	3.3-6.7	19	5.7	3.2-9.1	17	4.1	2.3-6.5
2008	105	12.3	10.0-15.0	64	17.4	13.1-22.6	41	8.7	6.2-11.9	16	1.9	1.1-3.2	9	2.5	1.0-4.8	7	1.5	0.6-3.0
2009	105	12.1	9.8-14.7	53	14.1	10.4-18.7	52	10.6	7.8-13.9	21	2.4	1.5-3.7	14	3.6	1.9-6.2	7	1.5	0.6-3.1
2010	105	11.7	9.5-14.3	58	15.5	11.5-20.4	47	9.2	6.7-12.3	23	2.6	1.6-3.9	12	3.1	1.4-5.7	11	2.2	1.1-4.0
2011	104	11.3	9.2-13.8	64	15.4	11.6-19.9	40	8.0	5.7-10.9	19	2.1	1.3-3.3	10	2.3	1.1-4.2	9	1.8	0.8-3.5
2012	104	10.5	8.5-12.8	56	12.6	9.3-16.6	48	8.8	6.5-11.7	13	1.6	0.9-2.8	9	3.4	1.5-6.3	4	0.7	0.2-1.9
2013	118	12.0	9.9-14.5	61	14.0	10.4-18.3	57	10.7	8.1-13.9	16	1.8	1.0-2.9	8	2.4	1.0-4.8	8	1.4	0.6-2.8
2014	120	11.7	9.6-14.1	64	14.3	10.8-18.6	56	9.7	7.3-12.6	24	2.5	1.6-3.7	17	4.3	2.4-6.9	7	1.2	0.5-2.5
2015	124	11.7	9.7-14.0	67	14.6	11.1-18.7	57	9.6	7.3-12.5	30	3.1	2.1-4.4	13	3.1	1.5-5.4	17	3.1	1.8-5.0
2016	112	10.7	8.7-12.9	68	15.0	11.5-19.3	44	7.5	5.4-10.1	16	1.7	1.0-2.7	9	2.2	1.0-4.2	7	1.2	0.5-2.5
2017	126	11.4	9.5-13.6	77	15.9	12.4-20.1	49	8.2	6.0-10.9	24	2.4	1.5-3.6	11	2.4	1.2-4.4	13	2.2	1.2-3.8
2018	125	10.9	9.0-13.0	67	13.2	10.1-16.9	58	9.3	7.0-12.0	29	2.7	1.8-3.9	15	3.3	1.7-5.5	14	2.4	1.3-4.0
2019	129	10.8	9.0-12.9	69	12.9	10.0-16.6	60	9.3	7.0-12.0	23	2.1	1.3-3.1	13	2.8	1.4-4.8	10	1.5	0.7-2.8
2020	113	9.5	7.8-11.4	71	13.4	10.3-17.0	42	6.5	4.7-8.8	32	2.6	1.7-3.6	20	3.6	2.1-5.8	12	1.9	1.0-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Liver / intrahepatic bile duct

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	16	4.1	2.3-6.9	14	8.2	4.2-14.6	2	0.9	0.1-3.4	18	5.0	2.7-8.4	15	8.3	4.4-14.5	3	2.2	0.3-6.3
1977	13	3.9	1.9-7.0	9	4.9	2.2-9.7	4	2.8	0.6-7.0	12	2.8	1.4-5.0	12	6.2	3.1-11.3	0	0.0	0.0-1.9
1978	20	5.6	3.2-9.0	14	9.0	4.3-16.1	6	3.2	1.1-6.9	11	3.3	1.5-6.0	5	3.9	1.0-9.6	6	2.9	1.0-6.4
1979	13	3.7	1.8-6.5	8	4.5	1.8-9.4	5	2.8	0.8-6.6	10	3.0	1.2-5.8	7	3.4	1.3-7.5	3	2.3	0.4-6.3
1980	21	5.9	3.5-9.2	14	8.0	4.0-14.2	7	4.1	1.5-8.3	16	4.7	2.5-7.8	12	7.4	3.5-13.5	4	2.6	0.7-6.4
1981	24	6.2	3.9-9.4	14	9.0	4.5-15.6	10	4.4	2.1-8.2	25	7.4	4.6-11.1	14	9.2	4.6-16.1	11	5.9	2.8-10.6
1982	24	6.5	4.0-9.8	21	13.8	7.9-21.9	3	1.5	0.3-4.4	20	5.7	3.4-8.9	15	10.2	5.2-17.5	5	2.7	0.8-6.1
1983	19	4.7	2.7-7.5	17	10.1	5.5-16.8	2	0.9	0.1-3.2	24	6.1	3.8-9.2	19	11.8	6.6-19.2	5	2.3	0.7-5.2
1984	15	3.3	1.8-5.4	11	5.4	2.6-9.9	4	1.6	0.4-4.1	19	4.8	2.8-7.7	14	8.7	4.4-15.1	5	2.2	0.7-5.1
1985	15	3.3	1.8-5.5	5	2.2	0.7-5.5	10	4.0	1.9-7.3	9	2.4	1.1-4.6	3	1.9	0.3-5.5	6	2.8	1.0-5.9
1986	18	3.6	2.1-5.8	11	5.5	2.5-10.2	7	2.4	0.9-5.0	15	2.8	1.5-4.6	9	3.8	1.7-7.5	6	1.9	0.7-4.3
1987	17	3.5	2.0-5.7	9	3.7	1.6-7.3	8	3.2	1.3-6.3	20	4.5	2.7-7.1	12	5.7	2.7-10.4	8	3.6	1.5-6.9
1988	24	5.0	3.1-7.5	16	7.9	4.1-13.5	8	3.0	1.2-5.9	25	5.4	3.4-8.0	21	12.0	6.9-19.0	4	1.2	0.3-3.2
1989	28	6.1	4.0-8.9	23	11.4	6.8-17.8	5	2.3	0.7-5.1	21	4.9	2.9-7.5	12	7.1	3.3-12.9	9	3.6	1.6-6.7
1990	31	6.4	4.3-9.2	21	9.9	5.9-15.6	10	3.7	1.7-6.9	25	4.6	3.0-6.9	19	8.3	4.8-13.3	6	2.0	0.7-4.3
1991	33	5.9	3.9-8.4	24	9.5	5.9-14.5	9	2.9	1.3-5.7	27	4.5	2.9-6.6	19	6.9	4.1-11.0	8	2.4	1.0-4.8
1992	23	4.4	2.7-6.6	15	6.1	3.2-10.3	8	2.8	1.2-5.5	29	5.2	3.4-7.5	18	7.1	4.1-11.5	11	3.5	1.7-6.3
1993	26	5.4	3.4-8.0	11	4.4	2.1-8.1	15	5.4	3.0-8.9	22	4.4	2.7-6.7	8	2.7	1.1-5.6	14	5.0	2.7-8.3
1994	25	4.5	2.9-6.8	19	8.7	4.7-14.2	6	2.0	0.7-4.2	28	5.4	3.5-7.8	19	9.5	5.4-15.2	9	2.8	1.2-5.3
1995	42	7.7	5.5-10.5	31	13.5	8.8-19.7	11	3.6	1.8-6.4	30	5.5	3.7-7.9	23	10.8	6.5-16.5	7	2.1	0.9-4.4
1996	40	7.2	5.1-9.9	25	11.0	6.6-17.0	15	4.9	2.7-8.1	39	7.1	5.0-9.7	25	10.8	6.5-16.5	14	4.6	2.5-7.7
1997	51	8.8	6.5-11.6	33	14.2	9.4-20.3	18	5.3	3.1-8.4	33	5.7	3.9-8.1	20	8.5	4.9-13.5	13	3.9	2.1-6.7
1998	54	8.4	6.2-11.0	37	11.8	8.2-16.4	17	5.2	3.0-8.2	47	8.1	5.8-10.8	30	11.4	7.4-16.7	17	5.3	3.1-8.5
1999	30	4.6	3.0-6.5	21	6.6	4.0-10.3	9	2.6	1.2-4.9	25	4.4	2.8-6.5	14	5.2	2.7-8.9	11	3.4	1.7-6.0
2000	34	5.1	3.5-7.1	25	8.6	5.4-13.0	9	2.4	1.1-4.6	39	6.0	4.2-8.2	26	9.0	5.6-13.7	13	3.8	2.0-6.5
2001	44	6.0	4.3-8.1	31	9.1	6.1-13.1	13	3.3	1.7-5.6	37	5.3	3.7-7.3	21	6.7	3.9-10.6	16	4.2	2.4-6.8
2002	62	8.4	6.4-10.8	50	14.6	10.6-19.5	12	3.3	1.7-5.7	33	4.7	3.2-6.6	22	7.2	4.3-11.2	11	2.8	1.4-5.0
2003	62	8.3	6.3-10.6	42	11.1	7.9-15.2	20	5.4	3.3-8.3	42	5.3	3.8-7.2	35	9.4	6.4-13.2	7	1.8	0.7-3.7
2004	69	8.8	6.8-11.2	49	14.5	10.4-19.4	20	4.6	2.7-7.1	48	6.5	4.7-8.6	30	9.1	6.0-13.3	18	4.4	2.6-7.0
2005	82	10.8	8.5-13.5	61	18.2	13.7-23.8	21	5.2	3.2-7.9	53	6.4	4.7-8.4	40	10.6	7.4-14.7	13	3.0	1.6-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Liver / intrahepatic bile duct

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	66	8.1	6.2-10.3	52	13.9	10.2-18.4	14	3.3	1.8-5.5	38	5.2	3.6-7.2	22	6.4	3.9-9.7	16	4.0	2.3-6.5
2007	99	11.5	9.2-14.0	72	18.2	14.0-23.3	27	5.9	3.9-8.6	53	6.9	5.1-9.0	35	10.0	6.8-14.0	18	4.2	2.5-6.7
2008	98	10.4	8.4-12.8	74	17.8	13.8-22.7	24	4.7	3.0-7.0	59	6.8	5.1-8.8	46	12.0	8.6-16.4	13	2.8	1.5-4.7
2009	112	11.7	9.6-14.1	91	20.9	16.7-25.9	21	4.1	2.5-6.3	73	8.0	6.2-10.2	64	15.9	12.0-20.6	9	1.9	0.9-3.6
2010	101	10.3	8.3-12.5	83	18.4	14.5-23.0	18	3.5	2.1-5.6	79	8.2	6.4-10.2	59	13.8	10.3-18.0	20	3.7	2.3-5.8
2011	98	9.3	7.5-11.4	81	16.3	12.8-20.5	17	3.3	1.9-5.4	77	7.7	6.0-9.7	60	13.0	9.8-16.9	17	3.2	1.9-5.2
2012	102	9.6	7.8-11.7	78	15.9	12.4-20.1	24	4.5	2.8-6.7	89	9.1	7.3-11.3	65	14.0	10.6-18.2	24	5.0	3.2-7.5
2013	106	10.1	8.2-12.3	69	14.3	11.0-18.3	37	6.7	4.7-9.2	75	7.5	5.8-9.4	61	14.2	10.6-18.4	14	2.6	1.4-4.4
2014	145	13.3	11.2-15.8	116	24.1	19.7-29.2	29	4.9	3.2-7.0	72	7.0	5.4-8.8	50	11.3	8.2-15.1	22	3.9	2.4-5.9
2015	113	10.1	8.3-12.2	78	15.7	12.3-19.8	35	5.7	4.0-8.0	86	8.4	6.7-10.5	61	13.7	10.3-17.8	25	4.6	2.9-6.7
2016	162	13.9	11.8-16.3	124	23.7	19.5-28.5	38	6.1	4.3-8.4	87	8.3	6.6-10.3	67	15.0	11.5-19.3	20	3.4	2.0-5.2
2017	140	12.1	10.1-14.3	97	19.2	15.4-23.6	43	6.6	4.7-8.9	88	7.9	6.3-9.8	57	11.0	8.2-14.4	31	5.3	3.5-7.5
2018	107	9.0	7.3-10.9	76	14.0	10.9-17.6	31	5.0	3.4-7.1	81	7.0	5.5-8.7	52	10.1	7.5-13.4	29	4.6	3.0-6.6
2019	117	9.8	8.0-11.8	86	17.1	13.5-21.4	31	4.6	3.1-6.5	82	6.9	5.5-8.6	56	10.4	7.8-13.6	26	4.1	2.7-6.0
2020	114	9.2	7.6-11.1	74	13.2	10.3-16.7	40	5.9	4.2-8.1	71	6.2	4.8-7.9	47	9.5	6.8-12.7	24	3.7	2.4-5.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Pancreas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	54	17.5	12.7-23.3	21	13.9	8.2-22.0	33	19.7	13.2-28.2	38	12.0	8.2-16.8	19	12.5	7.4-19.9	19	11.3	6.5-17.9
1977	55	16.3	12.0-21.6	33	19.9	13.3-28.6	22	12.8	7.7-19.6	57	17.1	12.7-22.6	34	21.4	14.3-30.7	23	13.4	8.2-20.3
1978	70	21.6	16.4-27.7	40	27.5	18.5-39.0	30	17.2	11.2-24.9	50	15.4	11.1-20.6	27	19.1	11.7-29.0	23	12.6	7.8-19.2
1979	70	23.3	17.7-30.0	41	33.1	22.2-46.8	29	16.9	11.0-24.6	57	17.8	13.1-23.6	31	20.9	13.1-31.1	26	15.1	9.6-22.4
1980	50	14.5	10.5-19.4	27	20.1	12.4-30.2	23	11.1	6.9-16.8	47	13.8	9.8-18.6	27	18.6	11.4-28.4	20	10.5	6.2-16.4
1981	59	17.8	13.3-23.3	28	21.2	13.3-31.5	31	15.9	10.6-22.8	54	16.0	11.7-21.2	26	18.1	10.8-27.9	28	14.7	9.6-21.4
1982	64	17.1	12.9-22.0	33	20.1	13.3-29.0	31	14.8	9.9-21.1	66	18.4	14.0-23.7	33	23.2	15.1-33.5	33	15.6	10.6-22.1
1983	73	19.6	15.1-24.9	41	26.4	18.2-36.7	32	15.0	10.1-21.3	65	17.7	13.4-22.7	36	23.7	15.9-33.8	29	13.7	9.0-19.7
1984	73	20.2	15.6-25.6	30	23.4	14.9-34.4	43	19.2	13.7-25.9	68	17.5	13.3-22.4	36	23.9	15.9-34.0	32	13.7	9.2-19.4
1985	67	17.9	13.6-22.9	33	21.1	13.8-30.5	34	15.6	10.7-21.8	57	15.6	11.6-20.4	28	20.8	13.0-31.0	29	13.0	8.6-18.7
1986	69	16.2	12.4-20.6	32	16.9	11.1-24.5	37	15.3	10.7-21.1	66	16.2	12.4-20.8	34	18.2	12.1-26.2	32	14.3	9.7-20.2
1987	67	16.2	12.4-20.7	21	14.2	8.0-22.7	46	18.4	13.4-24.6	67	16.5	12.6-21.1	26	16.4	9.9-25.1	41	16.9	12.0-23.0
1988	65	17.5	13.3-22.4	34	22.2	14.8-31.6	31	14.2	9.5-20.1	67	17.5	13.4-22.3	38	24.4	16.6-34.1	29	12.9	8.5-18.5
1989	60	14.2	10.7-18.4	25	15.4	9.4-23.5	35	13.7	9.4-19.1	54	13.6	10.1-17.8	24	15.2	9.2-23.3	30	12.6	8.4-18.0
1990	92	22.2	17.7-27.5	35	22.9	15.2-32.6	57	22.5	16.9-29.2	72	17.2	13.2-21.7	20	13.7	7.7-21.9	52	20.1	14.9-26.4
1991	76	16.5	12.8-20.8	42	21.9	15.0-30.6	34	12.7	8.7-17.8	56	13.2	9.8-17.2	29	16.1	10.2-23.8	27	10.9	7.1-15.8
1992	70	15.5	11.9-19.7	33	18.6	12.0-27.0	37	13.8	9.6-19.0	62	14.6	11.0-18.8	28	17.7	11.0-26.5	34	13.2	9.1-18.4
1993	51	11.1	8.2-14.7	22	11.4	6.7-17.7	29	10.7	7.1-15.3	56	11.8	8.8-15.4	26	13.9	8.6-21.0	30	10.6	7.1-15.1
1994	79	16.6	13.0-20.8	36	18.7	12.7-26.4	43	15.2	10.9-20.4	66	14.6	11.2-18.6	33	18.5	12.2-26.6	33	12.2	8.3-17.0
1995	77	14.8	11.6-18.6	36	15.6	10.7-21.9	41	13.5	9.6-18.3	67	13.2	10.1-16.9	33	16.6	10.9-23.8	34	11.2	7.7-15.6
1996	87	15.6	12.4-19.4	43	19.7	13.8-27.1	44	13.4	9.7-17.9	75	14.0	10.9-17.7	33	17.7	11.7-25.3	42	12.7	9.1-17.1
1997	79	14.7	11.5-18.4	32	14.7	9.7-21.1	47	14.5	10.6-19.3	63	11.9	9.0-15.3	31	14.3	9.4-20.6	32	10.2	6.9-14.4
1998	82	14.5	11.4-18.1	30	13.4	8.6-19.8	52	15.6	11.6-20.5	86	16.2	12.8-20.1	33	15.6	10.3-22.4	53	16.6	12.4-21.7
1999	87	14.6	11.6-18.1	39	14.1	9.8-19.6	48	14.2	10.4-18.8	90	15.0	12.0-18.5	37	13.6	9.3-19.2	53	15.5	11.5-20.2
2000	83	14.3	11.3-17.8	32	12.0	7.9-17.4	51	15.0	11.1-19.7	72	12.4	9.6-15.7	31	12.7	8.3-18.3	41	11.9	8.5-16.2
2001	117	19.1	15.6-22.9	50	18.6	13.3-25.0	67	18.9	14.6-24.0	84	14.3	11.3-17.8	38	15.2	10.4-21.2	46	13.3	9.7-17.7
2002	100	15.4	12.5-18.8	50	18.3	13.3-24.4	50	13.3	9.8-17.5	81	13.0	10.3-16.3	42	15.3	10.7-21.1	39	11.0	7.8-15.0
2003	93	14.2	11.4-17.5	49	18.7	13.4-25.2	44	11.6	8.3-15.5	87	13.8	11.0-17.1	40	16.4	11.3-22.7	47	12.4	9.1-16.5
2004	99	15.0	12.1-18.3	42	14.7	10.2-20.4	57	14.8	11.2-19.2	84	13.0	10.3-16.1	42	15.9	11.0-22.0	42	11.0	7.9-14.9
2005	126	18.1	15.0-21.7	59	19.2	14.4-25.1	67	16.6	12.8-21.1	84	12.3	9.7-15.2	44	15.0	10.5-20.6	40	10.0	7.1-13.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Pancreas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	114	15.4	12.7-18.6	53	17.5	12.9-23.2	61	14.1	10.7-18.1	83	11.1	8.8-13.9	32	10.5	7.0-15.0	51	11.7	8.7-15.4
2007	106	14.5	11.8-17.6	42	13.4	9.4-18.4	64	14.9	11.4-19.1	76	10.3	8.1-13.0	34	10.7	7.3-15.2	42	9.7	7.0-13.2
2008	125	16.2	13.4-19.4	58	17.1	12.8-22.4	67	15.1	11.7-19.2	90	12.0	9.6-14.9	35	11.7	7.9-16.5	55	12.6	9.4-16.4
2009	124	15.4	12.7-18.5	69	18.7	14.4-24.0	55	12.4	9.3-16.1	112	14.2	11.6-17.2	50	14.2	10.3-19.0	62	13.7	10.5-17.6
2010	112	13.5	11.0-16.3	55	15.3	11.2-20.3	57	12.2	9.2-15.8	89	10.9	8.7-13.5	43	11.9	8.4-16.2	46	9.9	7.2-13.3
2011	147	16.9	14.2-20.0	69	17.5	13.3-22.5	78	16.0	12.6-20.0	99	12.2	9.8-14.9	43	12.4	8.7-17.0	56	11.8	8.9-15.3
2012	142	15.4	12.9-18.2	66	16.7	12.7-21.6	76	14.4	11.3-18.1	115	12.5	10.2-15.0	55	13.1	9.7-17.3	60	11.9	9.0-15.3
2013	128	13.8	11.5-16.5	63	15.6	11.7-20.3	65	12.4	9.5-15.8	97	11.0	8.9-13.5	46	12.8	9.1-17.2	51	9.7	7.2-12.8
2014	164	17.3	14.7-20.3	77	18.5	14.4-23.4	87	16.2	12.9-20.1	103	11.3	9.2-13.8	48	12.3	8.9-16.5	55	10.4	7.8-13.6
2015	167	17.0	14.4-19.8	68	15.4	11.7-19.8	99	17.9	14.5-21.8	112	11.8	9.7-14.3	50	11.5	8.4-15.4	62	11.7	8.9-15.0
2016	164	16.4	13.9-19.2	75	17.0	13.2-21.5	89	15.7	12.5-19.4	132	13.5	11.2-16.1	63	15.1	11.4-19.5	69	12.3	9.5-15.6
2017	177	16.9	14.5-19.7	81	17.0	13.3-21.4	96	16.5	13.3-20.2	126	12.1	10.0-14.4	61	13.4	10.1-17.4	65	10.9	8.4-13.9
2018	180	16.9	14.4-19.6	86	18.9	14.9-23.5	94	15.4	12.4-18.9	129	12.5	10.4-14.9	60	14.1	10.6-18.4	69	11.4	8.8-14.5
2019	205	18.6	16.1-21.4	92	19.8	15.7-24.4	113	17.8	14.6-21.4	148	13.6	11.4-16.0	72	14.7	11.4-18.8	76	12.3	9.6-15.4
2020	178	15.5	13.3-18.0	86	17.2	13.6-21.5	92	14.1	11.3-17.3	125	11.3	9.4-13.5	59	12.8	9.6-16.7	66	10.4	8.0-13.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	36	9.3	6.5-13.0	32	18.6	12.6-26.7	4	1.9	0.5-4.9	16	5.0	2.7-8.4	13	10.1	4.9-18.1	3	1.4	0.3-4.2
1977	22	6.0	3.6-9.4	20	12.8	7.0-21.1	2	1.1	0.1-3.8	11	2.6	1.3-4.8	9	4.7	2.1-9.5	2	0.9	0.1-3.4
1978	25	6.2	4.0-9.3	20	11.4	6.8-18.2	5	2.2	0.7-5.1	19	5.1	2.9-8.1	16	10.4	5.2-18.3	3	1.3	0.3-3.8
1979	41	10.0	7.1-13.7	32	17.9	11.9-25.9	9	3.9	1.8-7.5	10	2.6	1.2-4.9	10	6.1	2.8-11.6	0	0.0	0.0-1.7
1980	38	9.1	6.4-12.5	34	18.8	12.7-26.7	4	1.6	0.4-4.2	14	3.3	1.8-5.6	11	6.0	3.0-11.1	3	1.2	0.2-3.6
1981	39	8.9	6.3-12.3	32	16.0	10.8-22.9	7	3.1	1.2-6.5	19	4.7	2.7-7.6	16	10.0	5.2-17.1	3	1.2	0.2-3.5
1982	36	8.0	5.6-11.2	34	17.3	11.7-24.7	2	0.8	0.1-2.9	19	4.7	2.7-7.4	16	9.2	4.7-15.7	3	1.4	0.3-4.0
1983	24	5.3	3.4-8.0	22	10.5	6.5-16.1	2	1.1	0.1-3.5	18	4.0	2.4-6.4	15	7.7	4.3-13.0	3	1.2	0.2-3.4
1984	36	8.0	5.6-11.2	27	14.1	8.9-21.2	9	3.5	1.6-6.7	11	2.9	1.4-5.1	10	6.3	2.7-12.2	1	0.6	0.0-2.7
1985	42	8.8	6.3-12.0	30	14.0	9.3-20.2	12	4.7	2.4-8.2	15	3.7	2.0-6.1	13	8.0	4.0-14.0	2	0.8	0.1-2.8
1986	39	8.3	5.8-11.4	32	16.2	10.8-23.4	7	2.5	1.0-5.1	16	3.5	2.0-5.7	11	5.8	2.8-10.6	5	1.9	0.6-4.4
1987	29	5.8	3.9-8.4	24	11.1	7.0-16.7	5	1.8	0.6-4.1	12	2.6	1.3-4.6	7	3.8	1.4-8.0	5	1.9	0.6-4.4
1988	24	5.0	3.1-7.5	17	8.1	4.5-13.3	7	2.7	1.1-5.5	17	3.5	2.0-5.6	17	8.2	4.6-13.3	0	0.0	0.0-1.2
1989	38	7.2	5.0-9.9	25	11.5	7.1-17.6	13	4.2	2.2-7.2	11	2.2	1.1-4.0	10	4.8	2.2-8.9	1	0.4	0.0-1.9
1990	33	6.1	4.2-8.7	28	12.0	7.7-17.8	5	1.7	0.6-4.0	16	3.2	1.8-5.2	14	6.8	3.3-12.0	2	0.7	0.1-2.6
1991	28	5.3	3.5-7.8	24	10.6	6.5-16.1	4	1.4	0.4-3.6	17	3.3	1.9-5.3	17	8.0	4.4-13.2	0	0.0	0.0-1.1
1992	39	7.1	5.0-9.8	28	12.1	7.9-17.8	11	3.5	1.7-6.3	17	3.1	1.8-4.9	12	5.3	2.6-9.6	5	1.5	0.5-3.5
1993	38	6.7	4.7-9.2	34	14.2	9.5-20.1	4	1.2	0.3-3.1	16	2.8	1.6-4.5	14	5.3	2.9-9.0	2	0.7	0.1-2.4
1994	41	6.9	4.9-9.5	38	15.4	10.6-21.6	3	0.9	0.2-2.5	18	3.4	2.0-5.5	14	6.8	3.3-11.8	4	1.3	0.4-3.3
1995	37	6.3	4.4-8.7	30	12.4	8.1-18.0	7	2.1	0.8-4.2	18	3.4	2.0-5.4	15	7.4	4.0-12.4	3	0.9	0.2-2.5
1996	40	6.7	4.7-9.1	34	13.9	9.2-20.0	6	1.7	0.6-3.7	17	2.9	1.7-4.7	15	6.8	3.4-11.7	2	0.6	0.1-2.1
1997	42	6.6	4.7-9.0	33	12.7	8.5-18.2	9	2.4	1.1-4.5	12	1.8	0.9-3.2	9	3.2	1.4-6.2	3	0.8	0.2-2.2
1998	49	7.4	5.5-9.9	35	12.7	8.5-18.0	14	3.8	2.0-6.3	15	2.4	1.3-4.0	14	5.3	2.7-9.1	1	0.3	0.0-1.6
1999	37	5.3	3.7-7.3	29	9.7	6.3-14.3	8	2.0	0.9-3.9	20	3.1	1.9-4.8	19	7.2	4.1-11.5	1	0.2	0.0-1.3
2000	33	4.6	3.1-6.4	23	7.4	4.6-11.3	10	2.4	1.1-4.4	13	2.1	1.1-3.6	7	3.3	1.2-6.9	6	1.5	0.6-3.3
2001	38	5.1	3.6-7.0	32	9.7	6.6-13.9	6	1.5	0.5-3.2	10	1.5	0.7-2.7	9	3.5	1.4-6.9	1	0.2	0.0-1.3
2002	37	4.8	3.4-6.7	28	8.3	5.5-12.2	9	2.0	0.9-3.9	24	3.3	2.1-5.0	18	5.4	3.2-8.6	6	1.6	0.6-3.4
2003	42	5.7	4.1-7.8	32	10.3	6.8-14.8	10	2.5	1.2-4.6	20	2.7	1.6-4.2	14	4.5	2.4-7.7	6	1.4	0.5-3.1
2004	40	5.0	3.6-6.9	36	10.5	7.2-14.7	4	0.9	0.2-2.3	14	1.8	1.0-3.0	12	3.5	1.8-6.2	2	0.5	0.1-1.7
2005	33	4.4	3.0-6.2	28	9.3	5.9-13.7	5	1.1	0.4-2.6	14	1.7	0.9-2.9	13	3.9	1.9-6.9	1	0.2	0.0-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	48	5.9	4.3-7.8	39	11.2	7.8-15.5	9	1.9	0.9-3.7	11	1.6	0.8-2.8	10	3.4	1.6-6.3	1	0.3	0.0-1.3
2007	46	5.8	4.2-7.7	31	9.5	6.2-13.8	15	3.2	1.8-5.3	11	1.4	0.7-2.6	7	2.1	0.8-4.5	4	0.9	0.2-2.3
2008	47	5.6	4.1-7.5	37	10.6	7.2-14.9	10	2.2	1.0-4.0	20	2.4	1.5-3.8	15	4.0	2.2-6.8	5	1.1	0.4-2.5
2009	34	4.0	2.8-5.7	24	6.7	4.2-10.2	10	2.1	1.0-3.9	18	2.2	1.3-3.4	13	3.9	1.9-6.9	5	1.0	0.3-2.4
2010	43	5.0	3.6-6.8	33	9.2	6.2-13.1	10	2.0	0.9-3.7	16	2.0	1.2-3.3	14	4.3	2.3-7.3	2	0.4	0.0-1.5
2011	42	4.3	3.1-5.9	36	8.5	5.8-11.9	6	1.1	0.4-2.4	11	1.1	0.5-2.0	9	2.0	0.9-3.9	2	0.4	0.0-1.5
2012	31	3.3	2.2-4.7	21	5.5	3.3-8.5	10	1.8	0.9-3.3	21	2.4	1.4-3.6	19	5.4	3.1-8.5	2	0.4	0.0-1.3
2013	33	3.3	2.2-4.7	23	5.7	3.5-8.7	10	1.6	0.8-3.0	15	1.7	0.9-2.8	14	4.1	2.2-7.0	1	0.2	0.0-1.0
2014	45	4.6	3.3-6.2	38	9.2	6.4-12.8	7	1.2	0.5-2.6	17	1.8	1.0-2.9	15	4.0	2.2-6.7	2	0.3	0.0-1.2
2015	45	4.3	3.1-5.8	39	8.8	6.1-12.2	6	1.0	0.4-2.3	13	1.4	0.7-2.4	12	3.3	1.6-5.9	1	0.2	0.0-1.0
2016	45	4.2	3.0-5.7	35	7.8	5.3-10.9	10	1.5	0.7-2.8	15	1.4	0.8-2.4	11	2.5	1.2-4.6	4	0.6	0.2-1.6
2017	32	2.9	2.0-4.2	24	5.1	3.2-7.8	8	1.3	0.6-2.6	9	1.0	0.4-1.8	7	1.8	0.7-3.6	2	0.4	0.0-1.4
2018	31	2.6	1.7-3.7	23	4.4	2.8-6.7	8	1.1	0.5-2.3	10	0.9	0.4-1.6	9	1.8	0.8-3.4	1	0.2	0.0-0.9
2019	32	2.6	1.8-3.7	27	4.9	3.2-7.3	5	0.7	0.2-1.7	13	1.0	0.5-1.7	9	1.7	0.7-3.3	4	0.5	0.1-1.3
2020	42	3.5	2.5-4.7	34	6.5	4.5-9.1	8	1.2	0.5-2.4	8	0.7	0.3-1.4	8	1.7	0.7-3.5	0	0.0	0.0-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Lung and bronchus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	224	62.9	54.5-72.0	174	111.8	94.7-130.9	50	24.6	18.1-32.7	156	44.2	37.2-52.1	122	80.3	65.3-97.6	34	16.6	11.4-23.3
1977	286	82.2	72.3-93.0	220	149.4	128.0-173.1	66	32.5	24.7-41.7	205	59.4	51.0-68.6	165	112.9	94.5-133.5	40	19.8	13.9-27.3
1978	270	74.4	65.3-84.4	203	128.7	110.0-149.5	67	33.7	25.7-43.2	216	60.8	52.4-70.1	169	111.8	93.6-132.1	47	23.6	17.0-31.7
1979	278	76.1	66.8-86.3	200	134.3	113.9-156.9	78	35.5	27.9-44.6	217	61.3	52.8-70.6	166	112.4	94.0-133.0	51	25.1	18.3-33.4
1980	334	87.3	77.7-97.7	254	160.6	139.3-183.9	80	35.4	27.9-44.3	262	71.2	62.3-81.0	193	129.2	109.3-151.4	69	31.6	24.4-40.2
1981	332	85.4	76.0-95.6	250	154.6	134.2-177.0	82	36.3	28.6-45.3	258	67.9	59.4-77.3	190	123.5	104.4-144.7	68	29.9	23.1-38.1
1982	326	80.1	71.2-89.7	244	145.1	125.6-166.5	82	34.5	27.2-43.0	265	65.4	57.4-74.2	197	115.5	98.5-134.4	68	29.5	22.8-37.6
1983	337	80.6	71.8-90.0	241	138.2	120.0-158.2	96	39.6	31.8-48.5	293	73.7	65.0-83.0	225	137.7	118.7-158.6	68	29.2	22.4-37.2
1984	325	76.4	67.9-85.6	228	124.2	107.4-142.8	97	41.2	33.2-50.6	273	63.9	56.3-72.3	197	110.4	94.2-128.4	76	31.1	24.4-39.1
1985	353	79.5	71.1-88.5	250	133.2	116.0-152.1	103	40.7	33.1-49.5	274	63.0	55.5-71.3	198	107.2	91.7-124.5	76	31.1	24.3-39.1
1986	371	84.8	76.0-94.3	262	149.1	129.6-170.5	109	42.4	34.7-51.3	293	67.0	59.3-75.5	199	112.5	95.9-130.8	94	36.7	29.6-45.1
1987	337	72.9	65.0-81.3	234	119.9	104.0-137.4	103	39.1	31.7-47.5	325	71.7	63.8-80.3	234	126.7	109.5-145.7	91	34.6	27.7-42.6
1988	394	85.9	77.3-95.1	273	148.4	129.7-168.9	121	44.9	37.2-53.8	292	63.7	56.4-71.7	204	110.0	94.2-127.4	88	32.8	26.2-40.5
1989	410	84.9	76.6-93.8	286	145.1	127.2-164.5	124	44.4	36.8-53.1	378	79.6	71.4-88.3	258	134.4	117.2-153.3	120	43.1	35.6-51.6
1990	384	80.5	72.3-89.3	273	140.2	122.6-159.5	111	40.3	33.0-48.6	331	69.6	62.0-77.8	224	115.3	99.3-133.0	107	39.2	32.0-47.4
1991	412	83.8	75.6-92.6	265	135.4	117.9-154.5	147	50.7	42.7-59.7	346	71.2	63.6-79.4	221	114.1	98.2-131.6	125	43.3	35.9-51.7
1992	420	81.8	73.9-90.3	295	145.1	127.2-164.6	125	41.1	34.1-49.1	326	65.6	58.4-73.5	227	116.6	100.3-134.6	99	33.4	27.0-40.8
1993	406	77.6	70.0-85.8	268	129.5	112.8-147.7	138	44.9	37.6-53.2	362	70.6	63.2-78.5	252	124.5	108.0-142.6	110	36.3	29.8-43.9
1994	440	79.9	72.3-87.9	256	118.8	103.3-135.8	184	56.3	48.4-65.2	369	67.9	60.9-75.4	260	118.9	103.5-135.7	109	34.4	28.2-41.5
1995	442	79.9	72.3-87.9	281	121.7	106.7-138.0	161	51.0	43.3-59.6	352	64.4	57.7-71.7	212	97.8	84.0-113.0	140	43.3	36.3-51.2
1996	456	79.7	72.3-87.6	277	118.5	103.8-134.6	179	53.9	46.2-62.5	359	63.7	57.1-70.9	215	91.1	78.3-105.2	144	44.7	37.6-52.7
1997	464	78.8	71.6-86.5	293	120.8	106.3-136.6	171	50.1	42.8-58.3	371	64.1	57.5-71.1	236	102.0	88.4-116.9	135	39.5	33.1-46.8
1998	467	76.9	69.8-84.4	273	108.1	94.6-122.9	194	55.6	47.9-64.0	375	63.6	57.1-70.6	239	99.5	86.4-114.0	136	39.4	33.0-46.7
1999	460	75.3	68.3-82.7	265	105.1	91.8-119.6	195	55.4	47.8-63.8	336	56.1	50.1-62.6	204	82.8	70.8-96.1	132	38.6	32.2-45.8
2000	482	74.7	68.0-81.9	306	113.1	99.8-127.5	176	48.1	41.2-55.9	374	58.9	52.9-65.3	231	89.7	77.6-102.9	143	38.6	32.5-45.5
2001	492	74.7	68.0-81.8	300	111.7	98.6-126.1	192	50.1	43.2-57.8	353	55.7	49.8-61.9	211	82.1	70.5-94.8	142	38.4	32.2-45.3
2002	465	69.9	63.5-76.7	273	101.3	88.6-115.0	192	49.5	42.7-57.1	371	55.8	50.1-62.0	233	86.1	74.6-98.9	138	35.8	30.0-42.4
2003	476	67.9	61.7-74.5	272	93.3	81.7-106.0	204	50.5	43.7-58.0	373	55.1	49.5-61.2	226	80.8	69.8-92.9	147	37.3	31.5-44.0
2004	506	69.0	62.9-75.5	299	97.0	85.5-109.5	207	49.3	42.8-56.6	372	54.1	48.6-60.1	224	81.9	70.7-94.2	148	36.3	30.6-42.8
2005	500	68.3	62.3-74.8	297	99.7	87.7-112.7	203	48.1	41.6-55.3	384	52.7	47.4-58.5	219	73.0	62.8-84.3	165	39.5	33.6-46.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Lung and bronchus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	496	67.3	61.3-73.7	289	94.0	82.7-106.3	207	49.4	42.8-56.7	324	44.6	39.7-49.9	186	61.1	52.0-71.2	138	32.9	27.6-39.0
2007	524	67.4	61.6-73.6	303	93.9	82.9-105.8	221	49.2	42.8-56.2	369	47.8	42.9-53.1	209	65.0	55.9-75.1	160	36.1	30.7-42.2
2008	503	63.3	57.7-69.2	271	82.0	71.8-93.1	232	50.8	44.4-57.9	326	42.1	37.5-47.1	182	56.4	47.9-65.9	144	32.0	26.9-37.8
2009	556	67.2	61.5-73.2	307	88.9	78.5-100.2	249	52.6	46.2-59.7	356	44.1	39.5-49.1	203	60.7	52.1-70.3	153	32.9	27.8-38.6
2010	564	66.1	60.6-72.0	298	83.5	73.6-94.3	266	54.7	48.2-61.8	378	46.4	41.7-51.4	223	68.3	59.0-78.5	155	32.6	27.6-38.3
2011	543	61.0	55.8-66.6	298	79.8	70.3-90.0	245	48.2	42.3-54.8	401	46.7	42.1-51.6	228	63.7	55.2-73.1	173	35.0	29.9-40.7
2012	578	63.4	58.2-69.0	338	88.2	78.5-98.8	240	46.1	40.4-52.4	379	42.9	38.6-47.6	215	59.3	51.2-68.4	164	32.0	27.2-37.4
2013	558	58.3	53.4-63.5	297	75.2	66.4-84.8	261	46.7	41.1-52.9	376	41.4	37.2-46.0	211	55.8	48.0-64.3	165	31.3	26.6-36.5
2014	564	58.1	53.3-63.3	308	75.1	66.4-84.5	256	46.1	40.5-52.2	361	37.4	33.5-41.6	205	50.8	43.7-58.7	156	27.9	23.7-32.8
2015	634	63.4	58.4-68.7	351	85.7	76.4-95.6	283	48.3	42.8-54.4	361	37.9	34.0-42.2	194	49.4	42.3-57.2	167	30.3	25.8-35.3
2016	618	60.8	55.9-65.9	333	77.2	68.7-86.5	285	48.9	43.3-55.1	348	34.5	30.8-38.4	202	49.5	42.5-57.2	146	24.4	20.6-28.8
2017	587	55.5	51.0-60.4	309	71.0	62.9-79.8	278	45.7	40.4-51.5	326	31.6	28.1-35.3	177	41.9	35.6-48.9	149	24.7	20.8-29.1
2018	578	52.8	48.4-57.4	291	64.7	57.1-73.0	287	45.4	40.2-51.1	326	31.1	27.7-34.7	176	41.8	35.5-48.7	150	24.1	20.4-28.4
2019	617	54.7	50.3-59.3	320	67.7	60.1-76.0	297	46.0	40.8-51.6	321	28.6	25.5-32.0	179	39.1	33.2-45.6	142	21.6	18.2-25.6
2020	548	46.5	42.6-50.7	285	56.1	49.4-63.4	263	39.7	35.0-44.9	282	24.5	21.7-27.7	160	33.6	28.3-39.5	122	18.6	15.4-22.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Melanoma of the skin

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2	0.6	0.1-2.2	1	0.9	0.0-4.7	1	0.4	0.0-2.7	4	1.7	0.4-4.4	1	0.5	0.0-3.8	3	2.4	0.4-6.6
1977	6	3.0	1.0-6.4	5	5.9	1.5-13.9	1	1.1	0.0-4.8	3	1.4	0.2-3.9	2	3.3	0.3-10.8	1	0.4	0.0-2.5
1978	4	1.4	0.3-3.7	2	2.8	0.2-9.6	2	0.8	0.1-3.2	1	0.4	0.0-2.1	0	0.0	0.0-2.9	1	0.7	0.0-3.4
1979	1	0.2	0.0-1.4	1	0.5	0.0-3.4	0	0.0	0.0-1.7	2	0.9	0.1-3.1	1	0.5	0.0-3.5	1	1.0	0.0-4.2
1980	4	1.4	0.3-3.6	2	1.0	0.1-4.1	2	1.5	0.2-4.7	0	0.0	0.0-1.0	0	0.0	0.0-2.6	0	0.0	0.0-1.6
1981	7	1.4	0.5-3.1	2	0.7	0.1-3.5	5	1.9	0.6-4.6	1	0.4	0.0-1.8	1	1.0	0.0-4.8	0	0.0	0.0-1.6
1982	9	1.7	0.8-3.3	6	2.6	0.9-6.0	3	0.9	0.2-3.0	3	0.7	0.2-2.2	2	1.2	0.1-4.5	1	0.4	0.0-2.2
1983	2	0.6	0.1-2.0	2	1.4	0.1-5.0	0	0.0	0.0-1.5	2	0.6	0.1-1.9	1	0.4	0.0-2.9	1	0.6	0.0-2.8
1984	5	1.5	0.4-3.4	2	1.7	0.1-5.9	3	1.4	0.3-4.0	1	0.1	0.0-1.0	1	0.2	0.0-2.5	0	0.0	0.0-1.4
1985	6	1.6	0.5-3.6	3	1.4	0.3-4.3	3	1.6	0.3-4.5	2	0.7	0.1-2.3	1	1.2	0.0-5.3	1	0.5	0.0-2.3
1986	6	1.2	0.4-2.7	3	1.1	0.2-3.7	3	1.2	0.2-3.5	7	1.3	0.5-2.7	5	2.1	0.7-5.2	2	0.7	0.1-2.5
1987	5	1.2	0.4-2.8	3	1.7	0.3-5.2	2	0.8	0.1-2.9	1	0.2	0.0-1.1	1	0.5	0.0-2.8	0	0.0	0.0-1.3
1988	5	1.0	0.3-2.3	2	1.1	0.1-4.0	3	1.0	0.2-2.9	2	0.4	0.0-1.5	2	1.0	0.1-3.8	0	0.0	0.0-1.2
1989	4	1.0	0.2-2.5	2	0.6	0.1-2.7	2	1.0	0.1-3.3	5	1.1	0.3-2.5	2	1.1	0.1-4.1	3	1.1	0.2-3.2
1990	4	0.8	0.2-2.0	1	0.4	0.0-2.5	3	1.0	0.2-3.0	0	0.0	0.0-0.7	0	0.0	0.0-1.9	0	0.0	0.0-1.2
1991	5	1.0	0.3-2.4	1	1.4	0.0-5.5	4	1.1	0.3-3.0	2	0.5	0.1-1.8	0	0.0	0.0-1.9	2	0.8	0.1-2.7
1992	3	0.5	0.1-1.4	2	0.7	0.1-2.8	1	0.3	0.0-1.7	1	0.1	0.0-0.9	1	0.3	0.0-2.2	0	0.0	0.0-1.1
1993	8	1.6	0.7-3.2	1	1.2	0.0-5.1	7	2.3	0.9-4.6	3	0.4	0.1-1.3	0	0.0	0.0-1.7	3	0.8	0.2-2.3
1994	5	0.9	0.3-2.1	2	0.5	0.1-2.4	3	1.0	0.2-2.9	3	0.6	0.1-1.6	0	0.0	0.0-1.7	3	1.0	0.2-2.8
1995	6	0.9	0.3-2.0	3	0.8	0.2-2.8	3	0.9	0.2-2.5	4	0.5	0.1-1.3	1	0.2	0.0-1.8	3	0.7	0.1-2.2
1996	5	0.8	0.2-2.0	3	1.6	0.2-5.0	2	0.5	0.1-1.9	4	0.6	0.2-1.5	1	0.3	0.0-2.0	3	0.8	0.2-2.4
1997	7	1.2	0.5-2.5	1	0.3	0.0-2.0	6	1.8	0.6-3.9	2	0.4	0.0-1.3	1	0.2	0.0-1.7	1	0.4	0.0-1.8
1998	8	1.3	0.5-2.6	3	0.9	0.2-2.8	5	1.5	0.5-3.4	3	0.4	0.1-1.2	1	0.2	0.0-1.6	2	0.6	0.1-2.0
1999	4	0.5	0.1-1.3	2	0.5	0.1-2.1	2	0.5	0.1-1.8	1	0.2	0.0-0.8	1	0.3	0.0-2.0	0	0.0	0.0-0.9
2000	0	0.0	0.0-0.5	0	0.0	0.0-1.3	0	0.0	0.0-0.9	3	0.6	0.1-1.7	2	1.1	0.1-3.6	1	0.3	0.0-1.6
2001	6	0.7	0.3-1.6	2	0.5	0.1-2.0	4	0.9	0.3-2.4	4	0.7	0.2-1.7	2	0.6	0.1-2.4	2	0.6	0.1-2.1
2002	8	1.3	0.5-2.5	4	1.5	0.4-3.9	4	1.1	0.3-2.8	2	0.4	0.0-1.3	1	0.9	0.0-3.7	1	0.3	0.0-1.4
2003	8	1.1	0.5-2.2	2	0.9	0.1-3.1	6	1.3	0.5-2.9	5	0.7	0.2-1.6	2	0.9	0.1-3.1	3	0.6	0.1-1.8
2004	9	1.1	0.5-2.0	5	1.3	0.4-3.2	4	0.9	0.2-2.2	1	0.1	0.0-0.6	1	0.2	0.0-1.5	0	0.0	0.0-0.8
2005	8	0.9	0.4-1.9	5	1.1	0.3-2.7	3	0.7	0.1-2.0	2	0.2	0.0-0.8	1	0.2	0.0-1.4	1	0.2	0.0-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Melanoma of the skin

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	11	1.5	0.7-2.6	7	1.9	0.7-4.1	4	1.1	0.3-2.6	4	0.6	0.1-1.4	2	0.7	0.1-2.5	2	0.5	0.1-1.7
2007	10	1.2	0.6-2.3	1	0.2	0.0-1.4	9	2.0	0.9-3.8	1	0.1	0.0-0.6	0	0.0	0.0-1.1	1	0.2	0.0-1.1
2008	12	1.4	0.7-2.5	4	0.9	0.2-2.4	8	1.8	0.8-3.5	2	0.3	0.0-0.9	0	0.0	0.0-1.0	2	0.5	0.1-1.6
2009	8	0.8	0.3-1.6	2	0.4	0.0-1.5	6	1.1	0.4-2.5	4	0.5	0.1-1.3	2	0.6	0.1-2.3	2	0.4	0.0-1.5
2010	10	1.0	0.5-1.8	3	0.7	0.1-2.2	7	1.2	0.5-2.6	5	0.6	0.2-1.4	1	0.2	0.0-1.3	4	0.8	0.2-2.1
2011	9	1.0	0.4-1.9	4	1.0	0.2-2.8	5	1.0	0.3-2.3	4	0.5	0.1-1.3	1	0.2	0.0-1.4	3	0.7	0.1-1.9
2012	7	0.7	0.3-1.4	4	1.1	0.2-2.8	3	0.5	0.1-1.5	4	0.5	0.1-1.2	0	0.0	0.0-0.9	4	0.8	0.2-2.0
2013	15	1.7	0.9-2.8	5	1.6	0.5-3.6	10	1.8	0.9-3.4	6	0.5	0.2-1.2	3	0.6	0.1-1.9	3	0.5	0.1-1.4
2014	13	1.2	0.6-2.1	6	1.1	0.4-2.6	7	1.3	0.5-2.6	1	0.1	0.0-0.6	0	0.0	0.0-0.8	1	0.2	0.0-1.0
2015	13	1.3	0.7-2.2	7	1.4	0.5-3.0	6	1.2	0.4-2.5	3	0.4	0.1-1.0	2	0.5	0.1-1.8	1	0.2	0.0-1.1
2016	11	1.0	0.5-1.8	6	1.1	0.4-2.5	5	0.8	0.3-2.0	1	0.1	0.0-0.6	1	0.3	0.0-1.4	0	0.0	0.0-0.6
2017	12	1.3	0.7-2.3	5	1.2	0.4-2.7	7	1.4	0.5-2.8	3	0.3	0.1-0.9	2	0.4	0.1-1.6	1	0.2	0.0-0.9
2018	12	1.1	0.6-1.9	7	1.7	0.6-3.5	5	0.8	0.2-1.8	3	0.3	0.1-0.8	3	0.6	0.1-1.9	0	0.0	0.0-0.6
2019	8	0.7	0.3-1.4	1	0.2	0.0-1.0	7	1.1	0.4-2.3	2	0.2	0.0-0.7	0	0.0	0.0-0.7	2	0.3	0.0-1.2
2020	13	1.2	0.6-2.0	4	0.8	0.2-2.2	9	1.4	0.6-2.7	3	0.2	0.0-0.7	1	0.1	0.0-0.9	2	0.3	0.0-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	150	78.6	65.7-93.2	67	37.3	28.3-48.1
1977	170	88.6	75.0-103.8	65	34.8	26.3-44.9
1978	185	90.7	77.5-105.4	55	26.1	19.4-34.3
1979	175	84.2	71.5-98.4	65	30.7	23.3-39.6
1980	204	91.0	78.4-104.9	70	32.5	25.0-41.4
1981	205	90.0	77.7-103.7	84	39.6	31.2-49.4
1982	204	85.8	74.1-98.8	90	38.1	30.4-47.1
1983	223	92.5	80.4-105.9	77	31.5	24.7-39.6
1984	222	89.5	77.7-102.5	97	43.2	34.7-53.0
1985	245	96.4	84.3-109.6	91	36.4	29.1-45.0
1986	257	96.9	85.1-109.8	89	34.6	27.6-42.7
1987	252	94.9	83.2-107.6	84	32.6	25.9-40.6
1988	279	99.0	87.5-111.5	106	38.3	31.3-46.5
1989	249	87.4	76.6-99.2	106	39.1	31.8-47.4
1990	303	102.4	91.0-114.8	90	30.8	24.6-38.0
1991	317	103.4	92.1-115.6	113	37.9	31.1-45.6
1992	330	104.8	93.6-116.9	114	36.6	30.1-44.0
1993	339	105.4	94.2-117.4	109	34.6	28.3-41.9
1994	351	104.0	93.3-115.7	116	36.2	29.8-43.5
1995	349	101.7	91.1-113.1	108	33.0	27.0-39.9
1996	397	110.9	100.1-122.5	108	31.5	25.7-38.1
1997	429	118.7	107.6-130.7	115	34.0	27.9-40.9
1998	440	115.6	104.9-127.1	122	33.7	27.9-40.3
1999	445	115.3	104.7-126.7	98	25.9	20.9-31.6
2000	400	100.6	90.8-111.1	127	34.2	28.4-40.7
2001	418	102.5	92.8-113.0	121	30.3	25.1-36.3
2002	464	108.4	98.6-118.8	123	29.8	24.7-35.6
2003	467	108.0	98.2-118.3	119	28.1	23.2-33.7
2004	510	114.9	105.0-125.4	128	30.5	25.4-36.3
2005	515	114.8	105.0-125.3	126	28.7	23.8-34.2
2006	553	118.7	108.9-129.1	121	26.7	22.1-31.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	557	117.1	107.5-127.4	135	29.2	24.4-34.6
2008	585	119.8	110.1-130.0	133	27.3	22.8-32.4
2009	595	120.5	110.9-130.7	151	31.5	26.6-37.0
2010	597	116.6	107.3-126.5	126	24.7	20.5-29.5
2011	687	130.0	120.3-140.2	148	28.7	24.2-33.8
2012	618	115.3	106.2-124.9	121	23.3	19.3-27.9
2013	662	121.6	112.4-131.4	140	26.0	21.8-30.7
2014	722	127.4	118.1-137.2	138	24.8	20.8-29.4
2015	716	124.0	114.9-133.5	124	21.2	17.6-25.3
2016	782	132.3	123.0-142.1	132	23.0	19.2-27.4
2017	836	139.5	130.0-149.5	123	20.7	17.1-24.7
2018	749	122.6	113.8-131.9	148	23.8	20.0-28.0
2019	862	139.5	130.1-149.3	153	24.4	20.6-28.7
2020	743	114.4	106.1-123.1	154	23.9	20.2-28.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Cervix uteri

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	70	35.8	27.3-46.0	29	16.5	10.6-24.3
1977	58	27.2	20.2-35.8	27	14.6	9.3-21.7
1978	57	25.7	19.2-33.7	21	10.3	6.1-16.0
1979	61	25.8	19.5-33.5	24	10.7	6.7-16.1
1980	58	26.1	19.5-34.1	25	12.2	7.7-18.3
1981	37	15.3	10.6-21.3	27	13.0	8.4-19.0
1982	43	16.4	11.7-22.2	16	7.0	3.9-11.4
1983	49	18.7	13.7-25.0	22	8.6	5.4-13.0
1984	52	19.4	14.3-25.7	13	6.9	3.6-11.8
1985	41	16.0	11.3-21.8	23	8.9	5.5-13.6
1986	53	18.8	13.9-24.8	20	8.0	4.8-12.3
1987	49	17.5	12.8-23.3	16	6.0	3.3-9.9
1988	46	15.8	11.5-21.2	18	6.6	3.8-10.5
1989	51	17.4	12.9-23.0	19	6.9	4.1-10.8
1990	63	19.0	14.5-24.4	18	6.2	3.6-9.9
1991	53	16.2	12.0-21.4	19	6.2	3.7-9.7
1992	41	11.2	8.0-15.3	14	4.2	2.3-7.1
1993	52	14.3	10.6-18.9	25	7.4	4.7-11.0
1994	48	12.9	9.4-17.2	15	4.2	2.4-7.0
1995	53	14.2	10.5-18.7	16	4.7	2.6-7.6
1996	73	19.3	15.0-24.3	19	5.2	3.1-8.2
1997	34	8.9	6.1-12.5	24	6.4	4.1-9.5
1998	51	12.0	8.9-15.9	17	4.6	2.7-7.4
1999	51	13.1	9.7-17.3	13	3.3	1.7-5.7
2000	42	10.2	7.3-13.8	19	4.6	2.7-7.1
2001	52	12.7	9.4-16.7	10	2.5	1.2-4.6
2002	46	10.6	7.7-14.1	16	4.0	2.3-6.5
2003	57	13.0	9.8-16.9	19	4.4	2.6-6.9
2004	50	10.8	8.0-14.2	14	3.0	1.6-5.0
2005	50	10.9	8.0-14.4	17	3.6	2.1-5.8
2006	49	10.2	7.5-13.5	15	3.3	1.9-5.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Cervix uteri

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	48	10.1	7.4-13.4	21	4.6	2.8-7.0
2008	52	10.3	7.7-13.5	15	3.1	1.7-5.1
2009	43	8.7	6.3-11.7	26	5.0	3.3-7.4
2010	45	8.7	6.3-11.7	13	2.5	1.3-4.3
2011	45	8.9	6.5-11.9	14	2.6	1.4-4.4
2012	44	8.1	5.8-10.9	19	3.6	2.1-5.6
2013	62	11.3	8.6-14.5	23	4.2	2.6-6.3
2014	51	8.7	6.4-11.4	15	2.5	1.4-4.1
2015	53	9.6	7.1-12.5	17	3.1	1.8-5.0
2016	47	7.8	5.7-10.4	6	1.2	0.4-2.5
2017	51	8.7	6.4-11.5	18	3.0	1.7-4.7
2018	43	7.0	5.1-9.5	16	2.6	1.5-4.3
2019	40	6.3	4.5-8.6	19	2.9	1.7-4.6
2020	41	6.7	4.7-9.1	12	2.0	1.0-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Corpus uteris and uterus, not otherwise specified**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	31	17.9	11.8-25.7	13	9.6	4.9-16.5
1977	36	19.4	13.2-27.3	15	8.2	4.4-13.9
1978	25	13.8	8.6-20.7	9	4.6	2.0-8.9
1979	38	16.7	11.7-23.2	14	7.6	4.0-12.9
1980	50	23.8	17.4-31.7	21	9.8	6.0-15.0
1981	29	12.6	8.3-18.2	21	9.9	6.0-15.3
1982	31	13.4	9.0-19.2	11	4.2	2.1-7.5
1983	32	13.3	9.0-18.8	13	6.0	3.1-10.3
1984	33	14.2	9.6-20.1	16	6.7	3.7-10.9
1985	49	19.8	14.6-26.3	19	9.5	5.6-14.8
1986	38	15.4	10.8-21.2	12	4.9	2.5-8.5
1987	50	19.3	14.2-25.5	11	4.7	2.3-8.4
1988	46	17.9	12.9-23.9	18	7.2	4.2-11.4
1989	35	12.6	8.7-17.6	24	9.4	5.9-14.0
1990	51	17.8	13.1-23.4	10	4.0	1.9-7.4
1991	42	14.5	10.4-19.6	17	5.8	3.3-9.2
1992	63	21.6	16.5-27.7	23	8.7	5.5-13.1
1993	60	19.4	14.8-25.0	16	6.0	3.4-9.6
1994	67	20.7	16.0-26.4	13	4.2	2.2-7.2
1995	52	15.8	11.7-20.7	18	5.9	3.4-9.2
1996	79	23.2	18.3-29.0	24	7.4	4.7-10.9
1997	70	20.8	16.1-26.2	12	3.7	1.9-6.5
1998	79	21.5	17.0-26.8	28	8.5	5.6-12.3
1999	84	22.7	18.1-28.2	24	7.0	4.4-10.4
2000	81	21.8	17.2-27.1	30	8.5	5.7-12.1
2001	80	20.8	16.4-25.9	25	7.0	4.5-10.3
2002	75	18.2	14.2-22.8	29	7.8	5.2-11.2
2003	84	20.1	16.0-25.0	36	9.1	6.3-12.6
2004	110	25.6	21.0-30.9	20	4.4	2.7-6.8
2005	102	23.8	19.4-29.0	33	8.3	5.7-11.6
2006	101	21.9	17.8-26.6	32	7.3	5.0-10.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
Black Individuals
Corpus uteris and uterus, not otherwise specified**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	126	26.7	22.2-31.9	41	9.7	6.9-13.1
2008	114	24.1	19.8-29.0	46	10.2	7.5-13.7
2009	137	28.6	23.9-33.9	30	6.7	4.5-9.5
2010	131	26.1	21.8-31.1	46	9.6	7.0-12.9
2011	136	25.2	21.1-29.9	36	7.3	5.1-10.2
2012	159	29.7	25.2-34.8	47	9.6	7.0-12.7
2013	136	23.8	19.9-28.2	45	8.5	6.2-11.4
2014	173	30.2	25.8-35.1	57	10.5	7.9-13.6
2015	168	28.7	24.4-33.4	40	7.2	5.1-9.8
2016	194	32.4	27.9-37.4	47	8.1	5.9-10.8
2017	165	26.5	22.5-30.9	56	9.1	6.8-11.8
2018	197	30.8	26.6-35.5	58	9.4	7.1-12.2
2019	215	32.7	28.4-37.5	57	8.9	6.7-11.6
2020	219	32.3	28.1-37.0	50	7.5	5.5-9.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Ovary

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	29	16.9	10.9-24.8	17	11.7	6.6-19.0
1977	29	13.2	8.5-19.4	17	8.4	4.7-13.7
1978	24	10.7	6.8-16.1	13	6.9	3.5-12.0
1979	34	16.4	11.0-23.2	22	10.1	6.1-15.6
1980	26	12.0	7.7-17.7	20	9.5	5.7-14.7
1981	28	12.3	7.9-18.0	14	6.2	3.3-10.5
1982	36	14.3	9.9-19.9	16	6.7	3.8-11.0
1983	29	12.2	8.0-17.7	14	6.1	3.2-10.3
1984	28	10.8	7.0-15.7	24	10.5	6.6-15.7
1985	35	14.3	9.9-20.0	13	5.4	2.8-9.2
1986	37	14.3	9.9-19.9	21	8.5	5.2-13.0
1987	28	11.1	7.3-16.2	27	10.9	7.1-15.9
1988	39	13.8	9.7-19.0	20	7.5	4.6-11.7
1989	38	13.2	9.2-18.2	23	8.3	5.2-12.4
1990	37	12.2	8.5-17.0	28	10.2	6.7-14.8
1991	23	8.8	5.5-13.2	20	7.2	4.3-11.1
1992	31	9.7	6.5-13.9	34	11.9	8.1-16.6
1993	41	12.0	8.5-16.3	19	6.5	3.9-10.1
1994	32	9.7	6.6-13.7	23	7.4	4.6-11.1
1995	42	11.8	8.4-16.0	23	7.9	5.0-11.8
1996	40	11.8	8.4-16.1	19	5.7	3.4-8.9
1997	51	14.2	10.5-18.7	26	7.6	4.9-11.1
1998	37	9.9	6.9-13.7	23	7.0	4.4-10.5
1999	47	12.6	9.2-16.8	26	7.2	4.7-10.6
2000	42	11.2	8.0-15.1	31	8.5	5.7-12.1
2001	41	10.1	7.2-13.7	24	6.6	4.2-9.7
2002	45	10.6	7.7-14.3	29	7.5	5.0-10.8
2003	28	6.4	4.2-9.3	35	8.5	5.9-11.9
2004	47	11.2	8.2-15.0	28	7.1	4.7-10.2
2005	40	8.7	6.2-11.9	26	6.0	3.9-8.8
2006	40	9.1	6.4-12.4	31	7.3	4.9-10.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	35	7.4	5.2-10.4	26	6.1	4.0-8.9
2008	49	10.6	7.8-14.1	21	4.9	3.0-7.5
2009	52	10.3	7.7-13.6	25	5.3	3.4-7.8
2010	42	8.1	5.8-11.0	26	5.3	3.4-7.8
2011	50	9.4	6.9-12.4	30	6.3	4.2-9.0
2012	64	11.7	9.0-15.0	34	6.4	4.4-8.9
2013	40	7.0	4.9-9.6	34	6.3	4.4-8.9
2014	62	10.8	8.2-13.9	37	6.5	4.5-9.0
2015	47	8.5	6.2-11.3	30	5.4	3.6-7.8
2016	54	8.5	6.4-11.2	28	4.9	3.2-7.1
2017	55	9.4	7.0-12.3	30	5.1	3.4-7.3
2018	42	6.8	4.8-9.2	29	4.7	3.2-6.8
2019	53	8.5	6.3-11.1	31	5.1	3.4-7.2
2020	41	6.3	4.5-8.5	23	3.7	2.4-5.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	123	126.0	101.5-153.8	65	71.9	53.1-94.3
1977	129	115.4	93.8-139.8	64	79.0	58.8-102.8
1978	143	128.1	104.9-154.2	59	54.2	39.8-71.5
1979	144	126.1	103.5-151.5	73	67.8	51.5-87.0
1980	176	142.7	119.9-168.0	56	57.7	42.1-76.3
1981	170	143.5	120.2-169.4	82	76.0	58.7-96.1
1982	160	123.3	103.3-145.6	74	64.2	49.1-82.0
1983	185	143.2	121.2-167.5	73	65.3	49.9-83.2
1984	171	123.2	103.6-144.9	67	51.9	39.0-67.2
1985	182	128.6	108.7-150.7	72	58.0	44.3-74.0
1986	206	146.6	125.2-170.0	92	74.5	58.5-92.8
1987	187	129.4	109.9-151.0	83	75.3	58.6-94.5
1988	212	139.2	119.2-161.2	100	83.0	66.2-102.1
1989	207	131.4	112.4-152.4	82	59.1	46.1-74.2
1990	258	162.1	141.0-185.2	96	67.9	53.7-84.2
1991	359	210.9	187.4-236.3	104	82.1	65.9-100.5
1992	434	238.6	214.3-264.7	87	60.9	47.7-76.1
1993	517	272.0	246.7-298.8	102	72.5	57.9-89.2
1994	516	255.6	232.0-280.7	121	83.3	67.9-100.6
1995	563	268.0	244.2-293.2	113	70.2	56.7-85.5
1996	629	269.3	247.1-293.0	116	70.6	57.4-85.5
1997	620	267.4	244.9-291.2	108	63.4	50.9-77.5
1998	672	273.9	252.1-296.9	114	70.9	57.7-85.8
1999	706	274.9	253.5-297.3	94	54.6	43.3-67.5
2000	774	305.1	282.3-329.0	129	74.3	61.2-89.0
2001	791	293.7	272.0-316.7	110	63.3	51.3-76.9
2002	828	300.1	278.2-323.1	114	61.5	49.9-74.7
2003	826	283.5	263.1-305.0	112	56.5	45.8-68.6
2004	767	245.7	227.5-265.0	97	47.0	37.4-58.0
2005	809	252.9	234.5-272.2	107	45.2	36.5-55.1
2006	892	270.4	251.9-290.0	83	35.1	27.6-43.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	971	286.8	267.9-306.5	99	43.3	34.7-53.0
2008	936	260.8	243.4-279.2	103	40.2	32.4-49.1
2009	958	252.4	235.7-270.0	111	44.8	36.4-54.3
2010	992	255.4	238.7-272.9	113	44.3	36.1-53.5
2011	1,032	252.7	236.7-269.6	88	30.8	24.4-38.1
2012	909	211.2	196.9-226.1	97	35.3	28.4-43.3
2013	966	219.6	205.2-234.7	97	34.2	27.5-41.9
2014	934	203.7	190.1-217.9	102	32.5	26.2-39.6
2015	963	202.2	188.9-216.1	77	24.4	19.1-30.6
2016	979	201.0	188.0-214.7	100	28.7	23.1-35.1
2017	1,075	216.3	202.9-230.3	126	36.0	29.8-43.0
2018	1,093	211.8	198.9-225.4	94	25.1	20.1-30.8
2019	1,136	214.5	201.6-228.0	95	24.0	19.2-29.5
2020	1,001	183.6	171.9-195.9	119	27.4	22.5-33.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Testis

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3	0.9	0.2-4.1	1	0.9	0.0-4.7
1977	5	1.5	0.5-4.7	1	0.4	0.0-3.5
1978	2	0.5	0.1-3.3	1	0.3	0.0-3.2
1980	7	2.1	0.8-5.2	2	0.6	0.1-3.4
1981	4	1.6	0.4-4.8	0	0.0	0.0-2.5
1982	3	1.2	0.2-4.1	0	0.0	0.0-2.4
1983	2	0.4	0.1-2.7	1	0.2	0.0-2.5
1984	5	1.5	0.4-4.2	0	0.0	0.0-2.2
1985	6	1.9	0.6-4.7	1	0.4	0.0-2.7
1986	6	1.8	0.6-4.5	0	0.0	0.0-2.1
1987	5	1.2	0.4-3.5	1	0.2	0.0-2.2
1988	5	1.2	0.4-3.5	0	0.0	0.0-2.0
1989	2	0.5	0.1-2.7	0	0.0	0.0-1.9
1990	3	0.7	0.1-2.8	0	0.0	0.0-1.9
1991	3	0.6	0.1-2.7	1	0.4	0.0-2.4
1992	8	2.6	0.6-6.5	1	0.3	0.0-2.1
1993	3	0.6	0.1-2.4	1	0.2	0.0-1.9
1994	4	0.8	0.2-2.6	0	0.0	0.0-1.7
1995	2	0.5	0.1-2.3	0	0.0	0.0-1.6
1996	4	0.9	0.2-2.7	0	0.0	0.0-1.5
1997	6	1.2	0.4-3.0	0	0.0	0.0-1.5
1998	6	1.2	0.4-3.1	0	0.0	0.0-1.4
1999	8	1.5	0.6-3.2	1	0.2	0.0-1.7
2000	6	1.7	0.6-4.0	0	0.0	0.0-1.3
2001	8	1.7	0.7-3.7	0	0.0	0.0-1.3
2002	6	1.2	0.4-2.9	0	0.0	0.0-1.3
2003	9	1.8	0.8-3.7	0	0.0	0.0-1.2
2004	5	1.0	0.3-2.5	0	0.0	0.0-1.2
2005	3	0.5	0.1-1.8	0	0.0	0.0-1.1
2006	6	1.2	0.4-2.8	0	0.0	0.0-1.1
2007	9	1.7	0.8-3.4	0	0.0	0.0-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Testis**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	6	1.2	0.4-2.7	1	0.1	0.0-1.2
2009	4	0.7	0.2-2.1	0	0.0	0.0-1.0
2010	5	1.0	0.3-2.5	1	0.2	0.0-1.3
2011	7	1.3	0.5-2.9	0	0.0	0.0-0.9
2012	6	1.2	0.4-2.7	0	0.0	0.0-0.9
2013	10	1.8	0.9-3.4	1	0.2	0.0-1.2
2014	8	1.4	0.6-3.0	0	0.0	0.0-0.8
2015	8	1.6	0.7-3.1	0	0.0	0.0-0.8
2016	9	1.4	0.6-2.8	0	0.0	0.0-0.8
2017	10	1.8	0.9-3.4	0	0.0	0.0-0.8
2018	10	1.7	0.8-3.3	0	0.0	0.0-0.8
2019	10	1.4	0.7-2.7	0	0.0	0.0-0.7
2020	12	1.8	0.9-3.3	0	0.0	0.0-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Urinary bladder (incl. in situ)

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	44	18.4	12.8-25.3	31	34.4	21.4-51.3	13	8.8	4.4-15.2	15	5.5	2.8-9.4	7	5.8	1.6-13.6	8	5.5	2.2-10.8
1977	29	10.9	6.9-16.1	13	10.4	5.1-18.4	16	10.6	5.7-17.6	14	6.7	3.4-11.3	8	11.1	4.3-22.0	6	4.2	1.4-9.2
1978	42	14.8	10.3-20.3	30	28.2	17.8-41.6	12	6.4	3.2-11.2	18	5.3	3.0-8.5	8	4.6	1.9-9.3	10	5.6	2.6-10.3
1979	55	18.2	13.3-24.1	30	23.0	14.9-33.6	25	14.6	9.1-21.8	23	9.9	6.0-15.0	14	17.3	8.7-29.5	9	5.9	2.5-11.1
1980	40	12.7	8.8-17.7	29	21.1	13.3-31.5	11	6.8	3.3-12.1	20	6.4	3.8-10.1	12	10.2	4.9-18.2	8	4.1	1.6-8.1
1981	42	12.7	8.9-17.4	23	16.2	9.6-25.1	19	10.2	5.9-15.9	16	5.0	2.7-8.4	13	11.7	5.4-21.0	3	1.3	0.3-3.7
1982	33	10.2	6.8-14.5	21	15.8	9.1-24.8	12	6.4	3.1-11.2	19	6.1	3.5-9.6	12	9.6	4.5-17.3	7	3.8	1.5-7.8
1983	36	9.9	6.8-13.9	18	11.1	6.1-18.2	18	8.9	5.1-14.0	18	5.4	3.1-8.5	7	5.3	2.0-10.8	11	5.4	2.6-9.7
1984	55	15.6	11.4-20.5	38	28.2	18.9-39.7	17	7.9	4.5-12.7	20	6.9	4.1-10.6	14	13.3	6.7-22.6	6	3.4	1.2-7.1
1985	45	12.0	8.5-16.2	27	18.2	11.4-27.0	18	8.1	4.7-12.9	16	4.8	2.6-7.8	10	8.4	3.8-15.6	6	2.8	1.0-6.0
1986	41	10.8	7.6-14.7	26	16.7	10.5-24.9	15	6.9	3.8-11.2	16	4.7	2.6-7.6	8	5.7	2.2-11.4	8	4.0	1.7-7.6
1987	34	9.1	6.1-12.8	20	12.6	7.3-19.9	14	6.6	3.6-11.0	13	3.9	2.0-6.6	6	4.9	1.7-10.4	7	3.3	1.3-6.7
1988	53	13.9	10.2-18.3	33	19.4	13.0-27.5	20	9.5	5.7-14.5	16	5.0	2.8-8.0	9	7.5	3.2-14.2	7	3.5	1.4-7.1
1989	54	12.1	9.0-15.9	34	19.3	12.7-27.6	20	7.6	4.6-11.8	22	6.7	4.1-10.0	15	13.0	6.9-21.5	7	3.4	1.3-6.7
1990	41	10.3	7.3-14.1	30	18.9	12.1-27.7	11	4.9	2.4-8.6	24	6.5	4.1-9.7	16	9.7	5.3-15.9	8	4.0	1.7-7.6
1991	64	14.1	10.7-18.0	47	27.8	19.6-37.9	17	6.0	3.5-9.6	24	6.1	3.8-9.0	12	7.5	3.4-13.7	12	5.1	2.6-8.8
1992	39	8.3	5.9-11.5	30	16.2	10.5-23.5	9	3.3	1.5-6.2	12	2.6	1.3-4.4	9	4.8	2.0-9.2	3	1.1	0.2-3.1
1993	49	10.0	7.3-13.4	24	10.2	6.2-15.6	25	9.2	5.9-13.5	21	5.0	3.0-7.6	10	7.0	3.2-12.6	11	4.0	1.9-7.0
1994	45	9.7	7.0-13.0	32	19.0	12.4-27.3	13	4.5	2.4-7.6	15	3.9	2.1-6.3	9	6.0	2.6-11.4	6	2.5	0.9-5.3
1995	48	9.5	6.9-12.7	34	17.2	11.5-24.3	14	4.6	2.5-7.7	14	3.5	1.9-5.8	8	6.8	2.7-13.2	6	2.3	0.8-4.8
1996	60	11.5	8.6-14.8	30	14.8	9.4-21.7	30	9.7	6.5-13.8	23	4.8	3.0-7.2	13	7.7	3.9-13.3	10	3.3	1.6-6.0
1997	65	12.7	9.7-16.3	48	26.6	18.8-36.2	17	5.6	3.2-8.8	16	3.0	1.7-4.9	9	4.5	1.8-8.8	7	2.3	0.9-4.6
1998	87	16.9	13.4-20.9	58	29.7	21.7-39.2	29	9.6	6.4-13.7	20	4.1	2.5-6.3	10	5.7	2.6-10.4	10	3.3	1.6-5.9
1999	64	11.3	8.6-14.5	40	18.8	12.9-26.1	24	7.1	4.5-10.5	20	3.8	2.3-5.9	12	5.9	2.7-10.5	8	2.6	1.1-4.9
2000	79	12.9	10.1-16.2	57	22.2	16.3-29.3	22	6.5	4.1-9.8	15	3.1	1.7-5.0	7	4.6	1.7-9.3	8	2.4	1.1-4.7
2001	77	12.8	10.0-16.1	50	21.5	15.3-29.0	27	7.7	5.1-11.2	19	3.5	2.1-5.4	10	5.3	2.2-10.0	9	2.8	1.3-5.1
2002	66	10.8	8.3-13.8	43	17.0	12.0-23.2	23	6.4	4.0-9.5	19	3.5	2.0-5.4	10	5.2	2.3-9.7	9	2.6	1.2-4.8
2003	84	12.4	9.8-15.4	56	20.5	14.9-27.2	28	7.2	4.8-10.4	24	4.1	2.6-6.1	13	6.3	3.2-10.8	11	3.0	1.5-5.3
2004	88	13.5	10.7-16.7	60	21.5	16.1-28.0	28	7.6	5.0-10.9	21	3.5	2.1-5.3	11	4.2	1.9-7.7	10	2.8	1.4-5.1
2005	85	12.4	9.9-15.4	62	23.4	17.4-30.5	23	5.7	3.6-8.6	23	4.2	2.6-6.2	13	7.1	3.6-11.9	10	2.8	1.3-5.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Urinary bladder (incl. in situ)

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	81	11.3	8.9-14.1	54	19.3	14.2-25.4	27	6.1	4.0-8.9	12	1.8	0.9-3.1	8	2.8	1.1-5.7	4	1.1	0.3-2.7
2007	85	11.6	9.2-14.4	58	19.1	14.2-24.8	27	6.4	4.2-9.3	31	4.7	3.1-6.6	15	5.7	3.1-9.4	16	3.9	2.2-6.3
2008	103	13.5	10.9-16.4	70	22.8	17.4-29.3	33	7.5	5.1-10.6	18	2.7	1.6-4.3	12	4.9	2.5-8.4	6	1.4	0.5-3.0
2009	100	13.1	10.6-16.0	67	21.4	16.2-27.5	33	7.5	5.1-10.5	27	3.8	2.5-5.5	20	7.6	4.4-11.8	7	1.7	0.7-3.4
2010	93	11.3	9.0-13.9	69	21.9	16.6-28.2	24	4.8	3.1-7.2	23	3.1	1.9-4.6	9	3.6	1.6-6.7	14	2.9	1.6-4.9
2011	132	15.3	12.7-18.2	102	29.5	23.6-36.2	30	5.9	3.9-8.4	27	3.6	2.3-5.2	19	6.6	3.8-10.4	8	1.9	0.8-3.6
2012	120	14.6	12.1-17.5	80	24.6	19.2-31.0	40	8.4	6.0-11.4	23	2.7	1.7-4.1	15	4.5	2.4-7.6	8	1.6	0.7-3.2
2013	123	14.2	11.7-17.0	83	23.1	18.2-29.0	40	7.8	5.6-10.7	30	3.6	2.4-5.1	19	5.8	3.4-9.2	11	2.2	1.1-3.9
2014	114	12.4	10.1-14.9	76	19.7	15.3-24.9	38	7.0	4.9-9.7	24	2.9	1.8-4.3	17	5.5	3.1-8.8	7	1.3	0.5-2.7
2015	128	13.7	11.3-16.3	87	22.7	17.9-28.3	41	7.7	5.5-10.4	26	2.7	1.7-4.0	14	3.6	1.9-6.1	12	2.1	1.1-3.6
2016	119	11.9	9.8-14.2	76	17.9	13.9-22.6	43	7.4	5.3-10.0	22	2.5	1.6-3.8	10	3.3	1.5-5.9	12	2.2	1.1-3.8
2017	142	14.0	11.7-16.5	100	24.0	19.3-29.5	42	7.1	5.1-9.6	24	2.6	1.6-3.9	16	4.4	2.4-7.1	8	1.4	0.6-2.7
2018	144	13.6	11.4-16.1	94	20.7	16.5-25.6	50	8.7	6.4-11.5	28	2.8	1.9-4.1	19	5.1	3.0-8.0	9	1.4	0.6-2.8
2019	142	12.9	10.8-15.2	102	22.3	18.0-27.2	40	6.3	4.5-8.6	24	2.5	1.6-3.7	14	3.9	2.1-6.6	10	1.8	0.8-3.2
2020	114	10.3	8.5-12.5	76	16.9	13.1-21.3	38	5.9	4.1-8.1	28	2.6	1.7-3.7	18	4.0	2.3-6.3	10	1.5	0.7-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Kidney and renal pelvis

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	31	9.8	6.4-14.2	20	12.7	7.3-20.5	11	7.3	3.5-13.0	10	2.8	1.2-5.4	6	3.4	1.2-7.9	4	2.2	0.4-5.9
1977	27	7.9	4.9-11.9	14	11.7	5.4-21.0	13	5.8	3.0-10.2	7	2.0	0.7-4.3	4	3.3	0.7-9.0	3	1.1	0.2-3.7
1978	30	9.4	6.0-13.9	17	10.7	5.6-18.1	13	8.0	4.0-14.0	15	4.5	2.3-7.7	8	3.5	1.4-7.6	7	4.7	1.8-9.6
1979	30	7.7	5.0-11.1	21	11.1	6.8-17.3	9	4.6	2.0-8.9	15	4.5	2.4-7.7	13	10.0	4.8-17.8	2	0.8	0.1-3.1
1980	21	5.5	3.2-8.6	11	7.0	3.0-13.6	10	4.5	2.0-8.5	10	2.7	1.2-5.1	7	4.8	1.4-11.0	3	1.3	0.3-3.9
1981	30	7.1	4.6-10.3	22	11.6	6.6-18.8	8	3.7	1.6-7.2	12	2.5	1.3-4.4	8	3.8	1.6-7.8	4	1.5	0.4-3.9
1982	31	7.9	5.1-11.4	18	10.0	5.4-16.6	13	6.1	3.1-10.6	19	5.6	3.2-8.9	9	4.5	1.9-9.0	10	5.8	2.7-10.5
1983	27	6.3	4.0-9.4	20	10.3	5.8-16.7	7	3.2	1.3-6.6	10	2.4	1.1-4.6	7	4.3	1.3-9.7	3	1.3	0.3-3.7
1984	44	10.0	7.1-13.6	28	14.8	8.9-22.7	16	6.7	3.8-10.9	12	3.0	1.5-5.3	7	4.7	1.7-9.9	5	2.0	0.6-4.6
1985	46	10.4	7.4-14.0	25	13.9	8.4-21.3	21	8.1	4.9-12.5	16	3.8	2.1-6.3	11	4.8	2.4-8.8	5	2.6	0.8-5.8
1986	39	8.6	6.1-11.9	20	10.3	6.1-16.2	19	7.3	4.3-11.5	20	4.5	2.7-7.1	11	6.1	2.6-11.6	9	3.5	1.6-6.7
1987	36	8.5	5.8-11.9	23	14.4	8.7-22.1	13	4.9	2.5-8.4	17	4.3	2.4-7.0	5	3.1	0.9-7.4	12	5.1	2.6-8.9
1988	40	8.4	5.8-11.5	18	9.9	5.4-16.2	22	7.7	4.7-11.8	17	4.1	2.3-6.7	9	6.4	2.7-12.3	8	2.9	1.2-5.8
1989	33	6.7	4.5-9.6	19	9.5	5.2-15.6	14	5.0	2.7-8.4	12	3.0	1.5-5.2	6	4.0	1.3-8.7	6	2.4	0.8-5.1
1990	62	11.5	8.7-14.8	37	15.8	10.8-22.2	25	8.3	5.3-12.2	16	3.2	1.7-5.3	9	4.1	1.6-8.2	7	2.5	0.9-5.1
1991	59	10.4	7.8-13.6	38	15.8	10.8-22.2	21	6.6	4.0-10.1	20	4.5	2.7-7.0	12	7.2	3.3-13.0	8	3.0	1.2-5.8
1992	60	10.6	8.0-13.8	40	16.9	11.5-23.9	20	6.3	3.8-9.8	19	3.4	1.9-5.4	12	5.4	2.6-9.9	7	2.0	0.7-4.3
1993	65	11.0	8.4-14.1	42	16.7	11.6-23.1	23	6.8	4.3-10.3	22	4.2	2.6-6.4	10	4.9	2.2-9.2	12	3.9	2.0-6.8
1994	75	12.2	9.4-15.4	37	12.8	8.8-18.1	38	11.3	7.9-15.6	20	3.6	2.1-5.6	13	5.9	2.9-10.6	7	2.1	0.8-4.3
1995	76	13.5	10.5-17.0	38	17.4	11.6-24.8	38	11.6	8.1-16.0	23	4.5	2.8-6.8	14	7.6	3.6-13.4	9	3.0	1.3-5.6
1996	71	11.3	8.7-14.4	48	18.4	13.0-25.2	23	6.4	4.0-9.7	22	3.8	2.3-5.8	10	5.1	2.1-9.8	12	3.4	1.7-5.9
1997	73	11.9	9.2-15.0	45	17.7	12.5-24.2	28	7.9	5.2-11.4	27	4.9	3.1-7.1	14	5.6	2.8-9.9	13	4.2	2.2-7.1
1998	103	15.6	12.6-19.0	63	22.8	17.1-29.6	40	10.5	7.5-14.4	14	2.3	1.3-3.9	7	2.7	1.1-5.7	7	2.0	0.8-4.1
1999	70	10.7	8.3-13.6	38	13.1	9.0-18.2	32	8.7	5.9-12.3	20	3.5	2.1-5.4	10	3.7	1.6-7.2	10	3.2	1.5-5.7
2000	73	10.3	8.0-13.0	38	12.6	8.6-17.7	35	8.7	6.0-12.2	22	3.2	2.0-4.9	12	4.2	2.1-7.6	10	2.6	1.2-4.7
2001	82	11.5	9.0-14.3	50	15.2	11.0-20.4	32	8.3	5.6-11.7	23	3.6	2.2-5.4	13	4.0	2.0-7.0	10	2.9	1.4-5.3
2002	104	13.9	11.3-17.0	60	19.2	14.3-25.2	44	10.4	7.5-14.0	10	1.3	0.6-2.4	7	2.2	0.9-4.6	3	0.6	0.1-1.8
2003	93	13.0	10.4-16.0	53	15.8	11.5-21.0	40	10.3	7.3-14.1	26	3.9	2.5-5.8	20	7.9	4.4-12.7	6	1.6	0.6-3.4
2004	115	14.0	11.5-16.9	68	17.7	13.5-22.6	47	10.7	7.8-14.3	21	3.0	1.8-4.6	11	3.8	1.7-7.0	10	2.5	1.2-4.6
2005	108	13.3	10.8-16.1	59	16.6	12.4-21.7	49	10.8	8.0-14.3	28	3.6	2.3-5.3	21	6.2	3.6-9.7	7	1.7	0.6-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Kidney and renal pelvis

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	131	15.8	13.1-18.8	89	24.5	19.2-30.8	42	9.4	6.7-12.7	21	2.6	1.6-4.0	12	3.7	1.7-6.7	9	1.9	0.9-3.6
2007	147	17.5	14.7-20.7	83	22.5	17.6-28.3	64	13.8	10.6-17.6	32	4.3	2.9-6.0	23	7.3	4.4-11.1	9	2.1	0.9-3.9
2008	148	17.1	14.4-20.2	85	20.8	16.4-26.1	63	13.6	10.4-17.5	28	3.2	2.1-4.7	19	4.7	2.7-7.4	9	2.0	0.9-3.7
2009	177	20.0	17.1-23.3	104	25.1	20.2-30.7	73	15.4	12.1-19.5	33	3.9	2.6-5.5	20	5.0	2.9-7.8	13	2.8	1.5-4.7
2010	172	18.9	16.1-22.1	108	25.9	21.0-31.6	64	13.1	10.1-16.8	25	3.2	2.0-4.7	11	3.0	1.4-5.5	14	3.2	1.7-5.3
2011	166	18.3	15.5-21.4	112	28.6	23.2-34.8	54	10.6	7.9-13.8	24	2.8	1.8-4.2	14	3.9	2.0-6.6	10	2.1	1.0-3.8
2012	163	16.7	14.2-19.6	103	24.6	19.8-30.2	60	11.0	8.4-14.2	29	3.4	2.2-4.9	19	6.2	3.6-9.8	10	1.8	0.9-3.4
2013	175	17.3	14.8-20.2	117	25.3	20.8-30.6	58	10.7	8.1-13.9	17	1.9	1.1-3.0	11	2.6	1.2-4.7	6	1.1	0.4-2.5
2014	195	18.6	16.0-21.5	126	25.9	21.4-31.1	69	12.6	9.8-16.0	28	3.1	2.0-4.4	23	6.2	3.8-9.4	5	1.0	0.3-2.2
2015	229	21.7	18.9-24.7	148	31.4	26.3-37.2	81	14.0	11.1-17.5	40	4.1	2.9-5.6	25	5.8	3.6-8.7	15	2.9	1.6-4.8
2016	205	19.2	16.6-22.1	137	28.7	23.9-34.2	68	11.8	9.1-15.0	25	2.4	1.5-3.5	20	4.4	2.6-6.8	5	0.8	0.3-1.9
2017	197	17.5	15.1-20.2	113	22.6	18.4-27.4	84	13.8	11.0-17.1	23	2.3	1.4-3.4	19	4.3	2.5-6.8	4	0.8	0.2-2.0
2018	249	21.8	19.1-24.8	157	30.3	25.5-35.7	92	15.1	12.1-18.5	25	2.3	1.5-3.4	14	2.6	1.4-4.5	11	1.9	1.0-3.4
2019	209	18.3	15.9-21.0	126	23.9	19.8-28.7	83	13.5	10.7-16.8	28	2.7	1.8-3.9	15	3.5	1.9-5.8	13	2.1	1.1-3.6
2020	190	16.0	13.7-18.5	130	24.5	20.3-29.3	60	9.2	7.0-12.0	24	2.2	1.4-3.3	12	2.6	1.3-4.6	12	2.0	1.0-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Brain and other central nervous system

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	20	4.4	2.6-7.2	10	4.9	2.3-9.7	10	4.0	1.7-7.9	16	3.3	1.8-5.8	8	2.9	1.2-6.8	8	3.5	1.4-7.3
1977	26	5.3	3.4-8.0	10	5.2	2.4-10.2	16	5.5	3.0-9.2	10	2.1	1.0-4.0	4	1.9	0.5-5.6	6	2.3	0.8-5.3
1978	19	4.1	2.3-6.6	6	3.4	1.2-7.8	13	4.7	2.3-8.4	10	2.3	1.0-4.4	5	2.3	0.7-6.0	5	2.2	0.6-5.3
1979	24	4.8	3.0-7.3	16	7.5	4.2-12.7	8	2.6	1.1-5.5	12	2.4	1.2-4.4	6	2.5	0.9-6.2	6	2.3	0.8-5.2
1980	19	3.8	2.2-6.2	16	6.8	3.7-11.9	3	1.3	0.3-3.9	9	1.8	0.8-3.5	8	3.4	1.4-7.1	1	0.4	0.0-2.4
1981	14	2.3	1.2-4.0	7	2.6	1.0-6.0	7	2.1	0.8-4.5	6	1.2	0.4-2.7	5	2.4	0.8-5.9	1	0.2	0.0-1.9
1982	14	2.5	1.3-4.3	10	3.7	1.7-7.4	4	1.4	0.4-3.6	9	2.1	0.9-4.1	5	3.3	0.9-8.1	4	1.3	0.4-3.6
1983	15	2.4	1.3-4.1	7	2.2	0.8-5.3	8	2.4	1.0-5.0	11	2.3	1.0-4.2	5	2.4	0.8-6.0	6	2.1	0.7-4.8
1984	22	3.4	2.1-5.2	14	4.5	2.4-8.2	8	2.3	1.0-4.8	18	2.8	1.6-4.6	11	3.8	1.8-7.4	7	1.9	0.8-4.2
1985	21	3.8	2.2-5.9	9	4.4	1.7-9.0	12	3.6	1.8-6.4	11	1.8	0.9-3.3	5	1.8	0.5-4.7	6	1.8	0.6-4.1
1986	19	4.1	2.3-6.6	8	4.2	1.4-9.2	11	4.0	1.9-7.4	9	2.1	0.9-4.0	3	1.3	0.3-4.0	6	2.5	0.9-5.4
1987	18	3.5	1.9-5.7	7	2.3	0.9-5.2	11	4.0	1.9-7.3	9	2.2	0.9-4.2	3	1.2	0.2-3.8	6	2.7	0.9-5.7
1988	24	3.9	2.4-5.9	14	4.5	2.4-8.0	10	3.0	1.4-5.8	11	1.7	0.8-3.1	7	2.6	1.0-5.9	4	0.9	0.2-2.6
1989	18	2.6	1.5-4.2	5	1.2	0.4-3.4	13	3.7	1.9-6.5	8	1.2	0.5-2.5	4	1.0	0.3-3.3	4	1.3	0.3-3.3
1990	21	3.3	1.9-5.2	10	4.7	1.8-9.4	11	2.7	1.3-5.1	9	1.5	0.6-2.9	3	1.2	0.2-3.9	6	1.7	0.6-3.8
1991	37	5.4	3.7-7.6	19	5.7	3.2-9.5	18	5.1	2.9-8.2	11	2.0	0.9-3.7	3	1.4	0.2-4.4	8	2.4	1.0-4.9
1992	38	5.8	4.0-8.2	19	6.0	3.4-9.8	19	5.6	3.3-8.8	16	2.6	1.4-4.3	5	2.1	0.6-5.2	11	3.1	1.5-5.7
1993	24	3.2	2.0-4.9	11	2.7	1.3-5.4	13	3.5	1.8-6.0	24	3.7	2.3-5.6	11	3.7	1.8-6.9	13	3.7	1.9-6.4
1994	20	3.0	1.7-4.8	9	2.4	0.9-5.2	11	3.3	1.6-6.0	8	1.1	0.5-2.2	4	1.0	0.3-3.0	4	1.1	0.3-2.9
1995	33	5.0	3.4-7.2	16	5.1	2.7-8.7	17	4.8	2.8-7.8	16	2.6	1.4-4.2	9	3.2	1.3-6.3	7	2.1	0.8-4.3
1996	39	5.3	3.7-7.4	22	6.7	4.0-10.4	17	4.2	2.4-6.9	18	2.9	1.6-4.6	12	4.2	2.0-7.8	6	1.8	0.7-3.9
1997	32	4.1	2.8-5.9	22	6.4	3.6-10.3	10	2.4	1.1-4.5	11	1.5	0.8-2.8	7	1.9	0.8-4.2	4	1.1	0.3-2.9
1998	21	2.4	1.5-3.8	14	3.3	1.7-5.8	7	1.6	0.7-3.4	13	1.8	0.9-3.1	10	3.0	1.4-5.8	3	0.8	0.2-2.2
1999	35	4.3	2.9-6.0	23	5.6	3.4-8.6	12	2.9	1.5-5.1	13	1.7	0.9-2.9	8	2.4	1.0-5.0	5	1.1	0.3-2.6
2000	27	3.1	2.0-4.6	11	2.3	1.1-4.4	16	3.6	2.0-5.9	13	1.7	0.9-2.9	8	2.3	1.0-4.6	5	1.2	0.4-2.8
2001	52	6.9	5.1-9.2	22	5.8	3.5-9.2	30	7.4	4.9-10.6	22	3.4	2.1-5.1	9	3.6	1.5-7.0	13	3.2	1.7-5.5
2002	42	4.8	3.4-6.5	19	5.0	2.9-8.1	23	4.7	3.0-7.1	20	2.4	1.4-3.7	10	2.0	1.0-4.0	10	2.4	1.1-4.5
2003	38	4.5	3.1-6.2	18	4.5	2.6-7.4	20	4.3	2.6-6.8	21	2.5	1.5-3.9	12	3.1	1.5-5.6	9	2.0	0.9-3.9
2004	42	4.6	3.2-6.3	19	4.3	2.5-6.9	23	4.7	2.9-7.1	13	1.6	0.8-2.8	4	1.3	0.3-3.5	9	1.9	0.9-3.7
2005	34	4.0	2.7-5.7	24	6.6	4.1-10.0	10	2.0	1.0-3.8	13	1.5	0.8-2.6	8	1.9	0.8-3.9	5	1.1	0.4-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Brain and other central nervous system

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	30	3.1	2.0-4.5	14	2.9	1.5-5.2	16	3.2	1.8-5.2	14	1.6	0.8-2.7	11	2.7	1.2-5.0	3	0.7	0.1-1.9
2007	28	3.1	2.0-4.5	14	3.1	1.6-5.4	14	2.9	1.6-4.9	16	2.1	1.2-3.5	7	2.6	1.0-5.3	9	2.0	0.9-3.7
2008	39	4.2	3.0-5.9	16	3.3	1.8-5.6	23	4.7	3.0-7.1	19	2.1	1.2-3.3	9	1.7	0.8-3.4	10	2.1	1.0-3.9
2009	39	4.3	3.0-5.9	22	5.3	3.1-8.3	17	3.5	2.0-5.7	17	2.3	1.3-3.6	8	2.6	1.1-5.2	9	2.0	0.9-3.8
2010	54	5.6	4.1-7.3	31	6.8	4.5-9.8	23	4.5	2.8-6.7	21	2.2	1.3-3.3	12	2.7	1.3-4.9	9	1.7	0.8-3.3
2011	46	4.4	3.2-5.9	24	4.4	2.8-6.6	22	4.1	2.5-6.2	23	2.3	1.4-3.4	15	3.2	1.7-5.5	8	1.5	0.6-3.0
2012	47	4.3	3.2-5.8	25	4.8	3.0-7.2	22	3.8	2.4-5.9	25	2.6	1.6-3.8	13	2.5	1.3-4.4	12	2.4	1.2-4.2
2013	43	3.9	2.8-5.3	19	3.4	2.0-5.5	24	4.2	2.7-6.3	26	2.4	1.6-3.6	15	3.1	1.7-5.3	11	1.9	0.9-3.4
2014	39	3.5	2.4-4.8	21	4.1	2.4-6.4	18	3.0	1.7-4.7	19	1.9	1.1-3.0	12	2.8	1.4-5.0	7	1.2	0.5-2.5
2015	45	4.1	3.0-5.5	25	4.6	2.9-7.0	20	3.6	2.2-5.6	28	2.8	1.9-4.1	11	2.7	1.3-4.9	17	3.0	1.7-4.8
2016	37	3.4	2.4-4.8	21	4.2	2.5-6.6	16	2.9	1.6-4.7	26	2.3	1.5-3.4	12	2.0	1.0-3.7	14	2.4	1.3-4.1
2017	45	4.0	2.9-5.3	21	4.5	2.7-7.0	24	3.7	2.4-5.6	21	1.9	1.1-2.9	10	1.9	0.9-3.6	11	1.7	0.8-3.1
2018	53	4.6	3.4-6.0	33	6.4	4.3-9.1	20	3.2	1.9-5.0	29	2.4	1.6-3.5	14	2.6	1.3-4.4	15	2.3	1.3-3.9
2019	46	4.0	2.9-5.4	20	3.6	2.2-5.6	26	4.3	2.8-6.3	23	2.1	1.3-3.1	13	2.5	1.3-4.3	10	1.7	0.8-3.1
2020	53	4.3	3.2-5.7	25	4.5	2.9-6.7	28	4.3	2.8-6.2	34	3.0	2.0-4.2	16	2.9	1.6-4.8	18	2.9	1.7-4.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Thyroid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	16	3.6	1.9-6.1	6	3.4	1.0-8.4	10	3.9	1.8-7.4	2	0.8	0.1-2.6	1	0.5	0.0-3.9	1	0.8	0.0-3.8
1977	12	3.1	1.5-5.8	2	0.9	0.1-4.2	10	4.7	2.1-9.1	2	0.9	0.1-3.0	1	0.5	0.0-3.7	1	1.0	0.0-4.4
1978	12	3.0	1.5-5.5	3	1.3	0.2-4.7	9	4.3	1.8-8.4	4	1.6	0.4-4.0	2	1.7	0.2-6.2	2	1.5	0.2-5.0
1979	11	2.5	1.1-4.9	2	0.5	0.1-3.2	9	4.0	1.6-7.9	3	1.5	0.3-4.0	0	0.0	0.0-2.8	3	2.3	0.4-6.3
1980	11	2.2	1.0-4.3	2	0.6	0.1-3.2	9	3.4	1.4-6.9	1	0.2	0.0-1.4	0	0.0	0.0-2.6	1	0.4	0.0-2.4
1981	7	1.1	0.4-2.4	2	0.7	0.1-3.5	5	1.4	0.4-3.6	2	0.5	0.1-1.7	1	0.5	0.0-3.2	1	0.4	0.0-2.3
1982	12	2.3	1.2-4.2	1	0.4	0.0-3.1	11	3.9	1.9-7.1	2	0.4	0.1-1.6	1	0.4	0.0-3.0	1	0.4	0.0-2.2
1983	16	2.9	1.6-4.9	2	0.8	0.1-3.5	14	4.7	2.5-8.1	0	0.0	0.0-0.9	0	0.0	0.0-2.3	0	0.0	0.0-1.5
1984	10	1.6	0.8-3.1	2	0.8	0.1-3.5	8	2.3	1.0-4.8	0	0.0	0.0-0.8	0	0.0	0.0-2.2	0	0.0	0.0-1.4
1985	20	3.9	2.3-6.1	3	1.2	0.2-3.9	17	6.1	3.5-9.9	1	0.2	0.0-1.2	1	0.5	0.0-2.9	0	0.0	0.0-1.4
1986	10	1.9	0.9-3.6	3	1.3	0.3-4.1	7	2.5	1.0-5.2	3	0.7	0.1-1.9	1	0.5	0.0-2.9	2	0.8	0.1-2.7
1987	12	2.2	1.1-3.9	3	1.4	0.3-4.4	9	2.9	1.3-5.6	4	0.8	0.2-2.0	1	0.4	0.0-2.6	3	1.1	0.2-3.1
1988	8	1.4	0.6-2.7	1	0.4	0.0-2.6	7	2.2	0.9-4.6	3	0.6	0.1-1.8	2	1.1	0.1-3.9	1	0.3	0.0-1.8
1989	11	1.8	0.9-3.3	1	0.2	0.0-2.2	10	3.1	1.5-5.8	3	0.8	0.1-2.2	0	0.0	0.0-1.9	3	1.3	0.3-3.5
1990	23	3.7	2.3-5.7	4	1.2	0.3-3.6	19	5.8	3.4-9.1	2	0.6	0.1-2.0	0	0.0	0.0-1.9	2	1.0	0.1-3.1
1991	26	4.5	2.9-6.8	4	2.3	0.6-5.8	22	6.6	4.1-10.1	3	0.7	0.1-1.9	1	0.8	0.0-3.8	2	0.7	0.1-2.3
1992	20	2.8	1.7-4.5	6	2.4	0.7-5.9	14	3.5	1.9-6.0	2	0.3	0.0-1.2	1	0.3	0.0-2.3	1	0.3	0.0-1.7
1993	26	3.7	2.3-5.6	7	1.8	0.7-4.2	19	5.1	3.0-8.2	3	0.6	0.1-1.8	2	0.9	0.1-3.2	1	0.4	0.0-2.0
1994	29	4.3	2.8-6.3	2	0.6	0.1-2.6	27	7.3	4.8-10.7	1	0.1	0.0-0.8	0	0.0	0.0-1.7	1	0.3	0.0-1.5
1995	32	4.8	3.2-6.9	4	1.5	0.4-4.0	28	7.5	4.9-10.9	3	0.6	0.1-1.7	1	0.3	0.0-2.0	2	0.7	0.1-2.3
1996	30	4.1	2.7-5.9	6	2.0	0.7-4.5	24	6.0	3.8-9.0	3	0.5	0.1-1.4	2	0.7	0.1-2.8	1	0.3	0.0-1.5
1997	34	4.6	3.1-6.5	8	2.3	0.9-4.8	26	6.5	4.2-9.6	5	0.7	0.2-1.7	4	1.3	0.3-3.5	1	0.3	0.0-1.4
1998	29	3.8	2.5-5.5	6	1.9	0.7-4.2	23	5.5	3.5-8.3	2	0.4	0.0-1.2	1	0.6	0.0-2.7	1	0.2	0.0-1.4
1999	27	3.5	2.2-5.1	3	0.9	0.2-2.8	24	5.6	3.6-8.4	4	0.6	0.2-1.6	0	0.0	0.0-1.4	4	1.1	0.3-2.8
2000	51	6.4	4.7-8.5	10	2.5	1.2-4.8	41	9.6	6.8-13.1	2	0.4	0.0-1.2	0	0.0	0.0-1.3	2	0.6	0.1-1.9
2001	51	6.0	4.4-8.0	12	3.4	1.5-6.3	39	8.6	6.1-11.8	2	0.3	0.0-1.1	0	0.0	0.0-1.3	2	0.6	0.1-1.9
2002	48	5.6	4.1-7.5	10	2.7	1.2-5.3	38	8.2	5.8-11.3	3	0.5	0.1-1.3	1	0.3	0.0-1.7	2	0.5	0.1-1.9
2003	54	6.2	4.6-8.2	15	4.1	2.2-6.9	39	8.3	5.9-11.4	4	0.6	0.2-1.6	1	0.2	0.0-1.6	3	0.8	0.2-2.3
2004	61	7.3	5.5-9.5	8	3.1	1.2-6.3	53	11.3	8.5-14.9	6	0.9	0.3-2.0	2	0.5	0.1-2.0	4	1.1	0.3-2.7
2005	70	7.5	5.8-9.5	12	2.8	1.4-4.9	58	11.8	8.9-15.3	5	0.7	0.2-1.6	4	1.3	0.3-3.4	1	0.3	0.0-1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Thyroid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	68	7.4	5.7-9.5	16	4.6	2.5-7.6	52	10.4	7.8-13.7	2	0.2	0.0-0.9	1	0.3	0.0-1.6	1	0.2	0.0-1.2
2007	81	9.0	7.1-11.2	18	4.6	2.6-7.4	63	12.9	9.9-16.6	3	0.5	0.1-1.3	2	0.7	0.1-2.5	1	0.3	0.0-1.4
2008	78	8.5	6.7-10.7	18	4.5	2.6-7.2	60	12.0	9.1-15.5	4	0.4	0.1-1.2	2	0.4	0.0-1.5	2	0.4	0.0-1.5
2009	94	10.0	8.0-12.3	22	5.1	3.2-7.9	72	14.1	11.0-17.8	5	0.7	0.2-1.5	1	0.3	0.0-1.5	4	0.9	0.2-2.2
2010	96	9.4	7.6-11.5	18	3.7	2.2-5.9	78	14.4	11.4-18.1	7	0.8	0.3-1.6	4	1.0	0.2-2.7	3	0.6	0.1-1.8
2011	116	11.4	9.4-13.7	25	5.7	3.6-8.6	91	16.5	13.3-20.4	4	0.5	0.1-1.3	0	0.0	0.0-0.9	4	0.8	0.2-2.1
2012	112	10.9	8.9-13.2	26	5.3	3.4-7.9	86	15.8	12.6-19.6	9	1.0	0.4-1.9	3	0.8	0.1-2.4	6	1.1	0.4-2.4
2013	121	11.7	9.6-14.0	17	3.5	2.0-5.7	104	18.5	15.1-22.5	7	0.9	0.3-1.8	0	0.0	0.0-0.9	7	1.4	0.6-2.9
2014	127	11.9	9.9-14.2	21	4.1	2.5-6.4	106	18.9	15.4-22.9	3	0.4	0.1-1.0	1	0.4	0.0-1.7	2	0.4	0.0-1.4
2015	146	13.4	11.3-15.8	34	7.2	4.9-10.2	112	19.4	15.9-23.4	6	0.7	0.2-1.4	3	0.8	0.1-2.4	3	0.6	0.1-1.7
2016	114	10.4	8.5-12.5	24	4.8	3.0-7.3	90	15.4	12.3-19.0	8	0.7	0.3-1.4	4	0.8	0.2-2.2	4	0.6	0.2-1.7
2017	113	10.2	8.4-12.3	27	6.0	3.8-8.7	86	14.4	11.5-17.9	5	0.6	0.2-1.3	3	1.0	0.2-2.7	2	0.3	0.0-1.1
2018	129	11.2	9.3-13.3	29	5.7	3.7-8.3	100	16.4	13.3-20.0	4	0.3	0.1-0.9	3	0.5	0.1-1.6	1	0.2	0.0-0.9
2019	133	11.5	9.6-13.7	24	4.4	2.7-6.6	109	18.2	14.9-22.0	6	0.6	0.2-1.4	1	0.3	0.0-1.4	5	0.8	0.3-2.0
2020	103	8.6	7.0-10.5	18	2.9	1.7-4.7	85	13.8	11.0-17.1	8	0.8	0.4-1.6	2	0.5	0.1-1.8	6	1.1	0.4-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Hodgkin lymphoma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	13	3.2	1.6-5.7	5	1.8	0.5-5.4	8	4.1	1.7-8.2	3	0.7	0.2-2.3	1	0.6	0.0-4.1	2	0.8	0.1-3.3
1977	8	1.4	0.6-3.1	4	1.4	0.3-4.7	4	1.5	0.4-4.0	4	0.9	0.2-2.4	3	1.6	0.3-5.2	1	0.2	0.0-2.2
1978	12	2.1	1.1-4.0	10	3.9	1.8-7.9	2	0.6	0.1-2.8	2	0.4	0.1-1.7	2	0.9	0.1-4.1	0	0.0	0.0-1.8
1979	8	0.9	0.4-2.1	5	1.1	0.3-3.8	3	0.7	0.1-2.6	4	1.0	0.2-2.9	3	0.8	0.1-3.8	1	0.9	0.0-4.0
1980	14	1.8	1.0-3.3	10	2.9	1.3-6.2	4	0.9	0.2-2.8	4	0.7	0.2-2.1	2	1.0	0.1-4.0	2	0.6	0.1-2.5
1981	13	1.9	0.9-3.6	8	2.0	0.8-5.0	5	1.6	0.4-4.1	5	1.1	0.3-2.7	4	1.4	0.3-4.5	1	0.6	0.0-3.0
1982	18	3.4	1.9-5.6	7	3.2	1.3-7.0	11	3.5	1.6-6.7	3	0.7	0.1-2.0	2	1.0	0.1-4.0	1	0.4	0.0-2.3
1983	12	1.7	0.9-3.2	7	2.0	0.8-4.9	5	1.4	0.4-3.5	2	0.2	0.0-1.1	2	0.4	0.1-2.7	0	0.0	0.0-1.5
1984	15	2.7	1.5-4.7	11	5.1	2.3-9.6	4	1.0	0.3-2.8	3	0.6	0.1-1.9	3	1.4	0.2-4.8	0	0.0	0.0-1.4
1985	17	2.7	1.5-4.4	9	3.2	1.4-6.6	8	2.2	0.9-4.6	6	0.9	0.3-2.2	2	1.0	0.1-3.8	4	1.0	0.3-2.9
1986	22	3.0	1.8-4.7	16	4.5	2.5-7.9	6	1.5	0.5-3.5	8	1.2	0.5-2.5	7	2.2	0.8-5.2	1	0.3	0.0-1.9
1987	25	3.3	2.1-5.0	10	3.0	1.4-6.1	15	3.7	2.0-6.3	4	0.6	0.2-1.8	3	1.2	0.3-3.9	1	0.2	0.0-1.6
1988	14	1.5	0.8-2.7	11	2.3	1.1-4.9	3	0.6	0.1-2.1	7	1.1	0.4-2.4	5	2.0	0.6-5.0	2	0.4	0.1-1.9
1989	19	2.4	1.3-4.0	10	2.0	0.9-4.4	9	2.5	1.1-5.0	3	0.7	0.1-2.1	1	0.4	0.0-2.5	2	0.9	0.1-3.0
1990	25	3.4	2.1-5.3	13	3.0	1.6-5.7	12	3.4	1.7-6.2	5	0.7	0.2-1.7	2	0.5	0.1-2.5	3	0.8	0.2-2.5
1991	25	2.9	1.8-4.5	13	3.0	1.5-5.6	12	2.8	1.4-5.1	7	0.8	0.3-1.8	3	0.7	0.1-2.8	4	0.9	0.2-2.5
1992	13	1.6	0.8-2.9	11	3.0	1.5-5.8	2	0.4	0.0-1.6	4	0.5	0.1-1.3	3	0.8	0.1-2.8	1	0.2	0.0-1.3
1993	26	2.8	1.8-4.3	17	3.6	2.0-6.3	9	2.1	0.9-4.0	3	0.4	0.1-1.3	2	0.5	0.1-2.5	1	0.3	0.0-1.6
1994	13	1.6	0.8-2.9	6	1.5	0.5-3.7	7	1.7	0.6-3.6	5	0.7	0.2-1.7	4	1.2	0.3-3.4	1	0.3	0.0-1.5
1995	26	2.8	1.8-4.3	18	3.8	2.2-6.4	8	1.9	0.8-3.7	8	0.9	0.4-1.8	8	1.8	0.7-3.9	0	0.0	0.0-1.0
1996	24	2.8	1.8-4.3	11	2.7	1.3-5.2	13	3.0	1.6-5.1	1	0.2	0.0-0.8	1	0.4	0.0-2.1	0	0.0	0.0-1.0
1997	18	1.9	1.1-3.0	14	2.9	1.6-5.2	4	0.8	0.2-2.2	4	0.4	0.1-1.2	2	0.4	0.0-2.0	2	0.5	0.1-1.8
1998	26	2.9	1.8-4.3	16	3.9	2.2-6.7	10	2.0	1.0-3.8	2	0.2	0.0-0.8	0	0.0	0.0-1.4	2	0.4	0.0-1.6
1999	29	3.2	2.1-4.6	16	3.5	2.0-5.9	13	2.8	1.5-4.9	4	0.6	0.2-1.5	2	0.6	0.1-2.2	2	0.6	0.1-2.0
2000	33	3.4	2.3-4.8	20	4.2	2.5-6.7	13	2.6	1.4-4.5	0	0.0	0.0-0.5	0	0.0	0.0-1.3	0	0.0	0.0-0.9
2001	32	3.4	2.3-4.9	20	4.2	2.6-6.7	12	2.6	1.3-4.5	4	0.5	0.1-1.3	2	0.6	0.1-2.4	2	0.4	0.0-1.6
2002	26	2.7	1.7-4.0	17	3.3	1.9-5.5	9	2.0	0.9-3.8	3	0.3	0.1-1.0	3	0.7	0.1-2.2	0	0.0	0.0-0.8
2003	23	2.2	1.4-3.3	10	1.8	0.9-3.6	13	2.5	1.3-4.3	1	0.1	0.0-0.6	1	0.2	0.0-1.5	0	0.0	0.0-0.8
2004	29	2.7	1.8-4.0	13	2.5	1.3-4.5	16	3.0	1.7-4.9	1	0.1	0.0-0.6	1	0.2	0.0-1.5	0	0.0	0.0-0.8
2005	25	2.5	1.6-3.8	15	3.4	1.8-5.8	10	1.9	0.9-3.6	6	0.6	0.2-1.3	5	1.1	0.3-2.7	1	0.2	0.0-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Hodgkin lymphoma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	33	3.7	2.5-5.2	19	4.3	2.5-7.0	14	3.0	1.6-5.0	4	0.4	0.1-1.1	4	0.9	0.2-2.5	0	0.0	0.0-0.8
2007	32	3.0	2.0-4.3	21	4.4	2.6-6.9	11	1.8	0.9-3.3	5	0.6	0.2-1.4	2	0.7	0.1-2.4	3	0.6	0.1-1.7
2008	30	2.9	2.0-4.2	19	4.1	2.4-6.5	11	1.9	1.0-3.5	3	0.3	0.1-0.9	2	0.4	0.1-1.7	1	0.2	0.0-1.1
2009	34	3.2	2.2-4.5	20	4.0	2.4-6.3	14	2.5	1.3-4.2	3	0.3	0.1-0.9	2	0.4	0.1-1.7	1	0.1	0.0-1.0
2010	38	3.5	2.4-4.8	21	4.0	2.5-6.3	17	3.0	1.7-4.9	1	0.1	0.0-0.5	1	0.2	0.0-1.2	0	0.0	0.0-0.7
2011	34	2.8	1.9-4.0	20	3.4	2.0-5.3	14	2.3	1.3-3.9	7	0.6	0.2-1.3	4	0.7	0.2-1.9	3	0.5	0.1-1.5
2012	39	3.6	2.6-5.0	17	3.6	2.0-5.8	22	3.8	2.4-5.8	1	0.1	0.0-0.5	1	0.2	0.0-1.1	0	0.0	0.0-0.7
2013	34	3.1	2.1-4.3	19	3.5	2.1-5.6	15	2.7	1.5-4.4	2	0.2	0.0-0.7	1	0.2	0.0-1.2	1	0.2	0.0-1.0
2014	30	2.6	1.8-3.8	15	2.8	1.5-4.7	15	2.6	1.4-4.3	1	0.1	0.0-0.5	1	0.2	0.0-1.1	0	0.0	0.0-0.7
2015	44	3.9	2.8-5.3	23	4.1	2.6-6.3	21	3.8	2.3-5.8	5	0.5	0.2-1.2	4	1.0	0.2-2.5	1	0.2	0.0-1.1
2016	31	2.9	1.9-4.1	12	2.6	1.3-4.7	19	3.2	1.9-5.1	2	0.2	0.0-0.7	2	0.4	0.0-1.5	0	0.0	0.0-0.6
2017	28	2.4	1.6-3.5	17	3.0	1.7-4.8	11	1.8	0.9-3.2	2	0.1	0.0-0.5	1	0.2	0.0-1.0	1	0.1	0.0-0.9
2018	40	3.3	2.4-4.6	23	4.0	2.5-6.0	17	2.8	1.6-4.5	3	0.3	0.1-0.9	3	0.8	0.1-2.2	0	0.0	0.0-0.6
2019	30	2.4	1.6-3.5	14	2.4	1.3-4.2	16	2.4	1.4-4.0	1	0.1	0.0-0.6	1	0.2	0.0-1.2	0	0.0	0.0-0.6
2020	29	2.3	1.5-3.3	15	2.6	1.4-4.4	14	2.1	1.1-3.5	3	0.2	0.0-0.7	2	0.3	0.0-1.1	1	0.2	0.0-0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Non-Hodgkin lymphomas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	29	7.7	4.9-11.5	17	9.6	5.3-16.1	12	6.0	2.8-10.9	12	4.7	2.2-8.4	6	5.4	1.6-12.5	6	4.0	1.3-8.8
1977	46	12.2	8.7-16.7	28	17.7	11.1-26.7	18	8.3	4.7-13.4	24	6.9	4.2-10.7	14	10.8	5.0-19.6	10	4.6	2.1-8.6
1978	47	12.1	8.6-16.6	24	12.0	7.5-18.4	23	11.6	7.0-17.9	13	3.3	1.6-5.9	8	4.0	1.7-8.5	5	2.6	0.7-6.3
1979	30	6.5	4.2-9.5	16	6.3	3.5-10.9	14	6.3	3.3-10.7	29	7.7	4.8-11.5	18	8.3	4.8-13.7	11	6.4	3.0-11.7
1980	26	6.9	4.2-10.5	16	11.0	5.3-19.5	10	4.5	2.1-8.5	18	5.1	2.8-8.2	8	5.9	2.0-12.6	10	4.8	2.2-9.0
1981	32	8.0	5.2-11.6	20	9.6	5.5-15.7	12	6.3	3.1-11.1	18	4.1	2.3-6.7	8	4.7	1.7-10.0	10	3.9	1.8-7.2
1982	35	7.6	5.2-10.7	24	12.3	7.5-19.0	11	4.1	2.0-7.5	8	1.6	0.7-3.3	6	3.0	1.1-6.9	2	0.6	0.1-2.5
1983	41	8.5	6.0-11.8	23	9.2	5.7-14.3	18	7.4	4.3-11.9	18	3.8	2.2-6.1	10	4.0	1.9-7.8	8	3.4	1.4-6.7
1984	41	8.6	6.0-12.0	24	9.7	6.0-15.0	17	7.3	4.1-11.7	35	7.9	5.4-11.2	20	9.1	5.4-14.4	15	6.5	3.5-10.9
1985	47	9.5	6.8-12.8	28	13.4	8.3-20.3	19	6.8	4.0-10.7	25	5.4	3.5-8.1	14	6.6	3.5-11.4	11	4.5	2.2-8.0
1986	51	10.2	7.4-13.7	25	10.5	6.2-16.4	26	9.7	6.2-14.4	20	4.1	2.4-6.4	12	5.3	2.6-9.8	8	2.9	1.2-5.9
1987	50	9.2	6.7-12.2	29	11.9	7.6-17.8	21	7.0	4.3-10.8	18	4.1	2.3-6.5	12	5.9	2.8-10.8	6	2.6	0.9-5.5
1988	57	10.9	8.0-14.3	35	15.8	10.2-23.1	22	7.5	4.6-11.5	32	7.2	4.7-10.4	21	13.1	7.3-21.0	11	3.8	1.8-7.0
1989	69	11.0	8.5-14.1	37	12.3	8.3-17.5	32	10.0	6.8-14.1	28	5.1	3.3-7.6	16	6.2	3.3-10.7	12	4.2	2.1-7.3
1990	79	14.0	10.9-17.7	46	15.9	11.3-21.9	33	11.5	7.8-16.3	31	5.7	3.8-8.1	16	6.7	3.6-11.3	15	4.9	2.7-8.2
1991	74	11.9	9.1-15.3	45	14.8	10.0-21.1	29	9.4	6.2-13.7	34	6.2	4.2-8.9	19	6.8	4.0-11.0	15	5.3	2.9-8.7
1992	77	12.5	9.7-15.9	55	19.2	13.9-25.7	22	7.0	4.3-10.7	29	5.6	3.6-8.1	19	8.4	4.7-13.5	10	3.5	1.6-6.4
1993	94	14.4	11.5-17.8	57	19.2	14.2-25.5	37	10.4	7.3-14.5	39	7.0	4.9-9.7	16	5.2	2.8-8.8	23	7.8	4.9-11.7
1994	86	13.1	10.4-16.4	47	14.8	10.6-20.2	39	11.4	8.0-15.6	28	5.4	3.5-7.8	19	8.3	4.6-13.6	9	3.2	1.4-6.0
1995	101	14.8	11.9-18.2	71	23.8	17.7-31.1	30	8.1	5.4-11.7	33	5.9	4.0-8.4	16	4.9	2.8-8.2	17	5.8	3.3-9.2
1996	116	17.0	13.9-20.6	70	20.1	15.4-25.9	46	13.3	9.6-17.8	32	5.5	3.7-7.9	19	7.9	4.3-13.0	13	4.0	2.1-6.8
1997	102	14.7	11.8-18.0	61	20.3	14.9-26.9	41	10.8	7.7-14.7	21	3.7	2.2-5.8	11	5.3	2.4-9.8	10	3.0	1.4-5.4
1998	118	16.3	13.4-19.6	73	22.5	17.1-29.0	45	11.6	8.4-15.5	34	5.4	3.7-7.6	20	7.6	4.3-12.2	14	4.0	2.2-6.7
1999	129	17.9	14.8-21.4	74	22.2	17.0-28.4	55	14.1	10.6-18.5	32	5.0	3.4-7.2	15	4.7	2.5-8.0	17	4.9	2.8-7.9
2000	101	13.0	10.5-15.9	55	16.1	11.8-21.4	46	10.9	7.9-14.6	42	6.1	4.4-8.4	31	10.6	6.8-15.5	11	2.9	1.5-5.2
2001	124	16.1	13.2-19.3	74	20.1	15.5-25.6	50	12.2	9.0-16.1	42	6.6	4.7-9.0	21	8.5	5.0-13.2	21	5.5	3.4-8.4
2002	129	16.3	13.5-19.5	74	20.2	15.6-25.8	55	13.0	9.7-16.9	48	6.6	4.8-8.9	26	7.7	4.9-11.6	22	5.5	3.4-8.4
2003	122	15.6	12.9-18.8	56	16.3	12.0-21.5	66	15.4	11.9-19.7	27	3.8	2.4-5.5	14	4.1	2.1-7.0	13	3.3	1.7-5.6
2004	112	14.3	11.7-17.4	63	19.3	14.2-25.4	49	11.2	8.2-14.9	36	5.2	3.6-7.3	19	5.8	3.3-9.4	17	4.5	2.6-7.1
2005	118	14.7	12.1-17.7	68	18.8	14.2-24.3	50	11.6	8.5-15.3	32	4.7	3.2-6.7	18	6.6	3.7-10.6	14	3.5	1.9-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Non-Hodgkin lymphomas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	140	16.6	13.9-19.7	75	19.7	15.2-25.1	65	14.1	10.8-18.0	34	4.2	2.9-5.9	19	4.8	2.8-7.8	15	3.5	2.0-5.8
2007	128	14.8	12.2-17.6	76	20.9	16.1-26.6	52	10.7	7.9-14.0	31	3.9	2.6-5.6	19	6.0	3.4-9.7	12	2.7	1.4-4.7
2008	161	17.9	15.2-21.0	96	24.3	19.4-30.1	65	13.2	10.1-16.8	32	3.8	2.6-5.5	21	6.3	3.7-9.8	11	2.2	1.1-4.0
2009	136	15.0	12.5-17.8	69	17.3	13.2-22.2	67	13.4	10.3-17.0	34	3.8	2.6-5.4	23	6.1	3.7-9.4	11	2.2	1.1-4.0
2010	144	15.5	13.0-18.4	78	18.1	14.1-22.9	66	13.1	10.1-16.8	42	5.0	3.6-6.8	22	5.9	3.6-9.2	20	4.4	2.7-6.8
2011	162	17.2	14.5-20.1	103	25.2	20.2-31.1	59	11.4	8.6-14.7	42	5.0	3.6-6.8	27	7.6	4.8-11.3	15	3.2	1.7-5.2
2012	142	14.4	12.1-17.1	74	16.9	13.1-21.5	68	12.4	9.6-15.8	33	3.7	2.5-5.3	16	4.8	2.6-7.9	17	3.2	1.8-5.1
2013	147	15.1	12.7-17.8	78	18.2	14.2-23.0	69	12.7	9.8-16.1	28	3.2	2.1-4.7	14	3.3	1.7-5.7	14	2.9	1.6-4.8
2014	154	15.2	12.9-17.9	89	20.1	15.9-25.0	65	11.5	8.9-14.8	43	4.9	3.5-6.6	25	7.3	4.6-10.8	18	3.4	2.0-5.4
2015	180	17.1	14.6-19.8	86	18.6	14.6-23.2	94	16.3	13.1-20.0	39	3.9	2.8-5.4	21	4.9	2.9-7.7	18	3.1	1.8-5.0
2016	174	16.5	14.1-19.3	92	19.9	15.8-24.6	82	14.0	11.1-17.4	36	3.7	2.6-5.1	21	5.5	3.3-8.5	15	2.5	1.4-4.1
2017	173	16.1	13.8-18.8	94	20.2	16.1-24.9	79	13.2	10.4-16.5	49	4.7	3.4-6.2	34	7.6	5.1-10.8	15	2.6	1.5-4.4
2018	166	15.0	12.8-17.6	93	18.9	15.0-23.3	73	12.2	9.5-15.4	37	3.2	2.2-4.4	21	4.0	2.4-6.2	16	2.6	1.5-4.3
2019	191	16.7	14.4-19.3	113	21.9	17.8-26.5	78	12.4	9.7-15.5	47	4.4	3.2-5.9	28	6.6	4.3-9.6	19	3.1	1.8-4.8
2020	153	13.3	11.3-15.7	79	15.5	12.1-19.5	74	11.8	9.2-14.8	41	3.6	2.6-4.9	20	3.7	2.2-5.9	21	3.3	2.0-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Myeloma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	31	9.2	6.1-13.2	13	8.6	4.3-15.3	18	9.7	5.6-15.5	22	6.4	3.9-9.9	11	7.1	3.3-13.3	11	5.8	2.8-10.6
1977	30	10.3	6.7-15.1	13	10.1	4.6-18.6	17	10.5	5.9-17.1	19	7.5	4.2-11.9	12	11.8	5.3-21.6	7	4.7	1.7-9.7
1978	29	8.7	5.6-12.7	12	10.1	4.5-18.7	17	8.3	4.8-13.4	16	5.4	2.9-9.1	9	8.8	3.4-17.4	7	3.6	1.4-7.6
1979	35	11.0	7.4-15.6	15	14.2	7.1-24.5	20	9.8	5.9-15.1	16	4.8	2.6-8.0	8	6.2	2.1-13.2	8	4.1	1.7-8.1
1980	33	9.5	6.4-13.6	13	8.3	4.3-14.6	20	10.1	6.0-15.8	22	6.4	3.9-9.8	9	5.6	2.4-10.9	13	6.8	3.5-11.7
1981	40	12.6	8.7-17.4	15	12.2	6.2-21.0	25	13.0	8.2-19.3	26	8.0	5.1-11.8	13	10.9	5.4-19.0	13	6.3	3.3-10.8
1982	47	13.1	9.4-17.7	26	17.1	10.6-25.8	21	10.1	6.1-15.6	28	7.1	4.6-10.4	16	9.9	5.4-16.4	12	5.2	2.6-9.1
1983	26	6.8	4.3-10.1	16	10.5	5.6-17.7	10	4.4	2.1-8.1	26	7.7	4.9-11.3	13	10.5	5.0-18.4	13	6.3	3.3-10.7
1984	49	12.5	9.1-16.7	27	14.9	9.3-22.4	22	10.4	6.4-15.8	27	7.6	4.8-11.2	13	9.9	4.8-17.4	14	6.4	3.4-10.7
1985	38	9.4	6.5-13.0	21	11.7	6.8-18.4	17	7.6	4.3-12.1	29	7.4	4.8-10.6	16	8.6	4.8-14.1	13	6.1	3.2-10.4
1986	52	14.1	10.3-18.7	25	16.4	10.0-24.9	27	12.5	8.1-18.3	35	9.9	6.7-13.9	22	14.7	8.7-22.7	13	6.6	3.5-11.2
1987	39	9.8	6.8-13.5	19	14.1	7.8-22.6	20	8.0	4.8-12.3	22	6.1	3.7-9.3	13	9.9	4.8-17.4	9	4.1	1.8-7.6
1988	45	10.6	7.6-14.3	20	11.5	6.6-18.2	25	9.9	6.4-14.7	20	4.7	2.8-7.3	10	5.4	2.5-10.0	10	4.1	1.9-7.5
1989	57	13.6	10.1-17.7	24	13.5	8.1-20.7	33	13.5	9.2-19.0	29	7.4	4.9-10.8	11	7.4	3.3-13.6	18	7.5	4.3-11.8
1990	54	12.3	9.1-16.1	29	16.8	10.8-24.7	25	9.5	6.1-14.0	33	7.6	5.1-10.8	16	9.0	4.7-15.2	17	6.8	3.9-10.8
1991	59	12.5	9.4-16.3	26	14.0	8.6-21.1	33	11.8	8.1-16.6	26	5.7	3.6-8.4	12	6.8	3.3-11.9	14	5.0	2.7-8.4
1992	58	12.0	9.0-15.6	27	13.8	8.6-20.7	31	10.9	7.4-15.5	50	11.2	8.2-14.9	23	14.2	8.4-22.0	27	9.9	6.5-14.4
1993	57	11.3	8.5-14.7	26	14.5	8.9-21.9	31	10.1	6.8-14.4	39	7.8	5.5-10.8	15	8.2	4.5-13.6	24	7.9	5.0-11.7
1994	61	12.9	9.7-16.6	27	13.1	8.1-19.5	34	12.1	8.3-16.9	41	8.7	6.1-11.9	16	8.4	4.5-13.9	25	8.6	5.5-12.7
1995	52	9.8	7.2-13.0	28	14.2	8.8-21.2	24	7.5	4.8-11.1	30	6.0	4.0-8.6	14	7.2	3.5-12.6	16	5.4	3.1-8.7
1996	61	10.8	8.2-14.0	30	12.6	8.0-18.7	31	9.6	6.5-13.7	28	5.2	3.4-7.6	17	8.2	4.3-13.8	11	3.5	1.7-6.2
1997	72	13.0	10.1-16.4	36	15.6	10.6-22.0	36	11.1	7.7-15.4	47	8.8	6.4-11.8	31	15.1	9.7-21.9	16	5.1	2.9-8.2
1998	67	11.6	8.9-14.8	31	13.2	8.6-19.2	36	10.6	7.4-14.6	36	6.7	4.7-9.3	15	7.4	3.8-12.6	21	6.6	4.1-10.0
1999	67	11.4	8.7-14.5	35	13.6	9.2-19.3	32	9.5	6.5-13.4	37	7.0	4.9-9.6	17	8.0	4.5-12.9	20	6.2	3.8-9.5
2000	87	13.8	11.0-17.2	41	15.5	10.6-21.6	46	12.9	9.4-17.2	45	7.7	5.6-10.4	25	10.3	6.4-15.5	20	5.9	3.6-9.1
2001	87	13.8	11.0-17.2	41	14.4	10.0-20.0	46	13.0	9.5-17.4	49	8.3	6.0-10.9	18	7.4	4.1-12.2	31	8.9	6.0-12.5
2002	72	11.4	8.8-14.4	30	12.0	7.7-17.5	42	11.2	8.0-15.1	40	6.1	4.3-8.3	16	5.4	2.9-9.1	24	6.5	4.1-9.7
2003	87	12.3	9.8-15.3	41	13.8	9.7-19.1	46	11.3	8.2-15.1	35	5.2	3.6-7.2	17	6.1	3.4-9.9	18	4.6	2.7-7.3
2004	94	14.1	11.3-17.3	42	16.1	11.1-22.3	52	13.3	9.9-17.5	42	6.9	4.9-9.4	24	10.6	6.6-16.0	18	4.8	2.8-7.6
2005	115	16.3	13.3-19.6	50	16.6	12.0-22.2	65	15.9	12.2-20.3	38	5.5	3.8-7.6	16	4.9	2.7-8.1	22	5.6	3.5-8.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Myeloma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	104	13.5	11.0-16.4	52	16.6	12.1-22.1	52	11.7	8.7-15.3	45	6.4	4.7-8.7	26	8.6	5.5-12.8	19	4.7	2.8-7.3
2007	93	11.6	9.3-14.3	43	13.0	9.2-17.8	50	11.0	8.1-14.5	44	6.4	4.6-8.6	22	7.7	4.7-11.8	22	5.4	3.3-8.1
2008	119	14.0	11.5-16.8	54	14.2	10.5-18.8	65	13.7	10.5-17.5	42	5.7	4.0-7.7	22	7.2	4.4-11.1	20	4.6	2.8-7.2
2009	126	15.1	12.5-18.1	66	17.9	13.6-23.1	60	12.6	9.6-16.3	36	4.5	3.1-6.2	19	5.1	3.0-8.1	17	3.7	2.2-6.0
2010	133	16.0	13.3-19.0	62	19.1	14.4-24.8	71	14.3	11.1-18.1	50	6.5	4.8-8.6	24	7.7	4.8-11.6	26	5.5	3.6-8.1
2011	116	13.5	11.1-16.2	44	11.5	8.2-15.6	72	14.6	11.4-18.4	43	5.2	3.7-7.0	15	4.7	2.5-7.8	28	5.7	3.8-8.2
2012	139	14.8	12.3-17.5	78	19.8	15.3-25.0	61	11.5	8.7-14.8	49	5.7	4.1-7.5	25	6.3	4.0-9.5	24	4.9	3.1-7.4
2013	161	17.8	15.1-20.9	88	22.3	17.6-27.8	73	14.6	11.4-18.3	48	5.7	4.2-7.6	23	7.2	4.4-10.9	25	4.9	3.2-7.3
2014	149	15.1	12.7-17.8	91	21.4	17.0-26.5	58	10.6	8.0-13.8	48	5.6	4.1-7.4	19	5.6	3.3-8.9	29	5.7	3.8-8.2
2015	150	15.0	12.6-17.7	74	17.6	13.6-22.3	76	13.2	10.4-16.6	54	5.6	4.2-7.3	28	6.5	4.2-9.6	26	4.7	3.1-7.0
2016	167	15.9	13.5-18.5	90	20.8	16.5-25.8	77	12.7	10.0-16.0	45	4.7	3.4-6.3	21	5.5	3.3-8.4	24	4.3	2.8-6.5
2017	153	14.7	12.4-17.3	74	16.8	13.0-21.4	79	13.2	10.4-16.5	45	4.7	3.4-6.3	20	5.6	3.3-8.6	25	4.3	2.7-6.3
2018	180	16.0	13.7-18.6	98	20.1	16.1-24.8	82	12.9	10.2-16.1	56	5.7	4.3-7.4	29	7.7	5.1-11.1	27	4.5	3.0-6.6
2019	179	16.2	13.9-18.9	90	18.6	14.8-23.1	89	14.5	11.6-18.0	47	4.5	3.3-6.1	28	6.3	4.1-9.3	19	3.3	2.0-5.1
2020	118	10.7	8.8-12.8	56	12.2	9.1-15.9	62	9.6	7.3-12.3	54	4.9	3.6-6.4	28	6.5	4.2-9.4	26	4.0	2.6-5.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Leukemias

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	31	7.1	4.5-10.6	19	10.8	5.3-19.0	12	4.6	2.3-8.5	18	4.8	2.6-8.0	12	6.8	2.9-13.3	6	3.1	1.0-7.1
1977	35	10.7	7.0-15.4	21	17.2	9.1-28.5	14	6.8	3.6-11.6	20	5.1	2.8-8.4	14	8.1	3.4-15.8	6	3.0	0.9-6.9
1978	39	10.0	6.8-14.2	16	7.8	4.1-13.4	23	11.4	6.9-17.4	32	9.7	6.2-14.2	15	10.1	4.8-18.2	17	9.2	5.0-15.2
1979	36	9.0	6.0-12.8	15	6.9	3.7-11.9	21	10.2	6.0-15.8	22	5.3	3.3-8.2	12	7.1	3.5-12.8	10	4.1	1.9-7.7
1980	43	10.0	6.9-13.8	21	11.6	6.4-19.1	22	9.0	5.5-13.9	31	6.9	4.5-10.1	13	6.7	3.1-12.5	18	7.2	4.2-11.6
1981	40	10.3	7.1-14.4	22	13.9	7.7-22.5	18	8.1	4.7-13.0	27	7.5	4.7-11.3	13	10.7	4.8-19.7	14	6.1	3.2-10.4
1982	32	6.7	4.4-9.8	15	7.4	3.8-12.9	17	6.3	3.6-10.4	18	3.5	2.0-5.7	8	3.3	1.2-7.3	10	3.6	1.7-6.9
1983	42	9.6	6.7-13.4	24	13.5	7.7-21.6	18	7.3	4.2-11.7	26	6.4	4.0-9.5	12	9.1	4.1-16.7	14	5.3	2.8-9.0
1984	33	6.6	4.3-9.7	16	7.2	3.6-12.6	17	6.2	3.4-10.3	22	5.3	3.1-8.3	13	6.9	3.2-12.9	9	4.2	1.8-7.9
1985	58	11.7	8.6-15.4	29	13.6	8.5-20.5	29	10.4	6.8-15.3	22	4.6	2.7-7.1	12	6.0	2.6-11.4	10	3.7	1.7-6.9
1986	42	8.4	5.9-11.7	28	13.0	7.9-19.9	14	5.2	2.7-8.9	35	7.0	4.7-10.0	21	7.6	4.5-12.3	14	5.9	3.1-10.0
1987	50	10.0	7.2-13.4	27	12.2	7.5-18.7	23	8.3	5.1-12.6	31	7.0	4.6-10.2	15	8.5	4.1-15.1	16	6.4	3.6-10.4
1988	45	9.4	6.7-12.7	19	11.0	6.2-17.6	26	8.9	5.8-13.2	40	8.6	6.0-12.0	20	11.3	6.2-18.4	20	7.2	4.3-11.3
1989	48	10.0	7.2-13.5	26	12.8	7.7-19.8	22	8.0	4.9-12.3	18	4.0	2.3-6.4	14	8.3	4.2-14.4	4	1.4	0.3-3.6
1990	64	12.4	9.3-16.0	33	14.4	9.2-21.3	31	10.9	7.3-15.5	33	7.2	4.8-10.2	14	8.1	3.9-14.3	19	6.9	4.1-10.8
1991	58	10.6	7.8-13.9	28	12.2	7.4-18.7	30	9.7	6.4-14.0	35	6.8	4.6-9.6	23	10.7	6.1-17.0	12	4.3	2.2-7.4
1992	57	9.6	7.1-12.6	30	11.6	7.2-17.6	27	8.3	5.4-12.2	21	4.2	2.5-6.5	12	5.5	2.4-10.2	9	3.3	1.5-6.2
1993	54	9.8	7.2-13.0	29	13.3	8.2-19.9	25	7.6	4.8-11.4	27	4.6	2.9-6.9	16	7.0	3.5-12.2	11	3.2	1.6-5.8
1994	56	9.7	7.1-12.8	30	13.1	8.3-19.4	26	7.5	4.8-11.2	43	8.6	6.1-11.7	23	11.5	6.8-17.8	20	6.6	4.0-10.2
1995	67	10.8	8.2-13.9	41	16.2	10.9-22.9	26	7.3	4.7-10.8	42	7.0	4.9-9.5	23	10.1	6.0-15.7	19	5.2	3.1-8.2
1996	74	11.3	8.6-14.4	44	15.8	10.9-21.9	30	7.9	5.3-11.5	42	7.1	5.0-9.7	21	8.4	4.9-13.3	21	6.1	3.7-9.3
1997	64	9.9	7.5-12.8	36	11.6	7.8-16.5	28	8.2	5.4-11.8	39	6.8	4.7-9.4	23	9.5	5.7-14.7	16	4.8	2.7-7.8
1998	73	11.3	8.7-14.3	36	13.8	8.9-20.0	37	10.3	7.2-14.2	47	7.9	5.7-10.7	25	10.6	6.4-16.0	22	6.4	4.0-9.7
1999	60	8.6	6.5-11.2	30	11.5	7.2-17.1	30	7.4	5.0-10.7	37	5.9	4.0-8.2	18	6.8	3.7-11.2	19	5.3	3.1-8.2
2000	66	9.5	7.2-12.2	39	12.7	8.7-17.8	27	7.0	4.6-10.3	32	4.6	3.0-6.5	18	5.9	3.2-9.7	14	3.6	1.9-6.1
2001	73	10.2	7.9-13.0	39	12.5	8.6-17.6	34	8.5	5.8-11.9	47	6.9	5.0-9.3	24	8.5	5.1-13.2	23	5.7	3.6-8.7
2002	85	12.3	9.7-15.3	46	15.0	10.7-20.3	39	10.0	7.0-13.7	31	5.0	3.4-7.2	14	4.5	2.3-7.9	17	5.0	2.9-7.9
2003	79	10.1	7.9-12.7	39	11.8	8.1-16.6	40	9.0	6.4-12.4	38	5.8	4.0-8.0	17	5.6	3.2-9.1	21	5.5	3.4-8.4
2004	98	12.7	10.2-15.7	60	18.6	13.6-24.7	38	8.8	6.2-12.1	52	7.8	5.8-10.3	28	11.0	6.9-16.3	24	6.1	3.8-9.0
2005	69	9.0	6.9-11.5	36	11.3	7.6-16.1	33	7.5	5.1-10.6	41	5.7	4.1-7.8	21	7.0	4.1-11.0	20	4.8	2.9-7.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
New York State Excl New York City
Black Individuals
Leukemias

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2006	100	12.0	9.7-14.7	57	15.3	11.2-20.3	43	9.5	6.8-12.8	39	5.1	3.6-7.0	19	5.9	3.3-9.4	20	4.6	2.8-7.2
2007	89	10.6	8.4-13.1	49	14.0	10.0-19.0	40	8.5	6.1-11.7	43	5.5	3.9-7.5	23	7.2	4.4-11.0	20	4.4	2.7-6.9
2008	92	10.1	8.1-12.5	52	12.3	8.9-16.4	40	8.2	5.8-11.3	44	5.3	3.8-7.2	23	6.8	4.1-10.6	21	4.5	2.8-6.9
2009	76	8.7	6.8-11.0	47	11.7	8.4-15.8	29	6.2	4.1-9.0	34	3.9	2.6-5.4	22	5.5	3.3-8.6	12	2.5	1.3-4.4
2010	105	11.1	9.1-13.6	66	15.5	11.7-20.0	39	7.7	5.4-10.5	35	4.1	2.8-5.8	23	6.9	4.2-10.6	12	2.3	1.2-4.1
2011	111	12.1	9.9-14.7	66	17.0	12.9-21.9	45	8.5	6.2-11.4	47	5.6	4.1-7.5	28	8.3	5.3-12.1	19	3.8	2.3-6.0
2012	126	13.5	11.2-16.2	61	15.9	11.9-20.7	65	12.1	9.3-15.5	47	5.2	3.8-6.9	31	8.1	5.3-11.8	16	3.0	1.7-4.9
2013	127	13.3	11.0-15.9	66	15.5	11.8-19.9	61	11.3	8.6-14.6	49	5.4	3.9-7.2	28	7.4	4.8-10.9	21	4.0	2.4-6.1
2014	119	11.9	9.8-14.3	74	18.2	14.0-23.2	45	7.8	5.7-10.5	49	5.3	3.8-7.0	24	6.4	4.0-9.7	25	4.5	2.9-6.7
2015	107	10.7	8.7-13.0	62	14.8	11.2-19.2	45	7.9	5.7-10.6	40	4.1	2.9-5.6	20	5.1	3.0-8.0	20	3.5	2.1-5.4
2016	137	13.4	11.2-15.9	78	18.3	14.2-23.0	59	10.1	7.6-13.0	61	6.3	4.8-8.1	32	7.9	5.2-11.3	29	5.3	3.5-7.6
2017	114	10.3	8.4-12.4	57	10.7	8.0-14.1	57	9.8	7.4-12.8	55	5.4	4.0-7.0	27	6.4	4.1-9.4	28	4.8	3.2-7.0
2018	108	9.7	7.9-11.7	67	13.7	10.4-17.6	41	6.6	4.7-9.0	47	4.5	3.3-6.1	28	6.8	4.4-10.0	19	3.2	1.9-5.0
2019	119	10.3	8.5-12.4	55	11.4	8.4-15.0	64	9.8	7.5-12.6	46	4.2	3.0-5.6	23	5.0	3.1-7.6	23	3.5	2.2-5.3
2020	106	9.1	7.4-11.1	55	10.8	8.0-14.2	51	8.0	5.9-10.6	50	4.4	3.2-5.8	25	5.0	3.1-7.5	25	3.9	2.5-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.