Expanded BRFSS, 2003

Rensselaer Locality Tables

TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days	
Core 2: Health Care Access	4
Core 3: Exercise	8
Core 3: Exercise	(
Core 5: Asthma	1(
Core 6: Arthritis	12
Core 6: Arthritis	14
Core 8: Tobacco ETS	11
Core 9: Alcohol Consumption	19
Core 10: Weight Status	2
Core 11: Mammography	23
Core 12: Sexual Behavior	2:
Core 13: Family Planning	28
Core 14: Cardiovascular Disease	29
Core 15: Prostate Cancer Screening	
	3

New York State Department Of Health

Self-rated Health Status

Demographic Groups	n¹	Fair or Poor	n	Good to Excellent %	C.I. ³
Total	95	12.3	543	87.7	2.6
Gender					
Male	37	12.2	209	87.8	4.1
Female	58	12.3	334	87.7	3.3
Age					
18-34	8	4.5	142	95.5	3.2
35-54	27	10.1	215	89.9	4.2
≥55	58	22.6	181	77.4	5.7
Education					
≤High School	65	21.4	197	78.6	5.4
>High School	30	6.4	343	93.6	2.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Reported Number of Physical Distress Days in the Past Month *Rensselaer*

Demographic Groups	n¹	≥14 days %2	n	< 14 days %	C.I. ³
Total	64	9.0	566	91.0	2.4
Gender					
Male	22	9.0	222	91.0	3.8
Female	42	9.1	344	90.9	2.9
Age					
18-34	5	2.4	145	97.6	2.2
35-54	28	11.4	212	88.6	4.4
≥55	30	12.7	202	87.3	4.8
Education					
≤High School	35	12.3	222	87.7	4.3
>High School	29	6.9	341	93.1	2.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Reported Number of Mental Distress Days in the Past Month Rensselaer

Demographic Groups	n¹	≥14 days %2	n	< 14 days %	C.I. ³
Total	63	9.4	563	90.6	2.5
Gender					
Male	23	9.1	221	90.9	3.9
Female	40	9.8	342	90.2	3.1
Age					
18-34	22	12.5	128	87.5	5.5
35-54	29	11.0	210	89.0	4.2
≥55	12	4.5	218	95.5	2.8
Education					
≤High School	36	13.4	220	86.6	4.6
>High School	27	6.9	340	93.1	2.8

¹ Percents based on row denominators of less than 50 are unstable and should be used with caution

² Weighted Percent³ 95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health **Health Insurance Coverage, 18-64 Years of Age** Rensselaer

Demographic		Yes		No	
Groups	n¹	% ²	n	%	C.I. ³
Total	434	89.7	49	10.3	3.1
Gender					
Male	171	89.5	19	10.5	5.0
Female	263	89.8	30	10.2	3.7
Age					
18-34	120	82.2	28	17.8	6.9
35-54	228	94.5	14	5.5	3.0
55-64	86	92.4	7	7.6	5.9
Education					
≤High School	147	83.9	30	16.1	5.7
>High School	286	92.8	19	7.2	3.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health **Health Insurance Coverage, Aged 18 and Older** Rensselaer

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	584	91.2	53	8.8	2.6
Gender					
Male	224	90.8	21	9.2	4.3
Female	360	91.7	32	8.3	3.0
Age					
18-34	120	82.2	28	17.8	6.9
35-54	228	94.5	14	5.5	3.0
≥55	229	96.0	10	4.0	2.7
Education					
≤High School	229	87.7	32	12.3	4.3
>High School	352	93.5	21	6.5	3.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Reported Having a Routine Physical Exam in Past Two Years Rensselaer

Demographic Groups	n¹	Yes %°	n	No %	C.I. ³
Total	579	90.3	55	9.7	2.7
Gender					
Male	214	86.5	30	13.5	4.9
Female	365	93.8	25	6.2	2.6
Age					
18-34	127	86.0	19	14.0	6.4
35-54	218	90.0	24	10.0	4.2
≥55	226	94.6	12	5.4	3.4
Education					
≤High School	232	88.2	27	11.8	4.6
>High School	345	91.8	27	8.2	3.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Time in Past 12 Months when Needed Medical Care but Could Not Get It Rensselaer

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Total	31	4.6	608	95.4	1.9
Gender					
Male	11	4.3	235	95.7	2.8
Female	20	5.0	373	95.0	2.4
Age					
18-34	13	8.2	137	91.8	4.9
35-54	10	3.0	232	97.0	2.0
≥55	8	3.3	231	96.7	2.5
Education					
≤High School	17	5.7	245	94.3	2.9
>High School	14	4.0	360	96.0	2.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days *Rensselaer*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	494	78.7	145	21.3	3.5
Gender					
Male	196	82.3	50	17.7	5.0
Female	298	75.3	95	24.7	4.8
Age					
18-34	125	84.6	25	15.4	6.1
35-54	193	81.0	49	19.0	5.2
≥55	170	70.0	69	30.0	6.8
Education					
≤High School	182	71.4	80	28.6	6.0
>High School	310	83.6	64	16.4	4.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Physician Diagnosed Diabetes

Demographic		Yes		No	G T 1
Groups	n¹	% ²	n	%	C.I. ³
Total	60	8.2	579	91.8	2.2
Gender					
Male	26	8.9	220	91.1	3.6
Female	34	7.5	359	92.5	2.6
Age					
18-34	2	0.9	148	99.1	1.3
35-54	22	9.4	220	90.6	4.0
≥55	33	13.3	206	86.7	4.6
Education					
≤High School	28	9.5	234	90.5	3.9
>High School	32	7.3	342	92.7	2.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Lifetime Asthma Diagnosed by a Medical Professional

Demographic		Yes		No	
Groups	n¹	%2	n	%	C.I. ³
Total	86	13.1	553	86.9	2.9
Gender					
Male	30	11.4	216	88.6	4.3
Female	56	14.6	337	85.4	3.8
Age					
18-34	32	18.9	118	81.1	6.6
35-54	22	8.9	220	91.1	3.9
≥55	32	12.8	207	87.2	4.5
Education					
≤High School	34	11.6	228	88.4	4.0
>High School	52	14.1	322	85.9	4.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health Current Asthma Diagnosed by a Medical Professional Rensselaer

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	66	10.0	572	90.0	2.6
Gender					
Male	22	8.3	223	91.7	3.7
Female	44	11.6	349	88.4	3.5
Age					
18-34	20	12.6	129	87.4	5.7
35-54	18	7.3	224	92.7	3.6
≥55	28	11.1	211	88.9	4.2

30 10.5

36

9.7

337

232 89.5 3.9

90.3 3.4

Education

≤High School
>High School

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Arthritis Diagnosed by a Medical Professional

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	239	33.8	396	66.2	4.1
Gender					
Male	85	30.6	160	69.4	6.2
Female	154	36.7	236	63.3	5.3
Age					
18-34	15	8.7	134	91.3	4.8
35-54	83	35.7	158	64.3	6.7
≥55	137	56.3	100	43.7	7.2
Education					
≤High School	116	41.3	143	58.7	6.8
>High School	122	28.8	251	71.2	5.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

Demographic Groups	n¹	Doctor Diagnosed %2	C.I. ³	n	Possible Arthritis (CJS Only) %	C.I.	n	No Arthritis or CJS %	C.I.
Total	239	33.9	4.1	77	12.8	3.3	316	53.2	4.5
Gender									
Male	85	30.8	6.2	29	12.7	5.6	130	56.4	7.1
Female	154	36.9	5.3	48	12.9	3.7	186	50.2	5.5
Age									
18-34	15	8.7	4.8	22	15.3	7.9	112	75.9	8.6
35-54	83	36.1	6.8	34	14.0	4.8	122	49.9	7.0
≥55	137	56.4	7.2	20	8.8	4.2	79	34.8	6.9
Education									
≤High School	116	41.4	6.8	33	15.4	6.5	109	43.3	7.0
>High School	122	29.1	5.1	43	11.1	3.5	206	59.9	5.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Ever Smoked 100 Cigarettes in Lifetime

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	333	51.0	305	49.0	4.5
Gender					
Male	139	53.8	107	46.2	7.1
Female	194	48.4	198	51.6	5.5
Age					
18-34	68	45.7	81	54.3	9.3
35-54	115	47.3	127	52.7	7.0
≥55	147	61.2	92	38.8	7.0
Education					
≤High School	169	62.9	93	37.1	6.7
>High School	162	43.0	211	57.0	5.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Current Smoker

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	151	24.3	486	75.7	4.0
Gender					
Male	52	23.2	193	76.8	6.6
Female	99	25.3	293	74.7	4.7
Age					
18-34	42	28.9	107	71.1	8.9
35-54	61	25.8	181	74.2	6.2
≥55	46	17.8	192	82.2	5.5
Education					
≤High School	85	32.4	176	67.6	7.0
>High School	65	18.9	308	81.1	4.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Status of Cigarette Smoking

Demographic Groups	n¹	Everyday Smoker %	C.I. ³	n	Someday Smoker %	C.I.	n	Former or Never Smoked	C.I.
Total	111	17.6	3.6	40	6.7	2.3	486	75.7	4.0
Gender									
Male	34	16.0	6.1	18	7.2	3.7	193	76.8	6.6
Female	77	19.1	4.2	22	6.3	2.8	293	74.7	4.7
Age									
18-34	27	18.0	8.1	15	10.9	5.6	107	71.1	8.9
35-54	49	21.0	5.8	12	4.8	2.8	181	74.2	6.2
≥55	35	13.3	4.7	11	4.5	3.3	192	82.2	5.5
Education									
≤High School	68	26.9	6.9	17	5.6	2.8	176	67.6	7.0
>High School	43	11.6	3.7	22	7.3	3.3	308	81.1	4.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Home Smoking Policies

Demographic Groups	n¹	Not Allowed	C.I. ³	n	Allowed Some Places %	C.I.	n	Allowed Anywhere	C.I.
Total	393	62.5	4.4	87	12.9	2.8	157	24.5	4.0
Gender									
Male	147	61.5	7.0	31	12.2	4.4	67	26.3	6.6
Female	246	63.5	5.3	56	13.6	3.6	90	22.9	4.7
Age									
18-34	101	67.0	9.0	23	13.3	5.5	26	19.8	8.5
35-54	161	67.3	6.6	39	15.3	4.8	41	17.5	5.5
≥55	126	52.1	7.2	24	9.5	4.3	88	38.4	7.1
Education									
≤High School	136	50.2	7.1	35	13.5	4.6	90	36.3	7.2
>High School	255	70.6	5.2	51	12.4	3.6	67	17.0	4.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Home Smoking

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	168	26.2	471	73.8	4.0
Gender					
Male	65	26.3	181	73.7	6.6
Female	103	26.2	290	73.8	4.8
Age					
18-34	37	26.2	113	73.8	8.8
35-54	66	26.6	176	73.4	6.1
≥55	63	25.8	176	74.2	6.3
Education					
≤High School	92	37.3	170	62.7	7.2
>High School	75	18.9	299	81.1	4.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Total	85	16.8	543	83.2	3.5
Gender					
Male	56	25.6	186	74.4	6.2
Female	29	8.6	357	91.4	3.3
Age					
18-34	43	32.5	105	67.5	8.5
35-54	33	16.0	205	84.0	5.3
≥55	9	2.9	226	97.1	2.0
Education					
≤High School	31	15.1	226	84.9	5.1
>High School	54	17.8	316	82.2	4.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Total	26	4.6	601	95.4	1.9
Gender					
Male	14	5.1	228	94.9	3.0
Female	12	4.1	373	95.9	2.5
Age					
18-34	10	8.0	138	92.0	5.1
35-54	6	2.9	230	97.1	2.4
≥55	10	3.4	226	96.6	2.3
Education					
≤High School	10	4.4	247	95.6	2.8
>High School	16	4.7	353	95.3	2.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health Overweight or Obese, BMI(Body Mass Index) 25 or More Rensselaer

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	354	59.1	243	40.9	4.5
Gender					
Male	171	68.6	73	31.4	6.6
Female	183	49.2	170	50.8	5.8
Age					
18-34	70	51.0	73	49.0	9.4
35-54	137	60.3	93	39.7	7.0
≥55	147	67.1	73	32.9	6.9
Education					
≤High School	150	61.8	91	38.2	7.1
>High School	204	57.5	152	42.5	5.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+) Rensselaer

Demographic Groups	n¹	Underweight\ Normal %2	C.I. ³	n	Overweight %	C.I.	n	Obese %	C.I.
Total	243	40.9	4.5	205	35.3	4.5	148	23.7	3.8
Gender									
Male	73	31.4	6.6	105	42.8	7.1	66	25.7	6.0
Female	170	51.0	5.8	100	27.4	5.1	82	21.6	4.6
Age									
18-34	73	49.3	9.5	41	31.1	9.3	28	19.6	7.3
35-54	93	39.7	7.0	79	37.7	7.1	58	22.6	5.8
≥55	73	32.9	6.9	85	37.2	7.1	62	29.9	6.9
Education									
≤High School	91	38.2	7.1	81	34.1	7.3	69	27.7	6.3
>High School	152	42.7	5.9	124	36.1	5.7	79	21.2	4.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Ever Had Mammography Screening, Women Aged 40 Years and Older *Rensselaer*

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Female	237	95.1	17	4.9	2.4
Age					
40-64	154	96.3	9	3.7	2.6
≥65	83	92.9	8	7.1	4.9
Education					
≤High School	103	93.6	9	6.4	4.3
>High School	133	96.2	8	3.8	2.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Female	204	82.6	49	17.4	4.9
Age					
40-64	136	86.4	27	13.6	5.2
≥65	68	74.9	22	25.1	10.0
Education					
≤High School	87	79.7	24	20.3	8.1
>High School	116	84.5	25	15.5	6.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	80	16.4	393	83.6	3.8
Gender					
Male	20	10.0	167	90.0	4.8
Female	60	22.9	226	77.1	5.6
Age					
18-34	47	29.7	100	70.3	8.3
35-54	29	10.3	205	89.7	3.9
55-64	4	2.8	88	97.2	2.8
Education					
≤High School	31	16.9	144	83.1	6.2
>High School	49	16.1	248	83.9	4.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health Sexually Active, Age 18-64 Years Old

Demographic		Yes		No	
Groups	n¹	% ²	n	%	C.I. ³
Total	359	82.4	89	17.6	4.4
Gender					
Male	142	80.7	34	19.3	7.3
Female	217	84.1	55	15.9	4.8
Age					
18-34	128	83.4	16	16.6	8.9
35-54	184	87.5	38	12.5	4.4
55-64	47	64.4	35	35.6	11.7
Education					
≤High School	127	76.2	39	23.8	8.8
>High School	231	85.9	50	14.1	4.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger Rensselaer

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Total	64	18.3	300	81.7	4.6
Gender					
Male	31	21.8	113	78.2	7.6
Female	33	14.8	187	85.2	5.2
Age					
18-34	37	28.8	93	71.2	8.8
35-54	26	13.9	162	86.1	5.8
55-64	1	1.4	45	98.6	2.8
Education					
≤High School	18	14.9	110	85.1	7.0
>High School	46	20.1	189	79.9	6.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

Demographic Groups	n¹	Use BC	C.I. ³	n	No BC at risk	C.I.	n	No BC	C.I.
Total	202	63.8		41	12.4	3.8	68	23.8	5.9
Gender									
Male	101	62.0	8.6	18	10.8	4.9	43	27.2	8.3
Female	101	66.8	8.4	23	14.9	6.1	25	18.3	7.2
Age									
18-34	95	63.2	9.6	21	13.4	5.8	26	23.3	9.3
35-59	107	64.4	8.0	20	11.4	5.1	42	24.2	7.3
Education									
≤High School	74	63.7	11.1	13	10.2	5.6	24	26.1	11.1
>High School	127	63.8	7.5	28	13.7	5.1	44	22.6	6.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health **Physician Diagnosed Heart Attack, Angina or Stroke**

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	50	6.2	576	93.8	1.8
Gender					
Male	23	6.3	219	93.7	2.7
Female	27	6.2	357	93.8	2.4
Age					
18-64	19	3.2	456	96.8	1.5
≥65	29	19.3	116	80.7	6.9
Education					
≤High School	29	8.9	229	91.1	3.5
>High School	21	4.5	344	95.5	2.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina) Rensselaer

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Total	50	6.2	576	93.8	1.8
Gender					
Male	23	6.3	219	93.7	2.7
Female	27	6.2	357	93.8	2.4
Age					
18-64	19	3.2	456	96.8	1.5
≥65	29	19.3	116	80.7	6.9
Education					
≤High School	29	8.9	229	91.1	3.5
>High School	21	4.5	344	95.5	2.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older *Rensselaer*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Male	97	62.5	52	37.5	8.8
Age					
40-64	52	56.5	42	43.5	10.9
≥65	45	78.9	10	21.1	12.6
Education					
≤High School	46	66.8	18	33.2	13.0
>High School	51	59.6	34	40.4	11.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Male	86	56.1	61	43.9	9.0
Age					
40-64	44	48.7	49	51.3	11.2
≥65	42	76.3	12	23.7	13.0
Education					
≤High School	40	57.7	23	42.3	13.5
>High School	46	54.9	38	45.1	12.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health Ever Had a Digital Rectal Exam, Men Aged 40 and Older Rensselaer

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Male	126	82.6	24	17.4	6.7
Age					
40-64	76	79.6	20	20.4	8.5
≥65	50	91.3	4	8.7	9.4
Education					
≤High School	49	70.6	16	29.4	12.5
>High School	77	91.0	8	9.0	6.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Ever Used a Blood Stool Test at Home, Aged 50 and Older *Rensselaer*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	133	47.4	160	52.6	6.6
Gender					
Male	42	42.4	63	57.6	10.8
Female	91	51.1	97	48.9	8.0
Age					
50-64	67	45.4	83	54.6	9.3
≥65	66	49.9	77	50.1	9.0
Education					
≤High School	55	42.5	83	57.5	9.2
>High School	78	51.7	76	48.3	9.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older *Rensselaer*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	56	22.6	231	77.4	6.0
Gender					
Male	19	22.0	84	78.0	9.7
Female	37	23.0	147	77.0	7.5
Age					
50-64	25	21.0	123	79.0	8.5
≥65	31	24.6	108	75.4	8.2
Education					
≤High School	24	20.3	109	79.7	8.0
>High School	32	24.5	121	75.5	8.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older *Rensselaer*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	87	31.6	200	68.4	6.3
Gender					
Male	29	30.6	74	69.4	10.3
Female	58	32.4	126	67.6	7.9
Age					
50-64	39	27.3	109	72.7	8.8
≥65	48	37.3	91	62.7	8.9
Education					
≤High School	37	29.5	96	70.5	8.7
>High School	50	33.5	103	66.5	9.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older *Rensselaer*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	161	55.3	133	44.7	6.5
Gender					
Male	63	60.9	42	39.1	10.6
Female	98	51.1	91	48.9	8.0
Age					
50-64	78	52.8	72	47.2	9.3
≥65	83	58.5	61	41.5	8.8
Education					
≤High School	74	52.4	64	47.6	9.3
>High School	87	58.1	68	41.9	9.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older *Rensselaer*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	149	51.0	144	49.0	6.5
Gender					
Male	58	56.0	47	44.0	10.7
Female	91	47.2	97	52.8	8.0
Age					
50-64	73	49.4	77	50.6	9.3
≥65	76	53.0	67	47.0	8.9
Education					
≤High School	70	49.0	67	51.0	9.3
>High School	79	52.9	76	47.1	9.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	168	57.7	127	42.3	6.4
Gender					
Male	61	58.4	44	41.6	10.7
Female	107	57.2	83	42.8	7.8
Age					
50-64	81	56.1	69	43.9	9.2
≥65	87	59.8	58	40.2	8.7
Education					
≤High School	79	55.2	60	44.8	9.2
>High School	89	60.2	66	39.8	8.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval