#### Expanded BRFSS, 2003

#### Saratoga Locality Tables

#### TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days	
Core 2: Health Care Access	
Core 3: Exercise	8
Core 3: Exercise	9
Core 5: Asthma	10
Core 6: Arthritis	
Core 7: Tobacco Use	
Core 8: Tobacco ETS	17
Core 9: Alcohol Consumption	19
Core 10: Weight Status	2
Core 11: Mammography	23
Core 12: Sexual Behavior	25
Core 13: Family Planning	28
Core 14: Cardiovascular Disease	29
Core 15: Prostate Cancer Screening	3
Core 16: Colorectal Cancer Screening	37

New York State Department Of Health

#### **Self-rated Health Status**

Demographic Groups	n¹	Fair or Poor	n	Good to Excellent	C.I. <sup>3</sup>
Total	67	8.5	577	91.5	2.2
Gender					
Male	21	8.5	201	91.5	3.7
Female	46	8.5	376	91.5	2.6
Age					
18-34	7	3.5	136	96.5	2.7
35-54	17	5.9	252	94.1	3.1
≥55	42	17.2	177	82.8	5.5
Education					
≤High School	37	13.9	195	86.1	4.8
>High School	30	5.5	380	94.5	2.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Reported Number of Physical Distress Days in the Past Month** Saratoga

Demographic Groups	n¹	≥14 days	n	< 14 days %	C.I. <sup>3</sup>
Total	62	8.5	568	91.5	2.5
Gender					
Male	17	8.7	202	91.3	4.2
Female	45	8.3	366	91.7	2.7
Age					
18-34	10	6.5	130	93.5	4.6
35-54	23	8.0	241	92.0	3.8
≥55	29	12.0	183	88.0	4.8
Education					
≤High School	31	14.1	189	85.9	5.5
>High School	31	5.5	377	94.5	2.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### **Reported Number of Mental Distress Days in the Past Month** Saratoga

Demographic Groups	n¹	≥14 days %2	n	< 14 days %	C.I. <sup>3</sup>
Total	55	7.4	587	92.6	2.3
Gender					
Male	16	6.5	206	93.5	3.6
Female	39	8.3	381	91.7	2.8
Age					
18-34	16	9.4	126	90.6	5.3
35-54	28	8.1	240	91.9	3.3
≥55	10	4.0	208	96.0	2.8
Education					
≤High School	27	11.1	205	88.9	4.8
>High School	28	5.4	380	94.6	2.2

<sup>&</sup>lt;sup>1</sup> Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup> Weighted Percent<sup>3</sup> 95% Confidence Interval

#### Expanded BRFSS, 2003 - New York State Department of Health **Health Insurance Coverage, 18-64 Years of Age** Saratoga

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	450	88.6	51	11.4	3.4
Gender					
Male	165	87.1	22	12.9	5.7
Female	285	90.2	29	9.8	3.6
Age					
18-34	120	82.8	23	17.2	7.3
35-54	245	90.3	24	9.7	4.2
55-64	85	96.5	4	3.5	3.5
Education					
≤High School	143	82.3	25	17.7	7.0
>High School	306	91.9	26	8.1	3.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### Expanded BRFSS, 2003 - New York State Department of Health Health Insurance Coverage, Aged 18 and Older Saratoga

Demographic		Yes		No	
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	590	90.2	56	9.8	2.8
Gender					
Male	200	89.0	22	11.0	4.9
Female	390	91.4	34	8.6	3.0
Age					
18-34	120	82.8	23	17.2	7.3
35-54	245	90.3	24	9.7	4.2
≥55	212	97.5	8	2.5	1.9
Education					
≤High School	205	85.4	28	14.6	5.7
>High School	383	92.9	28	7.1	3.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### Reported Having a Routine Physical Exam in Past Two Years Saratoga

Demographic		Yes		No	
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	582	89.9	58	10.1	2.9
Gender					
Male	193	86.3	28	13.7	5.2
Female	389	93.4	30	6.6	2.5
Age					
18-34	125	85.5	18	14.5	6.8
35-54	238	88.3	28	11.7	4.7
≥55	206	96.2	12	3.8	2.4
Education					
≤High School	205	85.9	26	14.1	5.9
>High School	375	92.2	32	7.8	3.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### Time in Past 12 Months when Needed Medical Care but Could Not Get It Saratoga

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	21	3.1	625	96.9	1.6
Gender					
Male	5	3.0	217	97.0	2.7
Female	16	3.2	408	96.8	1.7
Age					
18-34	7	4.8	136	95.2	3.8
35-54	9	2.9	260	97.1	2.4
≥55	5	2.0	215	98.0	2.1
Education					
≤High School	9	4.5	224	95.5	3.4
>High School	12	2.4	399	97.6	1.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

## Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days *Saratoga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	509	79.8	136	20.2	3.5
Gender					
Male	178	80.4	44	19.6	5.7
Female	331	79.2	92	20.8	4.3
Age					
18-34	115	80.8	28	19.2	7.2
35-54	224	81.1	45	18.9	5.5
≥55	158	76.5	61	23.5	6.2
Education					
≤High School	158	67.4	75	32.6	7.1
>High School	349	86.7	61	13.3	3.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### **Physician Diagnosed Diabetes**

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	44	5.5	602	94.5	1.9
Gender					
Male	14	5.3	208	94.7	2.9
Female	30	5.7	394	94.3	2.3
Age					
18-34	1	0.4	142	99.6	0.8
35-54	11	3.0	258	97.0	2.2
≥55	28	12.0	192	88.0	4.8
Education					
≤High School	19	6.2	214	93.8	3.1
>High School	25	5.1	386	94.9	2.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### Expanded BRFSS, 2003 - New York State Department of Health **Lifetime Asthma Diagnosed by a Medical Professional** *Saratoga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	86	12.2	558	87.8	2.9
Gender					
Male	21	9.9	201	90.1	4.4
Female	65	14.4	357	85.6	3.6
Age					
18-34	23	15.5	120	84.5	6.6
35-54	37	11.8	232	88.2	4.2
≥55	25	9.7	193	90.3	4.2
Education					
≤High School	34	14.0	198	86.0	5.3
>High School	52	11.3	358	88.7	3.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

## Expanded BRFSS, 2003 - New York State Department of Health Current Asthma Diagnosed by a Medical Professional Saratoga

Demographic		Yes		No	
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	58	7.8	584	92.2	2.4
Gender					
Male	13	6.5	209	93.5	3.8
Female	45	9.1	375	90.9	2.8
Age					
18-34	16	11.0	127	89.0	5.9
35-54	24	6.7	243	93.3	3.0
≥55	18	6.8	200	93.2	3.6
Education					
≤High School	25	10.1	207	89.9	4.8
>High School	33	6.6	375	93.4	2.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Expanded BRFSS, 2003 - New York State Department of Health **Arthritis Diagnosed by a Medical Professional**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	211	28.6	433	71.4	3.9
Gender					
Male	63	26.7	158	73.3	6.3
Female	148	30.3	275	69.7	4.7
Age					
18-34	15	8.6	128	91.4	5.1
35-54	76	27.5	193	72.5	6.0
≥55	117	51.1	101	48.9	7.8
Education					
≤High School	86	32.6	147	67.4	7.0
>High School	125	26.5	284	73.5	4.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

Demographic Groups	n¹	Doctor Diagnosed %2	C.I. <sup>3</sup>	n	Possible Arthritis (CJS Only) %	C.I.	n	No Arthritis or CJS %	C.I.
Total	211	28.7	3.9	95	17.0	3.6	336	54.3	4.5
Gender									
Male	63	26.9	6.3	38	19.6	6.2	119	53.4	7.3
Female	148	30.4	4.7	57	14.5	3.7	217	55.2	5.2
Age									
18-34	15	8.8	5.1	25	19.5	7.7	101	71.7	8.6
35-54	76	27.5	6.0	48	19.5	5.4	145	52.9	6.7
≥55	117	51.1	7.8	19	10.9	6.0	82	38.0	7.4
Education									
≤High School	86	32.9	7.0	30	13.7	5.4	116	53.4	7.6
>High School	125	26.5	4.7	64	18.4	4.6	219	55.1	5.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health **Ever Smoked 100 Cigarettes in Lifetime** 

Demographic		Yes		No	
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	324	49.8	321	50.2	4.5
Gender					
Male	109	50.6	113	49.4	7.3
Female	215	49.1	208	50.9	5.3
Age					
18-34	60	46.0	83	54.0	9.4
35-54	131	47.1	138	52.9	6.7
≥55	125	57.7	94	42.3	7.6
Education					
≤High School	143	59.2	90	40.8	7.6
>High School	180	44.2	230	55.8	5.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Current Smoker**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	126	20.4	518	79.6	3.8
Gender					
Male	37	19.5	185	80.5	6.1
Female	89	21.3	333	78.7	4.5
Age					
18-34	40	30.8	103	69.2	9.1
35-54	50	19.6	218	80.4	5.3
≥55	34	11.2	185	88.8	4.2
Education					
≤High School	71	30.3	161	69.7	7.1
>High School	55	14.9	355	85.1	4.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### **Status of Cigarette Smoking**

Demographic Groups	n¹	Everyday Smoker %	C.I. <sup>3</sup>	n	Someday Smoker	C.I.	n	Former or Never Smoked %	C.I.
Total	98	14.5	3.0	28	5.9	2.7	518	79.6	3.8
Gender									
Male	27	12.4	4.7	10	7.1	4.5	185	80.5	6.1
Female	71	16.5	3.9	18	4.8	2.8	333	78.7	4.5
Age									
18-34	30	18.2	6.7	10	12.6	7.9	103	69.2	9.1
35-54	41	16.5	5.0	9	3.2	2.2	218	80.4	5.3
≥55	25	7.8	3.4	9	3.4	2.5	185	88.8	4.2
Education									
≤High School	57	23.2	6.2	14	7.1	4.8	161	69.7	7.1
>High School	41	9.7	3.2	14	5.2	3.2	355	85.1	4.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### **Home Smoking Policies**

Demographic Groups	n¹	Not Allowed %2	C.I. <sup>3</sup>	n	Allowed Some Places %	C.I.	n	Allowed Anywhere	C.I.
Total	454	71.6	4.0	62	10.2	2.7	128	18.2	3.4
Gender									
Male	164	73.6	6.6	19	9.4	4.3	38	17.0	5.7
Female	290	69.8	4.8	43	10.9	3.3	90	19.3	4.1
Age									
18-34	107	74.1	8.2	17	12.4	6.1	19	13.5	6.4
35-54	195	73.4	5.9	28	11.6	4.4	45	15.0	4.6
≥55	140	66.6	7.4	17	6.6	3.5	62	26.8	7.1
Education									
≤High School	133	57.7	7.6	30	13.9	5.4	69	28.4	6.9
>High School	319	79.3	4.4	32	8.2	3.0	59	12.5	3.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	134	21.6	512	78.4	3.8
Gender					
Male	38	19.6	184	80.4	6.1
Female	96	23.6	328	76.4	4.7
Age					
18-34	35	26.5	108	73.5	8.4
35-54	56	21.0	213	79.0	5.5
≥55	41	16.7	179	83.3	6.2
Education					
≤High School	75	33.9	158	66.1	7.3
>High School	59	14.7	352	85.3	4.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	77	16.3	559	83.7	3.7
Gender					
Male	54	25.9	162	74.1	6.6
Female	23	7.3	397	92.7	3.4
Age					
18-34	33	30.4	107	69.6	9.3
35-54	34	15.3	231	84.7	5.1
≥55	9	3.2	209	96.8	2.2
Education					
≤High School	31	18.7	198	81.3	6.7
>High School	46	15.1	359	84.9	4.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	29	4.7	610	95.3	2.0
Gender					
Male	12	4.9	205	95.1	3.1
Female	17	4.5	405	95.5	2.7
Age					
18-34	9	7.7	133	92.3	5.7
35-54	9	2.8	255	97.2	1.9
≥55	10	4.5	209	95.5	3.2
Education					
≤High School	11	5.5	219	94.5	3.8
>High School	18	4.3	389	95.7	2.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

## Expanded BRFSS, 2003 - New York State Department of Health Overweight or Obese, BMI(Body Mass Index) 25 or More Saratoga

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	338	58.6	263	41.4	4.5
Gender					
Male	157	71.6	60	28.4	6.8
Female	181	45.2	203	54.8	5.5
Age					
18-34	63	47.9	70	52.1	9.7
35-54	135	57.8	117	42.2	6.7
≥55	136	70.7	70	29.3	7.0
Education					
≤High School	136	62.7	83	37.3	7.7
>High School	202	56.3	179	43.7	5.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+) Saratoga

Demographic Groups	n¹	Underweight\ Normal %2	C.I. <sup>3</sup>	n	Overweight %	C.I.	n	Obese %	C.I.
Total	262	41.4	4.6	221	37.6	4.5	116	20.9	3.9
Gender									
Male	60	28.4	6.8	105	46.4	7.4	52	25.1	6.5
Female	202	54.8	5.5	116	28.6	4.9	64	16.6	4.1
Age									
18-34	69	51.9	9.8	39	28.8	8.7	24	19.3	8.1
35-54	117	42.4	6.7	78	32.9	6.7	56	24.7	6.2
≥55	70	29.3	7.0	100	52.4	8.0	36	18.3	6.2
Education									
≤High School	82	37.3	7.7	80	34.1	7.3	55	28.6	7.4
>High School	179	43.7	5.6	141	39.7	5.7	61	16.5	4.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Ever Had Mammography Screening, Women Aged 40 Years and Older** *Saratoga*

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Female	251	93.5	18	6.5	3.3
Age					
40-64	160	93.5	11	6.5	4.1
≥65	91	93.4	7	6.6	5.3
Education					
≤High School	94	90.6	8	9.4	6.8
>High School	157	95.1	10	4.9	3.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Female	214	80.3	53	19.7	5.4
Age					
40-64	137	80.1	33	19.9	6.8
≥65	77	80.8	20	19.2	8.4
Education					
≤High School	78	75.1	22	24.9	10.1
>High School	136	83.1	31	16.9	6.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	81	16.5	402	83.5	4.0
Gender					
Male	22	12.6	157	87.4	5.8
Female	59	20.5	245	79.5	5.3
Age					
18-34	49	33.7	89	66.3	9.1
35-54	28	8.3	232	91.7	3.5
55-64	4	3.6	81	96.4	3.8
Education					
≤High School	37	24.6	124	75.4	8.3
>High School	44	12.3	278	87.7	4.1

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health **Sexually Active, Age 18-64 Years Old** *Saratoga* 

Demographic		Yes		No	G I 2
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	366	85.3	85	14.7	3.5
Gender					
Male	132	83.9	32	16.1	6.0
Female	234	86.8	53	13.2	3.7
Age					
18-34	112	87.5	18	12.5	6.5
35-54	211	89.3	33	10.7	4.1
55-64	43	66.8	34	33.2	11.5
Education					
≤High School	115	81.1	36	18.9	7.0
>High School	251	87.6	49	12.4	3.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger** Saratoga

Demographic		Yes		No	~ - 4
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	69	18.3	305	81.7	4.6
Gender					
Male	29	21.2	106	78.8	7.9
Female	40	15.5	199	84.5	4.8
Age					
18-34	33	28.8	80	71.2	9.8
35-54	33	14.0	182	86.0	5.1
55-64	3	5.5	43	94.5	6.5
Education					
≤High School	28	26.6	92	73.4	9.8
>High School	41	14.1	213	85.9	4.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

Demographic Groups	n¹	Use BC	C.I. <sup>3</sup>	n	No BC at risk	C.I.	n	No BC	C.I.
Total	200	61.5	6.0	48	15.7	4.4	84	22.9	5.1
Gender									
Male	91	58.3	8.4	20	14.3	6.1	51	27.3	7.5
Female	109	66.3	7.6	28	17.7	6.2	33	15.9	5.7
Age									
18-34	73	57.0	9.6	26	20.0	7.7	34	22.9	8.1
35-59	127	65.0	7.5	22	12.2	5.1	50	22.8	6.6
Education									
≤High School	60	56.9	10.6	13	13.2	7.5	37	29.8	9.5
>High School	140	64.0	7.2	35	17.0	5.5	47	19.0	6.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health **Physician Diagnosed Heart Attack, Angina or Stroke** *Saratoga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	52	7.5	583	92.5	2.2
Gender					
Male	23	9.7	194	90.3	4.0
Female	29	5.4	389	94.6	2.1
Age					
18-64	17	3.3	475	96.7	1.7
≥65	34	29.8	97	70.2	9.4
Education					
≤High School	33	12.5	195	87.5	4.6
>High School	19	4.7	387	95.3	2.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina) Saratoga

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	46	6.8	589	93.2	2.2
Gender					
Male	20	8.7	197	91.3	3.9
Female	26	5.0	392	95.0	2.1
Age					
18-64	16	3.2	476	96.8	1.7
≥65	29	26.0	102	74.0	9.1
Education					
≤High School	29	11.2	199	88.8	4.4
>High School	17	4.3	389	95.7	2.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### **Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older** *Saratoga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	81	61.4	46	38.6	9.5
Age					
40-64	54	54.3	42	45.7	11.1
≥65	27	83.6	4	16.4	14.5
Education					
≤High School	26	50.2	21	49.8	15.8
>High School	55	67.8	25	32.2	11.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	70	54.0	56	46.0	9.6
Age					
40-64	45	45.6	50	54.4	11.1
≥65	25	80.0	6	20.0	15.1
Education					
≤High School	22	43.5	25	56.5	15.4
>High School	48	60.2	31	39.8	12.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

## Expanded BRFSS, 2003 - New York State Department of Health Ever Had a Digital Rectal Exam, Men Aged 40 and Older Saratoga

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	107	81.5	25	18.5	7.2
Age					
40-64	80	79.8	20	20.2	8.7
≥65	27	86.9	5	13.1	11.7
Education					
≤High School	36	71.8	13	28.2	14.4
>High School	71	87.2	12	12.8	7.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Used a Blood Stool Test at Home, Aged 50 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	150	54.4	127	45.6	6.8
Gender					
Male	53	59.2	30	40.8	11.9
Female	97	50.5	97	49.5	7.8
Age					
50-64	84	56.5	65	43.5	9.2
≥65	66	51.3	62	48.7	10.0
Education					
≤High School	49	45.2	64	54.8	10.6
>High School	101	60.7	62	39.3	8.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older *Saratoga*

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	77	31.2	196	68.8	6.5
Gender					
Male	34	39.4	49	60.6	11.4
Female	43	24.5	147	75.5	7.0
Age					
50-64	42	30.7	106	69.3	8.6
≥65	35	32.1	90	67.9	9.8
Education					
≤High School	19	20.6	90	79.4	9.0
>High School	58	37.8	105	62.2	8.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older *Saratoga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	102	40.6	171	59.4	6.8
Gender					
Male	42	48.2	41	51.8	11.8
Female	60	34.3	130	65.7	7.7
Age					
50-64	58	41.6	90	58.4	9.1
≥65	44	39.2	81	60.8	10.1
Education					
≤High School	30	30.8	79	69.2	10.1
>High School	72	46.9	91	53.1	8.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older *Saratoga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	155	57.4	125	42.6	6.7
Gender					
Male	54	63.4	31	36.6	11.5
Female	101	52.5	94	47.5	7.8
Age					
50-64	80	54.1	71	45.9	9.2
≥65	75	62.3	54	37.7	9.5
Education					
≤High School	65	63.4	49	36.6	9.7
>High School	90	54.9	75	45.1	8.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older *Saratoga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	143	53.9	137	46.1	6.8
Gender					
Male	51	59.7	34	40.3	11.6
Female	92	49.2	103	50.8	7.8
Age					
50-64	73	50.4	78	49.6	9.2
≥65	70	59.1	59	40.9	9.7
Education					
≤High School	61	59.0	53	41.0	10.1
>High School	82	51.9	83	48.1	8.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	168	62.9	113	37.1	6.6
Gender					
Male	61	68.9	24	31.1	11.3
Female	107	58.0	89	42.0	7.6
Age					
50-64	88	60.7	63	39.3	9.0
≥65	80	66.2	50	33.8	9.3
Education					
≤High School	67	64.1	48	35.9	9.8
>High School	101	63.3	64	36.7	8.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval