Expanded BRFSS, 2003

Clinton, Essex and Franklin Locality Tables

TABLE OF CONTENTS:

| Core 1: Health Status / Healthy Days | |
|--------------------------------------|---|
| Core 2: Health Care Access | |
| Core 3: Exercise | |
| Core 4: Diabetes | 9 |
| Core 5: Asthma | |
| Core 6: Arthritis | |
| Core 7: Tobacco Use | |
| Core 8: Tobacco ETS | |
| Core 9: Alcohol Consumption | |
| Core 10: Weight Status | |
| Core 11: Mammography | |
| Core 12: Sexual Behavior | |
| Core 13: Family Planning | |
| Core 14: Cardiovascular Disease | |
| Core 15: Prostate Cancer Screening | |
| Core 16: Colorectal Cancer Screening | |

New York State Department Of Health

| Demographic Groups | n1 | Fair or Poor | n | Good to Excellent % | C.I. ³ |
|-----------------------|----|--------------|-----|---------------------|-------------------|
| Total | 96 | 13.4 | 550 | 86.6 | 2.9 |
| Gender | | | | | |
| Male | 47 | 14.6 | 227 | 85.4 | 4.6 |
| Female | 49 | 12.1 | 323 | 87.9 | 3.6 |
| Age | | | | | |
| 18-34 | 8 | 6.1 | 141 | 93.9 | 5.0 |
| 35-54 | 28 | 10.6 | 231 | 89.4 | 4.1 |
| ≥55 | 59 | 25.3 | 173 | 74.7 | 6.2 |
| Education | | | | | |
| ≤High School | 55 | 15.5 | 249 | 84.5 | 4.3 |
| >High School | 41 | 11.5 | 301 | 88.5 | 4.1 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health **Reported Number of Physical Distress Days in the Past Month** Clinton, Essex, Franklin

| Demographic Groups | n ¹ | ≥14 days % | n | < 14 days % | C.I. ³ |
|-----------------------|----------------|---------------|-----|----------------|-------------------|
| Total | 70 | 9.1 | 562 | 90.9 | 2.5 |
| Gender | | | | | |
| Male | 26 | 8.3 | 242 | 91.7 | 3.8 |
| Female | 44 | 10.1 | 320 | 89.9 | 3.2 |
| Age | | | | | |
| 18-34 | 8 | 4.9 | 139 | 95.1 | 4.5 |
| 35-54 | 25 | 9.0 | 234 | 91.0 | 3.8 |
| ≥55 | 37 | 14.3 | 183 | 85.7 | 4.8 |
| Education | | | | | |
| ≤High School | 41 | 11.0 | 254 | 89.0 | 3.7 |
| >High School | 29 | 7.5 | 308 | 92.5 | 3.4 |

Expanded BRFSS, 2003 - New York State Department of Health **Reported Number of Mental Distress Days in the Past Month** Clinton, Essex, Franklin

| Demographic Groups | n1 | ≥14 days % | n | < 14 days % | C.I. ³ |
|-----------------------|----|---------------|-----|----------------|-------------------|
| Total | 65 | 9.8 | 573 | 90.2 | 2.7 |
| Gender | | | | | |
| Male | 28 | 9.7 | 246 | 90.3 | 4.2 |
| Female | 37 | 9.9 | 327 | 90.1 | 3.4 |
| Age | | | | | |
| 18-34 | 17 | 10.2 | 132 | 89.8 | 5.7 |
| 35-54 | 29 | 10.7 | 226 | 89.3 | 4.4 |
| ≥55 | 19 | 8.5 | 209 | 91.5 | 4.0 |
| Education | | | | | |
| ≤High School | 35 | 12.1 | 263 | 87.9 | 4.5 |
| >High School | 30 | 7.8 | 310 | 92.2 | 3.2 |

¹ Percents based on row denominators of less than 50 are unstable and should be used with caution

² Weighted Percent³ 95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health Health Insurance Coverage, 18-64 Years of Age Clinton, Essex, Franklin

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|-----|------------|----|---------|-------------------|
| Total | 414 | 82.1 | 75 | 17.9 | 4.2 |
| Gender | | | | | |
| Male | 176 | 77.7 | 45 | 22.3 | 6.5 |
| Female | 238 | 87.4 | 30 | 12.6 | 4.8 |
| Age | | | | | |
| 18-34 | 116 | 76.1 | 31 | 23.9 | 8.3 |
| 35-54 | 223 | 84.8 | 36 | 15.2 | 5.1 |
| 55-64 | 75 | 88.7 | 8 | 11.3 | 8.2 |
| Education | | | | | |
| ≤High School | 170 | 75.6 | 43 | 24.4 | 7.0 |
| >High School | 244 | 87.2 | 32 | 12.8 | 4.9 |

Expanded BRFSS, 2003 - New York State Department of Health Health Insurance Coverage, Aged 18 and Older Clinton, Essex, Franklin

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|-----|------------|----|---------|-------------------|
| Total | 564 | 84.6 | 82 | 15.4 | 3.5 |
| Gender | | | | | |
| Male | 228 | 80.7 | 46 | 19.3 | 5.7 |
| Female | 336 | 88.8 | 36 | 11.2 | 3.9 |
| Age | | | | | |
| 18-34 | 116 | 76.1 | 31 | 23.9 | 8.3 |
| 35-54 | 223 | 84.8 | 36 | 15.2 | 5.1 |
| ≥55 | 220 | 93.2 | 13 | 6.8 | 4.0 |
| Education | | | | | |
| ≤High School | 258 | 80.3 | 47 | 19.7 | 5.6 |
| >High School | 306 | 88.4 | 35 | 11.6 | 4.3 |

Expanded BRFSS, 2003 - New York State Department of Health **Reported Having a Routine Physical Exam in Past Two Years** *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes % | n | No % | C.I. ³ |
|-----------------------|-----|----------|----|---------|-------------------|
| Total | 546 | 81.1 | 99 | 18.9 | 3.9 |
| Gender | | | | | |
| Male | 209 | 72.8 | 65 | 27.2 | 6.5 |
| Female | 337 | 90.3 | 34 | 9.7 | 3.5 |
| Age | | | | | |
| 18-34 | 114 | 72.3 | 33 | 27.7 | 9.0 |
| 35-54 | 211 | 81.7 | 48 | 18.3 | 5.4 |
| ≥55 | 215 | 89.7 | 17 | 10.3 | 4.9 |
| Education | | | | | |
| ≤High School | 254 | 80.4 | 51 | 19.6 | 5.6 |
| >High School | 292 | 81.8 | 48 | 18.2 | 5.4 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Time in Past 12 Months when Needed Medical Care but Could Not Get It *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes % | n | No % | C.I. ³ |
|-----------------------|----|----------|-----|---------|-------------------|
| Total | 32 | 4.2 | 618 | 95.8 | 1.6 |
| Gender | | | | | |
| Male | 13 | 4.3 | 264 | 95.7 | 2.5 |
| Female | 19 | 4.1 | 354 | 95.9 | 2.0 |
| Age | | | | | |
| 18-34 | 8 | 3.9 | 141 | 96.1 | 3.0 |
| 35-54 | 18 | 5.4 | 243 | 94.6 | 2.7 |
| ≥55 | 6 | 3.2 | 227 | 96.8 | 2.9 |
| Education | | | | | |
| ≤High School | 16 | 4.3 | 291 | 95.7 | 2.3 |
| >High School | 16 | 4.2 | 327 | 95.8 | 2.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health

Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes % | n | No % | C.I. ³ |
|-----------------------|-----|----------|-----|---------|-------------------|
| Total | 479 | 73.6 | 171 | 26.4 | 4.0 |
| Gender | | | | | |
| Male | 201 | 71.1 | 76 | 28.9 | 6.4 |
| Female | 278 | 76.4 | 95 | 23.6 | 4.6 |
| Age | | | | | |
| 18-34 | 123 | 78.8 | 26 | 21.2 | 8.4 |
| 35-54 | 202 | 77.9 | 59 | 22.1 | 5.4 |
| ≥55 | 149 | 62.5 | 84 | 37.5 | 7.1 |
| Education | | | | | |
| ≤High School | 190 | 60.6 | 117 | 39.4 | 6.5 |
| >High School | 289 | 85.3 | 54 | 14.7 | 4.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health **Physician Diagnosed Diabetes** Clinton, Essex, Franklin

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|----|------------|-----|---------|-------------------|
| Total | 53 | 6.5 | 596 | 93.5 | 1.9 |
| Gender | | | | | |
| Male | 23 | 5.8 | 254 | 94.2 | 2.5 |
| Female | 30 | 7.2 | 342 | 92.8 | 2.8 |
| Age | | | | | |
| 18-34 | 2 | 0.8 | 147 | 99.2 | 1.1 |
| 35-54 | 15 | 6.1 | 246 | 93.9 | 3.2 |
| ≥55 | 36 | 13.4 | 196 | 86.6 | 4.5 |
| Education | | | | | |
| ≤High School | 30 | 7.7 | 277 | 92.3 | 2.9 |
| >High School | 23 | 5.4 | 319 | 94.6 | 2.4 |

Expanded BRFSS, 2003 - New York State Department of Health Lifetime Asthma Diagnosed by a Medical Professional *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|----|------------|-----|---------|-------------------|
| Total | 83 | 11.3 | 566 | 88.7 | 2.7 |
| Gender | | | | | |
| Male | 23 | 7.0 | 253 | 93.0 | 3.2 |
| Female | 60 | 16.1 | 313 | 83.9 | 4.3 |
| Age | | | | | |
| 18-34 | 20 | 10.4 | 128 | 89.6 | 5.1 |
| 35-54 | 28 | 10.6 | 233 | 89.4 | 4.1 |
| ≥55 | 34 | 13.4 | 199 | 86.6 | 4.9 |
| Education | | | | | |
| ≤High School | 43 | 12.9 | 263 | 87.1 | 4.2 |
| >High School | 40 | 9.9 | 303 | 90.1 | 3.4 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health **Current Asthma Diagnosed by a Medical Professional** *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|----|------------|-----|---------|-------------------|
| Total | 59 | 7.4 | 588 | 92.6 | 2.1 |
| Gender | | | | | |
| Male | 16 | 4.2 | 259 | 95.8 | 2.3 |
| Female | 43 | 11.0 | 329 | 89.0 | 3.5 |
| Age | | | | | |
| 18-34 | 11 | 4.7 | 137 | 95.3 | 3.0 |
| 35-54 | 21 | 7.6 | 239 | 92.4 | 3.5 |
| ≥55 | 26 | 10.1 | 206 | 89.9 | 4.4 |
| Education | | | | | |
| ≤High School | 34 | 8.9 | 272 | 91.1 | 3.2 |
| >High School | 25 | 6.1 | 316 | 93.9 | 2.7 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health Arthritis Diagnosed by a Medical Professional *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes % | n | No % | C.I. ³ |
|-----------------------|-----|----------|-----|---------|-------------------|
| Total | 197 | 25.9 | 452 | 74.1 | 3.7 |
| Gender | | | | | |
| Male | 68 | 21.0 | 209 | 79.0 | 5.2 |
| Female | 129 | 31.3 | 243 | 68.7 | 5.1 |
| Age | | | | | |
| 18-34 | 15 | 8.0 | 134 | 92.0 | 4.4 |
| 35-54 | 54 | 20.2 | 206 | 79.8 | 5.3 |
| ≥55 | 128 | 53.5 | 105 | 46.5 | 7.2 |
| Education | | | | | |
| ≤High School | 96 | 27.2 | 211 | 72.8 | 5.5 |
| >High School | 101 | 24.7 | 241 | 75.3 | 4.9 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

Clinton, Essex, Franklin

| Demographic Groups | n1 | Doctor Diagnosed % | C.I. ³ | n | Possible Arthritis (CJS Only) % | C.I. | n | No Arthritis or CJS % | C.I. |
|-----------------------|-----|--------------------------|-------------------|-----|---------------------------------------|------|-----|-----------------------------|------|
| Total | 197 | 25.9 | 3.7 | 109 | 16.8 | 3.5 | 341 | 57.2 | 4.4 |
| Gender | | | | | | | | | |
| Male | 68 | 21.0 | 5.2 | 57 | 19.6 | 5.5 | 152 | 59.4 | 6.7 |
| Female | 129 | 31.5 | 5.1 | 52 | 13.8 | 3.9 | 189 | 54.8 | 5.6 |
| Age | | | | | | | | | |
| 18-34 | 15 | 8.0 | 4.4 | 24 | 15.1 | 7.2 | 110 | 76.8 | 8.0 |
| 35-54 | 54 | 20.3 | 5.3 | 46 | 17.4 | 5.2 | 159 | 62.3 | 6.5 |
| ≥55 | 128 | 53.8 | 7.3 | 36 | 16.7 | 5.4 | 68 | 29.6 | 6.7 |
| Education | | | | | | | | | |
| ≤High School | 96 | 27.2 | 5.5 | 52 | 17.3 | 5.2 | 159 | 55.5 | 6.4 |
| >High School | 101 | 24.8 | 4.9 | 57 | 16.4 | 4.6 | 182 | 58.8 | 6.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health Ever Smoked 100 Cigarettes in Lifetime *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes % | n | No % | C.I. ³ |
|-----------------------|-----|----------|-----|---------|-------------------|
| Total | 358 | 54.7 | 290 | 45.3 | 4.5 |
| Gender | | | | | |
| Male | 178 | 61.0 | 98 | 39.0 | 7.0 |
| Female | 180 | 47.6 | 192 | 52.4 | 5.6 |
| Age | | | | | |
| 18-34 | 71 | 46.8 | 78 | 53.2 | 9.5 |
| 35-54 | 148 | 58.9 | 112 | 41.1 | 6.6 |
| ≥55 | 136 | 58.5 | 96 | 41.5 | 7.1 |
| Education | | | | | |
| ≤High School | 183 | 60.1 | 123 | 39.9 | 6.4 |
| >High School | 175 | 49.9 | 167 | 50.1 | 6.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health **Current Smoker** Clinton, Essex, Franklin

| Demographic Groups | n ¹ | Yes % | n | No % | C.I. ³ |
|-----------------------|----------------|----------|-----|---------|-------------------|
| Total | 151 | 25.3 | 497 | 74.7 | 4.0 |
| Gender | | | | | |
| Male | 78 | 29.8 | 198 | 70.2 | 6.3 |
| Female | 73 | 20.2 | 299 | 79.8 | 4.6 |
| Age | | | | | |
| 18-34 | 45 | 28.6 | 104 | 71.4 | 8.6 |
| 35-54 | 81 | 33.8 | 179 | 66.2 | 6.4 |
| ≥55 | 25 | 11.3 | 207 | 88.7 | 4.7 |
| Education | | | | | |
| ≤High School | 88 | 31.8 | 218 | 68.2 | 6.2 |
| >High School | 63 | 19.4 | 279 | 80.6 | 5.0 |

Expanded BRFSS, 2003 - New York State Department of Health Status of Cigarette Smoking *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Everyday Smoker % | C.I. ³ | n | Someday Smoker % | C.I. | n | Former or Never Smoked % | C.I. |
|-----------------------|-----|----------------------|-------------------|----|---------------------|------|-----|--------------------------------|------|
| Total | 122 | 20.0 | 3.7 | 29 | 5.2 | 2.1 | 497 | 74.7 | 4.0 |
| Gender | | | | | | | | | |
| Male | 58 | 21.9 | 5.8 | 20 | 7.9 | 3.6 | 198 | 70.2 | 6.3 |
| Female | 64 | 17.9 | 4.4 | 9 | 2.3 | 1.6 | 299 | 79.8 | 4.6 |
| Age | | | | | | | | | |
| 18-34 | 38 | 23.8 | 8.1 | 7 | 4.7 | 4.0 | 104 | 71.4 | 8.6 |
| 35-54 | 62 | 24.5 | 5.8 | 19 | 9.2 | 4.2 | 179 | 66.2 | 6.4 |
| ≥55 | 22 | 10.5 | 4.7 | 3 | 0.8 | 0.9 | 207 | 88.7 | 4.7 |
| Education | | | | | | | | | |
| ≤High School | 74 | 26.5 | 5.9 | 14 | 5.3 | 3.1 | 218 | 68.2 | 6.2 |
| >High School | 48 | 14.2 | 4.5 | 15 | 5.2 | 2.8 | 279 | 80.6 | 5.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health Home Smoking Policies *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Not Allowed | C.I. ³ | n | Allowed Some Places % | C.I. | n | Allowed Anywhere % | C.I. |
|-----------------------|-----|-------------|-------------------|----|-----------------------------|------|-----|--------------------|------|
| Total | 398 | 62.7 | 4.4 | 94 | 13.7 | 3.1 | 156 | 23.6 | 3.8 |
| Gender | | | | | | | | | |
| Male | 164 | 62.0 | 6.7 | 40 | 13.8 | 4.8 | 71 | 24.2 | 5.8 |
| Female | 234 | 63.4 | 5.4 | 54 | 13.7 | 3.8 | 85 | 22.9 | 4.8 |
| Age | | | | | | | | | |
| 18-34 | 97 | 65.1 | 9.1 | 25 | 16.5 | 7.2 | 27 | 18.4 | 7.3 |
| 35-54 | 160 | 62.6 | 6.5 | 40 | 13.6 | 4.3 | 59 | 23.8 | 5.9 |
| ≥55 | 137 | 60.6 | 7.0 | 27 | 10.8 | 4.2 | 69 | 28.6 | 6.4 |
| Education | | | | | | | | | |
| ≤High School | 165 | 54.7 | 6.5 | 50 | 15.3 | 4.5 | 91 | 30.0 | 5.9 |
| >High School | 233 | 69.9 | 5.8 | 44 | 12.3 | 4.3 | 65 | 17.8 | 4.8 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health **Home Smoking** Clinton, Essex, Franklin

| Demographic Groups | n1 | Yes % | n | No % | C.I. ³ |
|-----------------------|-----|----------|-----|---------|-------------------|
| Total | 160 | 25.6 | 487 | 74.4 | 4.1 |
| Gender | | | | | |
| Male | 71 | 26.9 | 204 | 73.1 | 6.3 |
| Female | 89 | 24.2 | 283 | 75.8 | 4.9 |
| Age | | | | | |
| 18-34 | 41 | 29.0 | 108 | 71.0 | 8.9 |
| 35-54 | 72 | 27.1 | 186 | 72.9 | 5.9 |
| ≥55 | 45 | 19.5 | 188 | 80.5 | 5.8 |
| Education | | | | | |
| ≤High School | 92 | 32.2 | 212 | 67.8 | 6.2 |
| >High School | 68 | 19.8 | 275 | 80.2 | 5.2 |

Expanded BRFSS, 2003 - New York State Department of Health

At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink **Occasion**)

Clinton, Essex, Franklin

| Demographic Groups | n1 | Yes | n | No % | C.I. ³ |
|-----------------------|-----|------|-----|---------|-------------------|
| Total | 100 | 20.4 | 534 | 79.6 | 4.0 |
| Gender | 100 | 20.4 | 554 | 77.0 | т.0 |
| Male | 69 | 29.4 | 200 | 70.6 | 6.6 |
| Female | 31 | 10.4 | 334 | 89.6 | 3.9 |
| Age | | | | | |
| 18-34 | 40 | 32.3 | 106 | 67.7 | 9.3 |
| 35-54 | 45 | 19.8 | 211 | 80.2 | 5.6 |
| ≥55 | 14 | 8.1 | 212 | 91.9 | 4.6 |
| Education | | | | | |
| ≤High School | 43 | 18.7 | 252 | 81.3 | 5.6 |
| >High School | 57 | 21.9 | 282 | 78.1 | 5.7 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 **Drink per Day in Past Month)**

Clinton, Essex, Franklin

| Demographic Groups | | Yes | n | No % | C.I. ³ |
|-----------------------|----|-----|-----|---------|-------------------|
| Total | 40 | 6.4 | 602 | 93.6 | 2.3 |
| Gender | | | | | |
| Male | 20 | 6.7 | 254 | 93.3 | 3.5 |
| Female | 20 | 6.0 | 348 | 94.0 | 2.9 |
| Age | | | | | |
| 18-34 | 11 | 7.8 | 137 | 92.2 | 5.3 |
| 35-54 | 16 | 5.9 | 243 | 94.1 | 2.9 |
| ≥55 | 13 | 5.6 | 215 | 94.4 | 3.6 |
| Education | | | | | |
| ≤High School | 18 | 5.2 | 282 | 94.8 | 2.5 |
| >High School | 22 | 7.4 | 320 | 92.6 | 3.7 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health Overweight or Obese, BMI(Body Mass Index) 25 or More Clinton, Essex, Franklin

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|-----|------------|-----|---------|-------------------|
| Total | 352 | 56.7 | 269 | 43.3 | 4.6 |
| Gender | | | | | |
| Male | 185 | 65.8 | 87 | 34.2 | 6.9 |
| Female | 167 | 46.2 | 182 | 53.8 | 5.8 |
| Age | | | | | |
| 18-34 | 60 | 43.4 | 80 | 56.6 | 9.7 |
| 35-54 | 152 | 63.2 | 101 | 36.8 | 6.4 |
| ≥55 | 139 | 63.0 | 87 | 37.0 | 7.0 |
| Education | | | | | |
| ≤High School | 171 | 56.7 | 124 | 43.3 | 6.6 |
| >High School | 181 | 56.7 | 145 | 43.3 | 6.4 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+) *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Underweight\ Normal %0 ² | C.I. ³ | n | Overweight % | C.I. | n | Obese % | C.I. |
|-----------------------|-----|---|-------------------|-----|-----------------|------|-----|------------|------|
| Total | 268 | 43.1 | 4.6 | 212 | 35.1 | 4.4 | 140 | 21.7 | 3.7 |
| Gender | | | | | | | | | |
| Male | 87 | 34.2 | 6.9 | 117 | 43.6 | 6.9 | 68 | 22.2 | 5.5 |
| Female | 181 | 53.6 | 5.8 | 95 | 25.3 | 4.9 | 72 | 21.1 | 4.9 |
| Age | | | | | | | | | |
| 18-34 | 80 | 56.6 | 9.7 | 36 | 29.1 | 9.1 | 24 | 14.3 | 6.3 |
| 35-54 | 101 | 36.8 | 6.4 | 86 | 35.8 | 6.6 | 66 | 27.5 | 6.2 |
| ≥55 | 86 | 36.6 | 7.0 | 90 | 41.4 | 7.4 | 49 | 22.1 | 6.2 |
| Education | | | | | | | | | |
| ≤High School | 123 | 43.0 | 6.6 | 99 | 34.8 | 6.4 | 72 | 22.2 | 5.1 |
| >High School | 145 | 43.3 | 6.4 | 113 | 35.4 | 6.1 | 68 | 21.3 | 5.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

| Demographic Groups | n1 | Yes % | n | No % | C.I. ³ |
|-----------------------|-----|----------|----|---------|-------------------|
| Female | 217 | 91.5 | 22 | 8.5 | 3.7 |
| Age | | | | | |
| 40-64 | 126 | 89.8 | 16 | 10.2 | 5.1 |
| ≥65 | 91 | 94.9 | 6 | 5.1 | 4.2 |
| Education | | | | | |
| ≤High School | 113 | 91.9 | 10 | 8.1 | 5.1 |
| >High School | 104 | 91.1 | 12 | 8.9 | 5.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

Clinton, Essex, Franklin

| Demographic Groups | n1 | Yes % | n | No % | C.I. ³ |
|-----------------------|-----|----------|----|---------|-------------------|
| Female | 193 | 80.8 | 45 | 19.2 | 5.8 |
| Age | | | | | |
| 40-64 | 112 | 78.8 | 29 | 21.2 | 7.9 |
| ≥65 | 81 | 84.6 | 16 | 15.4 | 7.5 |
| Education | | | | | |
| ≤High School | 98 | 80.4 | 25 | 19.6 | 7.5 |
| >High School | 95 | 81.1 | 20 | 18.9 | 9.0 |

Expanded BRFSS, 2003 - New York State Department of Health

Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger

Clinton, Essex, Franklin

| Demographic Groups | n1 | Yes % | n | No % | C.I. ³ |
|-----------------------|----|----------|-----|---------|-------------------|
| Total | 53 | 11.6 | 423 | 88.4 | 3.5 |
| Gender | | | | | |
| Male | 12 | 7.2 | 205 | 92.8 | 4.7 |
| Female | 41 | 17.0 | 218 | 83.0 | 5.3 |
| Age | | | | | |
| 18-34 | 36 | 24.1 | 108 | 75.9 | 8.2 |
| 35-54 | 13 | 3.4 | 239 | 96.6 | 2.0 |
| 55-64 | 4 | 4.4 | 76 | 95.6 | 4.8 |
| Education | | | | | |
| ≤High School | 20 | 11.8 | 182 | 88.2 | 5.6 |
| >High School | 33 | 11.4 | 241 | 88.6 | 4.5 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health Sexually Active, Age 18-64 Years Old *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|-----|------------|----|---------|-------------------|
| Total | 383 | 86.0 | 70 | 14.0 | 3.9 |
| Gender | | | | | |
| Male | 175 | 83.9 | 34 | 16.1 | 6.1 |
| Female | 208 | 88.7 | 36 | 11.3 | 4.3 |
| Age | | | | | |
| 18-34 | 125 | 85.7 | 17 | 14.3 | 7.7 |
| 35-54 | 206 | 88.3 | 32 | 11.7 | 4.6 |
| 55-64 | 52 | 79.8 | 21 | 20.2 | 9.2 |
| Education | | | | | |
| ≤High School | 163 | 84.0 | 32 | 16.0 | 6.5 |
| >High School | 220 | 87.5 | 38 | 12.5 | 4.8 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health

Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|----|------------|-----|---------|-------------------|
| Total | 51 | 14.4 | 339 | 85.6 | 4.5 |
| Gender | | | | | |
| Male | 29 | 17.6 | 150 | 82.4 | 7.3 |
| Female | 22 | 10.5 | 189 | 89.5 | 4.5 |
| Age | | | | | |
| 18-34 | 28 | 25.8 | 98 | 74.2 | 9.6 |
| 35-54 | 19 | 6.9 | 190 | 93.1 | 3.3 |
| 55-64 | 4 | 6.7 | 51 | 93.3 | 7.8 |
| Education | | | | | |
| ≤High School | 23 | 13.0 | 144 | 87.0 | 6.5 |
| >High School | 28 | 15.5 | 195 | 84.5 | 6.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

Clinton, Essex, Franklin

| Demographic Groups | n1 | Use BC | C.I. ³ | n | No BC at risk % | C.I. | n | No BC % | C.I. |
|-----------------------|-----|--------|-------------------|----|--------------------|------|----|---------|------|
| Total | 220 | 64.7 | 6.1 | 44 | 11.7 | 3.7 | 75 | 23.6 | 5.6 |
| Gender | | | | | | | | | |
| Male | 122 | 65.5 | 8.1 | 22 | 10.5 | 4.7 | 43 | 24.0 | 7.5 |
| Female | 98 | 63.1 | 8.6 | 22 | 14.1 | 5.8 | 32 | 22.8 | 7.8 |
| Age | | | | | | | | | |
| 18-34 | 96 | 66.8 | 9.3 | 22 | 13.4 | 5.9 | 25 | 19.8 | 8.5 |
| 35-59 | 124 | 62.5 | 7.7 | 22 | 10.0 | 4.3 | 50 | 27.5 | 7.3 |
| Education | | | | | | | | | |
| ≤High School | 84 | 54.7 | 9.9 | 18 | 13.0 | 6.4 | 36 | 32.3 | 9.9 |
| >High School | 136 | 71.9 | 7.0 | 26 | 10.8 | 4.3 | 39 | 17.3 | 6.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health Physician Diagnosed Heart Attack, Angina or Stroke *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes % | n | No % | C.I. ³ |
|-----------------------|----|----------|-----|---------|-------------------|
| Total | 61 | 8.0 | 571 | 92.0 | 2.3 |
| Gender | | | | | |
| Male | 27 | 8.3 | 240 | 91.7 | 3.6 |
| Female | 34 | 7.6 | 331 | 92.4 | 2.7 |
| Age | | | | | |
| 18-64 | 20 | 4.1 | 458 | 95.9 | 2.1 |
| ≥65 | 41 | 26.8 | 108 | 73.2 | 7.7 |
| Education | | | | | |
| ≤High School | 37 | 9.9 | 255 | 90.1 | 3.4 |
| >High School | 24 | 6.3 | 316 | 93.7 | 3.0 |

Expanded BRFSS, 2003 - New York State Department of Health Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina) Clinton, Essex, Franklin

| Demographic Groups | n1 | Yes % | n | No % | C.I. ³ |
|-----------------------|----|----------|-----|---------|-------------------|
| Total | 47 | 6.2 | 584 | 93.8 | 2.1 |
| Gender | | | | | |
| Male | 23 | 7.6 | 244 | 92.4 | 3.5 |
| Female | 24 | 4.7 | 340 | 95.3 | 2.0 |
| Age | | | | | |
| 18-64 | 19 | 4.0 | 459 | 96.0 | 2.1 |
| ≥65 | 28 | 16.9 | 120 | 83.1 | 6.3 |
| Education | | | | | |
| ≤High School | 25 | 6.6 | 266 | 93.4 | 2.8 |
| >High School | 22 | 5.8 | 318 | 94.2 | 3.0 |

Expanded BRFSS, 2003 - New York State Department of Health Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|----|------------|----|---------|-------------------|
| Male | 95 | 54.2 | 75 | 45.8 | 8.4 |
| Age | | | | | |
| 40-64 | 56 | 44.4 | 66 | 55.6 | 9.8 |
| ≥65 | 39 | 84.7 | 9 | 15.3 | 10.4 |
| Education | | | | | |
| ≤High School | 37 | 48.5 | 43 | 51.5 | 12.2 |
| >High School | 58 | 59.2 | 32 | 40.8 | 11.7 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

Clinton, Essex, Franklin

| Demographic Groups | n ¹ | Yes %02 | n | No % | C.I. ³ |
|-----------------------|----------------|------------|----|---------|-------------------|
| Male | 78 | 44.1 | 89 | 55.9 | 8.3 |
| Age | | | | | |
| 40-64 | 45 | 33.8 | 76 | 66.2 | 9.0 |
| ≥65 | 33 | 76.5 | 13 | 23.5 | 12.8 |
| Education | | | | | |
| ≤High School | 27 | 35.1 | 51 | 64.9 | 11.7 |
| >High School | 51 | 52.0 | 38 | 48.0 | 11.7 |

Expanded BRFSS, 2003 - New York State Department of Health Ever Had a Digital Rectal Exam, Men Aged 40 and Older *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes % | n | No % | C.I. ³ |
|-----------------------|-----|----------|----|---------|-------------------|
| Male | 136 | 74.6 | 43 | 25.4 | 7.3 |
| Age | | | | | |
| 40-64 | 93 | 72.5 | 35 | 27.5 | 8.7 |
| ≥65 | 43 | 81.1 | 8 | 18.9 | 13.1 |
| Education | | | | | |
| ≤High School | 55 | 68.3 | 27 | 31.7 | 11.2 |
| >High School | 81 | 79.9 | 16 | 20.1 | 9.6 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

Expanded BRFSS, 2003 - New York State Department of Health Ever Used a Blood Stool Test at Home, Aged 50 and Older *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|-----|------------|-----|---------|-------------------|
| Total | 135 | 47.3 | 152 | 52.7 | 6.6 |
| Gender | | | | | |
| Male | 53 | 46.1 | 62 | 53.9 | 10.3 |
| Female | 82 | 48.5 | 90 | 51.5 | 8.3 |
| Age | | | | | |
| 50-64 | 55 | 38.2 | 87 | 61.8 | 9.0 |
| ≥65 | 80 | 59.5 | 65 | 40.5 | 8.6 |
| Education | | | | | |
| ≤High School | 58 | 42.1 | 85 | 57.9 | 9.1 |
| >High School | 77 | 52.3 | 67 | 47.7 | 9.4 |

Expanded BRFSS, 2003 - New York State Department of Health Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|----|------------|-----|---------|-------------------|
| Total | 51 | 20.2 | 227 | 79.8 | 5.5 |
| Gender | | | | | |
| Male | 17 | 17.9 | 96 | 82.1 | 8.3 |
| Female | 34 | 22.4 | 131 | 77.6 | 7.4 |
| Age | | | | | |
| 50-64 | 19 | 14.6 | 119 | 85.4 | 6.7 |
| ≥65 | 32 | 27.9 | 108 | 72.1 | 8.9 |
| Education | | | | | |
| ≤High School | 27 | 22.3 | 112 | 77.7 | 8.2 |
| >High School | 24 | 18.2 | 115 | 81.8 | 7.5 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

35

Expanded BRFSS, 2003 - New York State Department of Health Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older

Clinton, Essex, Franklin

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|----|------------|-----|---------|-------------------|
| Total | 77 | 29.4 | 201 | 70.6 | 6.1 |
| Gender | | | | | |
| Male | 30 | 29.2 | 83 | 70.8 | 9.5 |
| Female | 47 | 29.6 | 118 | 70.4 | 7.9 |
| Age | | | | | |
| 50-64 | 31 | 23.8 | 107 | 76.2 | 8.0 |
| ≥65 | 46 | 37.1 | 94 | 62.9 | 9.2 |
| Education | | | | | |
| ≤High School | 38 | 29.6 | 101 | 70.4 | 8.7 |
| >High School | 39 | 29.2 | 100 | 70.8 | 8.7 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

36

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|-----|------------|-----|---------|-------------------|
| Total | 157 | 50.9 | 132 | 49.1 | 6.6 |
| Gender | | | | | |
| Male | 59 | 43.8 | 56 | 56.2 | 10.0 |
| Female | 98 | 57.3 | 76 | 42.7 | 8.2 |
| Age | | | | | |
| 50-64 | 59 | 40.2 | 82 | 59.8 | 9.0 |
| ≥65 | 98 | 64.7 | 50 | 35.3 | 8.7 |
| Education | | | | | |
| ≤High School | 73 | 49.2 | 71 | 50.8 | 9.2 |
| >High School | 84 | 52.4 | 61 | 47.6 | 9.4 |

Expanded BRFSS, 2003 - New York State Department of Health

Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older *Clinton, Essex, Franklin*

| Demographic Groups | n1 | Yes %02 | n | No % | C.I. ³ |
|-----------------------|-----|------------|-----|---------|-------------------|
| Total | 145 | 47.5 | 141 | 52.5 | 6.6 |
| Gender | | | | | |
| Male | 55 | 40.7 | 59 | 59.3 | 9.8 |
| Female | 90 | 53.8 | 82 | 46.2 | 8.3 |
| Age | | | | | |
| 50-64 | 56 | 38.6 | 84 | 61.4 | 9.0 |
| ≥65 | 89 | 59.2 | 57 | 40.8 | 8.9 |
| Education | | | | | |
| ≤High School | 67 | 45.4 | 75 | 54.6 | 9.2 |
| >High School | 78 | 49.4 | 66 | 50.6 | 9.4 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

38

Expanded BRFSS, 2003 - New York State Department of Health

Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older

Clinton, Essex, Franklin

| Demographic Groups | n1 | Yes % | n | No % | C.I. ³ |
|-----------------------|-----|----------|-----|---------|-------------------|
| Total | 162 | 54.0 | 126 | 46.0 | 6.6 |
| Gender | | | | | |
| Male | 62 | 47.2 | 54 | 52.8 | 10.2 |
| Female | 100 | 60.4 | 72 | 39.6 | 8.1 |
| Age | | | | | |
| 50-64 | 63 | 42.2 | 78 | 57.8 | 9.1 |
| ≥65 | 99 | 69.5 | 48 | 30.5 | 8.0 |
| Education | | | | | |
| ≤High School | 77 | 54.4 | 66 | 45.6 | 9.2 |
| >High School | 85 | 53.6 | 60 | 46.4 | 9.4 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent