### Expanded BRFSS, 2003

### Hamilton and Herkimer Locality Tables

#### TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days	
Core 2: Health Care Access	
Core 3: Exercise	8
Core 4: Diabetes	9
Core 5: Asthma	10
Core 6: Arthritis	12
Core 7: Tobacco Use	
Core 8: Tobacco ETS	17
Core 9: Alcohol Consumption	19
Core 10: Weight Status	2
Core 11: Mammography	23
Core 12: Sexual Behavior	25
Core 13: Family Planning	28
Core 14: Cardiovascular Disease	29
Core 15: Prostate Cancer Screening	3
Core 16: Colorectal Cancer Screening	37

New York State Department Of Health

### **Self-rated Health Status**

Demographic Groups	n¹	Fair or Poor	n	Good to Excellent %	C.I. <sup>3</sup>
Total	118	17.2	499	82.8	3.3
Gender					
Male	56	19.3	198	80.7	5.2
Female	62	15.4	301	84.6	4.0
Age					
18-34	10	9.0	98	91.0	5.9
35-54	33	14.8	205	85.2	5.2
≥55	74	25.1	192	74.9	5.7
Education					
≤High School	77	21.8	227	78.2	5.0
>High School	40	12.9	271	87.1	4.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Reported Number of Physical Distress Days in the Past Month** Hamilton, Herkimer

Demographic Groups	n¹	≥14 days %2	n	< 14 days %	C.I. <sup>3</sup>
Total	79	11.9	523	88.1	2.8
Gender					
Male	31	10.8	216	89.2	4.1
Female	48	13.0	307	87.0	4.0
Age					
18-34	10	8.6	98	91.4	5.9
35-54	26	11.4	209	88.6	4.5
≥55	43	15.1	211	84.9	4.7
Education					
≤High School	48	15.0	248	85.0	4.5
>High School	31	9.1	274	90.9	3.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### **Reported Number of Mental Distress Days in the Past Month** Hamilton, Herkimer

Demographic Groups	n¹	≥14 days %2	n	< 14 days %	C.I. <sup>3</sup>
Total	72	10.4	531	89.6	2.6
Gender					
Male	29	9.0	221	91.0	3.6
Female	43	11.7	310	88.3	3.7
Age					
18-34	18	13.3	90	86.7	6.5
35-54	32	11.2	202	88.8	4.0
≥55	21	7.3	235	92.7	3.2
Education					
≤High School	39	12.5	256	87.5	4.2
>High School	31	8.2	275	91.8	3.0

<sup>&</sup>lt;sup>1</sup> Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup> Weighted Percent<sup>3</sup> 95% Confidence Interval

### Expanded BRFSS, 2003 - New York State Department of Health **Health Insurance Coverage, 18-64 Years of Age** Hamilton, Herkimer

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	387	83.5	66	16.5	4.4
Gender					
Male	163	78.5	40	21.5	7.0
Female	224	88.5	26	11.5	5.3
Age					
18-34	79	68.1	30	31.9	10.6
35-54	211	90.4	27	9.6	3.7
55-64	97	93.7	9	6.3	4.3
Education					
≤High School	163	76.9	36	23.1	7.9
>High School	224	88.8	30	11.2	4.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health **Health Insurance Coverage, Aged 18 and Older** *Hamilton, Herkimer*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	546	86.5	71	13.5	3.5
Gender					
Male	211	81.7	43	18.3	5.9
Female	335	91.0	28	9.0	4.0
Age					
18-34	79	68.1	30	31.9	10.6
35-54	211	90.4	27	9.6	3.7
≥55	251	95.8	13	4.2	2.4
Education					
≤High School	263	81.9	41	18.1	6.0
>High School	282	90.7	30	9.3	3.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Reported Having a Routine Physical Exam in Past Two Years *Hamilton*, *Herkimer*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	552	89.9	61	10.1	2.8
Gender					
Male	217	86.2	37	13.8	4.9
Female	335	93.3	24	6.7	2.9
Age					
18-34	94	87.2	12	12.8	7.7
35-54	199	84.9	38	15.1	4.8
≥55	253	96.2	11	3.8	2.4
Education					
≤High School	275	90.7	25	9.3	4.4
>High School	275	89.1	36	10.9	3.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Time in Past 12 Months when Needed Medical Care but Could Not Get It *Hamilton, Herkimer*

Demographic		Yes		No	CI.
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	37	6.4	581	93.6	2.3
Gender					
Male	13	5.8	242	94.2	3.6
Female	24	7.0	339	93.0	2.9
Age					
18-34	16	13.8	93	86.2	7.1
35-54	14	4.6	223	95.4	2.6
≥55	7	3.2	259	96.8	2.5
Education					
≤High School	16	6.0	288	94.0	3.2
>High School	21	6.9	291	93.1	3.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days *Hamilton, Herkimer*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	437	72.4	182	27.6	4.0
Gender					
Male	184	72.7	72	27.3	6.2
Female	253	72.1	110	27.9	5.2
Age					
18-34	86	76.9	23	23.1	9.6
35-54	181	75.9	57	24.1	6.1
≥55	165	65.6	101	34.4	6.2
Education					
≤High School	186	62.9	119	37.1	6.3
>High School	251	81.5	61	18.5	4.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Physician Diagnosed Diabetes**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	59	8.4	560	91.6	2.3
Gender	37	0.4	300	71.0	2.3
Male	31	10.4	225	89.6	3.8
Female	28	6.4	335	93.6	2.6
Age					
18-34	1	0.5	108	99.5	0.9
35-54	15	7.1	223	92.9	3.8
≥55	43	15.3	223	84.7	4.6
Education					
≤High School	36	10.1	269	89.9	3.4
>High School	23	6.7	289	93.3	3.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### Lifetime Asthma Diagnosed by a Medical Professional

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	69	10.7	549	89.3	2.9
Gender					
Male	24	8.7	231	91.3	4.0
Female	45	12.5	318	87.5	4.0
Age					
18-34	17	15.8	92	84.2	8.0
35-54	22	7.7	215	92.3	3.5
≥55	30	10.3	236	89.7	3.9
Education					
≤High School	34	11.4	270	88.6	4.3
>High School	34	9.9	278	90.1	3.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Current Asthma Diagnosed by a Medical Professional**

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	47	6.8	569	93.2	2.2
Gender					
Male	15	5.6	240	94.4	3.4
Female	32	7.9	329	92.1	2.9
Age					
18-34	10	8.0	98	92.0	5.7
35-54	18	6.5	219	93.5	3.3
≥55	19	6.3	246	93.7	3.2
Education					
≤High School	22	7.3	281	92.7	3.5
>High School	24	6.1	287	93.9	2.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### **Arthritis Diagnosed by a Medical Professional**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	231	34.0	388	66.0	4.2
Gender					
Male	80	29.7	176	70.3	6.4
Female	151	38.0	212	62.0	5.6
Age					
18-34	12	10.7	97	89.3	7.1
35-54	75	30.9	163	69.1	6.4
≥55	141	52.7	125	47.3	6.9
Education					
≤High School	130	38.2	175	61.8	6.2
>High School	101	30.3	211	69.7	5.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

Demographic Groups	n¹	Doctor Diagnosed %2	C.I. <sup>3</sup>	n	Possible Arthritis (CJS Only) %	rthritis (CJS Only)		No Arthritis or CJS %	C.I.
Total	231	34.4	4.3	82	14.9	3.5	301	50.7	4.6
Gender									
Male	80	30.3	6.5	41	15.7	5.1	133	54.0	7.0
Female	151	38.3	5.6	41	14.1	4.7	168	47.6	5.9
Age									
18-34	12	10.8	7.2	19	20.8	9.1	76	68.4	10.4
35-54	75	31.5	6.4	40	17.4	5.4	122	51.0	7.0
≥55	141	52.9	6.9	22	8.3	3.9	102	38.9	6.7
Education									
≤High School	130	38.5	6.2	37	14.4	5.2	136	47.1	6.6
>High School	101	30.8	5.9	43	15.0	4.6	165	54.2	6.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Ever Smoked 100 Cigarettes in Lifetime**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	342	55.1	276	44.9	4.5
Gender	3.2	00.1	2,0	,	
Male	154	59.4	101	40.6	7.0
Female	188	51.2	175	48.8	5.9
Age					
18-34	57	53.2	52	46.8	10.7
35-54	132	53.6	106	46.4	7.1
≥55	152	58.7	113	41.3	6.7
Education					
≤High School	193	65.7	111	34.3	6.1
>High School	148	45.4	164	54.6	6.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### **Current Smoker**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	162	25.7	456	74.3	4.0
Gender					
Male	69	25.3	186	74.7	6.0
Female	93	26.0	270	74.0	5.4
Age					
18-34	39	36.4	70	63.6	10.4
35-54	86	32.3	152	67.7	6.4
≥55	37	12.5	228	87.5	4.2
Education					
≤High School	90	30.5	214	69.5	6.3
>High School	72	21.3	240	78.7	5.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### **Status of Cigarette Smoking**

Demographic Groups	n¹	Everyday Smoker %	C.I. <sup>3</sup>	n	Someday Smoker %	C.I.	n	Former or Never Smoked %	C.I.
Total	134	21.5	3.8	28	4.2	1.7	456	74.3	4.0
Gender									
Male	58	21.2	5.7	11	4.1	2.6	186	74.7	6.0
Female	76	21.7	5.2	17	4.3	2.2	270	74.0	5.4
Age									
18-34	28	29.1	10.1	11	7.3	4.7	70	63.6	10.4
35-54	75	28.0	6.2	11	4.2	2.6	152	67.7	6.4
≥55	31	10.4	3.9	6	2.1	1.8	228	87.5	4.2
Education									
≤High School	76	25.7	6.1	14	4.8	2.7	214	69.5	6.3
>High School	58	17.7	4.7	14	3.7	2.0	240	78.7	5.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### **Home Smoking Policies**

Demographic Groups	n¹	Not Allowed	C.I. <sup>3</sup>	n	Allowed Some Places	C.I.	n	Allowed Anywhere	C.I.
Total	360	60.6	4.4	79	11.8	2.8	179	27.6	4.1
Gender									
Male	148	63.5	6.5	30	10.4	4.0	78	26.1	5.9
Female	212	57.9	5.9	49	13.1	3.9	101	29.0	5.6
Age									
18-34	64	59.7	10.5	15	12.1	6.6	30	28.2	9.9
35-54	139	62.1	6.7	37	15.1	4.9	62	22.9	5.7
≥55	154	60.4	6.6	26	8.3	3.3	85	31.3	6.3
Education									
≤High School	155	52.2	6.6	43	14.7	4.6	106	33.1	6.2
>High School	204	68.4	5.7	36	9.2	3.1	72	22.4	5.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Home Smoking**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	186	29.7	432	70.3	4.2
Gender					
Male	79	28.1	176	71.9	6.1
Female	107	31.2	256	68.8	5.7
Age					
18-34	42	38.7	67	61.3	10.4
35-54	89	33.5	149	66.5	6.5
≥55	55	20.4	211	79.6	5.4
Education					
≤High School	106	36.1	198	63.9	6.5
>High School	80	23.9	232	76.1	5.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)

Demographic		Yes		No	
Groups	n¹	%2	n	%	C.I. <sup>3</sup>
Total	77	17.0	525	83.0	4.0
Gender					
Male	51	26.2	192	73.8	7.0
Female	26	8.9	333	91.1	3.8
Age					
18-34	29	33.1	76	66.9	10.6
35-54	32	13.9	199	86.1	4.9
≥55	16	9.3	245	90.7	5.4
Education					
≤High School	36	16.7	257	83.3	5.7
>High School	41	17.3	266	82.7	5.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	27	4.6	582	95.4	1.9
Gender					
Male	15	5.8	232	94.2	3.1
Female	12	3.5	350	96.5	2.1
Age					
18-34	6	4.5	102	95.5	4.1
35-54	13	5.5	221	94.5	3.1
≥55	8	3.9	254	96.1	2.8
Education					
≤High School	13	4.6	284	95.4	2.6
>High School	14	4.7	296	95.3	2.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health Overweight or Obese, BMI(Body Mass Index) 25 or More *Hamilton, Herkimer*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	358	62.9	230	37.1	4.5
Gender					
Male	174	69.6	76	30.4	6.5
Female	184	56.3	154	43.7	6.1
Age					
18-34	59	58.8	46	41.2	10.6
35-54	138	62.2	89	37.8	7.0
≥55	159	66.2	94	33.8	6.5
Education					
≤High School	184	66.5	104	33.5	6.3
>High School	174	59.6	125	40.4	6.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+) *Hamilton, Herkimer*

Demographic Groups	n¹	Underweight\ Normal %2	C.I. <sup>3</sup>	n	Overweight %	C.I.	n	Obese %	C.I.
Total	230	37.2	4.5	226	40.6	4.7	131	22.3	3.8
Gender									
Male	76	30.4	6.5	127	52.0	7.1	47	17.6	5.2
Female	154	43.8	6.1	99	29.4	5.7	84	26.8	5.5
Age									
18-34	46	41.2	10.6	39	37.8	10.8	20	21.0	8.9
35-54	89	37.8	7.0	81	36.2	7.1	57	26.0	6.4
≥55	94	33.9	6.5	104	46.3	7.1	54	19.8	5.3
Education									
≤High School	104	33.5	6.3	110	39.3	6.7	73	27.1	6.1
>High School	125	40.4	6.3	116	41.8	6.5	58	17.9	4.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# **Ever Had Mammography Screening, Women Aged 40 Years and Older** *Hamilton, Herkimer*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Female	245	91.6	21	8.4	4.1
Age					
40-64	149	93.9	8	6.1	5.0
≥65	96	87.7	13	12.3	7.3
Education					
≤High School	113	87.2	14	12.8	7.7
>High School	132	95.6	6	4.4	3.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Female	203	75.9	62	24.1	5.9
Age					
40-64	126	79.2	30	20.8	7.5
≥65	77	70.3	32	29.7	9.5
Education					
≤High School	92	70.3	35	29.7	9.2
>High School	111	80.9	26	19.1	7.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	55	12.3	389	87.7	3.8
Gender					
Male	20	8.2	181	91.8	4.1
Female	35	16.5	208	83.5	6.2
Age					
18-34	28	23.5	80	76.5	9.2
35-54	22	8.1	211	91.9	3.8
55-64	5	3.0	98	97.0	3.0
Education					
≤High School	33	19.7	162	80.3	7.3
>High School	22	6.5	227	93.5	2.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### Sexually Active, Age 18-64 Years Old

Dama awan bia		Vac		Na	
Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	350	87.0	69	13.0	3.5
Gender					
Male	158	86.4	29	13.6	5.7
Female	192	87.6	40	12.4	4.1
Age					
18-34	94	90.7	8	9.3	7.1
35-54	195	90.0	30	10.0	3.9
55-64	61	71.8	31	28.2	9.5
Education					
≤High School	146	84.9	39	15.1	5.5
>High School	204	88.7	30	11.3	4.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger *Hamilton, Herkimer*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	58	15.9	300	84.1	4.5
Gender					
Male	35	18.2	128	81.8	6.4
Female	23	13.7	172	86.3	6.2
Age					
18-34	27	24.5	71	75.5	9.7
35-54	30	13.6	169	86.4	5.1
55-64	1	2.1	60	97.9	4.1
Education					
≤High School	27	15.9	121	84.1	6.6
>High School	31	15.9	179	84.1	6.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

Demographic Groups	n¹	Use BC	C.I. <sup>3</sup>	n	No BC at risk	C.I.	n	No BC	C.I.
Total	195	67.3	6.2	44	14.6	4.8	62	18.1	4.9
Gender									
Male	101	59.7	8.5	29	17.7	7.1	48	22.6	6.9
Female	94	78.6	7.9	15	10.0	5.4	14	11.3	6.4
Age									
18-34	76	70.4	10.0	13	11.8	6.9	17	17.8	8.5
35-59	119	64.8	7.6	31	16.9	6.6	45	18.3	5.5
Education									
≤High School	82	66.5	9.4	19	14.6	7.2	31	18.9	7.5
>High School	113	68.0	8.2	25	14.6	6.5	31	17.4	6.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Physician Diagnosed Heart Attack, Angina or Stroke

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	64	9.3	546	90.7	2.4
Gender					
Male	35	11.7	217	88.3	4.0
Female	29	7.1	329	92.9	2.7
Age					
18-64	19	4.1	427	95.9	2.0
≥65	45	27.8	115	72.2	7.5
Education					
≤High School	36	10.6	263	89.4	3.6
>High School	27	8.0	282	92.0	3.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina) *Hamilton, Herkimer*

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	60	8.8	549	91.2	2.3
Gender					
Male	34	11.4	218	88.6	4.0
Female	26	6.4	331	93.6	2.6
Age					
18-64	19	4.1	427	95.9	2.0
≥65	41	25.6	118	74.4	7.4
Education					
≤High School	34	10.0	264	90.0	3.5
>High School	25	7.6	284	92.4	3.1

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# **Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older** *Hamilton, Herkimer*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	101	63.8	65	36.2	8.1
Age					
40-64	68	59.0	52	41.0	9.9
≥65	33	75.1	13	24.9	14.0
Education					
≤High School	48	56.5	40	43.5	11.5
>High School	53	70.8	25	29.2	10.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	83	53.8	81	46.2	8.6
Age					
40-64	54	48.4	64	51.6	10.5
≥65	29	66.3	17	33.7	15.0
Education					
≤High School	37	44.1	49	55.9	11.5
>High School	46	62.8	32	37.2	11.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had a Digital Rectal Exam, Men Aged 40 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	141	84.6	30	15.4	5.6
Age					
40-64	95	79.7	28	20.3	7.5
≥65	46	95.9	2	4.1	5.6
Education					
≤High School	74	80.4	19	19.6	8.7
>High School	67	88.7	11	11.3	6.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# **Ever Used a Blood Stool Test at Home, Aged 50 and Older** *Hamilton, Herkimer*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	143	43.1	184	56.9	6.1
Gender					
Male	49	37.1	82	62.9	9.5
Female	94	48.9	102	51.1	7.7
Age					
50-64	74	42.4	98	57.6	8.5
≥65	69	44.0	86	56.0	8.8
Education					
≤High School	64	36.5	109	63.5	7.9
>High School	79	49.6	74	50.4	9.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older *Hamilton, Herkimer*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	69	19.1	257	80.9	4.6
Gender					
Male	25	16.7	105	83.3	6.7
Female	44	21.5	152	78.5	6.1
Age					
50-64	34	17.3	138	82.7	5.9
≥65	35	21.4	119	78.6	7.1
Education					
≤High School	28	14.7	145	85.3	5.7
>High School	41	23.5	111	76.5	7.1

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older *Hamilton, Herkimer*

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	97	28.6	229	71.4	5.6
Gender					
Male	35	26.1	95	73.9	8.8
Female	62	30.9	134	69.1	7.0
Age					
50-64	52	29.2	120	70.8	7.8
≥65	45	27.8	109	72.2	7.9
Education					
≤High School	41	22.5	132	77.5	6.7
>High School	56	34.6	96	65.4	8.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older *Hamilton, Herkimer*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	159	48.6	172	51.4	6.2
Gender					
Male	67	50.9	66	49.1	9.8
Female	92	46.4	106	53.6	7.6
Age					
50-64	77	44.4	96	55.6	8.6
≥65	82	53.6	76	46.4	8.8
Education					
≤High School	81	45.8	94	54.2	8.2
>High School	78	51.6	76	48.4	9.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older *Hamilton, Herkimer*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	144	43.9	185	56.1	6.2
Gender					
Male	61	45.3	72	54.7	9.8
Female	83	42.7	113	57.3	7.6
Age					
50-64	72	41.2	101	58.8	8.5
≥65	72	47.3	84	52.7	8.9
Education					
≤High School	74	42.5	99	57.5	8.2
>High School	70	45.6	84	54.4	9.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	180	54.7	152	45.3	6.1
Gender					
Male	74	55.3	59	44.7	9.7
Female	106	54.1	93	45.9	7.6
Age					
50-64	89	50.5	84	49.5	8.6
≥65	91	59.7	68	40.3	8.6
Education					
≤High School	87	49.4	89	50.6	8.2
>High School	93	60.2	61	39.8	9.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval