Expanded BRFSS, 2003

Jefferson, Lewis and St. Lawrence Locality Tables

TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days	
Core 1: Health Status / Healthy Days Core 2: Health Care Access	
Core 3: Exercise	
Core 4: Diabetes	
Core 5: Asthma	12
Core 7: Tobacco Use	
Core 8: Tobacco ETS	17
Core 9: Alcohol Consumption	19
Core 9: Alcohol Consumption Core 10: Weight Status Core 11: Mammography	2
Core 11: Mammography	23
Core 12: Sexual Behavior.	25
Core 13: Family Planning	
Core 14: Cardiovascular Disease	
Core 15: Prostate Cancer Screening	
Core 16: Colorectal Cancer Screening	

New York State Department Of Health

Self-rated Health Status

Demographic Groups	n¹	Fair or Poor	n	Good to Excellent %	C.I. ³
Total	114	15.7	525	84.3	3.5
Gender					
Male	41	16.5	205	83.5	5.9
Female	73	14.9	320	85.1	3.6
Age					
18-34	6	4.8	127	95.2	5.9
35-54	44	18.1	217	81.9	5.7
≥55	64	25.8	180	74.2	6.4
Education					
≤High School	72	19.1	263	80.9	4.9
>High School	42	12.2	260	87.8	5.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Reported Number of Physical Distress Days in the Past Month Jefferson, Lewis, St. Lawrence

Demographic Groups	n¹	≥14 days	n	< 14 days %	C.I. ³
Total	89	11.9	542	88.1	2.8
Gender					
Male	32	11.8	210	88.2	4.5
Female	57	12.1	332	87.9	3.3
Age					
18-34	8	4.0	125	96.0	3.1
35-54	37	14.6	220	85.4	5.2
≥55	44	18.2	196	81.8	5.7
Education					
≤High School	58	16.0	272	84.0	4.7
>High School	31	7.7	268	92.3	2.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

Reported Number of Mental Distress Days in the Past Month

Demographic Groups	n¹	≥14 days ⁰ / ₀ ²	n	< 14 days %	C.I. ³
Total	52	7.3	576	92.7	2.3
Gender					
Male	18	6.6	221	93.4	3.5
Female	34	8.0	355	92.0	3.0
Age					
18-34	8	4.2	122	95.8	3.3
35-54	26	10.3	229	89.7	4.7
≥55	18	7.3	224	92.7	3.6
Education					
≤High School	34	9.0	293	91.0	3.7
>High School	18	5.5	283	94.5	2.8

¹ Percents based on row denominators of less than 50 are unstable and should be used with caution

² Weighted Percent³ 95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health Health Insurance Coverage, 18-64 Years of Age *Jefferson, Lewis, St. Lawrence*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	417	82.6	79	17.4	4.4
Gender					
Male	165	79.9	36	20.1	7.1
Female	252	85.6	43	14.4	4.9
Age					
18-34	102	76.0	30	24.0	8.8
35-54	222	86.1	38	13.9	5.0
55-64	93	90.7	11	9.3	5.7
Education					
≤High School	192	75.6	51	24.4	7.3
>High School	224	89.8	27	10.2	4.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health Health Insurance Coverage, Aged 18 and Older *Jefferson, Lewis, St. Lawrence*

Demographic		Yes		No	
Groups	n¹	%2	n	%	C.I. ³
Total	556	85.2	84	14.8	3.7
Gender					
Male	207	82.4	38	17.6	6.2
Female	349	88.2	46	11.8	4.0
Age					
18-34	102	76.0	30	24.0	8.8
35-54	222	86.1	38	13.9	5.0
≥55	231	94.9	16	5.1	2.7
Education					
≤High School	281	80.2	54	19.8	6.1
>High School	274	90.9	29	9.1	3.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Reported Having a Routine Physical Exam in Past Two Years

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	558	83.3	82	16.7	4.2
Gender					
Male	196	75.7	50	24.3	7.1
Female	362	91.3	32	8.7	3.4
Age					
18-34	106	75.0	26	25.0	9.7
35-54	223	84.0	38	16.0	5.3
≥55	229	92.4	17	7.6	3.9
Education					
≤High School	296	83.6	42	16.4	6.0
>High School	261	83.5	39	16.5	5.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Time in Past 12 Months when Needed Medical Care but Could Not Get It *Jefferson, Lewis, St. Lawrence*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	36	4.9	604	95.1	1.8
Gender					
Male	12	3.7	233	96.3	2.2
Female	24	6.1	371	93.9	2.9
Age					
18-34	10	5.1	123	94.9	3.8
35-54	17	6.0	243	94.0	3.0
≥55	9	3.2	237	96.8	2.3
Education					
≤High School	17	4.8	319	95.2	2.8
>High School	19	4.9	283	95.1	2.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days *Jefferson, Lewis, St. Lawrence*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	480	78.3	163	21.7	
Gender	100	70.5	103	21.7	3.7
Male	196	81.0	50	19.0	5.8
Female	284	75.6	113	24.4	4.7
Age					
18-34	113	86.6	20	13.4	6.5
35-54	198	75.4	64	24.6	6.3
≥55	168	72.2	79	27.8	6.2
Education					
≤High School	226	71.2	112	28.8	5.9
>High School	254	86.9	49	13.1	4.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Physician Diagnosed Diabetes

Demographic		Yes		No	
Groups	n¹	% 2	n	%	C.I. ³
Total	73	10.2	569	89.8	2.7
Gender					
Male	29	10.9	216	89.1	4.3
Female	44	9.4	353	90.6	3.1
Age					
18-34	5	3.9	127	96.1	3.9
35-54	25	9.7	237	90.3	4.0
≥55	43	18.0	204	82.0	5.8
Education					
≤High School	45	12.0	292	88.0	3.8
>High School	28	8.3	275	91.7	3.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Lifetime Asthma Diagnosed by a Medical Professional

Demographic		Yes		No	
Groups	n¹	% ²	n	%	C.I. ³
Total	75	13.0	568	87.0	3.4
Gender					
Male	21	10.1	225	89.9	4.8
Female	54	16.0	343	84.0	4.7
Age					
18-34	20	15.0	113	85.0	6.9
35-54	31	13.1	231	86.9	5.2
≥55	24	10.5	223	89.5	4.9
Education					
≤High School	47	15.9	291	84.1	5.1
>High School	28	10.0	275	90.0	4.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Current Asthma Diagnosed by a Medical Professional

Demographic		Yes		No	
Groups	n¹	% 2	n	%	C.I. ³
Total	61	11.2	580	88.8	3.2
Gender					
Male	17	8.4	228	91.6	4.4
Female	44	14.2	352	85.8	4.6
Age					
18-34	17	14.1	116	85.9	6.9
35-54	27	10.8	234	89.2	4.6
≥55	17	8.3	229	91.7	4.7
Education					
≤High School	40	14.7	297	85.3	5.1
>High School	21	7.5	281	92.5	3.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Arthritis Diagnosed by a Medical Professional

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	254	35.4	384	64.6	4.4
Gender					
Male	96	35.3	148	64.7	7.0
Female	158	35.5	236	64.5	5.4
Age					
18-34	13	9.8	119	90.2	5.8
35-54	88	37.0	173	63.0	7.0
≥55	152	63.1	92	36.9	7.1
Education					
≤High School	147	39.4	189	60.6	6.4
>High School	107	31.3	193	68.7	6.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

Demographic Groups	n¹	Doctor Diagnosed	C.I. ³	n	Possible Arthritis (CJS Only) %	C.I.	n	No Arthritis or CJS %	C.I.
Total	254	35.4	4.4	73	11.8	3.4	310	52.8	4.8
Gender									
Male	96	35.3	7.0	31	13.9	5.8	117	50.8	7.6
Female	158	35.6	5.4	42	9.5	3.2	193	54.9	5.7
Age									
18-34	13	9.8	5.8	14	11.3	7.5	105	78.9	8.8
35-54	88	37.0	7.0	34	14.5	5.0	139	48.6	7.1
≥55	152	63.4	7.1	25	9.0	3.8	66	27.5	6.7
Education									
≤High School	147	39.5	6.4	36	11.1	4.9	152	49.4	6.8
>High School	107	31.3	6.1	36	12.2	4.6	157	56.5	6.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Ever Smoked 100 Cigarettes in Lifetime

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Groups	11	/0		/0	C.1.
Total	328	52.1	311	47.9	4.8
Gender					
Male	139	57.7	104	42.3	7.6
Female	189	46.4	207	53.6	5.8
Age					
18-34	56	44.8	76	55.2	10.2
35-54	136	52.0	125	48.0	7.1
≥55	136	60.9	109	39.1	7.0
Education					
≤High School	187	57.3	151	42.7	6.6
>High School	141	46.9	158	53.1	6.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Current Smoker

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	147	22.2	492	77.8	4.1
Gender					
Male	56	23.3	187	76.7	6.7
Female	91	21.1	305	78.9	4.8
Age					
18-34	35	28.0	97	72.0	9.4
35-54	74	23.1	187	76.9	5.5
≥55	38	14.4	207	85.6	5.2
Education					
≤High School	95	26.9	243	73.1	6.1
>High School	52	17.3	247	82.7	5.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Status of Cigarette Smoking

Demographic Groups	n¹	Everyday Smoker %	C.I. ³	n	Someday Smoker %	C.I.	n	Former or Never Smoked	C.I.
Total	120	17.9	3.8	27	4.4	2.1	492	77.8	4.1
Gender									
Male	44	17.8	6.1	12	5.5	3.6	187	76.7	6.7
Female	76	17.9	4.5	15	3.2	2.0	305	78.9	4.8
Age									
18-34	24	20.9	8.7	11	7.1	5.1	97	72.0	9.4
35-54	66	19.9	5.1	8	3.3	2.7	187	76.9	5.5
≥55	30	11.9	4.9	8	2.5	1.9	207	85.6	5.2
Education									
≤High School	81	23.5	5.9	14	3.4	2.2	243	73.1	6.1
>High School	39	11.8	4.5	13	5.5	3.7	247	82.7	5.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Home Smoking Policies

Demographic Groups	n¹	Not Allowed	C.I. ³	n	Allowed Some Places %	C.I.	n	Allowed Anywhere	C.I.
Total	375	63.9	4.5	80	12.2	3.0	188	23.8	3.8
Gender									
Male	146	64.9	7.0	28	11.0	4.4	72	24.1	6.1
Female	229	63.0	5.4	52	13.5	4.1	116	23.5	4.4
Age									
18-34	98	74.0	8.6	10	8.3	5.3	25	17.7	7.5
35-54	145	59.5	6.9	47	18.9	5.8	70	21.5	5.2
≥55	131	57.5	7.1	23	8.7	3.7	93	33.8	6.7
Education									
≤High School	173	57.9	6.5	43	12.7	4.4	122	29.4	5.6
>High School	200	70.2	6.1	37	11.9	4.2	66	17.9	5.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Home Smoking

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Groups	11	70	11	/0	C.1.
Total	169	24.1	474	75.9	4.0
Gender					
Male	61	23.0	185	77.0	6.2
Female	108	25.1	289	74.9	5.0
Age					
18-34	28	20.5	105	79.5	7.9
35-54	85	28.7	177	71.3	6.3
≥55	56	22.6	191	77.4	6.1
Education					
≤High School	105	28.1	233	71.9	5.7
>High School	64	19.9	239	80.1	5.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	90	18.0	540	82.0	4.2
Gender					
Male	61	28.5	176	71.5	7.4
Female	29	7.3	364	92.7	2.9
Age					
18-34	31	28.4	99	71.6	9.8
35-54	48	17.5	207	82.5	5.2
≥55	11	6.4	233	93.6	3.9
Education					
≤High School	49	20.3	279	79.7	6.5
>High School	41	15.7	259	84.3	5.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	29	5.2	606	94.8	2.8
Gender					
Male	16	8.1	224	91.9	5.2
Female	13	2.2	382	97.8	1.3
Age					
18-34	5	7.3	126	92.7	7.2
35-54	18	5.3	239	94.7	2.6
≥55	6	2.7	240	97.3	2.4
Education					
≤High School	16	5.5	316	94.5	4.3
>High School	13	5.0	288	95.0	3.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Overweight or Obese, BMI(Body Mass Index) 25 or More *Jefferson, Lewis, St. Lawrence*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	392	67.7	221	32.3	4.5
Gender					
Male	185	76.4	58	23.6	6.3
Female	207	58.1	163	41.9	5.9
Age					
18-34	75	63.9	48	36.1	9.7
35-54	170	74.3	82	25.7	5.8
≥55	147	64.1	90	35.9	7.1
Education					
≤High School	209	68.3	113	31.7	6.1
>High School	183	67.1	108	32.9	6.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+) *Jefferson, Lewis, St. Lawrence*

Demographic Groups	n¹	Underweight\ Normal %2	C.I. ³	n	Overweight %	C.I.	n	Obese %	C.I.
Total	221	32.3	4.5	221	37.7	4.8	171	30.0	4.7
Gender									
Male	58	23.6	6.3	111	44.5	7.6	74	32.0	7.5
Female	163	41.9	5.9	110	30.2	5.6	97	27.9	5.5
Age									
18-34	48	36.1	9.7	45	36.7	10.1	30	27.3	10.1
35-54	82	25.7	5.8	92	39.9	7.2	78	34.3	7.0
≥55	90	35.9	7.1	84	36.1	7.1	63	28.0	6.9
Education									
≤High School	113	31.7	6.1	106	32.2	6.3	103	36.1	7.1
>High School	108	32.9	6.5	115	43.6	7.1	68	23.5	5.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Ever Had Mammography Screening, Women Aged 40 Years and Older *Jefferson, Lewis, St. Lawrence*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Female	251	91.4	29	8.6	3.5
Age					
40-64	162	91.2	19	8.8	4.5
≥65	89	91.8	10	8.2	5.5
Education					
≤High School	132	89.5	20	10.5	4.9
>High School	119	93.5	9	6.5	5.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Female	215	79.5	63	20.5	5.2
Age					
40-64	138	78.0	42	22.0	6.8
≥65	77	82.4	21	17.6	7.7
Education					
≤High School	113	76.5	38	23.5	7.6
>High School	102	82.7	25	17.3	7.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	73	19.3	415	80.7	4.8
Gender					
Male	28	18.3	169	81.7	7.4
Female	45	20.5	246	79.5	6.1
Age					
18-34	44	36.4	88	63.6	9.9
35-54	28	8.9	227	91.1	3.8
55-64	1	1.7	100	98.3	3.4
Education					
≤High School	43	24.6	196	75.4	7.7
>High School	30	14.3	217	85.7	5.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Sexually Active, Age 18-64 Years Old

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	358	82.7	111	17.3	3.7
Gender					
Male	149	84.8	39	15.2	5.3
Female	209	80.3	72	19.7	5.0
Age					
18-34	115	90.9	16	9.1	5.0
35-54	197	82.5	47	17.5	5.4
55-64	46	59.8	48	40.2	11.2
Education					
≤High School	172	83.2	53	16.8	5.4
>High School	185	82.7	57	17.3	5.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger *Jefferson, Lewis, St. Lawrence*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	61	19.2	303	80.8	5.7
Gender					
Male	30	20.7	123	79.3	8.9
Female	31	17.5	180	82.5	6.5
Age					
18-34	38	33.6	76	66.4	10.7
35-54	23	8.8	178	91.2	4.0
55-64	0	0.0	49	100	0.0
Education					
≤High School	29	18.2	150	81.8	7.8
>High School	32	20.2	153	79.8	8.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

Demographic Groups	n¹	Use BC	C.I. ³	n	No BC at risk	C.I.	n	No BC	C.I.
Total	222	66.7	6.2	32	9.4	3.7	83	23.9	5.7
Gender									
Male	113	64.0	8.4	17	9.5	5.0	54	26.5	7.8
Female	109	71.6	8.4	15	9.1	5.0	29	19.3	7.6
Age									
18-34	90	69.6	9.4	14	10.4	5.9	27	19.9	8.3
35-59	132	63.5	7.9	18	8.2	4.1	56	28.3	7.6
Education									
≤High School	106	67.0	8.8	16	10.7	5.9	39	22.3	7.6
>High School	116	67.0	8.7	16	8.1	4.4	43	24.9	8.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Physician Diagnosed Heart Attack, Angina or Stroke

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	65	8.7	567	91.3	2.5
Gender					
Male	26	9.8	214	90.2	4.1
Female	39	7.6	353	92.4	2.6
Age					
18-64	28	4.1	460	95.9	1.8
≥65	37	31.4	106	68.6	9.5
Education					
≤High School	41	10.7	288	89.3	3.8
>High School	24	6.6	277	93.4	3.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina) *Jefferson, Lewis, St. Lawrence*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	53	7.1	577	92.9	2.2
Gender					
Male	22	8.1	217	91.9	3.7
Female	31	6.0	360	94.0	2.4
Age					
18-64	23	3.5	465	96.5	1.7
≥65	30	25.2	111	74.8	9.1
Education					
≤High School	32	8.4	296	91.6	3.4
>High School	21	5.7	279	94.3	2.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older *Jefferson, Lewis, St. Lawrence*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Male	89	63.6	55	36.4	8.9
Age					
40-64	57	56.4	47	43.6	10.9
≥65	32	83.9	8	16.1	11.3
Education					
≤High School	49	63.4	30	36.6	12.2
>High School	40	64.8	24	35.2	13.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Male	75	54.2	69	45.8	9.3
Age					
40-64	47	47.6	57	52.4	11.0
≥65	28	73.2	12	26.8	15.5
Education					
≤High School	43	58.4	36	41.6	12.4
>High School	32	50.5	32	49.5	13.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had a Digital Rectal Exam, Men Aged 40 and Older *Jefferson, Lewis, St. Lawrence*

Demographic Groups	n¹	Yes %°	n	No %	C.I. ³
Male	118	80.0	33	20.0	7.0
Age					
40-64	79	74.7	29	25.3	9.1
≥65	39	94.9	4	5.1	5.3
Education					
≤High School	63	78.0	21	22.0	9.5
>High School	54	82.0	12	18.0	10.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Ever Used a Blood Stool Test at Home, Aged 50 and Older

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Total	149	50.5	150	49.5	6.8
Gender					
Male	49	51.1	47	48.9	11.5
Female	100	50.0	103	50.0	7.8
Age					
50-64	72	46.3	88	53.7	9.3
≥65	77	56.0	62	44.0	9.6
Education					
≤High School	72	42.0	98	58.0	8.8
>High School	77	60.5	52	39.5	9.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older *Jefferson, Lewis, St. Lawrence*

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Total	61	21.8	236	78.2	5.7
Gender					
Male	22	23.5	73	76.5	9.9
Female	39	20.4	163	79.6	6.5
Age					
50-64	28	17.9	132	82.1	7.2
≥65	33	27.0	104	73.0	9.0
Education					
≤High School	30	16.3	139	83.7	6.2
>High School	31	28.3	97	71.7	9.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older *Jefferson, Lewis, St. Lawrence*

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Total	84	29.4	213	70.6	6.3
Gender					
Male	29	31.6	66	68.4	10.9
Female	55	27.5	147	72.5	7.0
Age					
50-64	38	25.4	122	74.6	8.4
≥65	46	34.6	91	65.4	9.4
Education					
≤High School	39	23.7	130	76.3	7.7
>High School	45	36.0	83	64.0	10.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	133	45.7	166	54.3	6.8
Gender					
Male	46	48.9	50	51.1	11.5
Female	87	43.0	116	57.0	7.7
Age					
50-64	63	40.1	97	59.9	9.2
≥65	70	53.1	69	46.9	9.6
Education					
≤High School	71	40.6	99	59.4	8.7
>High School	62	51.7	67	48.3	10.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older *Jefferson, Lewis, St. Lawrence*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	114	40.5	180	59.5	6.8
Gender					
Male	40	43.9	55	56.1	11.6
Female	74	37.6	125	62.4	7.7
Age					
50-64	54	36.3	106	63.7	9.2
≥65	60	46.2	74	53.8	9.9
Education					
≤High School	63	37.1	104	62.9	8.7
>High School	51	44.5	76	55.5	10.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	143	49.6	158	50.4	6.7
Gender					
Male	50	54.3	46	45.7	11.4
Female	93	45.7	112	54.3	7.8
Age					
50-64	67	43.2	93	56.8	9.3
≥65	76	57.9	65	42.1	9.3
Education					
≤High School	78	45.3	93	54.7	8.9
>High School	65	54.6	65	45.4	10.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval