### Expanded BRFSS, 2003

### Onondaga Locality Tables

#### TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days	
Core 2: Health Care Access	
Core 3: Exercise	
Core 5: Asthma	
Core 6: Arthritis	
Core 7: Tobacco Use	
Core 8: Tobacco ETS	17
Core 9: Alcohol Consumption	
Core 10: Weight Status	
Core 11: Mammography	23
Core 12: Sexual Behavior	
Core 13: Family Planning	28
Core 14: Cardiovascular Disease	29
Core 15: Prostate Cancer Screening	
Core 16: Colorectal Cancer Screening.	37

New York State Department Of Health

#### **Self-rated Health Status**

Demographic Groups	n¹	Fair or Poor	n	Good to Excellent	C.I. <sup>3</sup>
Total	89	11.9	557	88.1	3.0
Gender					
Male	37	13.1	206	86.9	5.4
Female	52	10.7	351	89.3	3.1
Age					
18-34	12	9.1	129	90.9	7.3
35-54	26	7.2	232	92.8	3.0
≥55	50	20.4	184	79.6	5.6
Education					
≤High School	55	18.3	180	81.7	5.0
>High School	34	8.6	374	91.4	3.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Reported Number of Physical Distress Days in the Past Month** Onondaga

Demographic Groups	n¹	≥14 days %2	n	< 14 days %	C.I. <sup>3</sup>
Total	56	7.5	578	92.5	2.1
Gender					
Male	22	7.9	217	92.1	3.6
Female	34	7.1	361	92.9	2.5
Age					
18-34	8	3.5	133	96.5	2.6
35-54	25	8.7	230	91.3	3.8
≥55	23	10.3	203	89.7	4.3
Education					
≤High School	23	9.0	204	91.0	4.0
>High School	33	6.8	371	93.2	2.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### **Reported Number of Mental Distress Days in the Past Month** Onondaga

Demographic Groups	n¹	≥14 days %2	n	< 14 days %	C.I. <sup>3</sup>
Total	62	8.9	573	91.1	2.4
Gender					
Male	14	5.1	223	94.9	2.8
Female	48	12.2	350	87.8	3.7
Age					
18-34	25	14.4	113	85.6	6.1
35-54	23	7.7	232	92.3	3.4
≥55	13	5.1	217	94.9	3.0
Education					
≤High School	29	10.8	202	89.2	4.2
>High School	33	7.9	369	92.1	3.0

<sup>&</sup>lt;sup>1</sup> Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup> Weighted Percent<sup>3</sup> 95% Confidence Interval

### Expanded BRFSS, 2003 - New York State Department of Health **Health Insurance Coverage, 18-64 Years of Age** Onondaga

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	436	85.3	57	14.7	4.1
Gender					
Male	157	80.3	28	19.7	7.3
Female	279	90.0	29	10.0	3.8
Age					
18-34	109	74.4	31	25.6	9.1
35-54	233	89.9	24	10.1	4.2
55-64	94	97.1	2	2.9	3.9
Education					
≤High School	141	80.0	26	20.0	7.8
>High School	292	87.6	31	12.4	4.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### Expanded BRFSS, 2003 - New York State Department of Health Health Insurance Coverage, Aged 18 and Older Onondaga

Demographic	n¹	Yes	<b>n</b>	No	C.I. <sup>3</sup>
Groups	11	70	n	%	C.1.
Total	584	87.8	60	12.2	3.4
Gender					
Male	211	82.8	30	17.2	6.3
Female	373	92.1	30	7.9	3.0
Age					
18-34	109	74.4	31	25.6	9.1
35-54	233	89.9	24	10.1	4.2
≥55	230	97.7	4	2.3	2.3
Education					
≤High School	205	83.9	29	16.1	6.0
>High School	376	89.7	31	10.3	4.1

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

# Reported Having a Routine Physical Exam in Past Two Years *Onondaga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
•					
Total	580	87.0	62	13.0	3.7
Gender					
Male	200	77.9	42	22.1	6.9
Female	380	95.2	20	4.8	2.3
Age					
18-34	119	78.6	19	21.4	9.7
35-54	221	84.7	36	15.3	5.0
≥55	227	97.2	7	2.8	2.8
Education					
≤High School	210	87.3	25	12.7	5.3
>High School	368	87.2	36	12.8	4.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Time in Past 12 Months when Needed Medical Care but Could Not Get It *Onondaga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	31	5.4	615	94.6	2.5
Gender					
Male	12	6.2	231	93.8	4.5
Female	19	4.7	384	95.3	2.4
Age					
18-34	14	11.6	127	88.4	7.3
35-54	15	4.9	243	95.1	2.6
≥55	2	0.4	232	99.6	0.6
Education					
≤High School	12	4.7	223	95.3	2.8
>High School	19	5.9	389	94.1	3.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days *Onondaga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	514	80.4	129	19.6	3.8
Gender					
Male	192	77.8	49	22.2	6.7
Female	322	82.7	80	17.3	3.9
Age					
18-34	117	80.6	23	19.4	9.1
35-54	215	85.4	43	14.6	4.6
≥55	172	74.1	60	25.9	6.2
Education					
≤High School	165	73.5	69	26.5	6.3
>High School	346	83.8	60	16.2	4.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### **Physician Diagnosed Diabetes**

,				
n¹	Yes	n	No %	C.I. <sup>3</sup>
48	6.2	597	93.8	1.9
29	8.4	214	91.6	3.3
19	4.3	383	95.7	2.0
1	0.4	139	99.6	0.8
10	3.8	248	96.2	2.5
37	14.9	197	85.1	4.8
28	10.4	207	89.6	4.1
20	4.1	387	95.9	1.9
	1 10 37 28	n¹     %²       48     6.2       29     8.4       19     4.3       1     0.4       10     3.8       37     14.9       28     10.4	n¹         %²         n           48         6.2         597           29         8.4         214           19         4.3         383           1         0.4         139           10         3.8         248           37         14.9         197           28         10.4         207	n¹         %o²         n         %o           48         6.2         597         93.8           29         8.4         214         91.6           19         4.3         383         95.7           1         0.4         139         99.6           10         3.8         248         96.2           37         14.9         197         85.1           28         10.4         207         89.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health **Lifetime Asthma Diagnosed by a Medical Professional** *Onondaga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	82	11.6	563	88.4	2.7
Gender					
Male	22	8.3	221	91.7	3.7
Female	60	14.4	342	85.6	3.8
Age					
18-34	20	12.5	121	87.5	5.9
35-54	36	12.5	222	87.5	4.2
≥55	23	9.1	210	90.9	3.9
Education					
≤High School	31	11.8	203	88.2	4.4
>High School	51	11.6	357	88.4	3.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Expanded BRFSS, 2003 - New York State Department of Health **Current Asthma Diagnosed by a Medical Professional** Onondaga

Demographic		Yes		No	
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	61	8.8	582	91.2	2.4
Gender					
Male	15	5.7	228	94.3	3.1
Female	46	11.5	354	88.5	3.5
Age					
18-34	15	9.9	125	90.1	5.3
35-54	28	9.9	230	90.1	3.9
≥55	16	6.4	216	93.6	3.2
Education					
≤High School	23	8.0	210	92.0	3.4
>High School	38	9.3	369	90.7	3.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health **Arthritis Diagnosed by a Medical Professional**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	211	28.9	432	71.1	3.8
Gender					
Male	76	26.1	166	73.9	5.9
Female	135	31.3	266	68.7	4.9
Age					
18-34	17	10.9	122	89.1	5.8
35-54	58	21.5	200	78.5	5.4
≥55	130	54.2	103	45.8	7.1
Education					
≤High School	94	36.4	140	63.6	6.9
>High School	117	25.2	289	74.8	4.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

## Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

Demographic Groups	n¹	Doctor Diagnosed	C.I. <sup>3</sup>	n	Possible Arthritis (CJS Only)	C.I.	n	No Arthritis or CJS %	C.I.
Total	211	29.0	3.8	96	15.4	3.2	335	55.6	4.4
Gender									
Male	76	26.5	6.0	38	16.3	5.3	127	57.3	7.1
Female	135	31.3	4.9	58	14.6	3.8	208	54.1	5.4
Age									
18-34	17	11.2	5.9	22	16.2	7.0	99	72.7	8.6
35-54	58	21.5	5.4	43	17.6	5.3	157	60.9	6.6
≥55	130	54.2	7.1	30	12.6	4.7	73	33.2	6.9
Education									
≤High School	94	36.4	6.9	27	12.3	4.8	113	51.3	7.3
>High School	117	25.5	4.6	68	16.8	4.2	220	57.7	5.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Ever Smoked 100 Cigarettes in Lifetime**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	361	54.3	284	45.7	4.5
Gender					
Male	152	58.7	90	41.3	7.4
Female	209	50.4	194	49.6	5.4
Age					
18-34	74	47.2	67	52.8	9.8
35-54	141	55.8	116	44.2	6.7
≥55	137	58.7	97	41.3	6.9
Education					
≤High School	151	65.8	84	34.2	6.8
>High School	210	48.8	197	51.2	5.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Current Smoker**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	163	25.8	481	74.2	4.0
Gender					
Male	65	28.5	176	71.5	6.7
Female	98	23.4	305	76.6	4.6
Age					
18-34	55	36.4	85	63.6	9.4
35-54	73	28.1	184	71.9	6.1
≥55	32	13.1	202	86.9	4.7
Education					
≤High School	81	37.7	154	62.3	7.3
>High School	82	19.7	324	80.3	4.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### **Status of Cigarette Smoking**

Demographic Groups	n¹	Everyday Smoker %2	C.I. <sup>3</sup>	n	Someday Smoker	C.I.	n	Former or Never Smoked %	C.I.
Total	122	19.8	3.6	41	6.0	2.1	481	74.2	4.0
Gender									
Male	51	22.4	6.2	14	6.1	3.5	176	71.5	6.7
Female	71	17.5	4.1	27	5.9	2.5	305	76.6	4.6
Age									
18-34	41	27.5	8.7	14	8.9	5.2	85	63.6	9.4
35-54	57	22.3	5.6	16	5.8	3.3	184	71.9	6.1
≥55	22	10.0	4.3	10	3.1	2.1	202	86.9	4.7
Education									
≤High School	62	30.1	7.0	19	7.6	4.1	154	62.3	7.3
>High School	60	14.5	4.0	22	5.2	2.4	324	80.3	4.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### **Home Smoking Policies**

Demographic Groups	n¹	Not Allowed	C.I. <sup>3</sup>	n	Allowed Some Places %	C.I.	n	Allowed Anywhere	C.I.
Total	414	64.4	4.4	100	15.1	3.3	131	20.5	3.8
Gender									
Male	150	60.9	7.3	32	13.8	5.5	61	25.2	6.6
Female	264	67.4	5.1	68	16.3	4.0	70	16.3	4.0
Age									
18-34	88	61.1	9.9	26	18.0	8.0	27	20.9	8.6
35-54	171	68.2	6.2	47	17.4	5.0	40	14.4	4.6
≥55	148	63.2	7.0	24	9.6	4.0	61	27.2	6.6
Education									
≤High School	116	51.8	7.3	50	20.5	5.7	68	27.8	6.6
>High School	296	70.9	5.5	50	12.5	4.1	62	16.5	4.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Expanded BRFSS, 2003 - New York State Department of Health **Home Smoking** Onondaga

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	161	26.9	484	73.1	4.2
Gender					
Male	68	31.0	174	69.0	7.2
Female	93	23.2	310	76.8	4.7
Age					
18-34	49	38.2	91	61.8	10.0
35-54	69	26.3	189	73.7	5.9
≥55	40	17.4	194	82.6	5.5
Education					
≤High School	84	38.2	151	61.8	7.3
>High School	77	21.2	330	78.8	5.1

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

## At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	91	18.5	548	81.5	4.0
Gender					
Male	60	29.6	180	70.4	7.2
Female	31	8.7	368	91.3	3.4
Age					
18-34	31	30.0	108	70.0	9.9
35-54	46	20.1	209	79.9	5.6
≥55	13	5.9	219	94.1	3.6
Education					
≤High School	36	20.4	196	79.6	6.8
>High School	55	17.6	351	82.4	5.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

## At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)

Demographic		Yes		No	
Groups	n¹	%2	n	%	C.I. <sup>3</sup>
Total	31	6.3	607	93.7	2.6
Gender					
Male	12	7.2	228	92.8	4.6
Female	19	5.4	379	94.6	2.8
Age					
18-34	11	11.5	128	88.5	7.3
35-54	12	5.0	242	95.0	3.0
≥55	7	2.5	225	97.5	2.0
Education					
≤High School	9	5.8	223	94.2	4.3
>High School	21	6.2	383	93.8	3.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health Overweight or Obese, BMI(Body Mass Index) 25 or More Onondaga

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	361	59.5	250	40.5	4.6
Gender					
Male	173	70.2	64	29.8	7.2
Female	188	49.7	186	50.3	5.6
Age					
18-34	59	46.5	75	53.5	10.1
35-54	140	59.2	108	40.8	6.7
≥55	158	72.7	64	27.3	6.4
Education					
≤High School	140	62.9	80	37.1	7.4
>High School	220	57.8	169	42.2	5.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+) *Onondaga*

Demographic Groups	n¹	Underweight\ Normal	C.I. <sup>3</sup>	n	Overweight %	C.I.	n	Obese %	C.I.
Total	250	40.5	4.6	235	39.2	4.5	126	20.3	3.6
Gender									
Male	64	29.8	7.2	121	48.1	7.4	52	22.1	5.9
Female	186	50.3	5.6	114	31.0	5.2	74	18.7	4.4
Age									
18-34	75	53.5	10.1	32	27.7	9.2	27	18.8	7.6
35-54	108	40.8	6.7	96	41.1	6.8	44	18.2	5.3
≥55	64	27.3	6.4	104	48.2	7.3	54	24.5	6.2
Education									
≤High School	80	37.1	7.4	90	41.1	7.4	50	21.8	6.1
>High School	169	42.2	5.8	144	38.1	5.6	76	19.7	4.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# **Ever Had Mammography Screening, Women Aged 40 Years and Older** *Onondaga*

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Female	246	96.0	10	4.0	2.6
Age					
40-64	165	96.3	6	3.7	3.1
≥65	81	95.4	4	4.6	4.9
Education					
≤High School	100	97.4	3	2.6	3.0
>High School	146	95.1	7	4.9	3.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Female	220	86.5	35	13.5	4.6
Age					
40-64	148	86.4	22	13.6	5.9
≥65	72	86.8	13	13.2	7.3
Education					
≤High School	91	90.2	11	9.8	5.9
>High School	129	84.3	24	15.7	6.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	79	14.9	401	85.1	3.6
Gender					
Male	15	7.6	168	92.4	4.4
Female	64	21.8	233	78.2	5.4
Age					
18-34	45	27.0	94	73.0	8.2
35-54	29	8.5	219	91.5	3.3
55-64	5	4.4	88	95.6	3.9
Education					
≤High School	38	22.5	122	77.5	7.5
>High School	41	11.6	276	88.4	3.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health **Sexually Active, Age 18-64 Years Old** *Onondaga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	354	84.5	93	15.5	3.6
Gender					
Male	138	85.8	31	14.2	5.7
Female	216	83.3	62	16.7	4.4
Age					
18-34	115	87.8	16	12.2	6.6
35-54	191	88.5	38	11.5	3.9
55-64	48	64.6	39	35.4	10.7
Education					
≤High School	112	77.8	40	22.2	7.9
>High School	241	87.5	53	12.5	3.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger** Onondaga

Demographic	1	Yes		No	CI3
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	87	23.8	279	76.2	5.2
Gender					
Male	34	23.7	110	76.3	8.3
Female	53	23.8	169	76.2	6.4
Age					
18-34	43	37.5	78	62.5	10.3
35-54	40	16.1	157	83.9	5.2
55-64	4	7.4	44	92.6	7.5
Education					
≤High School	31	27.4	86	72.6	9.6
>High School	55	22.0	192	78.0	6.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

## Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

Demographic	n¹	Use BC	C.I. <sup>3</sup>		No BC at risk	C.I.	_	No BC	C.I.
Groups	111-	70-	C.I.	n	%	C.I.	n	70	C.I.
Total	223	69.6	5.7	35	10.8	3.7	72	19.6	5.0
Gender									
Male	109	71.8	8.0	13	7.6	4.3	36	20.6	7.4
Female	114	66.5	7.9	22	15.2	6.2	36	18.3	6.3
Age									
18-34	91	67.1	9.2	18	12.7	5.9	29	20.2	8.1
35-59	132	72.1	6.8	17	8.9	4.3	43	19.0	6.0
Education									
≤High School	69	68.3	11.1	8	8.2	5.9	20	23.6	10.6
>High School	153	69.9	6.7	27	11.9	4.6	52	18.1	5.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health **Physician Diagnosed Heart Attack, Angina or Stroke** *Onondaga*

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	46	6.4	582	93.6	2.0
Gender					
Male	27	8.7	206	91.3	3.5
Female	19	4.5	376	95.5	2.2
Age					
18-64	16	3.0	463	97.0	1.6
≥65	29	21.0	108	79.0	7.4
Education					
≤High School	19	7.3	206	92.7	3.5
>High School	27	6.0	374	94.0	2.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina) *Onondaga*

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	38	5.4	590	94.6	1.8
Gender					
Male	22	7.2	211	92.8	3.2
Female	16	3.9	379	96.1	2.0
Age					
18-64	15	2.6	464	97.4	1.5
≥65	22	16.9	115	83.1	6.9
Education					
≤High School	12	4.6	213	95.4	2.9
>High School	26	5.9	375	94.1	2.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# **Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older** *Onondaga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	109	68.0	45	32.0	8.2
Age					
40-64	61	57.9	42	42.1	10.4
≥65	48	93.8	3	6.2	6.9
Education					
≤High School	37	69.7	12	30.3	14.7
>High School	72	69.0	31	31.0	9.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

## Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	95	59.1	58	40.9	8.6
Age					
40-64	52	49.2	50	50.8	10.5
≥65	43	84.5	8	15.5	10.3
Education					
≤High School	33	63.5	15	36.5	15.2
>High School	62	58.7	41	41.3	10.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health Ever Had a Digital Rectal Exam, Men Aged 40 and Older Onondaga

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Male	149	94.3	11	5.7	3.5
Age					
40-64	100	93.9	9	6.1	4.1
≥65	49	95.4	2	4.6	6.2
Education					
≤High School	45	91.9	5	8.1	7.3
>High School	102	95.3	6	4.7	3.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Ever Used a Blood Stool Test at Home, Aged 50 and Older *Onondaga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	164	56.3	125	43.7	6.4
Gender					
Male	63	55.0	50	45.0	10.2
Female	101	57.4	75	42.6	8.1
Age					
50-64	82	52.0	70	48.0	8.9
≥65	82	61.3	55	38.7	9.0
Education					
≤High School	68	58.9	52	41.1	9.7
>High School	96	55.2	72	44.8	8.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older *Onondaga*

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	71	24.2	214	75.8	5.4
Gender					
Male	31	23.8	81	76.2	8.2
Female	40	24.5	133	75.5	7.1
Age					
50-64	34	21.1	117	78.9	6.9
≥65	37	27.8	97	72.2	8.3
Education					
≤High School	28	24.7	90	75.3	8.6
>High School	43	24.1	123	75.9	6.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older *Onondaga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	106	36.6	179	63.4	6.1
Gender					
Male	44	38.4	68	61.6	9.8
Female	62	35.1	111	64.9	7.7
Age					
50-64	55	34.4	96	65.6	8.3
≥65	51	39.2	83	60.8	9.0
Education					
≤High School	44	38.4	74	61.6	9.7
>High School	62	35.8	104	64.2	7.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older *Onondaga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	160	57.3	128	42.7	6.3
Gender					
Male	65	61.9	45	38.1	10.0
Female	95	53.6	83	46.4	8.1
Age					
50-64	75	52.1	76	47.9	8.9
≥65	85	63.1	52	36.9	8.9
Education					
≤High School	64	56.2	54	43.8	9.8
>High School	96	58.5	73	41.5	8.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older *Onondaga*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	151	54.7	135	45.3	6.4
Gender					
Male	62	60.0	47	40.0	10.1
Female	89	50.7	88	49.3	8.1
Age					
50-64	71	49.9	80	50.1	9.0
≥65	80	60.4	55	39.6	9.0
Education					
≤High School	60	53.6	57	46.4	9.9
>High School	91	56.0	77	44.0	8.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	186	65.2	105	34.8	6.1
Gender					
Male	79	70.7	34	29.3	9.3
Female	107	60.8	71	39.2	7.9
Age					
50-64	87	58.7	67	41.3	8.7
≥65	99	72.9	38	27.1	8.2
Education					
≤High School	77	64.7	44	35.3	9.4
>High School	109	66.2	60	33.8	8.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval