#### Expanded BRFSS, 2003

#### Cortland, Tompkins Locality Tables

#### TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days	
Core 2: Health Care Access	4
Core 3: Exercise	8
Core 4: Diabetes	
Core 5: Asthma	10
Core 6: Arthritis	12
Core 7: Tobacco Use	14
Core 8: Tobacco ETS	17
Core 9: Alcohol Consumption	19
Core 9: Alcohol Consumption	21
Core 11: Mammography	23
Core 12: Sexual Behavior	
Core 13: Family Planning	28
Core 14: Cardiovascular Disease	29
Core 15: Prostate Cancer Screening	
Core 16: Colorectal Cancer Screening	

New York State Department Of Health

#### **Self-rated Health Status**

Demographic Groups	n¹	Fair or Poor	n	Good to Excellent	C.I. <sup>3</sup>
Total	78	9.8	565	90.2	2.6
Gender					
Male	28	10.0	234	90.0	4.2
Female	50	9.7	331	90.3	3.1
Age					
18-34	8	5.2	168	94.8	4.0
35-54	18	5.7	228	94.3	2.8
≥55	50	22.5	166	77.5	6.1
Education					
≤High School	49	18.4	163	81.6	5.5
>High School	29	6.0	402	94.0	2.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### **Reported Number of Physical Distress Days in the Past Month** Cortland, Tompkins

Demographic Groups	n¹	≥14 days %2	n	< 14 days %	C.I. <sup>3</sup>
Total	58	7.4	570	92.6	2.1
Gender					
Male	11	4.0	244	96.0	2.7
Female	47	10.5	326	89.5	3.2
Age					
18-34	8	4.2	166	95.8	3.1
35-54	21	8.0	221	92.0	3.4
≥55	28	12.2	178	87.8	4.7
Education					
≤High School	34	13.1	171	86.9	4.7
>High School	24	4.8	399	95.2	2.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### **Reported Number of Mental Distress Days in the Past Month** Cortland, Tompkins

Demographic Groups	n¹	≥14 days	n	< 14 days %	C.I. <sup>3</sup>
Total	50	6.9	581	93.1	2.2
Gender					
Male	15	4.1	243	95.9	2.4
Female	35	9.5	338	90.5	3.6
Age					
18-34	13	6.8	162	93.2	3.9
35-54	23	8.2	218	91.8	3.6
≥55	13	5.2	196	94.8	3.2
Education					
≤High School	26	10.8	184	89.2	4.7
>High School	24	5.1	397	94.9	2.4

<sup>&</sup>lt;sup>1</sup> Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup> Weighted Percent<sup>3</sup> 95% Confidence Interval

#### Expanded BRFSS, 2003 - New York State Department of Health **Health Insurance Coverage, 18-64 Years of Age** Cortland, Tompkins

Demographic		Yes		No	G T 1
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	464	86.6	57	13.4	3.8
Gender					
Male	193	84.4	32	15.6	5.5
Female	271	88.7	25	11.3	5.2
Age					
18-34	143	81.2	32	18.8	6.6
35-54	228	92.2	18	7.8	3.7
55-64	93	93.0	7	7.0	5.5
Education					
≤High School	119	70.5	33	29.5	9.6
>High School	345	93.1	24	6.9	3.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### Expanded BRFSS, 2003 - New York State Department of Health Health Insurance Coverage, Aged 18 and Older Cortland, Tompkins

Demographic		Yes		No	
Groups	n¹	%2	n	%	C.I. <sup>3</sup>
Total	585	88.3	58	11.7	3.3
Gender					
Male	229	86.2	32	13.8	4.9
Female	356	90.2	26	9.8	4.5
Age					
18-34	143	81.2	32	18.8	6.6
35-54	228	92.2	18	7.8	3.7
≥55	208	96.0	8	4.0	2.9
Education					
≤High School	179	76.7	33	23.3	7.9
>High School	406	93.6	25	6.4	2.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### Reported Having a Routine Physical Exam in Past Two Years *Cortland, Tompkins*

Demographic		Yes		No	
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	550	84.7	87	15.3	3.4
Gender					
Male	196	75.9	62	24.1	6.0
Female	354	93.0	25	7.0	3.0
Age					
18-34	141	80.9	34	19.1	6.3
35-54	206	83.6	36	16.4	5.2
≥55	197	92.7	17	7.3	3.7
Education					
≤High School	184	85.7	25	14.3	5.8
>High School	366	84.3	62	15.7	4.1

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Time in Past 12 Months when Needed Medical Care but Could Not Get It *Cortland, Tompkins*

Demographic		Yes		No	CI3
Groups	n¹	%2	n	%	C.I. <sup>3</sup>
Total	34	4.7	609	95.3	1.8
Gender					
Male	10	4.1	252	95.9	2.7
Female	24	5.4	357	94.6	2.4
Age					
18-34	12	5.8	163	94.2	3.4
35-54	13	4.5	233	95.5	2.7
≥55	8	3.0	208	97.0	2.3
Education					
≤High School	19	8.9	192	91.1	4.4
>High School	15	2.9	417	97.1	1.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days *Cortland, Tompkins*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	505	79.9	138	20.1	3.5
Gender					
Male	209	82.1	53	17.9	4.9
Female	296	77.8	85	22.2	5.0
Age					
18-34	147	83.6	29	16.4	5.8
35-54	196	79.5	50	20.5	5.5
≥55	157	73.2	58	26.8	6.5
Education					
≤High School	132	62.9	79	37.1	7.7
>High School	373	87.5	59	12.5	3.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### **Physician Diagnosed Diabetes**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Groups	11	/0	11	/0	C.I.
Total	33	4.0	611	96.0	1.5
Gender					
Male	15	4.8	247	95.2	2.5
Female	18	3.3	364	96.7	1.6
Age					
18-34	1	0.4	175	99.6	0.8
35-54	9	3.0	237	97.0	2.0
≥55	23	11.7	193	88.3	4.9
Education					
≤High School	19	7.4	193	92.6	3.6
>High School	14	2.4	418	97.6	1.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### **Lifetime Asthma Diagnosed by a Medical Professional**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	88	13.9	551	86.1	3.2
Gender					
Male	25	10.2	234	89.8	4.3
Female	63	17.2	317	82.8	4.7
Age					
18-34	27	15.4	148	84.6	6.0
35-54	34	14.6	210	85.4	4.9
≥55	25	9.6	189	90.4	3.9
Education					
≤High School	34	14.7	177	85.3	5.6
>High School	54	13.5	374	86.5	3.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Current Asthma Diagnosed by a Medical Professional**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	59	8.6	575	91.4	2.5
Gender					
Male	15	5.9	242	94.1	3.3
Female	44	11.0	333	89.0	3.6
Age					
18-34	15	7.9	158	92.1	4.2
35-54	23	10.2	218	89.8	4.3
≥55	19	7.2	195	92.8	3.4
Education					
≤High School	28	11.5	182	88.5	4.8
>High School	31	7.2	393	92.8	2.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Arthritis Diagnosed by a Medical Professional**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	190	23.8	452	76.2	3.7
Gender					
Male	59	18.3	203	81.7	4.9
Female	131	28.9	249	71.1	5.4
Age					
18-34	16	10.5	160	89.5	5.6
35-54	58	22.8	187	77.2	5.6
≥55	114	48.9	101	51.1	7.4
Education					
≤High School	90	37.8	122	62.2	7.9
>High School	100	17.4	330	82.6	3.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

Demographic Groups	n¹	Doctor Diagnosed %2	C.I. <sup>3</sup>	n	Possible Arthritis (CJS Only) %	C.I.	n	No Arthritis or CJS %	C.I.
Total	190	23.9	3.7	90	14.3	3.2	359	61.8	4.4
Gender									
Male	59	18.4	5.0	40	15.3	4.9	162	66.3	6.4
Female	131	29.0	5.5	50	13.4	4.3	197	57.6	6.0
Age									
18-34	16	10.5	5.6	33	17.2	6.0	127	72.4	7.5
35-54	58	22.8	5.6	39	14.5	4.8	147	62.7	6.5
≥55	114	49.5	7.4	17	8.4	4.0	82	42.0	7.3
Education									
≤High School	90	38.0	7.9	25	12.9	5.2	96	49.2	8.0
>High School	100	17.5	3.7	65	15.0	4.1	263	67.5	5.1

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Ever Smoked 100 Cigarettes in Lifetime**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	280	41.0	362	59.0	4.5
Gender					
Male	129	44.8	133	55.2	7.0
Female	151	37.4	229	62.6	5.8
Age					
18-34	56	32.8	119	67.2	7.8
35-54	117	47.2	129	52.8	6.8
≥55	104	48.1	111	51.9	7.4
Education					
≤High School	129	59.0	83	41.0	8.0
>High School	151	32.8	279	67.2	5.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Current Smoker**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	128	20.4	514	79.6	3.8
Gender					
Male	54	21.7	208	78.3	6.1
Female	74	19.3	306	80.7	4.6
Age					
18-34	41	23.2	134	76.8	7.0
35-54	57	23.3	189	76.7	5.8
≥55	28	11.7	187	88.3	4.5
Education					
≤High School	73	34.9	139	65.1	7.6
>High School	55	13.9	375	86.1	4.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### **Status of Cigarette Smoking**

Demographic Groups	n¹	Everyday Smoker %	C.I. <sup>3</sup>	n	Someday Smoker	C.I.	n	Former or Never Smoked %	C.I.
Total	95	14.6	3.1	33	5.8	2.5	514	79.6	3.8
Gender									
Male	42	15.0	4.8	12	6.7	4.5	208	78.3	6.1
Female	53	14.3	4.1	21	5.0	2.4	306	80.7	4.6
Age									
18-34	28	15.1	5.6	13	8.1	5.1	134	76.8	7.0
35-54	45	18.3	5.3	12	5.1	3.1	189	76.7	5.8
≥55	21	9.4	4.2	7	2.4	1.8	187	88.3	4.5
Education									
≤High School	57	27.2	7.0	16	7.7	4.2	139	65.1	7.6
>High School	38	8.9	3.1	17	4.9	3.2	375	86.1	4.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### **Home Smoking Policies**

Demographic Groups	n¹	Not Allowed	C.I. <sup>3</sup>	n	Allowed Some Places	C.I.	n	Allowed Anywhere	C.I.
Total	440	70.5	4.3	63	10.6	3.2	135	19.0	3.5
Gender									
Male	181	70.7	6.6	18	8.0	4.5	61	21.3	5.6
Female	259	70.2	5.7	45	13.0	4.6	74	16.7	4.2
Age									
18-34	128	71.9	7.8	16	11.0	6.2	32	17.1	6.1
35-54	175	71.9	6.1	27	11.3	4.4	44	16.7	5.0
≥55	136	67.6	6.8	18	7.9	3.7	56	24.5	6.2
Education									
≤High School	106	51.9	8.1	37	18.9	7.0	66	29.2	7.1
>High School	334	78.8	4.8	26	6.8	3.4	69	14.4	3.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Home Smoking**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	125	20.6	519	79.4	4.1
Gender					
Male	56	22.4	206	77.6	6.4
Female	69	19.0	313	81.0	5.1
Age					
18-34	35	22.8	141	77.2	7.7
35-54	52	20.5	194	79.5	5.4
≥55	34	15.3	182	84.7	5.2
Education					
≤High School	72	34.9	140	65.1	7.8
>High School	53	14.2	379	85.8	4.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)

Demographic		Yes		No	
Groups	n¹	%2	n	%	C.I. <sup>3</sup>
Total	83	18.0	554	82.0	4.1
Gender					
Male	58	27.4	200	72.6	6.9
Female	25	9.2	354	90.8	4.1
Age					
18-34	47	29.8	128	70.2	7.9
35-54	24	10.2	219	89.8	4.3
≥55	11	5.2	203	94.8	3.2
Education					
≤High School	19	9.0	191	91.0	4.3
>High School	64	22.0	363	78.0	5.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)

Demographic		Yes		No	
Groups	n¹	<b>0</b> ∕₀²	n	%	C.I. <sup>3</sup>
Total	39	7.4	599	92.6	2.9
Gender					
Male	18	9.1	241	90.9	5.2
Female	21	5.9	358	94.1	2.9
Age					
18-34	13	9.9	161	90.1	5.9
35-54	14	5.9	230	94.1	3.2
≥55	11	4.6	203	95.4	2.9
Education					
≤High School	12	5.0	197	95.0	3.1
>High School	27	8.5	402	91.5	4.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health Overweight or Obese, BMI(Body Mass Index) 25 or More Cortland, Tompkins

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	326	51.6	294	48.4	4.7
Gender					
Male	163	61.6	99	38.4	6.8
Female	163	41.7	195	58.3	6.2
Age					
18-34	67	43.2	103	56.8	8.6
35-54	134	59.3	104	40.7	6.7
≥55	125	58.3	85	41.7	7.4
Education					
≤High School	125	61.8	77	38.2	7.9
>High School	201	47.2	217	52.8	5.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+) *Cortland, Tompkins*

Demographic Groups	n¹	Underweight\ Normal %2	C.I. <sup>3</sup>	n	Overweight %	C.I.	n	Obese %	C.I.
Total	294	48.4	4.7	204	32.0	4.4	122	19.6	3.8
Gender									
Male	99	38.4	6.8	112	41.0	6.9	51	20.5	6.0
Female	195	58.3	6.2	92	23.0	5.3	71	18.7	4.6
Age									
18-34	103	56.8	8.6	42	26.6	7.8	25	16.6	6.8
35-54	104	40.7	6.7	77	34.8	6.7	57	24.5	6.0
≥55	85	41.7	7.4	85	38.9	7.2	40	19.4	5.9
Education									
≤High School	77	38.2	7.9	66	33.6	8.0	59	28.2	7.0
>High School	217	52.8	5.8	138	31.4	5.3	63	15.8	4.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Ever Had Mammography Screening, Women Aged 40 Years and Older** *Cortland, Tompkins*

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Female	224	90.0	26	10.0	4.0
Age					
40-64	149	88.7	20	11.3	5.2
≥65	75	93.1	6	6.9	5.5
Education					
≤High School	84	91.5	8	8.5	5.9
>High School	140	89.2	18	10.8	5.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Female	193	79.3	55	20.7	5.3
Age					
40-64	128	77.0	40	23.0	6.8
≥65	65	84.5	15	15.5	7.6
Education					
≤High School	70	78.5	21	21.5	8.8
>High School	123	79.7	34	20.3	6.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	60	15.2	451	84.8	4.2
Gender					
Male	13	8.7	210	91.3	5.0
Female	47	21.6	241	78.4	6.4
Age					
18-34	44	25.3	130	74.7	7.4
35-54	15	5.4	223	94.6	3.0
55-64	1	0.5	98	99.5	0.9
Education					
≤High School	20	14.0	131	86.0	6.5
>High School	40	15.6	320	84.4	5.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### Sexually Active, Age 18-64 Years Old

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	389	82.7	102	17.3	4.0
Gender					
Male	175	83.7	40	16.3	5.7
Female	214	81.7	62	18.3	5.5
Age					
18-34	141	82.2	32	17.8	6.5
35-54	190	88.2	36	11.8	4.1
55-64	58	71.2	34	28.8	9.5
Education					
≤High School	113	82.1	33	17.9	7.0
>High School	276	82.9	69	17.1	4.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger** Cortland, Tompkins

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	77	23.7	314	76.3	5.6
Gender	, ,	23.7	311	70.5	3.0
Male	39	25.6	138	74.4	8.4
Female	38	21.7	176	78.3	7.2
Age					
18-34	51	37.1	92	62.9	9.1
35-54	23	8.9	168	91.1	3.9
55-64	3	4.1	54	95.9	4.9
Education					
≤High School	13	11.0	103	89.0	6.4
>High School	64	28.9	211	71.1	7.1

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

Demographic Groups	n¹	Use BC	C.I. <sup>3</sup>	n	No BC at risk	C.I.	n	No BC	C.I.
Total	219	61.5	5.9	41	13.4	4.5	106	25.1	5.1
Gender									
Male	114	61.9	7.7	24	13.4	5.4	62	24.7	6.6
Female	105	61.0	9.2	17	13.4	7.4	44	25.6	8.0
Age									
18-34	108	61.8	8.2	19	14.0	6.3	46	24.2	7.0
35-59	111	60.9	7.4	22	12.4	5.1	60	26.7	6.6
Education									
≤High School	61	59.3	11.8	12	19.2	11.3	26	21.6	8.8
>High School	158	62.3	6.8	29	11.3	4.3	80	26.3	6.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### Physician Diagnosed Heart Attack, Angina or Stroke

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	44	5.3	592	94.7	1.7
Gender					
Male	23	6.7	237	93.3	2.9
Female	21	3.9	355	96.1	2.0
Age					
18-64	16	2.4	500	97.6	1.3
≥65	26	23.2	90	76.8	8.6
Education					
≤High School	24	8.6	187	91.4	4.0
>High School	20	3.8	405	96.2	1.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina) *Cortland, Tompkins*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	36	4.3	600	95.7	1.6
Gender					
Male	21	5.9	239	94.1	2.7
Female	15	2.7	361	97.3	1.7
Age					
18-64	14	2.1	502	97.9	1.3
≥65	20	17.2	96	82.8	7.7
Education					
≤High School	20	7.5	191	92.5	3.8
>High School	16	2.8	409	97.2	1.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# **Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older** *Cortland, Tompkins*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	91	61.0	62	39.0	8.3
Age					
40-64	61	53.1	57	46.9	9.7
≥65	30	86.1	5	13.9	11.8
Education					
≤High School	25	49.9	24	50.1	14.8
>High School	66	66.4	38	33.6	9.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	77	51.7	75	48.3	8.5
Age					
40-64	53	46.9	65	53.1	9.8
≥65	24	67.6	10	32.4	16.8
Education					
≤High School	21	41.1	28	58.9	14.5
>High School	56	57.0	47	43.0	10.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had a Digital Rectal Exam, Men Aged 40 and Older

Cortland, Tompkins

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	128	82.4	29	17.6	6.3
Age					
40-64	96	79.2	26	20.8	7.7
≥65	32	92.7	3	7.3	8.4
Education					
≤High School	36	70.6	15	29.4	13.2
>High School	92	88.3	14	11.7	6.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Ever Used a Blood Stool Test at Home, Aged 50 and Older *Cortland, Tompkins*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	125	44.4	154	55.6	6.4
Gender					
Male	51	44.0	60	56.0	9.9
Female	74	44.9	94	55.1	8.2
Age					
50-64	61	36.7	102	63.3	8.2
≥65	64	56.0	52	44.0	9.8
Education					
≤High School	48	42.8	61	57.2	10.1
>High School	77	45.4	93	54.6	8.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older *Cortland, Tompkins*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	52	19.9	224	80.1	5.3
Gender					
Male	26	24.5	85	75.5	8.7
Female	26	15.4	139	84.6	5.9
Age					
50-64	26	18.0	135	82.0	6.8
≥65	26	22.7	89	77.3	8.4
Education					
≤High School	17	14.7	90	85.3	7.1
>High School	35	23.1	134	76.9	7.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older *Cortland, Tompkins*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	76	27.9	200	72.1	5.9
Gender					
Male	36	31.3	75	68.7	9.2
Female	40	24.5	125	75.5	7.2
Age					
50-64	36	22.8	125	77.2	7.3
≥65	40	35.5	75	64.5	9.5
Education					
≤High School	27	23.8	80	76.2	8.6
>High School	49	30.3	120	69.7	7.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older *Cortland, Tompkins*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	138	49.9	142	50.1	6.5
Gender					
Male	58	53.4	54	46.6	9.9
Female	80	46.5	88	53.5	8.2
Age					
50-64	70	43.0	95	57.0	8.4
≥65	68	60.4	47	39.6	9.7
Education					
≤High School	48	44.3	60	55.7	10.2
>High School	90	53.3	82	46.7	8.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older *Cortland, Tompkins*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	128	46.2	151	53.8	6.5
Gender					
Male	55	50.2	57	49.8	10.0
Female	73	42.2	94	57.8	8.1
Age					
50-64	66	40.6	99	59.4	8.3
≥65	62	54.8	52	45.2	9.9
Education					
≤High School	46	43.0	61	57.0	10.2
>High School	82	48.1	90	51.9	8.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	150	55.2	131	44.8	6.4
Gender					
Male	66	61.5	46	38.5	9.6
Female	84	48.9	85	51.1	8.2
Age					
50-64	77	48.8	88	51.2	8.5
≥65	73	64.8	43	35.2	9.3
Education					
≤High School	55	51.3	54	48.7	10.2
>High School	95	57.5	77	42.5	8.1

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval