#### Expanded BRFSS, 2003

#### **Broome Locality Tables**

#### TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days	
Core 2: Health Care Access	
Core 3: Exercise	
Core 4: Diabetes	
Core 5: Asthma	10
Core 6: Arthritis	
Core 7: Tobacco Use	14
Core 8: Tobacco ETS	
Core 9: Alcohol Consumption	19
Core 10: Weight Status	
Core 11: Mammography	23
Core 12: Sexual Behavior	25
Core 13: Family Planning	28
Core 14: Cardiovascular Disease	
Core 15: Prostate Cancer Screening	31
Core 16: Colorectal Cancer Screening	

New York State Department Of Health

#### **Self-rated Health Status**

Demographic Groups	n¹	Fair or Poor	n	Good to Excellent %	C.I. <sup>3</sup>
Total	111	16.1	529	83.9	3.1
Gender					
Male	44	16.2	223	83.8	5.0
Female	67	15.9	306	84.1	3.9
Age					
18-34	4	2.8	138	97.2	3.0
35-54	31	15.6	202	84.4	5.7
≥55	76	29.5	177	70.5	6.1
Education					
≤High School	73	25.1	196	74.9	5.8
>High School	38	9.9	331	90.1	3.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### **Reported Number of Physical Distress Days in the Past Month** Broome

Demographic Groups	n¹	≥14 days %2	n	< 14 days %	C.I. <sup>3</sup>
Total	87	12.6	540	87.4	2.8
Gender					
Male	30	10.4	234	89.6	3.9
Female	57	14.7	306	85.3	4.0
Age					
18-34	9	6.4	132	93.6	4.3
35-54	27	12.3	205	87.7	4.9
≥55	51	19.6	191	80.4	5.3
Education					
≤High School	48	17.0	211	83.0	4.9
>High School	39	9.8	327	90.2	3.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### **Reported Number of Mental Distress Days in the Past Month** Broome

Demographic Groups	n¹	≥14 days %2	n	< 14 days %	C.I. <sup>3</sup>
Total	69	11.3	563	88.7	3.0
Gender					
Male	23	8.8	241	91.2	4.0
Female	46	13.6	322	86.4	4.4
Age					
18-34	12	10.6	130	89.4	6.5
35-54	31	13.1	202	86.9	5.0
≥55	25	9.3	221	90.7	3.8
Education					
≤High School	39	17.3	225	82.7	5.8
>High School	30	7.2	336	92.8	3.0

<sup>&</sup>lt;sup>1</sup> Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup> Weighted Percent<sup>3</sup> 95% Confidence Interval

#### Expanded BRFSS, 2003 - New York State Department of Health **Health Insurance Coverage, 18-64 Years of Age Broome**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	399	84.1	66	15.9	4.1
Gender					
Male	180	85.0	33	15.0	5.5
Female	219	83.1	33	16.9	6.0
Age					
18-34	109	75.5	33	24.5	8.3
35-54	205	87.9	28	12.1	4.8
55-64	85	94.1	5	5.9	5.3
Education					
≤High School	134	79.5	28	20.5	7.9
>High School	264	86.5	38	13.5	4.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### Expanded BRFSS, 2003 - New York State Department of Health **Health Insurance Coverage, Aged 18 and Older Broome**

Demographic	1	Yes		No	CI.
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	569	87.3	71	12.7	3.2
Gender					
Male	232	87.8	34	12.2	4.5
Female	337	86.8	37	13.2	4.6
Age					
18-34	109	75.5	33	24.5	8.3
35-54	205	87.9	28	12.1	4.8
≥55	245	96.9	8	3.1	2.3
Education					
≤High School	237	85.5	31	14.5	5.6
>High School	330	88.4	40	11.6	3.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### Reported Having a Routine Physical Exam in Past Two Years **Broome**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	549	87.2	86	12.8	2.8
Gender					
Male	208	80.8	58	19.2	5.0
Female	341	93.1	28	6.9	2.7
Age					
18-34	119	87.4	22	12.6	5.4
35-54	190	83.1	41	16.9	5.3
≥55	229	91.0	22	9.0	4.0
Education					
≤High School	242	90.4	27	9.6	3.7
>High School	305	85.0	59	15.0	4.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### Time in Past 12 Months when Needed Medical Care but Could Not Get It **Broome**

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	38	6.6	603	93.4	2.5
Gender					
Male	15	5.6	253	94.4	3.5
Female	23	7.6	350	92.4	3.5
Age					
18-34	10	8.3	133	91.7	5.8
35-54	21	9.0	213	91.0	4.2
≥55	6	1.8	246	98.2	1.6
Education					
≤High School	12	5.2	257	94.8	3.6
>High School	26	7.6	344	92.4	3.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days **Broome**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	519	83.2	123	16.8	3.1
Gender					
Male	223	85.0	45	15.0	4.6
Female	296	81.5	78	18.5	4.2
Age					
18-34	128	92.4	15	7.6	4.1
35-54	192	81.9	42	18.1	5.5
≥55	189	75.4	64	24.6	5.9
Education					
≤High School	205	79.0	65	21.0	5.1
>High School	313	86.1	57	13.9	3.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Physician Diagnosed Diabetes**

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	50	7.3	591	92.7	2.2
Gender					
Male	22	8.1	246	91.9	3.5
Female	28	6.6	345	93.4	2.7
Age					
18-34	1	0.4	142	99.6	0.8
35-54	13	6.8	221	93.2	3.9
≥55	36	14.6	217	85.4	4.8
Education					
≤High School	28	9.8	242	90.2	3.8
>High School	21	5.5	348	94.5	2.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### **Lifetime Asthma Diagnosed by a Medical Professional Broome**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	60	8.9	581	91.1	2.5
Gender					
Male	22	7.8	246	92.2	3.5
Female	38	10.0	335	90.0	3.4
Age					
18-34	18	12.8	125	87.2	6.0
35-54	21	6.9	213	93.1	3.1
≥55	18	6.5	234	93.5	3.1
Education					
≤High School	21	7.5	249	92.5	3.5
>High School	39	9.9	330	90.1	3.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Current Asthma Diagnosed by a Medical Professional Broome**

Demographic	1	Yes		No	CI3
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	43	6.5	597	93.5	2.2
Gender					
Male	13	4.5	254	95.5	2.8
Female	30	8.2	343	91.8	3.2
Age					
18-34	14	10.2	129	89.8	5.6
35-54	13	4.0	221	96.0	2.3
≥55	13	4.4	238	95.6	2.4
Education					
≤High School	15	5.7	255	94.3	3.1
>High School	28	7.0	340	93.0	3.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### **Arthritis Diagnosed by a Medical Professional Broome**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	233	32.4	406	67.6	4.0
Gender					
Male	73	25.6	194	74.4	5.6
Female	160	38.5	212	61.5	5.5
Age					
18-34	12	8.2	131	91.8	4.9
35-54	66	28.9	168	71.1	6.5
≥55	150	58.0	100	42.0	6.8
Education					
≤High School	120	39.7	149	60.3	6.5
>High School	113	27.4	255	72.6	5.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

Demographic Groups	n¹	Doctor Diagnosed %2	C.I. <sup>3</sup>	n	Possible Arthritis (CJS Only) %	C.I.	n	No Arthritis or CJS %	C.I.
Total	233	32.5	4.0	93	15.6	3.4	311	51.9	4.4
Gender									
Male	73	25.7	5.7	53	20.7	5.6	140	53.6	6.7
Female	160	38.6	5.5	40	11.0	3.9	171	50.4	5.8
Age									
18-34	12	8.2	4.9	26	20.8	7.9	105	71.0	8.6
35-54	66	29.0	6.5	39	16.3	5.3	128	54.7	7.1
≥55	150	58.1	6.8	26	10.6	4.3	73	31.3	6.5
Education									
≤High School	120	39.8	6.5	33	13.2	5.2	115	47.1	6.8
>High School	113	27.5	5.0	60	17.3	4.5	194	55.1	5.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Ever Smoked 100 Cigarettes in Lifetime**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	313	49.2	329	50.8	4.4
Gender					
Male	148	54.4	120	45.6	6.7
Female	165	44.4	209	55.6	5.7
Age					
18-34	61	39.9	82	60.1	8.8
35-54	121	52.5	113	47.5	7.1
≥55	125	53.6	128	46.4	6.8
Education					
≤High School	150	58.6	120	41.4	6.6
>High School	162	42.6	208	57.4	5.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Current Smoker**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	149	23.2	493	76.8	3.7
Gender					
Male	64	23.2	204	76.8	5.6
Female	85	23.3	289	76.7	5.0
Age					
18-34	42	27.0	101	73.0	7.9
35-54	70	29.7	164	70.3	6.6
≥55	35	13.7	218	86.3	4.7
Education					
≤High School	83	31.7	187	68.3	6.5
>High School	65	17.3	305	82.7	4.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### **Status of Cigarette Smoking**

Demographic Groups	n¹	Everyday Smoker %2	C.I. <sup>3</sup>	n	Someday Smoker	C.I.	n	Former or Never Smoked %	C.I.
Total	120	18.3	3.4	29	5.0	2.0	493	76.8	3.7
Gender									
Male	53	18.5	5.0	11	4.7	3.1	204	76.8	5.6
Female	67	18.1	4.6	18	5.2	2.6	289	76.7	5.0
Age									
18-34	33	21.6	7.4	9	5.4	3.7	101	73.0	7.9
35-54	58	23.1	5.9	12	6.6	4.2	164	70.3	6.6
≥55	27	10.6	4.2	8	3.1	2.3	218	86.3	4.7
Education									
≤High School	71	26.9	6.2	12	4.8	3.2	187	68.3	6.5
>High School	48	12.2	3.6	17	5.1	2.5	305	82.7	4.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### **Home Smoking Policies**

Demographic Groups	n¹	Not Allowed	C.I. <sup>3</sup>	n	Allowed Some Places	C.I.	n	Allowed Anywhere	C.I.
Total	417	67.2	4.1	80	12.4	2.9	141	20.4	3.5
Gender									
Male	175	66.8	6.3	24	9.2	3.9	67	24.1	5.7
Female	242	67.6	5.4	56	15.3	4.1	74	17.1	4.3
Age									
18-34	107	73.0	8.4	18	12.8	6.1	18	14.2	7.1
35-54	151	66.2	6.7	38	15.2	5.0	44	18.6	5.6
≥55	152	62.1	6.6	23	9.4	4.0	76	28.5	6.0
Education									
≤High School	151	57.5	6.7	38	12.8	4.2	81	29.7	6.3
>High School	266	74.1	5.0	42	12.2	3.9	58	13.7	3.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	159	24.8	483	75.2	3.9
Gender					
Male	61	22.5	207	77.5	5.7
Female	98	26.9	276	73.1	5.3
Age					
18-34	32	24.4	111	75.6	8.3
35-54	69	28.8	165	71.2	6.5
≥55	54	21.2	199	78.8	5.5
Education					
≤High School	85	31.2	185	68.8	6.4
>High School	73	20.3	297	79.7	4.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	78	13.2	559	86.8	3.0
Gender					
Male	55	20.4	210	79.6	5.3
Female	23	6.7	349	93.3	2.9
Age					
18-34	34	23.4	107	76.6	7.6
35-54	28	10.6	205	89.4	4.1
≥55	15	7.2	237	92.8	3.8
Education					
≤High School	26	10.0	240	90.0	4.0
>High School	52	15.4	317	84.6	4.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	35	5.6	605	94.4	2.1
Gender					
Male	22	8.2	246	91.8	3.7
Female	13	3.3	359	96.7	2.0
Age					
18-34	12	8.5	131	91.5	5.1
35-54	11	3.7	223	96.3	2.4
≥55	12	5.5	240	94.5	3.3
Education					
≤High School	14	4.5	256	95.5	2.5
>High School	21	6.5	347	93.5	3.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health Overweight or Obese, BMI(Body Mass Index) 25 or More Broome

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	373	60.3	235	39.7	4.4
Gender					
Male	193	70.6	73	29.4	6.3
Female	180	50.4	162	49.6	6.0
Age					
18-34	63	44.7	75	55.3	9.2
35-54	149	67.0	71	33.0	6.9
≥55	160	68.6	83	31.4	6.3
Education					
≤High School	161	64.7	91	35.3	6.7
>High School	211	57.4	143	42.6	5.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+) *Broome*

Demographic Groups	n¹	Underweight\ Normal %2	C.I. <sup>3</sup>	n	Overweight %	C.I.	n	Obese %	C.I.
Total	234	39.6	4.4	231	37.5	4.3	142	23.0	3.8
Gender									
Male	73	29.4	6.3	125	46.0	6.7	68	24.5	5.7
Female	161	49.4	6.0	106	29.1	5.3	74	21.4	5.0
Age									
18-34	75	55.3	9.2	42	28.7	8.2	21	16.0	7.0
35-54	71	33.0	6.9	98	43.1	7.2	51	23.9	6.4
≥55	82	31.0	6.3	90	39.7	6.9	70	29.3	6.4
Education									
≤High School	90	35.0	6.7	94	39.5	7.0	67	25.6	6.1
>High School	143	42.6	5.8	136	36.1	5.5	75	21.3	4.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Ever Had Mammography Screening, Women Aged 40 Years and Older Broome**

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Female	231	92.3	22	7.7	3.4
Age					
40-64	127	91.2	14	8.8	4.8
≥65	104	94.1	8	5.9	4.2
Education					
≤High School	123	94.0	10	6.0	4.1
>High School	106	90.4	12	9.6	5.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Female	200	79.9	52	20.1	5.6
Age					
40-64	109	78.4	32	21.6	7.4
≥65	91	82.2	20	17.8	8.3
Education					
≤High School	103	78.3	29	21.7	8.1
>High School	95	81.4	23	18.6	7.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	55	11.3	401	88.7	3.2
Gender					
Male	15	5.8	196	94.2	3.2
Female	40	16.8	205	83.2	5.3
Age					
18-34	29	18.7	111	81.3	6.8
35-54	23	8.2	205	91.8	3.6
55-64	3	2.0	85	98.0	2.3
Education					
≤High School	24	12.3	136	87.7	5.1
>High School	30	10.6	265	89.4	4.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health **Sexually Active, Age 18-64 Years Old Broome**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	317	77.7	106	22.3	4.5
Gender					
Male	145	74.5	53	25.5	6.8
Female	172	81.0	53	19.0	5.8
Age					
18-34	102	77.1	30	22.9	8.2
35-54	171	82.6	41	17.4	5.7
55-64	44	64.3	35	35.7	11.3
Education					
≤High School	101	70.6	47	29.4	8.5
>High School	216	81.9	58	18.1	4.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### **Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger Broome**

Demographic		Yes		No	~ - 4
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	62	18.3	265	81.7	4.7
Gender					
Male	34	21.3	115	78.7	7.2
Female	28	15.5	150	84.5	6.0
Age					
18-34	31	28.8	75	71.2	9.5
35-54	26	12.2	148	87.8	4.8
55-64	5	8.8	42	91.2	8.4
Education					
≤High School	15	13.8	90	86.2	7.1
>High School	47	20.5	175	79.5	6.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

Demographic		Use BC			No BC at risk			No BC	
Groups	n¹	% <sup>2</sup>	C.I. <sup>3</sup>	n	%	C.I.	n	%	C.I.
Total	193	57.3	5.9	34	11.7	3.9	105	30.9	5.5
Gender									
Male	99	52.0	7.8	18	11.2	5.1	77	36.8	7.5
Female	94	65.4	8.9	16	12.6	5.9	28	22.1	7.9
Age									
18-34	78	56.9	9.1	16	12.5	5.9	44	30.6	8.4
35-59	115	57.7	7.7	18	11.0	5.1	61	31.3	7.2
Education									
≤High School	60	58.2	10.8	7	6.6	4.9	36	35.2	10.4
>High School	133	56.9	7.1	27	14.2	5.1	69	28.9	6.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health **Physician Diagnosed Heart Attack, Angina or Stroke**

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	59	7.9	573	92.1	2.2
Gender					
Male	27	9.3	237	90.7	3.6
Female	32	6.5	336	93.5	2.6
Age					
18-64	18	3.5	443	96.5	1.8
≥65	41	24.1	121	75.9	7.1
Education					
≤High School	36	11.6	230	88.4	4.1
>High School	23	5.3	341	94.7	2.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

## Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina) **Broome**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	45	6.3	586	93.7	2.0
Gender					
Male	24	8.1	239	91.9	3.3
Female	21	4.7	347	95.3	2.3
Age					
18-64	12	2.4	448	97.6	1.6
≥65	33	20.8	129	79.2	6.9
Education					
≤High School	29	9.4	237	90.6	3.7
>High School	16	4.2	347	95.8	2.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### **Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older** *Broome*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	98	56.9	72	43.1	8.2
Age					
40-64	55	41.6	69	58.4	9.6
≥65	43	94.8	3	5.2	6.2
Education					
≤High School	42	58.0	30	42.0	12.4
>High School	56	56.1	42	43.9	10.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	81	48.6	88	51.4	8.3
Age					
40-64	44	33.9	79	66.1	9.2
≥65	37	84.4	9	15.6	10.2
Education					
≤High School	31	45.9	41	54.1	12.7
>High School	50	50.6	47	49.4	11.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Ever Had a Digital Rectal Exam, Men Aged 40 and Older** *Broome*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	141	81.3	34	18.7	6.3
Age					
40-64	99	78.0	28	22.0	8.0
≥65	42	89.7	6	10.3	8.4
Education					
≤High School	54	76.1	20	23.9	10.2
>High School	87	85.3	14	14.7	7.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Ever Used a Blood Stool Test at Home, Aged 50 and Older *Broome*

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	132	43.8	164	56.2	6.2
Gender					
Male	55	46.3	54	53.7	10.1
Female	77	41.9	110	58.1	7.8
Age					
50-64	54	39.8	84	60.2	9.0
≥65	78	47.4	80	52.6	8.6
Education					
≤High School	69	42.6	93	57.4	8.3
>High School	62	45.4	70	54.6	9.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older *Broome*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	39	12.9	256	87.1	4.2
Gender					
Male	17	13.2	91	86.8	6.6
Female	22	12.6	165	87.4	5.3
Age					
50-64	17	10.6	120	89.4	5.3
≥65	22	14.9	136	85.1	6.2
Education					
≤High School	20	12.6	142	87.4	5.6
>High School	18	12.9	113	87.1	6.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

## Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older **Broome**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	62	20.3	233	79.7	4.9
Gender					
Male	24	17.9	84	82.1	7.2
Female	38	22.2	149	77.8	6.7
Age					
50-64	27	18.2	110	81.8	6.8
≥65	35	22.2	123	77.8	7.1
Education					
≤High School	33	19.8	129	80.2	6.6
>High School	28	20.8	103	79.2	7.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

## Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older **Broome**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	151	49.2	148	50.8	6.3
Gender					
Male	52	48.8	58	51.2	10.2
Female	99	49.5	90	50.5	7.8
Age					
50-64	54	38.8	84	61.2	9.0
≥65	97	58.5	64	41.5	8.5
Education					
≤High School	76	44.3	86	55.7	8.3
>High School	74	55.1	61	44.9	9.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older *Broome*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	137	45.1	161	54.9	6.2
Gender					
Male	48	44.8	62	55.2	10.1
Female	89	45.3	99	54.7	7.8
Age					
50-64	51	37.1	87	62.9	8.9
≥65	86	52.3	74	47.7	8.6
Education					
≤High School	69	41.4	92	58.6	8.3
>High School	67	49.5	68	50.5	9.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	152	50.3	149	49.7	6.3
Gender					
Male	54	49.1	57	50.9	10.1
Female	98	51.1	92	48.9	7.8
Age					
50-64	56	41.3	83	58.7	9.0
≥65	96	58.3	66	41.7	8.5
Education					
≤High School	80	48.4	84	51.6	8.4
>High School	71	52.6	64	47.4	9.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval