#### Expanded BRFSS, 2003

### Allegany, Wyoming Locality Tables

#### TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days	
Core 2: Health Care Access	4
Core 3: Exercise	8
Core 4: Diabetes	
Core 5: Asthma	10
Core 6: Arthritis	12
Core 7: Tobacco Use	14
Core 8: Tobacco ETS	
Core 9: Alcohol Consumption	19
Core 9: Alcohol Consumption	21
Core 11: Mammography	23
Core 12: Sexual Behavior	
Core 13: Family Planning	28
Core 14: Cardiovascular Disease	29
Core 15: Prostate Cancer Screening	
Core 16: Colorectal Cancer Screening	

New York State Department Of Health

#### **Self-rated Health Status**

Demographic Groups	n¹	Fair or Poor	n	Good to Excellent	C.I. <sup>3</sup>
Total	112	15.5	524	84.5	3.1
Gender					
Male	43	15.1	204	84.9	4.7
Female	69	15.8	320	84.2	3.9
Age					
18-34	13	8.9	122	91.1	4.8
35-54	40	15.0	223	85.0	5.2
≥55	57	23.2	173	76.8	5.9
Education					
≤High School	73	19.0	266	81.0	4.6
>High School	38	11.3	258	88.7	3.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### **Reported Number of Physical Distress Days in the Past Month** Allegany, Wyoming

Demographic Groups	n¹	≥14 days %2	n	< 14 days %	C.I. <sup>3</sup>
Total	76	11.2	546	88.8	2.7
Gender					
Male	27	10.7	214	89.3	4.2
Female	49	11.8	332	88.2	3.5
Age					
18-34	13	9.7	123	90.3	5.3
35-54	26	8.2	229	91.8	3.4
≥55	36	16.5	187	83.5	5.5
Education					
≤High School	40	11.6	288	88.4	3.9
>High School	36	10.7	257	89.3	3.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### **Reported Number of Mental Distress Days in the Past Month** Allegany, Wyoming

Demographic Groups	n¹	≥14 days %2	n	< 14 days %	C.I. <sup>3</sup>
Total	72	10.0	548	90.0	2.7
Gender					
Male	15	6.7	225	93.3	3.6
Female	57	13.7	323	86.3	3.9
Age					
18-34	14	11.0	120	89.0	6.0
35-54	38	12.3	220	87.7	4.2
≥55	20	6.5	200	93.5	3.0
Education					
≤High School	38	10.9	291	89.1	4.0
>High School	34	9.1	256	90.9	3.4

<sup>&</sup>lt;sup>1</sup> Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup> Weighted Percent<sup>3</sup> 95% Confidence Interval

#### Expanded BRFSS, 2003 - New York State Department of Health **Health Insurance Coverage, 18-64 Years of Age** Allegany, Wyoming

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	433	85.4	62	14.6	3.8
Gender					
Male	169	83.4	31	16.6	5.9
Female	264	87.7	31	12.3	4.6
Age					
18-34	111	81.3	25	18.7	7.1
35-54	235	87.3	28	12.7	5.1
55-64	87	90.4	9	9.6	6.9
Education					
≤High School	199	79.6	43	20.4	6.2
>High School	234	91.4	19	8.6	4.1

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### Expanded BRFSS, 2003 - New York State Department of Health Health Insurance Coverage, Aged 18 and Older Allegany, Wyoming

Demographic		Yes		No	
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	573	87.6	65	12.4	3.2
Gender					
Male	217	85.4	32	14.6	5.1
Female	356	90.0	33	10.0	3.7
Age					
18-34	111	81.3	25	18.7	7.1
35-54	235	87.3	28	12.7	5.1
≥55	220	94.7	11	5.3	3.5
Education					
≤High School	297	83.7	44	16.3	5.0
>High School	275	92.2	21	7.8	3.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

# Reported Having a Routine Physical Exam in Past Two Years *Allegany, Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	547	84.2	87	15.8	3.5
Gender					
Male	198	78.6	48	21.4	6.0
Female	349	90.3	39	9.7	3.2
Age					
18-34	109	77.6	27	22.4	8.0
35-54	224	84.5	39	15.5	5.2
≥55	209	91.9	18	8.1	3.9
Education					
≤High School	294	85.5	44	14.5	4.6
>High School	252	82.6	43	17.4	5.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Time in Past 12 Months when Needed Medical Care but Could Not Get It *Allegany, Wyoming*

Demographic	n¹	Yes		No	C.I. <sup>3</sup>
Groups	11	70-	n	%	C.I.
Total	35	5.1	603	94.9	2.0
Gender					
Male	10	3.8	238	96.2	2.7
Female	25	6.5	365	93.5	2.8
Age					
18-34	9	6.4	127	93.6	4.2
35-54	18	6.1	245	93.9	3.5
≥55	8	2.7	223	97.3	2.0
Education					
≤High School	19	5.8	322	94.2	3.0
>High School	16	4.3	280	95.7	2.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days *Allegany, Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	462	74.1	176	25.9	3.8
Gender					
Male	175	71.9	74	28.1	6.0
Female	287	76.4	102	23.6	4.5
Age					
18-34	109	81.2	27	18.8	6.7
35-54	188	71.9	75	28.1	6.1
≥55	159	68.5	72	31.5	6.7
Education					
≤High School	227	67.0	114	33.0	5.6
>High School	235	82.5	61	17.5	4.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Physician Diagnosed Diabetes**

Demographic		Yes		No	
Groups	n¹	%2	n	%	C.I. <sup>3</sup>
Total	51	6.2	588	93.8	1.8
Gender					
Male	18	4.8	231	95.2	2.3
Female	33	7.7	357	92.3	2.8
Age					
18-34	2	1.1	134	98.9	1.6
35-54	14	4.7	249	95.3	2.6
≥55	33	13.1	199	86.9	4.6
Education					
≤High School	28	6.2	314	93.8	2.5
>High School	22	6.1	274	93.9	2.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health **Lifetime Asthma Diagnosed by a Medical Professional** *Allegany, Wyoming* 

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	71	10.6	565	89.4	2.7
Gender					
Male	28	11.1	220	88.9	4.3
Female	43	10.0	345	90.0	3.3
Age					
18-34	17	12.7	119	87.3	6.0
35-54	27	9.1	235	90.9	3.7
≥55	27	10.3	203	89.7	4.3
Education					
≤High School	43	11.6	297	88.4	3.7
>High School	28	9.3	267	90.7	4.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Current Asthma Diagnosed by a Medical Professional**

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	50	7.1	583	92.9	2.3
Gender					
Male	15	6.2	230	93.8	3.4
Female	35	8.2	353	91.8	3.0
Age					
18-34	13	9.6	122	90.4	5.4
35-54	17	5.0	244	95.0	2.6
≥55	20	7.1	209	92.9	3.5
Education					
≤High School	33	8.8	305	91.2	3.3
>High School	17	5.2	277	94.8	3.1

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Arthritis Diagnosed by a Medical Professional**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	221	29.1	415	70.9	3.8
Gender					
Male	74	25.6	174	74.4	5.6
Female	147	32.8	241	67.2	5.0
Age					
18-34	16	9.2	120	90.8	4.7
35-54	87	30.7	175	69.3	6.3
≥55	116	48.4	114	51.6	7.1
Education					
≤High School	127	30.9	212	69.1	5.3
>High School	94	26.9	202	73.1	5.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

Demographic Groups	n¹	Doctor Diagnosed %2	C.I. <sup>3</sup>	n	Possible Arthritis (CJS Only) %	C.I.	n	No Arthritis or CJS %	C.I.
Total	221	29.2	3.8	97	16.9	3.6	314	53.9	4.4
Gender									
Male	74	25.8	5.7	39	18.5	5.8	133	55.7	6.8
Female	147	32.8	5.0	58	15.2	4.1	181	51.9	5.5
Age									
18-34	16	9.2	4.7	28	24.1	8.2	92	66.7	8.7
35-54	87	30.8	6.3	40	14.5	4.7	134	54.7	6.8
≥55	116	49.0	7.2	28	12.2	4.8	84	38.9	7.0
Education									
≤High School	127	31.0	5.3	50	16.0	4.6	160	52.9	6.0
>High School	94	27.1	5.5	47	18.0	5.6	153	54.9	6.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Ever Smoked 100 Cigarettes in Lifetime**

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	343	52.1	294	47.9	4.5
Gender					
Male	148	57.3	99	42.7	6.9
Female	195	46.6	195	53.4	5.5
Age					
18-34	65	46.6	71	53.4	9.0
35-54	153	54.2	110	45.8	6.9
≥55	122	56.8	108	43.2	7.0
Education					
≤High School	194	56.4	146	43.6	6.0
>High School	148	47.1	148	52.9	6.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Current Smoker**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	177	27.0	460	73.0	4.0
Gender					
Male	66	28.1	181	71.9	6.3
Female	111	25.9	279	74.1	4.7
Age					
18-34	48	35.3	88	64.7	8.7
35-54	94	32.2	169	67.8	6.3
≥55	34	12.3	196	87.7	4.5
Education					
≤High School	109	32.3	231	67.7	5.8
>High School	68	20.9	228	79.1	5.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

#### **Status of Cigarette Smoking**

Demographic Groups	n¹	Everyday Smoker %2	C.I. <sup>3</sup>	n	Someday Smoker	C.I.	n	Former or Never Smoked %	C.I.
Total	137	21.3	3.7	40	5.7	2.1	460	73.0	4.0
Gender									
Male	54	22.0	5.7	12	6.1	3.6	181	71.9	6.3
Female	83	20.6	4.4	28	5.3	2.2	279	74.1	4.7
Age									
18-34	36	26.3	7.9	12	9.0	5.4	88	64.7	8.7
35-54	75	27.1	6.0	19	5.0	2.6	169	67.8	6.3
≥55	25	9.3	4.0	9	3.0	2.2	196	87.7	4.5
Education									
≤High School	85	24.7	5.3	24	7.6	3.5	231	67.7	5.8
>High School	52	17.5	4.9	16	3.5	2.1	228	79.1	5.2

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### **Home Smoking Policies**

Demographic Groups	n¹	Not Allowed	C.I. <sup>3</sup>	n	Allowed Some Places %	C.I.	n	Allowed Anywhere	C.I.
Total	361	58.6	4.4	100	15.8	3.2	175	25.6	3.8
Gender									
Male	139	57.8	6.7	32	14.1	4.7	77	28.1	6.1
Female	222	59.4	5.4	68	17.7	4.3	98	22.9	4.5
Age									
18-34	79	57.8	8.9	30	21.1	7.1	27	21.1	7.6
35-54	139	56.0	6.8	48	15.5	4.6	76	28.5	6.1
≥55	137	61.5	7.0	22	11.0	4.7	70	27.5	6.3
Education									
≤High School	172	52.4	6.1	51	16.0	4.5	116	31.5	5.6
>High School	188	65.6	6.1	49	15.6	4.6	59	18.7	5.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### **Home Smoking**

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	207	31.9	432	68.1	4.2
Gender					
Male	76	31.4	173	68.6	6.5
Female	131	32.4	259	67.6	5.1
Age					
18-34	44	34.5	92	65.5	8.9
35-54	110	38.5	153	61.5	6.6
≥55	51	21.4	181	78.6	5.9
Education					
≤High School	129	37.4	213	62.6	5.9
>High School	78	25.5	218	74.5	5.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)

Demographic	n¹	Yes		No o/	C.I. <sup>3</sup>
Groups	11-	70-	n	%	C.I.
Total	101	20.0	531	80.0	4.0
Gender					
Male	67	30.8	176	69.2	6.7
Female	34	8.6	355	91.4	3.0
Age					
18-34	42	35.2	91	64.8	9.1
35-54	47	20.0	213	80.0	5.6
≥55	10	4.0	221	96.0	2.8
Education					
≤High School	59	21.1	279	78.9	5.4
>High School	42	18.8	251	81.2	5.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	48	9.2	582	90.8	3.0
Gender					
Male	29	13.1	213	86.9	5.2
Female	19	5.1	369	94.9	2.4
Age					
18-34	16	13.6	118	86.4	7.2
35-54	24	10.3	237	89.7	4.2
≥55	7	3.3	221	96.7	2.6
Education					
≤High School	31	10.6	304	89.4	4.0
>High School	17	7.7	277	92.3	4.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Expanded BRFSS, 2003 - New York State Department of Health Overweight or Obese, BMI(Body Mass Index) 25 or More Allegany, Wyoming

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	351	57.4	266	42.6	4.5
Gender					
Male	156	62.5	87	37.5	6.8
Female	195	51.9	179	48.1	5.7
Age					
18-34	65	49.0	68	51.0	9.2
35-54	146	59.9	105	40.1	6.9
≥55	137	63.1	91	36.9	6.8
Education					
≤High School	197	57.8	136	42.2	6.1
>High School	154	57.1	130	42.9	6.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

#### Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+) Allegany, Wyoming

Demographic Groups	n¹	Underweight\ Normal %2	C.I. <sup>3</sup>	n	Overweight %	C.I.	n	Obese %	C.I.
Total	266	42.6	4.5	190	33.0	4.3	160	24.4	3.7
Gender									
Male	87	37.5	6.8	99	41.4	6.8	57	21.1	5.3
Female	179	48.3	5.7	91	23.8	4.8	103	27.9	5.1
Age									
18-34	68	51.4	9.2	38	30.2	8.6	26	18.4	6.7
35-54	105	40.1	6.9	66	28.8	6.5	80	31.1	6.4
≥55	91	36.9	6.8	85	40.4	7.2	52	22.7	6.0
Education									
≤High School	136	42.4	6.1	102	31.2	5.6	94	26.3	5.2
>High School	130	42.9	6.6	88	35.1	6.7	66	22.0	5.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# **Ever Had Mammography Screening, Women Aged 40 Years and Older** *Allegany, Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Female	245	89.2	28	10.8	4.0
Age					
40-64	165	87.5	20	12.5	5.4
≥65	80	92.5	8	7.5	5.4
Education					
≤High School	127	85.9	19	14.1	6.3
>High School	118	93.2	9	6.8	4.6

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Female	198	74.3	75	25.7	5.5
Age					
40-64	134	73.3	51	26.7	6.8
≥65	64	76.2	24	23.8	9.2
Education					
≤High School	98	68.7	48	31.3	8.0
>High School	100	81.0	27	19.0	7.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	62	14.7	422	85.3	3.9
Gender					
Male	21	12.0	177	88.0	5.3
Female	41	18.0	245	82.0	5.6
Age					
18-34	35	26.7	100	73.3	8.2
35-54	23	7.5	232	92.5	3.4
55-64	4	3.8	90	96.2	3.8
Education					
≤High School	36	17.7	197	82.3	5.9
>High School	26	11.7	225	88.3	4.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health **Sexually Active, Age 18-64 Years Old** 

Demographic		Yes		No	
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	355	81.8	106	18.2	3.9
Gender					
Male	143	80.5	43	19.5	6.2
Female	212	83.3	63	16.7	4.5
Age					
18-34	112	84.9	20	15.1	6.7
35-54	194	84.2	49	15.8	5.0
55-64	49	66.2	37	33.8	10.7
Education					
≤High School	172	79.6	50	20.4	6.1
>High School	183	84.0	56	16.0	4.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### **Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger** Allegany, Wyoming

Demographic		Yes		No	G T 1
Groups	n¹	% <sup>2</sup>	n	%	C.I. <sup>3</sup>
Total	49	14.9	316	85.1	4.5
Gender					
Male	28	16.3	123	83.7	6.6
Female	21	13.2	193	86.8	5.8
Age					
18-34	29	25.2	86	74.8	8.9
35-54	18	8.0	183	92.0	3.9
55-64	2	3.6	47	96.4	5.2
Education					
≤High School	26	14.2	154	85.8	5.6
>High School	23	15.6	162	84.4	6.9

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent <sup>3</sup>95% Confidence Interval

### Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

Demographic Groups	n¹	Use BC	C.I. <sup>3</sup>	n	No BC at risk	C.I.	n	No BC	C.I.
Total	205	65.6	5.8	34	12.1	4.0	74	22.3	5.1
Gender									
Male	99	61.9	7.8	23	14.5	5.7	49	23.5	6.7
Female	106	71.8	8.4	11	8.0	4.7	25	20.2	7.7
Age									
18-34	85	65.1	8.7	22	15.6	6.3	24	19.3	7.4
35-59	120	66.1	7.7	12	8.3	4.9	50	25.5	6.9
Education									
≤High School	110	65.6	8.2	20	14.1	6.1	34	20.3	6.8
>High School	95	65.6	8.3	14	9.9	5.1	40	24.6	7.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Physician Diagnosed Heart Attack, Angina or Stroke

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	51	7.5	574	92.5	2.2
Gender					
Male	27	9.4	217	90.6	3.7
Female	24	5.5	357	94.5	2.3
Age					
18-64	18	4.2	465	95.8	2.1
≥65	33	23.8	101	76.2	7.8
Education					
≤High School	38	9.5	293	90.5	3.3
>High School	13	5.2	280	94.8	3.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

#### Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina) Allegany, Wyoming

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	41	6.1	584	93.9	2.0
Gender					
Male	22	7.9	222	92.1	3.4
Female	19	4.2	362	95.8	2.0
Age					
18-64	16	3.7	467	96.3	1.9
≥65	25	17.9	109	82.1	6.9
Education					
≤High School	33	8.5	298	91.5	3.1
>High School	8	3.3	285	96.7	2.4

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### **Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older** *Allegany, Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	91	64.3	53	35.7	8.4
Age					
40-64	59	58.5	46	41.5	10.1
≥65	32	83.8	7	16.2	12.1
Education					
≤High School	55	66.0	31	34.0	10.7
>High School	36	61.7	22	38.3	13.5

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

### Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	77	53.3	67	46.7	8.8
Age					
40-64	48	46.4	57	53.6	10.3
≥65	29	76.3	10	23.7	14.0
Education					
≤High School	46	52.9	40	47.1	11.5
>High School	31	54.0	27	46.0	13.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had a Digital Rectal Exam, Men Aged 40 and Older

Allegany, Wyoming

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Male	123	79.1	35	20.9	6.7
Age					
40-64	83	76.0	30	24.0	8.2
≥65	40	88.7	5	11.3	10.1
Education					
≤High School	75	79.6	22	20.4	8.4
>High School	48	78.2	13	21.8	11.1

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval

# Ever Used a Blood Stool Test at Home, Aged 50 and Older *Allegany, Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	110	38.1	176	61.9	6.2
Gender					
Male	41	38.4	60	61.6	10.0
Female	69	37.9	116	62.1	7.5
Age					
50-64	54	35.2	101	64.8	8.4
≥65	56	41.8	75	58.2	9.2
Education					
≤High School	60	36.2	104	63.8	8.0
>High School	50	40.9	72	59.1	9.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older *Allegany, Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	44	14.8	241	85.2	4.5
Gender					
Male	17	15.3	83	84.7	7.3
Female	27	14.3	158	85.7	5.4
Age					
50-64	20	13.3	135	86.7	6.1
≥65	24	16.6	106	83.4	6.6
Education					
≤High School	24	14.5	139	85.5	5.8
>High School	20	15.1	102	84.9	7.0

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

### Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older *Allegany, Wyoming*

Demographic Groups	n¹	Yes %2	n	No %	C.I. <sup>3</sup>
Total	66	22.4	219	77.6	5.2
Gender					
Male	24	21.4	76	78.6	8.3
Female	42	23.3	143	76.7	6.5
Age					
50-64	34	22.2	121	77.8	7.2
≥65	32	22.7	98	77.3	7.5
Education					
≤High School	34	20.7	129	79.3	6.7
>High School	32	24.7	90	75.3	8.3

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older *Allegany, Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	129	44.7	158	55.3	6.4
Gender					
Male	46	43.6	55	56.4	10.2
Female	83	45.6	103	54.4	7.7
Age					
50-64	54	33.3	101	66.7	8.2
≥65	75	58.8	57	41.2	9.2
Education					
≤High School	72	46.1	93	53.9	8.4
>High School	57	42.5	65	57.5	9.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

# Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older *Allegany, Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	117	41.6	167	58.4	6.3
Gender					
Male	44	41.8	56	58.2	10.2
Female	73	41.4	111	58.6	7.7
Age					
50-64	49	31.2	104	68.8	8.1
≥65	68	54.5	63	45.5	9.3
Education					
≤High School	64	42.3	99	57.7	8.4
>High School	53	40.6	68	59.4	9.7

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>395%</sup> Confidence Interval

Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. <sup>3</sup>
Total	136	46.9	152	53.1	6.4
Gender					
Male	49	44.9	53	55.1	10.2
Female	87	48.7	99	51.3	7.8
Age					
50-64	57	35.6	98	64.4	8.3
≥65	79	60.8	54	39.2	9.1
Education					
≤High School	77	48.3	89	51.7	8.4
>High School	59	44.8	63	55.2	9.8

<sup>&</sup>lt;sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>&</sup>lt;sup>2</sup>Weighted Percent

<sup>&</sup>lt;sup>3</sup>95% Confidence Interval