### Bronx Locality Tables

### TABLE OF CONTENTS:

| Core 1: Health Status / Healthy Days | 1  |
|--------------------------------------|----|
| Core 2: Health Care Access           | 4  |
| Core 3: Exercise                     | 8  |
| Core 4: Diabetes                     |    |
| Core 5: Asthma                       |    |
| Core 6: Arthritis                    | 12 |
| Core 7: Tobacco Use                  | 14 |
| Core 8: Tobacco ETS                  | 17 |
| Core 9: Alcohol Consumption          | 19 |
| fore 10: Weight Status               | 21 |
|                                      | 23 |
|                                      | 25 |
| Core 13: Family Planning             | 28 |
| Core 14: Cardiovascular Disease      | 29 |
| Core 15: Prostate Cancer Screening   | 31 |
| Core 16: Colorectal Cancer Screening | 37 |

New York State Department Of Health

## Expanded BRFSS, 2003 - New York State Department of Health **Self-rated Health Status**

### Bronx

| Demographic<br>Groups | n1  | Fair or Poor | n   | Good to Excellent | C.I. <sup>3</sup> |
|-----------------------|-----|--------------|-----|-------------------|-------------------|
| Total                 | 184 | 31.5         | 441 | 68.5              | 4.6               |
| Gender                |     |              |     |                   |                   |
| Male                  | 50  | 31.3         | 139 | 68.7              | 8.0               |
| Female                | 134 | 31.7         | 302 | 68.3              | 5.1               |
| Age                   |     |              |     |                   |                   |
| 18-34                 | 35  | 24.7         | 148 | 75.3              | 8.2               |
| 35-54                 | 71  | 27.3         | 200 | 72.7              | 6.5               |
| ≥55                   | 77  | 49.6         | 85  | 50.4              | 9.1               |
| Education             |     |              |     |                   |                   |
| ≤High School          | 136 | 43.2         | 196 | 56.8              | 6.6               |
| >High School          | 44  | 16.1         | 243 | 83.9              | 5.2               |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

1

### Expanded BRFSS, 2003 - New York State Department of Health Reported Number of Physical Distress Days in the Past Month **Bronx**

| Demographic<br>Groups | n1 | ≥14 days<br>%0 <sup>2</sup> | n   | < 14 days<br>% | C.I. <sup>3</sup> |
|-----------------------|----|-----------------------------|-----|----------------|-------------------|
| Total                 | 91 | 13.6                        | 513 | 86.4           | 3.0               |
| Gender                |    |                             |     |                |                   |
| Male                  | 23 | 9.7                         | 161 | 90.3           | 4.3               |
| Female                | 68 | 16.9                        | 352 | 83.1           | 4.1               |
| Age                   |    |                             |     |                |                   |
| 18-34                 | 14 | 6.3                         | 166 | 93.7           | 3.5               |
| 35-54                 | 39 | 13.6                        | 224 | 86.4           | 4.5               |
| ≥55                   | 38 | 26.1                        | 113 | 73.9           | 8.3               |
| Education             |    |                             |     |                |                   |
| ≤High School          | 54 | 15.8                        | 260 | 84.2           | 4.5               |
| >High School          | 36 | 11.1                        | 249 | 88.9           | 3.9               |

### Expanded BRFSS, 2003 - New York State Department of Health **Reported Number of Mental Distress Days in the Past Month Bronx**

| Demographic<br>Groups | n1 | ≥14 days<br>%0 <sup>2</sup> | n   | < 14 days | C.I. <sup>3</sup> |
|-----------------------|----|-----------------------------|-----|-----------|-------------------|
| Total                 | 86 | 13.0                        | 531 | 87.0      | 3.0               |
| Gender                |    |                             |     |           |                   |
| Male                  | 19 | 9.3                         | 167 | 90.7      | 4.5               |
| Female                | 67 | 16.0                        | 364 | 84.0      | 4.1               |
| Age                   |    |                             |     |           |                   |
| 18-34                 | 21 | 10.4                        | 163 | 89.6      | 5.1               |
| 35-54                 | 48 | 16.4                        | 220 | 83.6      | 4.8               |
| ≥55                   | 17 | 12.2                        | 138 | 87.8      | 6.4               |
| Education             |    |                             |     |           |                   |
| ≤High School          | 57 | 15.8                        | 269 | 84.2      | 4.6               |
| >High School          | 29 | 9.7                         | 256 | 90.3      | 3.7               |

<sup>1</sup> Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup> Weighted Percent<sup>3</sup> 95% Confidence Interval

### Expanded BRFSS, 2003 - New York State Department of Health Health Insurance Coverage, 18-64 Years of Age **Bronx**

| Demographic<br>Groups | n1  | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|-----|------------|-----|---------|-------------------|
| Total                 | 431 | 72.1       | 113 | 27.9    | 5.1               |
| Gender                |     |            |     |         |                   |
| Male                  | 122 | 63.5       | 45  | 36.5    | 9.0               |
| Female                | 309 | 79.6       | 68  | 20.4    | 5.0               |
| Age                   |     |            |     |         |                   |
| 18-34                 | 137 | 64.1       | 49  | 35.9    | 9.0               |
| 35-54                 | 215 | 74.5       | 56  | 25.5    | 6.5               |
| 55-64                 | 79  | 91.6       | 8   | 8.4     | 7.8               |
| Education             |     |            |     |         |                   |
| ≤High School          | 204 | 65.5       | 69  | 34.5    | 7.7               |
| >High School          | 223 | 79.4       | 42  | 20.6    | 6.4               |

### Expanded BRFSS, 2003 - New York State Department of Health Health Insurance Coverage, Aged 18 and Older **Bronx**

| Demographic<br>Groups | n1  | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|-----|------------|-----|---------|-------------------|
| Total                 | 512 | 75.2       | 119 | 24.8    | 4.6               |
| Gender                |     |            |     |         |                   |
| Male                  | 144 | 66.7       | 47  | 33.3    | 8.3               |
| Female                | 368 | 82.1       | 72  | 17.9    | 4.3               |
| Age                   |     |            |     |         |                   |
| 18-34                 | 137 | 64.1       | 49  | 35.9    | 9.0               |
| 35-54                 | 215 | 74.5       | 56  | 25.5    | 6.5               |
| ≥55                   | 152 | 93.3       | 12  | 6.7     | 4.6               |
| Education             |     |            |     |         |                   |
| ≤High School          | 262 | 71.0       | 74  | 29.0    | 6.6               |
| >High School          | 246 | 80.6       | 43  | 19.4    | 6.0               |

### Expanded BRFSS, 2003 - New York State Department of Health **Reported Having a Routine Physical Exam in Past Two Years Bronx**

| Demographic<br>Groups | n1  | Yes<br>%02 | n  | No<br>% | C.I. <sup>3</sup> |
|-----------------------|-----|------------|----|---------|-------------------|
| Total                 | 589 | 92.9       | 38 | 7.1     | 2.6               |
| Gender                |     |            |    |         |                   |
| Male                  | 168 | 88.8       | 22 | 11.2    | 5.1               |
| Female                | 421 | 96.2       | 16 | 3.8     | 2.0               |
| Age                   |     |            |    |         |                   |
| 18-34                 | 174 | 92.6       | 10 | 7.4     | 4.9               |
| 35-54                 | 243 | 88.8       | 27 | 11.2    | 4.6               |
| ≥55                   | 163 | 99.1       | 1  | 0.9     | 1.7               |
| Education             |     |            |    |         |                   |
| ≤High School          | 316 | 94.7       | 17 | 5.3     | 3.1               |
| >High School          | 268 | 90.8       | 20 | 9.2     | 4.3               |

Time in Past 12 Months when Needed Medical Care but Could Not Get It *Bronx* 

| Demographic<br>Groups | n1 | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|------------|-----|---------|-------------------|
| Total                 | 80 | 14.0       | 548 | 86.0    | 3.6               |
| Gender                |    |            |     |         |                   |
| Male                  | 24 | 14.9       | 167 | 85.1    | 6.4               |
| Female                | 56 | 13.2       | 381 | 86.8    | 3.8               |
| Age                   |    |            |     |         |                   |
| 18-34                 | 23 | 14.5       | 163 | 85.5    | 6.8               |
| 35-54                 | 49 | 18.6       | 222 | 81.4    | 5.6               |
| ≥55                   | 8  | 6.9        | 153 | 93.1    | 5.5               |
| Education             |    |            |     |         |                   |
| ≤High School          | 45 | 16.7       | 289 | 83.3    | 5.5               |
| >High School          | 35 | 10.8       | 253 | 89.2    | 4.0               |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days **Bronx** 

| Demographic<br>Groups | n1  | Yes<br>% | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|-----|----------|-----|---------|-------------------|
| Total                 | 393 | 62.9     | 238 | 37.1    | 4.5               |
| Gender                |     |          |     |         |                   |
| Male                  | 130 | 68.6     | 60  | 31.4    | 7.5               |
| Female                | 263 | 58.3     | 178 | 41.7    | 5.4               |
| Age                   |     |          |     |         |                   |
| 18-34                 | 119 | 64.9     | 67  | 35.1    | 8.2               |
| 35-54                 | 174 | 65.4     | 98  | 34.6    | 6.6               |
| ≥55                   | 93  | 56.0     | 70  | 44.0    | 9.1               |
| Education             |     |          |     |         |                   |
| ≤High School          | 182 | 55.6     | 154 | 44.4    | 6.5               |
| >High School          | 209 | 73.0     | 80  | 27.0    | 5.9               |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

### Expanded BRFSS, 2003 - New York State Department of Health **Physician Diagnosed Diabetes Bronx**

| Demographic<br>Groups | n1 | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|------------|-----|---------|-------------------|
| Total                 | 68 | 10.0       | 562 | 90.0    | 2.7               |
| Gender                |    |            |     |         |                   |
| Male                  | 24 | 11.1       | 166 | 88.9    | 4.8               |
| Female                | 44 | 9.2        | 396 | 90.8    | 3.0               |
| Age                   |    |            |     |         |                   |
| 18-34                 | 2  | 0.8        | 184 | 99.2    | 1.2               |
| 35-54                 | 25 | 8.1        | 246 | 91.9    | 3.6               |
| ≥55                   | 40 | 27.0       | 123 | 73.0    | 8.3               |
| Education             |    |            |     |         |                   |
| ≤High School          | 49 | 13.7       | 286 | 86.3    | 4.2               |
| >High School          | 18 | 5.4        | 271 | 94.6    | 3.0               |

### Expanded BRFSS, 2003 - New York State Department of Health Lifetime Asthma Diagnosed by a Medical Professional **Bronx**

| Demographic<br>Groups | n1 | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|------------|-----|---------|-------------------|
| Total                 | 93 | 14.0       | 539 | 86.0    | 3.2               |
| Gender                |    |            |     |         |                   |
| Male                  | 17 | 9.1        | 174 | 90.9    | 4.7               |
| Female                | 76 | 18.0       | 365 | 82.0    | 4.2               |
| Age                   |    |            |     |         |                   |
| 18-34                 | 29 | 15.2       | 157 | 84.8    | 5.8               |
| 35-54                 | 46 | 14.8       | 226 | 85.2    | 4.6               |
| ≥55                   | 17 | 11.0       | 147 | 89.0    | 6.3               |
| Education             |    |            |     |         |                   |
| ≤High School          | 47 | 12.0       | 290 | 88.0    | 4.0               |
| >High School          | 46 | 17.1       | 243 | 82.9    | 5.1               |

### Expanded BRFSS, 2003 - New York State Department of Health **Current Asthma Diagnosed by a Medical Professional Bronx**

| Demographic<br>Groups | n1 | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|------------|-----|---------|-------------------|
| Total                 | 70 | 10.9       | 562 | 89.1    | 2.9               |
| Gender                |    |            |     |         |                   |
| Male                  | 11 | 6.2        | 180 | 93.8    | 4.0               |
| Female                | 59 | 14.8       | 382 | 85.2    | 4.0               |
| Age                   |    |            |     |         |                   |
| 18-34                 | 19 | 10.6       | 167 | 89.4    | 5.0               |
| 35-54                 | 38 | 12.7       | 234 | 87.3    | 4.4               |
| ≥55                   | 13 | 9.1        | 151 | 90.9    | 6.1               |
| Education             |    |            |     |         |                   |
| ≤High School          | 40 | 10.3       | 297 | 89.7    | 3.8               |
| >High School          | 30 | 12.1       | 259 | 87.9    | 4.5               |

### Expanded BRFSS, 2003 - New York State Department of Health Arthritis Diagnosed by a Medical Professional **Bronx**

| Demographic<br>Groups | n1  | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|-----|------------|-----|---------|-------------------|
| Total                 | 143 | 20.7       | 485 | 79.3    | 3.7               |
| Gender                |     |            |     |         |                   |
| Male                  | 29  | 14.9       | 160 | 85.1    | 5.8               |
| Female                | 114 | 25.4       | 325 | 74.6    | 4.7               |
| Age                   |     |            |     |         |                   |
| 18-34                 | 6   | 3.4        | 179 | 96.6    | 3.2               |
| 35-54                 | 53  | 20.2       | 218 | 79.8    | 5.8               |
| ≥55                   | 80  | 47.2       | 82  | 52.8    | 9.2               |
| Education             |     |            |     |         |                   |
| ≤High School          | 91  | 25.4       | 243 | 74.6    | 5.5               |
| >High School          | 51  | 15.0       | 237 | 85.0    | 4.7               |

## Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

### **Bronx**

| Demographic<br>Groups | n1  | Doctor<br>Diagnosed<br>% | C.I. <sup>3</sup> | n  | Possible<br>Arthritis (CJS Only)<br>% | C.I. | n   | No Arthritis<br>or CJS<br>% | C.I. |
|-----------------------|-----|--------------------------|-------------------|----|---------------------------------------|------|-----|-----------------------------|------|
| Total                 | 143 | 20.7                     | 3.7               | 93 | 14.7                                  | 3.3  | 391 | 64.6                        | 4.5  |
| Gender                |     |                          |                   |    |                                       |      |     |                             |      |
| Male                  | 29  | 14.9                     | 5.8               | 29 | 15.5                                  | 5.9  | 131 | 69.6                        | 7.6  |
| Female                | 114 | 25.5                     | 4.7               | 64 | 14.0                                  | 3.7  | 260 | 60.5                        | 5.3  |
| Age                   |     |                          |                   |    |                                       |      |     |                             |      |
| 18-34                 | 6   | 3.4                      | 3.2               | 23 | 12.5                                  | 5.7  | 156 | 84.1                        | 6.3  |
| 35-54                 | 53  | 20.2                     | 5.8               | 52 | 18.6                                  | 5.3  | 165 | 61.2                        | 6.9  |
| ≥55                   | 80  | 47.2                     | 9.2               | 17 | 12.3                                  | 6.5  | 65  | 40.5                        | 9.0  |
| Education             |     |                          |                   |    |                                       |      |     |                             |      |
| ≤High School          | 91  | 25.4                     | 5.5               | 50 | 14.1                                  | 4.5  | 192 | 60.4                        | 6.4  |
| >High School          | 51  | 15.0                     | 4.7               | 42 | 15.8                                  | 5.0  | 195 | 69.3                        | 6.3  |

### Expanded BRFSS, 2003 - New York State Department of Health **Ever Smoked 100 Cigarettes in Lifetime Bronx**

| Demographic<br>Groups | n1  | Yes<br>% | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|-----|----------|-----|---------|-------------------|
| Total                 | 226 | 35.3     | 402 | 64.7    | 4.5               |
| Gender                |     |          |     |         |                   |
| Male                  | 86  | 42.1     | 103 | 57.9    | 8.1               |
| Female                | 140 | 29.8     | 299 | 70.2    | 4.8               |
| Age                   |     |          |     |         |                   |
| 18-34                 | 40  | 20.5     | 145 | 79.5    | 7.0               |
| 35-54                 | 113 | 44.5     | 158 | 55.5    | 7.0               |
| ≥55                   | 67  | 42.8     | 95  | 57.2    | 9.1               |
| Education             |     |          |     |         |                   |
| ≤High School          | 124 | 37.2     | 210 | 62.8    | 6.4               |
| >High School          | 101 | 33.8     | 187 | 66.2    | 6.5               |

### Expanded BRFSS, 2003 - New York State Department of Health Current Smoker Bronx

### Demographic Yes No C.I.<sup>3</sup> Groups n1 **%**⁰ % n Total 130 18.4 498 81.6 3.5 Gender Male 39 17.1 150 82.9 5.8 80.5 4.2 Female 91 19.5 348 Age 18-34 14.1 85.9 5.7 31 154 35-54 71 25.4 200 74.6 6.0 ≥55 26 14.3 136 85.7 6.0 Education 70 18.1 264 81.9 4.6 ≤High School 59 19.3 229 80.7 5.4 >High School

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

# Expanded BRFSS, 2003 - New York State Department of Health **Status of Cigarette Smoking** *Bronx*

| Demographic<br>Groups | n1 | Everyday Smoker | C.I. <sup>3</sup> | n  | Someday Smoker<br>% | C.I. | n   | Former or<br>Never Smoked<br>% | C.I. |
|-----------------------|----|-----------------|-------------------|----|---------------------|------|-----|--------------------------------|------|
| Total                 | 89 | 12.2            | 2.8               | 41 | 6.3                 | 2.3  | 498 | 81.6                           | 3.5  |
| Gender                |    |                 |                   |    |                     |      |     |                                |      |
| Male                  | 28 | 12.0            | 4.8               | 11 | 5.1                 | 3.7  | 150 | 82.9                           | 5.8  |
| Female                | 61 | 12.3            | 3.3               | 30 | 7.2                 | 2.8  | 348 | 80.5                           | 4.2  |
| Age                   |    |                 |                   |    |                     |      |     |                                |      |
| 18-34                 | 17 | 5.9             | 3.1               | 14 | 8.2                 | 5.0  | 154 | 85.9                           | 5.7  |
| 35-54                 | 53 | 19.8            | 5.6               | 18 | 5.6                 | 2.8  | 200 | 74.6                           | 6.0  |
| ≥55                   | 17 | 9.6             | 5.2               | 9  | 4.7                 | 3.4  | 136 | 85.7                           | 6.0  |
| Education             |    |                 |                   |    |                     |      |     |                                |      |
| ≤High School          | 50 | 12.7            | 3.9               | 20 | 5.4                 | 2.6  | 264 | 81.9                           | 4.6  |
| >High School          | 38 | 11.7            | 4.1               | 21 | 7.6                 | 4.1  | 229 | 80.7                           | 5.4  |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

### Expanded BRFSS, 2003 - New York State Department of Health Home Smoking Policies Bronx

| Demographic<br>Groups | n1  | Not Allowed | C.I. <sup>3</sup> | n  | Allowed Some<br>Places<br>% | C.I. | n   | Allowed Anywhere % | C.I. |
|-----------------------|-----|-------------|-------------------|----|-----------------------------|------|-----|--------------------|------|
| Total                 | 435 | 69.9        | 4.3               | 80 | 12.1                        | 3.0  | 114 | 18.0               | 3.7  |
| Gender                |     |             |                   |    |                             |      |     |                    |      |
| Male                  | 123 | 68.0        | 7.5               | 22 | 11.2                        | 5.0  | 44  | 20.8               | 6.5  |
| Female                | 312 | 71.4        | 4.9               | 58 | 12.9                        | 3.5  | 70  | 15.7               | 4.0  |
| Age                   |     |             |                   |    |                             |      |     |                    |      |
| 18-34                 | 132 | 70.9        | 8.0               | 24 | 10.7                        | 4.9  | 29  | 18.4               | 7.1  |
| 35-54                 | 188 | 69.9        | 6.4               | 41 | 15.4                        | 5.0  | 42  | 14.7               | 4.9  |
| ≥55                   | 109 | 69.2        | 8.2               | 14 | 9.2                         | 5.4  | 40  | 21.6               | 7.0  |
| Education             |     |             |                   |    |                             |      |     |                    |      |
| ≤High School          | 228 | 67.9        | 6.1               | 44 | 12.0                        | 4.0  | 63  | 20.1               | 5.4  |
| >High School          | 203 | 72.2        | 6.0               | 36 | 12.8                        | 4.5  | 49  | 15.0               | 4.5  |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

### Expanded BRFSS, 2003 - New York State Department of Health Home Smoking Bronx

### Demographic Yes No C.I.<sup>3</sup> Groups n1 **%**⁰ % n 77.4 4.0 Total 142 22.6 489 Gender Male 53 26.5 138 73.5 7.3 Female 89 19.4 351 80.6 4.2 Age 18-34 24.7 75.3 7.9 42 144 35-54 24.1 67 204 75.9 5.8 ≥55 31 17.6 133 82.4 6.8 Education 74 22.9 262 77.1 5.7 ≤High School 66 20.9 223 79.1 5.3 >High School

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink **Occasion**)

**Bronx** 

| Demographic  |    | Yes                   |     | No   |                   |
|--------------|----|-----------------------|-----|------|-------------------|
| Groups       | n1 | <b>%</b> <sup>2</sup> | n   | %    | C.I. <sup>3</sup> |
| Total        | 52 | 9.5                   | 568 | 90.5 | 2.9               |
| Gender       |    |                       |     |      |                   |
| Male         | 28 | 13.6                  | 156 | 86.4 | 5.5               |
| Female       | 24 | 6.3                   | 412 | 93.7 | 2.8               |
| Age          |    |                       |     |      |                   |
| 18-34        | 19 | 11.8                  | 164 | 88.2 | 5.7               |
| 35-54        | 22 | 9.3                   | 243 | 90.7 | 4.3               |
| ≥55          | 9  | 6.3                   | 153 | 93.7 | 4.8               |
| Education    |    |                       |     |      |                   |
| ≤High School | 28 | 10.2                  | 301 | 89.8 | 4.3               |
| >High School | 24 | 9.0                   | 261 | 91.0 | 3.9               |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 **Drink per Day in Past Month)** 

### **Bronx**

| Demographic<br>Groups | n <sup>1</sup> | Yes<br>% | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----------------|----------|-----|---------|-------------------|
| Total                 | 20             | 3.1      | 602 | 96.9    | 1.5               |
| Gender                |                |          |     |         |                   |
| Male                  | 9              | 3.4      | 176 | 96.6    | 2.5               |
| Female                | 11             | 2.9      | 426 | 97.1    | 1.9               |
| Age                   |                |          |     |         |                   |
| 18-34                 | 5              | 2.5      | 179 | 97.5    | 2.3               |
| 35-54                 | 11             | 4.5      | 257 | 95.5    | 2.9               |
| ≥55                   | 4              | 1.9      | 157 | 98.1    | 2.4               |
| Education             |                |          |     |         |                   |
| ≤High School          | 10             | 3.4      | 318 | 96.6    | 2.3               |
| >High School          | 10             | 2.8      | 278 | 97.2    | 1.9               |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

### Expanded BRFSS, 2003 - New York State Department of Health Overweight or Obese, BMI(Body Mass Index) 25 or More *Bronx*

| Demographic<br>Groups | n1  | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|-----|------------|-----|---------|-------------------|
| Total                 | 366 | 62.7       | 224 | 37.3    | 4.7               |
| Gender                |     |            |     |         |                   |
| Male                  | 127 | 67.8       | 60  | 32.2    | 7.8               |
| Female                | 239 | 58.2       | 164 | 41.8    | 5.6               |
| Age                   |     |            |     |         |                   |
| 18-34                 | 88  | 49.4       | 84  | 50.6    | 9.2               |
| 35-54                 | 169 | 68.3       | 91  | 31.7    | 6.4               |
| ≥55                   | 105 | 72.9       | 46  | 27.1    | 8.0               |
| Education             |     |            |     |         |                   |
| ≤High School          | 200 | 65.7       | 107 | 34.3    | 6.5               |
| >High School          | 162 | 58.4       | 116 | 41.6    | 6.9               |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

### Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+) Bronx

| Demographic<br>Groups | n1  | Underweight\<br>Normal<br>%02 | C.I. <sup>3</sup> | n   | Overweight % | C.I. | n   | Obese<br>% | C.I. |
|-----------------------|-----|-------------------------------|-------------------|-----|--------------|------|-----|------------|------|
| Total                 | 224 | 37.3                          | 4.7               | 213 | 38.1         | 4.9  | 153 | 24.6       | 4.1  |
| Gender                |     |                               |                   |     |              |      |     |            |      |
| Male                  | 60  | 32.2                          | 7.8               | 81  | 44.8         | 8.3  | 46  | 23.0       | 6.7  |
| Female                | 164 | 41.8                          | 5.6               | 132 | 32.2         | 5.2  | 107 | 26.0       | 5.0  |
| Age                   |     |                               |                   |     |              |      |     |            |      |
| 18-34                 | 84  | 50.6                          | 9.2               | 53  | 29.8         | 8.6  | 35  | 19.6       | 7.0  |
| 35-54                 | 91  | 31.7                          | 6.4               | 95  | 41.3         | 7.2  | 74  | 27.0       | 6.2  |
| ≥55                   | 46  | 27.1                          | 8.0               | 62  | 44.1         | 9.6  | 43  | 28.8       | 8.6  |
| Education             |     |                               |                   |     |              |      |     |            |      |
| ≤High School          | 107 | 34.3                          | 6.5               | 112 | 38.9         | 6.9  | 88  | 26.7       | 5.8  |
| >High School          | 116 | 41.6                          | 6.9               | 98  | 36.8         | 6.8  | 64  | 21.6       | 5.6  |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

Ever Had Mammography Screening, Women Aged 40 Years and Older *Bronx* 

| Demographic<br>Groups | n1  | Yes<br>% | n  | No<br>% | C.I. <sup>3</sup> |
|-----------------------|-----|----------|----|---------|-------------------|
| Female                | 215 | 91.1     | 21 | 8.9     | 4.0               |
| Age                   |     |          |    |         |                   |
| 40-64                 | 168 | 90.3     | 16 | 9.7     | 5.1               |
| ≥65                   | 47  | 93.0     | 5  | 7.0     | 6.2               |
| Education             |     |          |    |         |                   |
| ≤High School          | 132 | 91.1     | 13 | 8.9     | 5.0               |
| >High School          | 81  | 91.0     | 8  | 9.0     | 6.7               |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

**Bronx** 

| Demographic<br>Groups | n1  | Yes<br>% | n  | No<br>% | C.I. <sup>3</sup> |
|-----------------------|-----|----------|----|---------|-------------------|
| Female                | 189 | 82.0     | 44 | 18.0    | 5.4               |
| Age                   |     |          |    |         |                   |
| 40-64                 | 150 | 82.6     | 32 | 17.4    | 6.2               |
| ≥65                   | 39  | 80.6     | 12 | 19.4    | 10.8              |
| Education             |     |          |    |         |                   |
| ≤High School          | 117 | 82.6     | 26 | 17.4    | 6.7               |
| >High School          | 70  | 80.5     | 18 | 19.5    | 9.2               |

Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger

**Bronx** 

| Demographic<br>Groups | n1  | Yes<br>% | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|-----|----------|-----|---------|-------------------|
| Total                 | 200 | 37.2     | 328 | 62.8    | 5.1               |
| Gender                |     |          |     |         |                   |
| Male                  | 46  | 29.5     | 118 | 70.5    | 8.3               |
| Female                | 154 | 43.9     | 210 | 56.1    | 5.9               |
| Age                   |     |          |     |         |                   |
| 18-34                 | 99  | 50.7     | 84  | 49.3    | 8.8               |
| 35-54                 | 82  | 29.1     | 181 | 70.9    | 6.3               |
| 55-64                 | 19  | 17.8     | 63  | 82.2    | 8.3               |
| Education             |     |          |     |         |                   |
| ≤High School          | 118 | 44.2     | 145 | 55.8    | 7.5               |
| >High School          | 80  | 30.2     | 179 | 69.8    | 6.4               |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

### Expanded BRFSS, 2003 - New York State Department of Health Sexually Active, Age 18-64 Years Old **Bronx**

| Demographic<br>Groups | n1  | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|-----|------------|-----|---------|-------------------|
| Total                 | 345 | 75.0       | 149 | 25.0    | 4.4               |
| Gender                |     |            |     |         |                   |
| Male                  | 127 | 85.1       | 23  | 14.9    | 6.4               |
| Female                | 218 | 66.4       | 126 | 33.6    | 5.7               |
| Age                   |     |            |     |         |                   |
| 18-34                 | 138 | 85.9       | 28  | 14.1    | 6.0               |
| 35-54                 | 169 | 70.5       | 79  | 29.5    | 6.6               |
| 55-64                 | 38  | 54.6       | 42  | 45.4    | 13.2              |
| Education             |     |            |     |         |                   |
| ≤High School          | 154 | 68.4       | 91  | 31.6    | 6.9               |
| >High School          | 190 | 83.1       | 55  | 16.9    | 5.1               |

Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger **Bronx** 

| Demographic<br>Groups | n1  | Yes<br>% | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|-----|----------|-----|---------|-------------------|
| Total                 | 137 | 40.3     | 220 | 59.7    | 6.4               |
| Gender                |     |          |     |         |                   |
| Male                  | 64  | 49.1     | 70  | 50.9    | 9.9               |
| Female                | 73  | 30.2     | 150 | 69.8    | 6.9               |
| Age                   |     |          |     |         |                   |
| 18-34                 | 66  | 50.1     | 78  | 49.9    | 10.0              |
| 35-54                 | 63  | 33.1     | 111 | 66.9    | 8.0               |
| 55-64                 | 8   | 19.7     | 31  | 80.3    | 15.5              |
| Education             |     |          |     |         |                   |
| ≤High School          | 61  | 41.9     | 103 | 58.1    | 9.7               |
| >High School          | 75  | 37.8     | 116 | 62.2    | 8.1               |

### **Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59)** and Women(18-44)

### **Bronx**

| Demographic  |     | Use BC                       |                   |    | No BC at risk |      |     | No BC |      |
|--------------|-----|------------------------------|-------------------|----|---------------|------|-----|-------|------|
| Groups       | n1  | <sup>0</sup> /0 <sup>2</sup> | C.I. <sup>3</sup> | n  | %             | C.I. | n   | %     | C.I. |
| Total        | 216 | 59.4                         | 6.0               | 56 | 14.5          | 4.2  | 104 | 26.1  | 5.3  |
| Gender       |     |                              |                   |    |               |      |     |       |      |
| Male         | 89  | 59.5                         | 9.1               | 22 | 15.1          | 6.4  | 38  | 25.5  | 8.1  |
| Female       | 127 | 59.2                         | 7.3               | 34 | 13.9          | 5.1  | 66  | 26.9  | 6.5  |
| Age          |     |                              |                   |    |               |      |     |       |      |
| 18-34        | 109 | 65.0                         | 8.5               | 22 | 12.4          | 5.9  | 44  | 22.6  | 7.4  |
| 35-59        | 107 | 52.9                         | 8.1               | 34 | 16.9          | 6.0  | 60  | 30.2  | 7.6  |
| Education    |     |                              |                   |    |               |      |     |       |      |
| ≤High School | 90  | 54.0                         | 9.0               | 30 | 15.7          | 6.0  | 60  | 30.3  | 8.1  |
| >High School | 126 | 66.6                         | 7.8               | 24 | 11.8          | 5.3  | 43  | 21.6  | 6.9  |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

### Expanded BRFSS, 2003 - New York State Department of Health **Physician Diagnosed Heart Attack, Angina or Stroke** *Bronx*

| Demographic<br>Groups | n1 | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|------------|-----|---------|-------------------|
| Total                 | 40 | 5.5        | 579 | 94.5    | 1.9               |
| Gender                |    |            |     |         |                   |
| Male                  | 10 | 4.4        | 179 | 95.6    | 3.0               |
| Female                | 30 | 6.5        | 400 | 93.5    | 2.5               |
| Age                   |    |            |     |         |                   |
| 18-64                 | 25 | 3.5        | 509 | 96.5    | 1.5               |
| ≥65                   | 14 | 18.1       | 62  | 81.9    | 9.6               |
| Education             |    |            |     |         |                   |
| ≤High School          | 31 | 7.8        | 298 | 92.2    | 3.1               |
| >High School          | 8  | 2.5        | 276 | 97.5    | 2.0               |

Expanded BRFSS, 2003 - New York State Department of Health **Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina)** *Bronx* 

| Demographic<br>Groups | n1 | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|------------|-----|---------|-------------------|
| Total                 | 27 | 4.1        | 591 | 95.9    | 1.7               |
| Gender                |    |            |     |         |                   |
| Male                  | 6  | 3.0        | 183 | 97.0    | 2.7               |
| Female                | 21 | 4.9        | 408 | 95.1    | 2.2               |
| Age                   |    |            |     |         |                   |
| 18-64                 | 17 | 2.5        | 516 | 97.5    | 1.3               |
| ≥65                   | 10 | 13.7       | 66  | 86.3    | 8.9               |
| Education             |    |            |     |         |                   |
| ≤High School          | 20 | 5.5        | 309 | 94.5    | 2.7               |
| >High School          | 7  | 2.3        | 276 | 97.7    | 1.9               |

### Expanded BRFSS, 2003 - New York State Department of Health Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older *Bronx*

| Demographic<br>Groups | n1 | Yes<br>%02 | n  | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|------------|----|---------|-------------------|
| Male                  | 65 | 57.0       | 51 | 43.0    | 10.2              |
| Age                   |    |            |    |         |                   |
| 40-64                 | 47 | 49.8       | 47 | 50.2    | 11.5              |
| ≥65                   | 18 | 83.5       | 4  | 16.5    | 16.1              |
| Education             |    |            |    |         |                   |
| ≤High School          | 33 | 56.0       | 27 | 44.0    | 14.1              |
| >High School          | 32 | 58.4       | 24 | 41.6    | 14.5              |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

**Bronx** 

| Demographic<br>Groups | n <sup>1</sup> | Yes<br>%02 | n  | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----------------|------------|----|---------|-------------------|
| Male                  | 55             | 50.1       | 61 | 49.9    | 10.4              |
| Age                   |                |            |    |         |                   |
| 40-64                 | 42             | 44.7       | 52 | 55.3    | 11.5              |
| ≥65                   | 13             | 69.8       | 9  | 30.2    | 19.6              |
| Education             |                |            |    |         |                   |
| ≤High School          | 27             | 49.9       | 33 | 50.1    | 14.3              |
| >High School          | 28             | 50.4       | 28 | 49.6    | 14.9              |

### Expanded BRFSS, 2003 - New York State Department of Health Ever Had a Digital Rectal Exam, Men Aged 40 and Older *Bronx*

| Demographic<br>Groups | n1 | Yes<br>% | n  | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|----------|----|---------|-------------------|
| Male                  | 78 | 64.8     | 43 | 35.2    | 9.7               |
| Age                   |    |          |    |         |                   |
| 40-64                 | 59 | 58.9     | 41 | 41.1    | 11.0              |
| ≥65                   | 19 | 88.4     | 2  | 11.6    | 15.3              |
| Education             |    |          |    |         |                   |
| ≤High School          | 35 | 59.8     | 25 | 40.2    | 14.0              |
| >High School          | 43 | 70.6     | 18 | 29.4    | 13.0              |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent

### Expanded BRFSS, 2003 - New York State Department of Health Ever Used a Blood Stool Test at Home, Aged 50 and Older *Bronx*

| Demographic<br>Groups | n1 | Yes<br>% | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|----------|-----|---------|-------------------|
| Total                 | 65 | 28.0     | 146 | 72.0    | 7.1               |
| Gender                |    |          |     |         |                   |
| Male                  | 18 | 22.3     | 45  | 77.7    | 10.9              |
| Female                | 47 | 31.7     | 101 | 68.3    | 9.1               |
| Age                   |    |          |     |         |                   |
| 50-64                 | 48 | 29.0     | 95  | 71.0    | 8.6               |
| ≥65                   | 17 | 26.6     | 51  | 73.4    | 12.1              |
| Education             |    |          |     |         |                   |
| ≤High School          | 33 | 22.3     | 99  | 77.7    | 8.0               |
| >High School          | 32 | 40.6     | 45  | 59.4    | 13.2              |

Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older *Bronx* 

| Demographic<br>Groups | n1 | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|------------|-----|---------|-------------------|
| Total                 | 30 | 14.2       | 179 | 85.8    | 5.7               |
| Gender                |    |            |     |         |                   |
| Male                  | 9  | 11.8       | 54  | 88.2    | 8.6               |
| Female                | 21 | 15.7       | 125 | 84.3    | 7.5               |
| Age                   |    |            |     |         |                   |
| 50-64                 | 26 | 18.5       | 115 | 81.5    | 7.7               |
| ≥65                   | 4  | 7.9        | 64  | 92.1    | 8.0               |
| Education             |    |            |     |         |                   |
| ≤High School          | 14 | 10.3       | 117 | 89.7    | 5.9               |
| >High School          | 16 | 22.5       | 60  | 77.5    | 12.1              |

Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older *Bronx* 

| Demographic<br>Groups | n1 | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|------------|-----|---------|-------------------|
| Total                 | 40 | 18.4       | 169 | 81.6    | 6.2               |
| Gender                |    |            |     |         |                   |
| Male                  | 13 | 17.1       | 50  | 82.9    | 9.9               |
| Female                | 27 | 19.2       | 119 | 80.8    | 7.9               |
| Age                   |    |            |     |         |                   |
| 50-64                 | 32 | 21.6       | 109 | 78.4    | 8.0               |
| ≥65                   | 8  | 13.8       | 60  | 86.2    | 9.6               |
| Education             |    |            |     |         |                   |
| ≤High School          | 21 | 14.5       | 110 | 85.5    | 6.6               |
| >High School          | 19 | 27.0       | 57  | 73.0    | 12.7              |

Expanded BRFSS, 2003 - New York State Department of Health Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older *Bronx* 

| Demographic<br>Groups | n1 | Yes<br>%02 | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|------------|-----|---------|-------------------|
| Total                 | 90 | 44.5       | 126 | 55.5    | 8.0               |
| Gender                |    |            |     |         |                   |
| Male                  | 23 | 41.4       | 40  | 58.6    | 14.4              |
| Female                | 67 | 46.5       | 86  | 53.5    | 9.3               |
| Age                   |    |            |     |         |                   |
| 50-64                 | 53 | 34.6       | 91  | 65.4    | 9.2               |
| ≥65                   | 37 | 57.8       | 35  | 42.2    | 12.7              |
| Education             |    |            |     |         |                   |
| ≤High School          | 54 | 43.6       | 83  | 56.4    | 9.9               |
| >High School          | 35 | 46.4       | 42  | 53.6    | 13.3              |

Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older *Bronx* 

| Demographic<br>Groups | n1 | Yes<br>% | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|----------|-----|---------|-------------------|
| Total                 | 84 | 43.0     | 128 | 57.0    | 8.1               |
| Gender                |    |          |     |         |                   |
| Male                  | 21 | 39.7     | 41  | 60.3    | 14.5              |
| Female                | 63 | 45.1     | 87  | 54.9    | 9.4               |
| Age                   |    |          |     |         |                   |
| 50-64                 | 50 | 33.4     | 92  | 66.6    | 9.2               |
| ≥65                   | 34 | 56.1     | 36  | 43.9    | 13.0              |
| Education             |    |          |     |         |                   |
| ≤High School          | 51 | 42.7     | 84  | 57.3    | 10.0              |
| >High School          | 32 | 43.7     | 43  | 56.3    | 13.5              |

Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older

**Bronx** 

| Demographic<br>Groups | n1 | Yes<br>% | n   | No<br>% | C.I. <sup>3</sup> |
|-----------------------|----|----------|-----|---------|-------------------|
| Total                 | 99 | 47.9     | 119 | 52.1    | 7.9               |
| Gender                |    |          |     |         |                   |
| Male                  | 25 | 42.2     | 40  | 57.8    | 14.2              |
| Female                | 74 | 51.6     | 79  | 48.4    | 9.2               |
| Age                   |    |          |     |         |                   |
| 50-64                 | 64 | 42.5     | 80  | 57.5    | 9.6               |
| ≥65                   | 35 | 55.0     | 39  | 45.0    | 12.7              |
| Education             |    |          |     |         |                   |
| ≤High School          | 58 | 46.5     | 81  | 53.5    | 9.9               |
| >High School          | 40 | 51.0     | 37  | 49.0    | 13.3              |

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution <sup>2</sup>Weighted Percent