Medicaid Disability Manual

- **G.** Absolute CD4 count of less than 200 cells/mm3 or CD4 percentage of less than 14 percent, and one of the following (values do not have to be measured on the same date) (see 14.00F5):
- 1. BMI measurement of less than 18.5; or
- 2. Hemoglobin measurement of less than 8.0 grams per deciliter (g/dL).

OR

H. Complication(s) of HIV infection requiring at least three hospitalizations within a 12-month period and at least 30 days apart (see 14.00F6). Each hospitalization must last at least 48 hours, including hours in a hospital emergency department immediately before the hospitalization.

OR

- I. Repeated (as defined in 14.00I3) manifestations of HIV infection, including those listed in 14.11A-H, but without the requisite findings for those listings (for example, Kaposi sarcoma not meeting the criteria in 14.11E), or other manifestations (including, but not limited to, cardiovascular disease (including myocarditis, pericardial effusion, pericarditis, endocarditis, or pulmonary arteritis), diarrhea, distal sensory polyneuropathy, glucose intolerance, gynecologic conditions (including cervical cancer or pelvic inflammatory disease, see 14.00F7), hepatitis, HIV-associated dementia, immune reconstitution inflammatory syndrome (IRIS), infections (bacterial, fungal, parasitic, or viral), lipodystrophy (lipoatrophy or lipohypertrophy), malnutrition, muscle weakness, myositis, neurocognitive or other mental limitations not meeting the criteria in 12.00, oral hairy leukoplakia, osteoporosis, pancreatitis, peripheral neuropathy) resulting in significant, documented symptoms or signs (for example, but not limited to, fever, headaches, insomnia, involuntary weight loss, malaise, nausea, night sweats, pain, severe fatigue, or vomiting) and one of the following at the marked level:
- 1. Limitation of activities of daily living.
- 2. Limitation in maintaining social functioning.
- 3. Limitation in completing tasks in a timely manner due to deficiencies in concentration, persistence, or pace.

Part B – Section Overview

The following sections provide medical criteria for the evaluation of impairments of children under age 18. In evaluation disability for a person under age 18, Part B will be used first. If medical criteria in Part B do not apply, then the medical criteria in Part A will be used.

100.00 Low Birth Weight and Failure to Thrive

101.00 Musculoskeletal Disorders

102.00 Special Senses and Speech

103.00 Respiratory Disorders

104.00 Cardiovascular System

Medicaid Disability Manual

- 105.00 Digestive System
- 106.00 Genitourinary Disorders
- 107.00 Hematological Disorders
- 108.00 Skin Disorders
- 109.00 Endocrine Disorders
- 110.00 Congenital Disorders that Affect Multiple Body Systems
- 111.00 Neurological Disorders
- 112.00 Mental Disorders
- 113.00 Cancer (Malignant Neoplastic Diseases)
- 114.00 Immune System Disorders

100.00 Low Birth Weight and Failure to Thrive

- **A.** What conditions do we evaluate under these listings? We evaluate low birth weight (LBW) in infants from birth to attainment of age 1 and failure to thrive (FTT) in infants and toddlers from birth to attainment of age 3.
- **B.** How do we evaluate disability based on LBW under 100.04? In 100.04A and 100.04B, we use an infant's birth weight as documented by an original or certified copy of the infant's birth certificate or by a medical record signed by a physician. *Birth weight* means the first weight recorded after birth. In 100.04B, *gestational age* is the infant's age based on the date of conception as recorded in the medical record. If the infant's impairment meets the requirements for listing 100.04A or 100.04B, we will follow the rules in § 416.990(b)(11) of this chapter.

C. How do we evaluate disability under 100.05?

- 1. General. We establish FTT with or without a known cause when we have documentation of an infant's or toddler's growth failure and developmental delay from an acceptable medical source(s) as defined in § 416.913(a) of this chapter. We require documentation of growth measurements in 100.05A and developmental delay in 100.05B or 100.05C within the same consecutive 12-month period. The dates of developmental testing and reports may be different from the dates of the growth measurements. After the attainment of age 3, we evaluate growth failure under the affected body system(s).
- 2. Growth failure. Under 100.05A, we use the appropriate table(s) under 105.08B in the digestive system to determine whether a child's growth is less than the third percentile. The child does not need to have a digestive disorder for the purposes of 100.05.
- a. For children from birth to attainment of age 2, we use the weight-for-length table corresponding to the child's gender (Table I or Table II).
- b. For children age 2 to the attainment of age 3, we use the body mass index (BMI)-forage table corresponding to the child's gender (Table III or Table IV).