

**COMMUNICABLE DISEASE IN NEW YORK CITY
REPORTED CASES OF SELECTED DISEASES BY DIAGNOSIS DATE
1987 - 1996**

	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
AIDS	5164	6351	6719	7456	8689	10207	11673	9636	6555	10463
AMEBIASIS	206	102	260	411	438	465	708	701	976	869
BABESIOSIS	1	4	2	1	5	3	2	4	7	8
CAMPYLOBACTERIOSIS	267	302	667	788	899	1165	1286	1340	1265	1248
ENCEPHALITIS OTHER	8	6	9	12	5	9	4	11	8	2
GIARDIASIS	213	290	496	810	944	1093	1937	2456	2486	2287
GONORRHEA	66545	54099	40533	35236	28945	21709	18477	19246	16361	12998
HAEMOPHILUS INFLUENZAE	79	58	147	128	54	33	39	34	36	57
HEPATITIS A	172	368	502	791	1283	883	1028	942	1008	609
HEPATITIS B	1213	1307	1418	674	440	439	472	544	524	491
KAWASAKI SYNDROME	19	9	11	11	11	12	10	8	7	7
LEGIONELLOSIS (2)	27	50	65	94	82	10	6	11	6	19
LYME DISEASE	108	157	107	93	137	103	214	95	455	403
MALARIA	27	94	99	114	133	169	143	104	222	268
MEASLES	469	57	135	1108	1909	68	19	15	5	11
MENINGITIS VIRAL	187	149	199	165	426	179	273	150	157	127
MENINGOCOCCAL (1)	57	66	87	79	30	28	40	40	54	56
MENINGITIS OTHER	297	134	320	195	201	143	195	205	254	235
MUMPS	40	104	23	65	14	12	2	12	17	15
PERTUSSIS	15	11	19	21	22	24	116	223	67	62
RMSF	6	1	3	4	1	7	1	3	6	19
RUBELLA	3	7	16	4	2	0	22	1	8	5
SALMONELLOSIS (3)	1802	1687	2022	1691	1709	1831	2024	1896	2165	1925
SHIGELLOSIS	1771	482	354	673	428	753	677	1006	845	630
SYPHILIS EARLY	8043	9557	10209	11750	9902	7619	4876	2986	2306	1215
TOXIC SHOCK SYNDROME	7	8	4	7	2	0	2	0	4	4
TUBERCULOSIS (4)	2197	2317	2545	3520	3673	3811	3235	2995	2445	2053
TYPHOID FEVER	29	55	56	84	79	50	102	79	66	64
YERSINIOSIS	18	48	46	33	41	51	45	49	36	42

(1) Case definition require positive culture from blood, spinal, or other sterile sites

(2) Beginning 1991 only confirmed cases reported

(3) Salmonellosis paratyphi A and B plus Salmonellosis nontyphoidal

(4) Revised in 1978 to include persons with verified disease in past or lost to supervision for more than 12 months and had disease again