

# Expanded BRFSS, 2003

## Nassau Locality Tables

### TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days .....	1
Core 2: Health Care Access .....	4
Core 3: Exercise .....	8
Core 4: Diabetes .....	9
Core 5: Asthma .....	10
Core 6: Arthritis .....	12
Core 7: Tobacco Use .....	14
Core 8: Tobacco ETS .....	17
Core 9: Alcohol Consumption .....	19
Core 10: Weight Status .....	21
Core 11: Mammography .....	23
Core 12: Sexual Behavior .....	25
Core 13: Family Planning .....	28
Core 14: Cardiovascular Disease .....	29
Core 15: Prostate Cancer Screening .....	31
Core 16: Colorectal Cancer Screening .....	37

New York State Department Of Health

Expanded BRFSS, 2003 - New York State Department of Health

## Self-rated Health Status

*Nassau*

Demographic Groups	n <sup>1</sup>	Fair or Poor % <sup>2</sup>	n	Good to Excellent %	C.I. <sup>3</sup>
Total	55	7.9	557	92.1	2.3
Gender					
Male	23	6.6	220	93.4	3.0
Female	32	9.0	337	91.0	3.5
Age					
18-34	3	2.2	115	97.8	3.0
35-54	9	3.1	250	96.9	2.5
≥55	41	17.6	178	82.4	5.5
Education					
≤High School	30	16.6	133	83.4	6.4
>High School	25	4.5	422	95.5	2.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Reported Number of Physical Distress Days in the Past Month

*Nassau*

Demographic Groups	n <sup>1</sup>	≥14 days % <sup>2</sup>	n	< 14 days %	C.I. <sup>3</sup>
Total	44	6.6	563	93.4	2.2
Gender					
Male	17	6.6	224	93.4	3.5
Female	27	6.6	339	93.4	2.7
Age					
18-34	2	0.9	117	99.1	1.4
35-54	16	6.3	243	93.7	3.4
≥55	25	11.4	189	88.6	4.9
Education					
≤High School	14	8.4	143	91.6	5.1
>High School	30	6.0	418	94.0	2.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Reported Number of Mental Distress Days in the Past Month

*Nassau*

Demographic Groups	n <sup>1</sup>	≥14 days % <sup>2</sup>	n	< 14 days %	C.I. <sup>3</sup>
Total	42	5.8	567	94.2	1.9
Gender					
Male	9	2.9	232	97.1	2.0
Female	33	8.5	335	91.5	3.2
Age					
18-34	9	6.6	111	93.4	4.5
35-54	14	4.0	242	96.0	2.2
≥55	17	7.2	200	92.8	3.8
Education					
≤High School	12	5.6	150	94.4	3.4
>High School	30	5.9	415	94.1	2.4

<sup>1</sup> Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup> Weighted Percent

<sup>3</sup> 95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Health Insurance Coverage, 18-64 Years of Age

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	434	90.5	32	9.5	3.7
Gender					
Male	167	88.3	14	11.7	6.3
Female	267	92.7	18	7.3	3.8
Age					
18-34	108	85.4	11	14.6	8.7
35-54	242	92.0	18	8.0	4.2
55-64	84	95.6	3	4.4	5.6
Education					
≤High School	87	79.1	16	20.9	10.2
>High School	346	94.1	16	5.9	3.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Health Insurance Coverage, Aged 18 and Older

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	580	92.1	36	7.9	3.0
Gender					
Male	229	90.0	15	10.0	5.3
Female	351	93.9	21	6.1	2.9
Age					
18-34	108	85.4	11	14.6	8.7
35-54	242	92.0	18	8.0	4.2
≥55	217	97.6	4	2.4	2.6
Education					
≤High School	143	83.8	20	16.2	7.4
>High School	435	95.2	16	4.8	2.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Reported Having a Routine Physical Exam in Past Two Years

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	567	92.5	47	7.5	2.4
Gender					
Male	217	90.4	25	9.6	4.1
Female	350	94.3	22	5.7	2.6
Age					
18-34	104	89.8	14	10.2	5.7
35-54	237	91.6	23	8.4	3.7
≥55	210	94.9	10	5.1	3.5
Education					
≤High School	149	90.8	15	9.2	4.9
>High School	416	93.0	32	7.0	2.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Time in Past 12 Months when Needed Medical Care but Could Not Get It

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	15	2.7	601	97.3	1.5
Gender					
Male	5	1.6	239	98.4	1.5
Female	10	3.7	362	96.3	2.5
Age					
18-34	4	4.6	114	95.4	4.5
35-54	7	2.6	254	97.4	2.1
≥55	2	0.8	219	99.2	1.3
Education					
≤High School	5	4.2	158	95.8	3.8
>High School	10	2.1	441	97.9	1.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

## Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days

*Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	485	75.6	132	24.4	4.3
Gender					
Male	201	77.4	44	22.6	7.0
Female	284	74.0	88	26.0	5.2
Age					
18-34	91	72.6	29	27.4	10.2
35-54	215	79.7	46	20.3	6.1
≥55	165	72.5	55	27.5	7.0
Education					
≤High School	108	63.4	56	36.6	8.7
>High School	376	80.4	75	19.6	4.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Physician Diagnosed Diabetes

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	43	5.8	575	94.2	1.9
Gender					
Male	22	6.2	223	93.8	2.9
Female	21	5.4	352	94.6	2.6
Age					
18-34	1	0.4	119	99.6	0.8
35-54	16	6.2	245	93.8	3.5
≥55	24	9.0	197	91.0	3.9
Education					
≤High School	17	9.6	148	90.4	5.1
>High School	26	4.3	425	95.7	1.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Lifetime Asthma Diagnosed by a Medical Professional

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	58	9.3	560	90.7	2.7
Gender					
Male	14	5.1	231	94.9	3.2
Female	44	13.0	329	87.0	4.1
Age					
18-34	19	13.4	101	86.6	6.8
35-54	17	7.6	244	92.4	3.9
≥55	21	8.2	200	91.8	3.9
Education					
≤High School	15	8.4	150	91.6	4.8
>High School	43	9.6	408	90.4	3.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Current Asthma Diagnosed by a Medical Professional

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	40	6.3	578	93.7	2.3
Gender					
Male	7	2.7	238	97.3	2.5
Female	33	9.6	340	90.4	3.6
Age					
18-34	11	8.4	109	91.6	5.7
35-54	13	5.7	248	94.3	3.4
≥55	16	5.9	205	94.1	3.2
Education					
≤High School	12	7.0	153	93.0	4.5
>High School	28	6.1	423	93.9	2.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Arthritis Diagnosed by a Medical Professional

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	177	25.8	437	74.2	3.9
Gender					
Male	62	20.2	181	79.8	5.6
Female	115	30.8	256	69.2	5.3
Age					
18-34	9	4.7	111	95.3	3.1
35-54	49	19.3	209	80.7	5.6
≥55	114	48.9	106	51.1	7.7
Education					
≤High School	60	34.2	104	65.8	8.4
>High School	117	22.7	331	77.3	4.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

*Nassau*

Demographic Groups	n <sup>1</sup>	Doctor Diagnosed % <sup>2</sup>	C.I. <sup>3</sup>	n	Possible Arthritis (CJS Only) %	C.I.	n	No Arthritis or CJS %	C.I.
Total	177	25.8	3.9	77	12.4	3.0	360	61.8	4.4
Gender									
Male	62	20.2	5.6	29	11.7	4.6	152	68.1	6.7
Female	115	30.8	5.3	48	13.0	3.9	208	56.1	5.7
Age									
18-34	9	4.7	3.1	15	13.2	7.2	96	82.1	7.7
35-54	49	19.3	5.6	42	15.4	4.8	167	65.3	6.6
≥55	114	48.9	7.7	18	8.4	4.1	88	42.7	7.7
Education									
≤High School	60	34.2	8.4	14	7.6	4.5	90	58.2	8.7
>High School	117	22.7	4.2	63	14.3	3.8	268	63.0	5.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Smoked 100 Cigarettes in Lifetime**

*Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	320	49.9	297	50.1	4.7
Gender					
Male	129	50.1	115	49.9	7.5
Female	191	49.8	182	50.2	5.7
Age					
18-34	42	34.6	78	65.4	9.9
35-54	136	52.5	124	47.5	7.0
≥55	134	58.8	87	41.2	7.6
Education					
≤High School	94	54.8	71	45.2	8.9
>High School	225	48.1	225	51.9	5.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Current Smoker

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	112	17.7	505	82.3	3.5
Gender					
Male	47	18.7	197	81.3	5.7
Female	65	16.9	308	83.1	4.3
Age					
18-34	25	20.8	95	79.2	8.1
35-54	59	22.4	201	77.6	6.0
≥55	26	10.5	195	89.5	4.6
Education					
≤High School	39	21.3	126	78.7	6.9
>High School	72	16.3	378	83.7	4.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

**Status of Cigarette Smoking***Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Everyday Smoker %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Someday Smoker %</b>	<b>C.I.</b>	<b>n</b>	<b>Former or Never Smoked %</b>	<b>C.I.</b>
Total	77	11.9	2.9	35	5.9	2.4	505	82.3	3.5
Gender									
Male	36	13.3	4.6	11	5.4	3.9	197	81.3	5.7
Female	41	10.6	3.5	24	6.3	2.8	308	83.1	4.3
Age									
18-34	16	13.6	6.8	9	7.2	5.1	95	79.2	8.1
35-54	42	14.4	4.5	17	8.0	4.7	201	77.6	6.0
≥55	19	8.3	4.3	7	2.2	1.8	195	89.5	4.6
Education									
≤High School	25	11.6	4.7	14	9.7	5.5	126	78.7	6.9
>High School	51	11.8	3.5	21	4.4	2.5	378	83.7	4.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution<sup>2</sup>Weighted Percent<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Home Smoking Policies***Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Not Allowed %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Allowed Some Places %</b>	<b>C.I.</b>	<b>n</b>	<b>Allowed Anywhere %</b>	<b>C.I.</b>
Total	451	73.3	4.1	66	10.3	2.7	100	16.4	3.5
Gender									
Male	175	71.9	6.6	22	8.9	4.0	48	19.2	5.9
Female	276	74.5	5.0	44	11.5	3.6	52	13.9	4.0
Age									
18-34	99	82.7	7.4	12	9.2	5.4	9	8.2	5.5
35-54	194	72.1	6.7	30	11.3	4.4	37	16.6	5.9
≥55	147	67.9	7.1	21	9.4	4.5	52	22.8	6.3
Education									
≤High School	110	69.0	8.0	26	14.9	6.0	29	16.0	6.3
>High School	340	75.0	4.8	40	8.5	3.0	70	16.5	4.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution<sup>2</sup>Weighted Percent<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Home Smoking

*Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	109	17.4	508	82.6	3.6
Gender					
Male	46	18.8	198	81.2	5.9
Female	63	16.2	310	83.8	4.2
Age					
18-34	18	14.5	102	85.5	7.0
35-54	47	17.5	214	82.5	5.4
≥55	40	19.3	180	80.7	6.7
Education					
≤High School	39	24.0	126	76.0	7.9
>High School	69	14.8	381	85.2	3.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)**

*Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	62	10.8	545	89.2	2.8
Gender					
Male	43	17.0	198	83.0	5.3
Female	19	5.1	347	94.9	2.5
Age					
18-34	22	18.1	96	81.9	7.7
35-54	26	10.3	229	89.7	4.2
≥55	12	5.6	206	94.4	3.5
Education					
≤High School	12	6.4	150	93.6	3.7
>High School	50	12.5	394	87.5	3.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	20	3.2	591	96.8	1.5
Gender					
Male	9	3.4	233	96.6	2.3
Female	11	3.0	358	97.0	2.0
Age					
18-34	6	3.7	114	96.3	3.2
35-54	4	1.7	252	98.3	1.8
≥55	8	4.1	211	95.9	3.0
Education					
≤High School	5	3.0	157	97.0	2.7
>High School	15	3.3	433	96.7	1.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Overweight or Obese, BMI(Body Mass Index) 25 or More**

*Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	300	52.1	283	47.9	4.8
Gender					
Male	158	61.7	83	38.3	7.4
Female	142	42.9	200	57.1	5.9
Age					
18-34	38	34.9	75	65.1	10.8
35-54	122	53.3	122	46.7	7.2
≥55	139	65.0	75	35.0	7.5
Education					
≤High School	88	58.1	62	41.9	9.3
>High School	211	49.8	221	50.2	5.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+)**

*Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Underweight\ Normal %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Overweight %</b>	<b>C.I.</b>	<b>n</b>	<b>Obese %</b>	<b>C.I.</b>
Total	283	47.9	4.8	209	35.2	4.5	91	16.8	3.8
Gender									
Male	83	38.3	7.4	113	43.3	7.4	45	18.4	6.2
Female	200	57.1	5.9	96	27.6	5.3	46	15.4	4.6
Age									
18-34	75	65.1	10.8	28	20.6	8.0	10	14.3	9.9
35-54	122	46.7	7.2	88	38.9	7.2	34	14.5	5.0
≥55	75	35.0	7.5	93	43.4	7.7	46	21.7	6.3
Education									
≤High School	62	41.9	9.3	62	39.1	9.0	26	19.0	7.1
>High School	221	50.2	5.6	146	33.7	5.2	65	16.1	4.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Had Mammography Screening, Women Aged 40 Years and Older**

*Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Female	221	91.5	21	8.5	3.8
Age					
40-64	152	88.4	18	11.6	5.5
≥65	69	97.6	3	2.4	2.8
Education					
≤High School	68	88.9	9	11.1	7.3
>High School	152	92.8	12	7.2	4.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

**Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older**

*Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Female	190	78.5	51	21.5	5.8
Age					
40-64	133	77.4	36	22.6	7.2
≥65	57	80.8	15	19.2	9.9
Education					
≤High School	53	70.1	24	29.9	11.1
>High School	136	82.6	27	17.4	6.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger**  
*Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	55	13.8	393	86.2	4.3
Gender					
Male	16	11.1	162	88.9	6.7
Female	39	16.6	231	83.4	5.5
Age					
18-34	31	28.0	84	72.0	10.3
35-54	20	8.6	229	91.4	4.4
55-64	4	2.4	80	97.6	2.4
Education					
≤High School	20	23.0	83	77.0	9.9
>High School	35	10.8	309	89.2	4.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Sexually Active, Age 18-64 Years Old**

*Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	356	83.5	65	16.5	4.4
Gender					
Male	129	78.5	33	21.5	7.7
Female	227	88.3	32	11.7	4.2
Age					
18-34	98	83.7	14	16.3	8.4
35-54	203	85.2	32	14.8	5.8
55-64	55	77.8	19	22.2	11.0
Education					
≤High School	77	77.6	22	22.4	9.6
>High School	279	85.5	43	14.5	4.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	91	28.0	263	72.0	5.9
Gender					
Male	47	34.3	89	65.7	9.8
Female	44	22.1	174	77.9	6.5
Age					
18-34	36	44.1	58	55.9	12.3
35-54	47	22.3	157	77.7	6.6
55-64	8	12.3	48	87.7	9.1
Education					
≤High School	14	17.7	64	82.3	9.9
>High School	77	31.4	199	68.6	7.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)**

*Nassau*

Demographic Groups	Use BC			No BC at risk			No BC		
	n <sup>1</sup>	% <sup>2</sup>	C.I. <sup>3</sup>	n	%	C.I.	n	%	C.I.
Total	199	62.4	6.4	56	17.7	4.8	59	19.9	5.5
Gender									
Male	98	60.0	9.1	25	16.6	6.6	35	23.4	8.1
Female	101	65.9	8.2	31	19.5	6.8	24	14.6	6.0
Age									
18-34	77	68.4	9.7	14	10.4	5.7	23	21.2	8.7
35-59	122	57.9	8.2	42	23.2	7.0	36	18.9	7.0
Education									
≤High School	43	55.7	13.1	16	22.6	11.1	17	21.7	10.8
>High School	156	64.8	7.2	40	16.0	5.1	42	19.2	6.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Physician Diagnosed Heart Attack, Angina or Stroke**

*Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	53	8.0	554	92.0	2.4
Gender					
Male	34	11.1	209	88.9	4.1
Female	19	5.2	345	94.8	2.6
Age					
18-64	12	2.9	446	97.1	1.8
≥65	38	27.9	95	72.1	8.5
Education					
≤High School	19	10.1	144	89.9	4.8
>High School	34	7.3	408	92.7	2.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina)

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	46	7.0	561	93.0	2.3
Gender					
Male	32	10.6	211	89.4	4.1
Female	14	3.7	350	96.3	2.2
Age					
18-64	9	2.2	449	97.8	1.7
≥65	34	25.1	99	74.9	8.3
Education					
≤High School	16	8.0	147	92.0	4.2
>High School	30	6.6	412	93.4	2.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Male	125	69.4	51	30.6	7.9
Age					
40-64	72	64.2	44	35.8	9.9
≥65	53	83.8	7	16.2	11.8
Education					
≤High School	25	61.7	19	38.3	16.4
>High School	100	72.5	31	27.5	9.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

## Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Male	119	65.1	57	34.9	8.3
Age					
40-64	66	58.4	50	41.6	10.3
≥65	53	83.8	7	16.2	11.8
Education					
≤High School	24	56.4	20	43.6	17.3
>High School	95	68.6	36	31.4	9.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Had a Digital Rectal Exam, Men Aged 40 and Older

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Male	138	77.7	41	22.3	6.9
Age					
40-64	84	75.0	34	25.0	8.5
≥65	54	85.2	7	14.8	10.6
Education					
≤High School	28	63.3	17	36.7	16.2
>High School	109	82.8	24	17.2	7.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Used a Blood Stool Test at Home, Aged 50 and Older

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	114	40.0	165	60.0	6.6
Gender					
Male	52	35.0	77	65.0	9.2
Female	62	45.7	88	54.3	9.1
Age					
50-64	52	33.5	96	66.5	8.8
≥65	62	48.1	69	51.9	9.6
Education					
≤High School	36	37.6	51	62.4	11.5
>High School	77	40.7	114	59.3	8.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older**

*Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	51	16.4	228	83.6	4.7
Gender					
Male	30	18.9	99	81.1	7.2
Female	21	13.5	129	86.5	5.7
Age					
50-64	24	15.3	124	84.7	6.5
≥65	27	17.8	104	82.2	6.8
Education					
≤High School	19	19.5	68	80.5	9.1
>High School	32	15.1	159	84.9	5.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	73	24.4	206	75.6	5.6
Gender					
Male	38	26.8	91	73.2	8.5
Female	35	21.5	115	78.5	7.0
Age					
50-64	36	23.3	112	76.7	7.8
≥65	37	25.6	94	74.4	8.0
Education					
≤High School	25	25.4	62	74.6	10.0
>High School	48	24.0	143	76.0	6.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older

*Nassau*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	164	54.0	116	46.0	6.9
Gender					
Male	82	53.4	47	46.6	10.2
Female	82	54.6	69	45.4	9.0
Age					
50-64	77	46.3	73	53.7	9.4
≥65	87	63.6	43	36.4	9.5
Education					
≤High School	45	48.1	42	51.9	12.1
>High School	118	56.4	74	43.6	8.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older

*Nassau*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	156	51.7	122	48.3	6.9
Gender					
Male	80	52.3	49	47.7	10.1
Female	76	51.1	73	48.9	9.1
Age					
50-64	73	44.2	76	55.8	9.4
≥65	83	61.1	46	38.9	9.6
Education					
≤High School	39	42.2	46	57.8	11.9
>High School	116	55.8	76	44.2	8.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older**

***Nassau***

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	168	54.7	115	45.3	6.8
Gender					
Male	85	55.2	45	44.8	10.2
Female	83	54.3	70	45.7	9.0
Age					
50-64	78	46.6	72	53.4	9.4
≥65	90	64.8	43	35.2	9.3
Education					
≤High School	44	46.1	44	53.9	12.0
>High School	123	58.5	71	41.5	8.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval