

**Expanded BRFSS, 2003**

**Queens Locality Tables**

TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days.....	1
Core 2: Health Care Access.....	4
Core 3: Exercise.....	8
Core 4: Diabetes.....	9
Core 5: Asthma.....	10
Core 6: Arthritis.....	12
Core 7: Tobacco Use.....	14
Core 8: Tobacco ETS.....	17
Core 9: Alcohol Consumption.....	19
Core 10: Weight Status.....	21
Core 11: Mammography.....	23
Core 12: Sexual Behavior.....	25
Core 13: Family Planning.....	28
Core 14: Cardiovascular Disease.....	29
Core 15: Prostate Cancer Screening.....	31
Core 16: Colorectal Cancer Screening.....	37

**New York State Department Of Health**

Expanded BRFSS, 2003 - New York State Department of Health

## Self-rated Health Status

### Queens

Demographic Groups	n <sup>1</sup>	Fair or Poor % <sup>2</sup>	n	Good to Excellent %	C.I. <sup>3</sup>
Total	132	22.1	475	77.9	3.9
Gender					
Male	53	23.0	194	77.0	6.3
Female	79	21.3	281	78.7	4.9
Age					
18-34	26	15.8	150	84.2	6.5
35-54	40	20.7	181	79.3	6.8
≥55	64	31.7	132	68.3	7.4
Education					
≤High School	71	30.6	155	69.4	7.1
>High School	61	16.5	311	83.5	4.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Reported Number of Physical Distress Days in the Past Month

### *Queens*

Demographic Groups	n <sup>1</sup>	≥14 days % <sup>2</sup>	n	< 14 days %	C.I. <sup>3</sup>
Total	71	11.7	523	88.3	3.1
Gender					
Male	23	9.3	217	90.7	4.5
Female	48	13.9	306	86.1	4.3
Age					
18-34	6	4.4	168	95.6	4.4
35-54	19	9.5	201	90.5	5.4
≥55	45	22.9	141	77.1	6.7
Education					
≤High School	38	15.0	182	85.0	5.2
>High School	32	8.7	333	91.3	3.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Reported Number of Mental Distress Days in the Past Month

### *Queens*

Demographic Groups	n <sup>1</sup>	≥14 days % <sup>2</sup>	n	< 14 days %	C.I. <sup>3</sup>
Total	59	9.7	540	90.3	2.7
Gender					
Male	19	7.8	225	92.2	3.9
Female	40	11.4	315	88.6	3.7
Age					
18-34	23	13.6	149	86.4	5.9
35-54	19	7.9	200	92.1	3.8
≥55	17	8.5	176	91.5	4.3
Education					
≤High School	21	8.5	202	91.5	4.0
>High School	38	10.8	331	89.2	3.6

<sup>1</sup> Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup> Weighted Percent

<sup>3</sup> 95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Health Insurance Coverage, 18-64 Years of Age

### *Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	393	74.9	98	25.1	5.0
Gender					
Male	149	70.2	50	29.8	7.8
Female	244	79.6	48	20.4	6.0
Age					
18-34	121	60.7	57	39.3	8.8
35-54	191	81.7	31	18.3	6.7
55-64	81	92.4	10	7.6	5.2
Education					
≤High School	118	64.1	49	35.9	9.3
>High School	271	82.4	47	17.6	5.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Health Insurance Coverage, Aged 18 and Older

### *Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	514	79.5	99	20.5	4.2
Gender					
Male	198	74.6	50	25.4	6.9
Female	316	83.9	49	16.1	4.8
Age					
18-34	121	60.7	57	39.3	8.8
35-54	191	81.7	31	18.3	6.7
≥55	188	96.6	10	3.4	2.3
Education					
≤High School	179	73.2	49	26.8	7.5
>High School	328	84.6	48	15.4	4.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Reported Having a Routine Physical Exam in Past Two Years

### *Queens*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	547	89.2	61	10.8	3.1
Gender					
Male	208	84.4	38	15.6	5.5
Female	339	93.5	23	6.5	2.9
Age					
18-34	151	85.5	25	14.5	6.0
35-54	194	85.9	28	14.1	6.0
≥55	189	96.5	8	3.5	2.7
Education					
≤High School	204	88.1	23	11.9	5.6
>High School	336	89.9	37	10.1	3.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Time in Past 12 Months when Needed Medical Care but Could Not Get It

### *Queens*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	56	10.6	556	89.4	2.9
Gender					
Male	18	8.5	230	91.5	4.0
Female	38	12.5	326	87.5	4.2
Age					
18-34	25	15.2	153	84.8	6.2
35-54	20	11.6	202	88.4	5.2
≥55	10	4.3	187	95.7	2.9
Education					
≤High School	27	14.3	200	85.7	5.5
>High School	28	8.0	348	92.0	3.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

## Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days

### *Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	413	68.2	200	31.8	4.3
Gender					
Male	176	73.4	73	26.6	6.3
Female	237	63.5	127	36.5	5.7
Age					
18-34	134	77.7	44	22.3	6.9
35-54	152	68.0	71	32.0	7.2
≥55	115	56.8	82	43.2	8.0
Education					
≤High School	133	62.6	95	37.4	7.3
>High School	275	73.1	101	26.9	5.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Physician Diagnosed Diabetes

### *Queens*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	50	8.8	564	91.2	2.7
Gender					
Male	25	9.2	224	90.8	3.9
Female	25	8.5	340	91.5	3.6
Age					
18-34	3	1.3	175	98.7	1.6
35-54	14	5.9	209	94.1	3.6
≥55	33	21.4	165	78.6	7.1
Education					
≤High School	23	11.0	206	89.0	4.7
>High School	26	7.2	350	92.8	3.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Lifetime Asthma Diagnosed by a Medical Professional

### *Queens*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	69	11.0	545	89.0	2.9
Gender					
Male	25	10.8	224	89.2	4.8
Female	44	11.2	321	88.8	3.5
Age					
18-34	21	10.6	157	89.4	4.8
35-54	22	9.8	201	90.2	5.1
≥55	23	11.6	175	88.4	4.9
Education					
≤High School	31	11.8	198	88.2	4.4
>High School	38	10.8	338	89.2	4.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Current Asthma Diagnosed by a Medical Professional

### Queens

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	43	6.7	569	93.3	2.4
Gender					
Male	13	6.3	236	93.7	4.0
Female	30	7.0	333	93.0	2.7
Age					
18-34	14	7.0	163	93.0	4.1
35-54	16	6.9	207	93.1	4.7
≥55	11	5.7	186	94.3	3.5
Education					
≤High School	19	5.9	209	94.1	2.9
>High School	24	7.4	351	92.6	3.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Arthritis Diagnosed by a Medical Professional

### *Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	145	22.2	467	77.8	3.8
Gender					
Male	49	17.1	200	82.9	5.5
Female	96	26.8	267	73.2	5.3
Age					
18-34	9	5.3	169	94.7	4.5
35-54	41	18.8	180	81.2	6.4
≥55	90	45.0	108	55.0	8.0
Education					
≤High School	59	23.2	170	76.8	6.2
>High School	83	20.7	291	79.3	4.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

### Queens

Demographic Groups	n <sup>1</sup>	Doctor Diagnosed % <sup>2</sup>	C.I. <sup>3</sup>	n	Possible Arthritis (CJS Only) %	C.I.	n	No Arthritis or CJS %	C.I.
Total	145	22.5	3.9	87	13.1	2.9	372	64.5	4.4
Gender									
Male	49	17.2	5.5	31	11.0	4.0	167	71.8	6.4
Female	96	27.2	5.4	56	15.0	4.2	205	57.8	5.9
Age									
18-34	9	5.4	4.6	27	12.2	5.0	139	82.4	6.4
35-54	41	18.8	6.4	34	14.1	5.0	145	67.1	7.3
≥55	90	45.5	8.0	25	12.9	5.5	81	41.6	8.0
Education									
≤High School	59	23.7	6.3	27	11.0	4.5	139	65.3	7.1
>High School	83	20.8	4.7	59	14.7	4.0	228	64.4	5.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Smoked 100 Cigarettes in Lifetime**

*Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	242	37.0	370	63.0	4.5
Gender					
Male	115	42.9	133	57.1	7.2
Female	127	31.6	237	68.4	5.3
Age					
18-34	52	28.0	126	72.0	7.5
35-54	87	38.3	135	61.7	7.7
≥55	97	45.8	100	54.2	8.0
Education					
≤High School	96	39.0	132	61.0	7.5
>High School	144	36.5	231	63.5	5.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Current Smoker

### *Queens*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	97	14.6	515	85.4	3.2
Gender					
Male	47	17.9	201	82.1	5.6
Female	50	11.8	314	88.2	3.5
Age					
18-34	35	17.4	143	82.6	6.1
35-54	36	15.9	186	84.1	6.0
≥55	22	9.7	175	90.3	4.3
Education					
≤High School	42	16.5	186	83.5	5.8
>High School	54	13.7	321	86.3	3.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

**Status of Cigarette Smoking***Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Everyday Smoker %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Someday Smoker %</b>	<b>C.I.</b>	<b>n</b>	<b>Former or Never Smoked %</b>	<b>C.I.</b>
Total	57	8.7	2.6	40	6.0	2.1	515	85.4	3.2
Gender									
Male	23	9.0	4.5	24	8.8	3.8	201	82.1	5.6
Female	34	8.4	3.0	16	3.4	1.8	314	88.2	3.5
Age									
18-34	16	7.0	3.7	19	10.4	5.1	143	82.6	6.1
35-54	20	10.0	5.5	16	5.9	3.1	186	84.1	6.0
≥55	18	8.4	4.1	4	1.3	1.4	175	90.3	4.3
Education									
≤High School	32	12.0	5.1	10	4.5	3.3	186	83.5	5.8
>High School	25	6.5	2.7	29	7.1	2.7	321	86.3	3.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution<sup>2</sup>Weighted Percent<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Home Smoking Policies***Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Not Allowed %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Allowed Some Places %</b>	<b>C.I.</b>	<b>n</b>	<b>Allowed Anywhere %</b>	<b>C.I.</b>
Total	444	74.2	4.0	56	8.0	2.3	108	17.8	3.6
Gender									
Male	178	76.4	6.0	21	6.3	3.0	45	17.3	5.5
Female	266	72.3	5.4	35	9.6	3.4	63	18.2	4.7
Age									
18-34	122	71.0	7.9	25	11.1	4.6	30	18.0	7.2
35-54	181	84.2	5.1	17	6.1	3.2	23	9.7	4.2
≥55	132	67.3	7.5	11	6.2	4.1	52	26.5	6.9
Education									
≤High School	156	73.6	6.4	23	8.4	3.8	47	18.0	5.6
>High School	282	76.1	4.9	33	8.0	2.9	58	15.9	4.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution<sup>2</sup>Weighted Percent<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Home Smoking

### *Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	101	16.0	513	84.0	3.4
Gender					
Male	38	14.6	211	85.4	5.3
Female	63	17.2	302	82.8	4.3
Age					
18-34	43	21.0	135	79.0	6.5
35-54	29	13.5	194	86.5	5.9
≥55	26	13.6	172	86.4	5.3
Education					
≤High School	44	18.7	185	81.3	6.1
>High School	57	14.5	319	85.5	3.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)**

*Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	68	11.8	535	88.2	3.2
Gender					
Male	50	20.5	196	79.5	6.1
Female	18	4.0	339	96.0	2.1
Age					
18-34	39	20.0	135	80.0	6.7
35-54	20	11.2	201	88.8	5.8
≥55	9	4.2	184	95.8	3.0
Education					
≤High School	23	11.9	201	88.1	5.6
>High School	45	12.2	325	87.8	3.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)**

*Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	16	2.8	588	97.2	1.9
Gender					
Male	9	4.4	236	95.6	3.9
Female	7	1.4	352	98.6	1.1
Age					
18-34	4	1.6	169	98.4	1.7
35-54	4	3.7	217	96.3	4.7
≥55	8	3.3	187	96.7	2.4
Education					
≤High School	4	2.8	217	97.2	4.0
>High School	12	3.0	362	97.0	1.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Overweight or Obese, BMI(Body Mass Index) 25 or More**

*Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	318	57.6	259	42.4	4.7
Gender					
Male	160	64.6	85	35.4	7.1
Female	158	50.9	174	49.1	6.1
Age					
18-34	70	45.5	93	54.5	9.1
35-54	131	62.8	83	37.2	7.7
≥55	114	64.7	76	35.3	7.5
Education					
≤High School	135	62.8	79	37.2	7.8
>High School	181	53.9	178	46.1	5.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+)

### Queens

Demographic Groups	n <sup>1</sup>	Underweight\ Normal % <sup>2</sup>	C.I. <sup>3</sup>	n	Overweight %	C.I.	n	Obese %	C.I.
Total	259	42.4	4.7	198	37.6	4.7	120	20.0	3.7
Gender									
Male	85	35.4	7.1	108	44.4	7.3	52	20.2	5.9
Female	174	49.1	6.1	90	31.0	6.0	68	19.9	4.7
Age									
18-34	93	54.5	9.1	50	33.9	8.8	20	11.7	5.6
35-54	83	37.2	7.7	76	36.8	7.6	55	26.1	7.0
≥55	76	35.3	7.5	69	42.2	8.3	45	22.5	6.5
Education									
≤High School	79	37.2	7.8	77	39.4	7.8	58	23.3	6.3
>High School	178	46.1	5.9	119	36.0	5.9	62	17.9	4.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Had Mammography Screening, Women Aged 40 Years and Older**

*Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Female	178	91.1	17	8.9	4.6
Age					
40-64	126	93.4	10	6.6	4.5
≥65	52	86.7	7	13.3	10.0
Education					
≤High School	68	83.6	13	16.4	9.3
>High School	107	97.1	4	2.9	3.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

**Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older**

*Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Female	152	76.2	40	23.8	7.1
Age					
40-64	110	80.0	24	20.0	8.0
≥65	42	69.2	16	30.8	13.5
Education					
≤High School	59	66.4	22	33.6	12.3
>High School	92	85.1	17	14.9	7.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger**  
*Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	109	24.6	361	75.4	4.8
Gender					
Male	28	17.4	161	82.6	6.7
Female	81	31.6	200	68.4	6.5
Age					
18-34	66	38.0	112	62.0	8.5
35-54	36	17.6	173	82.4	6.1
55-64	7	6.1	76	93.9	4.9
Education					
≤High School	48	32.4	110	67.6	9.0
>High School	61	20.5	245	79.5	5.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Sexually Active, Age 18-64 Years Old**

*Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	310	79.5	104	20.5	4.3
Gender					
Male	133	83.2	33	16.8	6.4
Female	177	76.0	71	24.0	5.8
Age					
18-34	133	88.5	21	11.5	5.8
35-54	144	79.6	44	20.4	6.3
55-64	33	53.6	39	46.4	13.9
Education					
≤High School	108	80.1	42	19.9	6.9
>High School	198	78.8	61	21.2	5.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger

### *Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	127	40.2	205	59.8	6.4
Gender					
Male	58	42.9	89	57.1	9.5
Female	69	37.1	116	62.9	8.2
Age					
18-34	68	51.9	78	48.1	9.5
35-54	52	33.4	97	66.6	9.0
55-64	7	14.1	30	85.9	11.1
Education					
≤High School	42	40.2	68	59.8	11.0
>High School	83	39.8	135	60.2	7.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)**

*Queens*

Demographic Groups	Use BC			No BC at risk			No BC		
	n <sup>1</sup>	% <sup>2</sup>	C.I. <sup>3</sup>	n	%	C.I.	n	%	C.I.
Total	191	59.7	6.1	62	17.4	4.6	90	22.9	5.1
Gender									
Male	90	58.8	8.5	41	22.7	7.0	40	18.5	6.4
Female	101	60.9	8.5	21	9.9	4.5	50	29.2	8.1
Age									
18-34	100	60.7	8.6	27	14.8	6.0	46	24.5	7.6
35-59	91	58.5	8.5	35	20.3	7.0	44	21.1	6.5
Education									
≤High School	59	58.8	10.3	27	18.9	7.8	32	22.2	8.4
>High School	131	62.3	7.2	33	15.1	5.2	57	22.6	6.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Physician Diagnosed Heart Attack, Angina or Stroke**

*Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	40	6.0	551	94.0	2.1
Gender					
Male	21	7.1	218	92.9	3.4
Female	19	5.1	333	94.9	2.4
Age					
18-64	14	2.5	459	97.5	1.5
≥65	26	23.6	80	76.4	8.9
Education					
≤High School	24	9.7	196	90.3	4.1
>High School	16	3.5	346	96.5	2.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina)

### *Queens*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	32	5.0	558	95.0	1.9
Gender					
Male	18	6.0	220	94.0	3.1
Female	14	4.0	338	96.0	2.2
Age					
18-64	9	1.5	463	98.5	1.1
≥65	23	21.9	83	78.1	8.8
Education					
≤High School	20	8.5	200	91.5	3.9
>High School	12	2.5	349	97.5	1.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older**

*Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Male	78	53.6	52	46.4	10.4
Age					
40-64	42	43.3	43	56.7	12.5
≥65	36	80.0	9	20.0	13.0
Education					
≤High School	28	45.3	24	54.7	16.1
>High School	50	61.6	27	38.4	12.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

## Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

### *Queens*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Male	69	47.7	60	52.3	10.3
Age					
40-64	35	36.7	49	63.3	12.1
≥65	34	75.7	11	24.3	13.8
Education					
≤High School	26	43.6	25	56.4	16.1
>High School	43	51.9	34	48.1	12.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Had a Digital Rectal Exam, Men Aged 40 and Older

### *Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Male	95	67.9	40	32.1	9.8
Age					
40-64	59	64.4	30	35.6	12.3
≥65	36	76.8	10	23.2	14.6
Education					
≤High School	40	68.2	16	31.8	15.9
>High School	55	68.4	23	31.6	11.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Used a Blood Stool Test at Home, Aged 50 and Older**

*Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	78	33.6	149	66.4	7.1
Gender					
Male	39	38.7	58	61.3	11.5
Female	39	29.5	91	70.5	8.7
Age					
50-64	40	32.0	84	68.0	9.8
≥65	38	35.3	65	64.7	10.1
Education					
≤High School	35	33.6	66	66.4	10.6
>High School	43	34.5	80	65.5	9.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older**

*Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	41	17.8	183	82.2	5.9
Gender					
Male	22	22.9	74	77.1	10.3
Female	19	13.8	109	86.2	6.4
Age					
50-64	21	18.5	102	81.5	8.7
≥65	20	17.1	81	82.9	7.7
Education					
≤High School	17	15.0	82	85.0	8.3
>High School	24	21.1	98	78.9	8.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older

### *Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	55	23.6	169	76.4	6.4
Gender					
Male	29	28.6	67	71.4	10.8
Female	26	19.5	102	80.5	7.5
Age					
50-64	31	26.2	92	73.8	9.5
≥65	24	20.6	77	79.4	8.2
Education					
≤High School	23	21.1	76	78.9	9.3
>High School	32	26.6	90	73.4	9.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older

### *Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	115	48.6	114	51.4	7.4
Gender					
Male	49	44.9	49	55.1	11.4
Female	66	51.5	65	48.5	9.7
Age					
50-64	59	44.9	66	55.1	10.3
≥65	56	52.6	48	47.4	10.7
Education					
≤High School	48	42.7	55	57.3	10.9
>High School	66	54.7	57	45.3	10.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older

### *Queens*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	108	46.0	120	54.0	7.4
Gender					
Male	47	43.5	51	56.5	11.3
Female	61	47.9	69	52.1	9.7
Age					
50-64	57	42.7	68	57.3	10.2
≥65	51	49.6	52	50.4	10.7
Education					
≤High School	43	38.6	59	61.4	10.7
>High School	64	53.5	59	46.5	10.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older**

***Queens***

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	118	50.0	112	50.0	7.5
Gender					
Male	52	48.7	47	51.3	11.7
Female	66	51.0	65	49.0	9.7
Age					
50-64	64	48.4	62	51.6	10.5
≥65	54	51.8	50	48.2	10.7
Education					
≤High School	47	43.4	56	56.6	11.1
>High School	70	56.6	54	43.4	10.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval