

Columbia and Greene Locality Tables

TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days 1
Core 2: Health Care Access 4
Core 3: Exercise 8
Core 4: Diabetes 9
Core 5: Asthma 10
Core 6: Arthritis 12
Core 7: Tobacco Use 14
Core 8: Tobacco ETS 17
Core 9: Alcohol Consumption 19
Core 10: Weight Status 21
Core 11: Mammography 23
Core 12: Sexual Behavior 25
Core 13: Family Planning 28
Core 14: Cardiovascular Disease 29
Core 15: Prostate Cancer Screening 31
Core 16: Colorectal Cancer Screening 37

Expanded BRFSS, 2003 - New York State Department of Health

Self-rated Health Status

Columbia, Greene

| Demographic Groups | n ¹ | Fair or Poor % ² | n | Good to Excellent % | C.I. ³ |
|--------------------|----------------|-----------------------------|-----|---------------------|-------------------|
| Total | 103 | 14.5 | 531 | 85.5 | 2.9 |
| Gender | | | | | |
| Male | 37 | 12.9 | 219 | 87.1 | 4.3 |
| Female | 66 | 16.2 | 312 | 83.8 | 3.9 |
| Age | | | | | |
| 18-34 | 10 | 7.5 | 104 | 92.5 | 4.8 |
| 35-54 | 32 | 10.9 | 229 | 89.1 | 3.9 |
| ≥55 | 60 | 23.1 | 191 | 76.9 | 5.7 |
| Education | | | | | |
| ≤High School | 69 | 21.2 | 220 | 78.8 | 5.1 |
| >High School | 34 | 9.2 | 309 | 90.8 | 3.2 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Reported Number of Physical Distress Days in the Past Month

Columbia, Greene

| Demographic Groups | n ¹ | ≥14 days % ² | n | < 14 days % | C.I. ³ |
|--------------------|----------------|-------------------------|-----|-------------|-------------------|
| Total | 76 | 10.8 | 535 | 89.2 | 2.7 |
| Gender | | | | | |
| Male | 25 | 8.1 | 225 | 91.9 | 3.5 |
| Female | 51 | 13.5 | 310 | 86.5 | 4.0 |
| Age | | | | | |
| 18-34 | 5 | 4.2 | 108 | 95.8 | 4.1 |
| 35-54 | 32 | 10.0 | 227 | 90.0 | 3.7 |
| ≥55 | 39 | 16.8 | 193 | 83.2 | 5.5 |
| Education | | | | | |
| ≤High School | 46 | 13.1 | 227 | 86.9 | 3.9 |
| >High School | 30 | 9.1 | 306 | 90.9 | 3.6 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Reported Number of Mental Distress Days in the Past Month

Columbia, Greene

| Demographic Groups | n ¹ | ≥14 days % ² | n | < 14 days % | C.I. ³ |
|--------------------|----------------|-------------------------|-----|-------------|-------------------|
| Total | 62 | 8.3 | 554 | 91.7 | 2.2 |
| Gender | | | | | |
| Male | 12 | 3.7 | 242 | 96.3 | 2.3 |
| Female | 50 | 13.1 | 312 | 86.9 | 3.7 |
| Age | | | | | |
| 18-34 | 9 | 6.4 | 102 | 93.6 | 4.3 |
| 35-54 | 33 | 10.5 | 226 | 89.5 | 3.8 |
| ≥55 | 20 | 7.5 | 220 | 92.5 | 3.4 |
| Education | | | | | |
| ≤High School | 36 | 10.2 | 242 | 89.8 | 3.5 |
| >High School | 26 | 6.8 | 310 | 93.2 | 2.7 |

¹ Percents based on row denominators of less than 50 are unstable and should be used with caution

² Weighted Percent

³ 95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Health Insurance Coverage, 18-64 Years of Age

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|----|------|-------------------|
| Total | 413 | 83.4 | 75 | 16.6 | 3.9 |
| Gender | | | | | |
| Male | 167 | 80.8 | 37 | 19.2 | 6.2 |
| Female | 246 | 86.3 | 38 | 13.7 | 4.5 |
| Age | | | | | |
| 18-34 | 86 | 74.5 | 27 | 25.5 | 9.1 |
| 35-54 | 227 | 88.9 | 34 | 11.1 | 4.0 |
| 55-64 | 100 | 85.0 | 14 | 15.0 | 7.3 |
| Education | | | | | |
| ≤High School | 166 | 76.8 | 43 | 23.2 | 6.8 |
| >High School | 246 | 88.3 | 32 | 11.7 | 4.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Health Insurance Coverage, Aged 18 and Older

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|----|------|-------------------|
| Total | 557 | 86.9 | 76 | 13.1 | 3.1 |
| Gender | | | | | |
| Male | 219 | 84.4 | 37 | 15.6 | 5.2 |
| Female | 338 | 89.5 | 39 | 10.5 | 3.4 |
| Age | | | | | |
| 18-34 | 86 | 74.5 | 27 | 25.5 | 9.1 |
| 35-54 | 227 | 88.9 | 34 | 11.1 | 4.0 |
| ≥55 | 236 | 93.1 | 15 | 6.9 | 3.4 |
| Education | | | | | |
| ≤High School | 243 | 82.3 | 44 | 17.7 | 5.3 |
| >High School | 312 | 90.5 | 32 | 9.5 | 3.6 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Reported Having a Routine Physical Exam in Past Two Years

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|----|------|-------------------|
| Total | 562 | 88.9 | 71 | 11.1 | 2.8 |
| Gender | | | | | |
| Male | 218 | 85.9 | 39 | 14.1 | 4.7 |
| Female | 344 | 92.0 | 32 | 8.0 | 3.0 |
| Age | | | | | |
| 18-34 | 94 | 84.3 | 19 | 15.7 | 7.5 |
| 35-54 | 226 | 88.6 | 34 | 11.4 | 4.1 |
| ≥55 | 235 | 92.4 | 17 | 7.6 | 3.6 |
| Education | | | | | |
| ≤High School | 255 | 88.1 | 33 | 11.9 | 4.4 |
| >High School | 306 | 89.7 | 37 | 10.3 | 3.6 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Time in Past 12 Months when Needed Medical Care but Could Not Get It

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 43 | 5.8 | 593 | 94.2 | 1.9 |
| Gender | | | | | |
| Male | 13 | 3.9 | 244 | 96.1 | 2.3 |
| Female | 30 | 7.6 | 349 | 92.4 | 2.9 |
| Age | | | | | |
| 18-34 | 8 | 5.9 | 106 | 94.1 | 4.3 |
| 35-54 | 27 | 8.1 | 235 | 91.9 | 3.3 |
| ≥55 | 8 | 3.5 | 244 | 96.5 | 2.4 |
| Education | | | | | |
| ≤High School | 24 | 7.7 | 265 | 92.3 | 3.3 |
| >High School | 19 | 4.2 | 326 | 95.8 | 2.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 477 | 75.7 | 158 | 24.3 | 3.7 |
| Gender | | | | | |
| Male | 198 | 77.4 | 58 | 22.6 | 5.7 |
| Female | 279 | 74.0 | 100 | 26.0 | 4.9 |
| Age | | | | | |
| 18-34 | 97 | 83.7 | 17 | 16.3 | 7.8 |
| 35-54 | 202 | 76.0 | 60 | 24.0 | 5.9 |
| ≥55 | 171 | 69.6 | 80 | 30.4 | 6.2 |
| Education | | | | | |
| ≤High School | 183 | 63.9 | 105 | 36.1 | 6.3 |
| >High School | 293 | 85.4 | 52 | 14.6 | 4.1 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Physician Diagnosed Diabetes

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 40 | 6.2 | 596 | 93.8 | 2.0 |
| Gender | | | | | |
| Male | 16 | 6.2 | 241 | 93.8 | 3.1 |
| Female | 24 | 6.2 | 355 | 93.8 | 2.6 |
| Age | | | | | |
| 18-34 | 0 | 0.0 | 114 | 100 | 0.0 |
| 35-54 | 13 | 5.3 | 249 | 94.7 | 3.1 |
| ≥55 | 26 | 11.4 | 226 | 88.6 | 4.4 |
| Education | | | | | |
| ≤High School | 29 | 9.1 | 260 | 90.9 | 3.4 |
| >High School | 11 | 3.9 | 334 | 96.1 | 2.4 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Lifetime Asthma Diagnosed by a Medical Professional

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 83 | 12.0 | 552 | 88.0 | 2.7 |
| Gender | | | | | |
| Male | 23 | 8.4 | 234 | 91.6 | 3.7 |
| Female | 60 | 15.6 | 318 | 84.4 | 3.9 |
| Age | | | | | |
| 18-34 | 14 | 10.7 | 100 | 89.3 | 5.9 |
| 35-54 | 41 | 15.1 | 221 | 84.9 | 4.7 |
| ≥55 | 27 | 10.0 | 224 | 90.0 | 3.8 |
| Education | | | | | |
| ≤High School | 36 | 11.5 | 252 | 88.5 | 4.0 |
| >High School | 47 | 12.4 | 298 | 87.6 | 3.7 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Current Asthma Diagnosed by a Medical Professional

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 64 | 9.1 | 569 | 90.9 | 2.4 |
| Gender | | | | | |
| Male | 15 | 5.9 | 241 | 94.1 | 3.2 |
| Female | 49 | 12.4 | 328 | 87.6 | 3.5 |
| Age | | | | | |
| 18-34 | 8 | 6.3 | 105 | 93.7 | 5.0 |
| 35-54 | 33 | 11.8 | 229 | 88.2 | 4.2 |
| ≥55 | 22 | 8.4 | 228 | 91.6 | 3.6 |
| Education | | | | | |
| ≤High School | 30 | 9.7 | 257 | 90.3 | 3.8 |
| >High School | 34 | 8.7 | 310 | 91.3 | 3.1 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Arthritis Diagnosed by a Medical Professional

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 220 | 32.7 | 415 | 67.3 | 4.1 |
| Gender | | | | | |
| Male | 76 | 27.4 | 180 | 72.6 | 5.9 |
| Female | 144 | 38.0 | 235 | 62.0 | 5.5 |
| Age | | | | | |
| 18-34 | 4 | 2.8 | 110 | 97.2 | 2.9 |
| 35-54 | 88 | 34.6 | 173 | 65.4 | 6.6 |
| ≥55 | 124 | 50.9 | 128 | 49.1 | 6.8 |
| Education | | | | | |
| ≤High School | 111 | 37.2 | 177 | 62.8 | 6.2 |
| >High School | 108 | 29.0 | 237 | 71.0 | 5.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

Columbia, Greene

| Demographic Groups | n ¹ | Doctor Diagnosed % ² | C.I. ³ | n | Possible Arthritis (CJS Only) % | C.I. | n | No Arthritis or CJS % | C.I. |
|--------------------|----------------|---------------------------------|-------------------|-----|---------------------------------|------|-----|-----------------------|------|
| Total | 220 | 32.9 | 4.1 | 107 | 17.2 | 3.4 | 304 | 49.9 | 4.4 |
| Gender | | | | | | | | | |
| Male | 76 | 27.4 | 5.9 | 45 | 17.5 | 5.2 | 135 | 55.1 | 6.7 |
| Female | 144 | 38.4 | 5.5 | 62 | 16.9 | 4.3 | 169 | 44.7 | 5.6 |
| Age | | | | | | | | | |
| 18-34 | 4 | 2.8 | 2.9 | 29 | 27.2 | 9.2 | 81 | 69.9 | 9.3 |
| 35-54 | 88 | 34.6 | 6.6 | 40 | 13.3 | 4.4 | 133 | 52.1 | 6.8 |
| ≥55 | 124 | 51.5 | 6.8 | 37 | 14.3 | 4.5 | 87 | 34.2 | 6.4 |
| Education | | | | | | | | | |
| ≤High School | 111 | 37.6 | 6.3 | 38 | 12.3 | 4.1 | 135 | 50.1 | 6.5 |
| >High School | 108 | 29.0 | 5.3 | 69 | 21.2 | 5.1 | 168 | 49.8 | 6.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Smoked 100 Cigarettes in Lifetime

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 371 | 56.5 | 263 | 43.5 | 4.4 |
| Gender | | | | | |
| Male | 152 | 56.5 | 105 | 43.5 | 6.8 |
| Female | 219 | 56.5 | 158 | 43.5 | 5.6 |
| Age | | | | | |
| 18-34 | 60 | 49.8 | 54 | 50.2 | 10.1 |
| 35-54 | 154 | 56.8 | 108 | 43.2 | 6.8 |
| ≥55 | 151 | 60.4 | 100 | 39.6 | 6.6 |
| Education | | | | | |
| ≤High School | 185 | 64.6 | 103 | 35.4 | 6.2 |
| >High School | 185 | 50.0 | 159 | 50.0 | 6.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Current Smoker

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 163 | 24.9 | 471 | 75.1 | 3.8 |
| Gender | | | | | |
| Male | 61 | 23.6 | 196 | 76.4 | 5.7 |
| Female | 102 | 26.3 | 275 | 73.7 | 4.8 |
| Age | | | | | |
| 18-34 | 41 | 34.3 | 73 | 65.7 | 9.4 |
| 35-54 | 75 | 25.8 | 187 | 74.2 | 5.7 |
| ≥55 | 45 | 17.7 | 206 | 82.3 | 5.2 |
| Education | | | | | |
| ≤High School | 96 | 33.4 | 192 | 66.6 | 6.1 |
| >High School | 67 | 18.1 | 277 | 81.9 | 4.5 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Status of Cigarette Smoking*Columbia, Greene*

| Demographic Groups | n¹ | Everyday Smoker %² | C.I.³ | n | Someday Smoker % | C.I. | n | Former or Never Smoked % | C.I. |
|---------------------------|----------------------|--------------------------------------|-------------------------|----------|-------------------------|-------------|----------|---------------------------------|-------------|
| Total | 135 | 20.9 | 3.5 | 28 | 4.1 | 1.7 | 471 | 75.1 | 3.8 |
| Gender | | | | | | | | | |
| Male | 50 | 19.4 | 5.4 | 11 | 4.1 | 2.7 | 196 | 76.4 | 5.7 |
| Female | 85 | 22.3 | 4.6 | 17 | 4.0 | 2.1 | 275 | 73.7 | 4.8 |
| Age | | | | | | | | | |
| 18-34 | 31 | 26.8 | 8.8 | 10 | 7.6 | 5.1 | 73 | 65.7 | 9.4 |
| 35-54 | 64 | 22.5 | 5.5 | 11 | 3.3 | 2.1 | 187 | 74.2 | 5.7 |
| ≥55 | 38 | 15.2 | 4.9 | 7 | 2.5 | 2.0 | 206 | 82.3 | 5.2 |
| Education | | | | | | | | | |
| ≤High School | 79 | 27.6 | 5.8 | 17 | 5.8 | 2.9 | 192 | 66.6 | 6.1 |
| >High School | 56 | 15.4 | 4.2 | 11 | 2.7 | 1.9 | 277 | 81.9 | 4.5 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution²Weighted Percent³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Home Smoking Policies*Columbia, Greene*

| Demographic Groups | n¹ | Not Allowed %² | C.I.³ | n | Allowed Some Places % | C.I. | n | Allowed Anywhere % | C.I. |
|---------------------------|----------------------|----------------------------------|-------------------------|----------|------------------------------|-------------|----------|---------------------------|-------------|
| Total | 400 | 63.1 | 4.3 | 91 | 13.4 | 2.9 | 143 | 23.5 | 3.8 |
| Gender | | | | | | | | | |
| Male | 162 | 64.1 | 6.5 | 26 | 8.2 | 3.2 | 68 | 27.7 | 6.2 |
| Female | 238 | 62.1 | 5.5 | 65 | 18.6 | 4.6 | 75 | 19.3 | 4.4 |
| Age | | | | | | | | | |
| 18-34 | 69 | 61.1 | 9.9 | 20 | 14.4 | 6.3 | 25 | 24.6 | 9.2 |
| 35-54 | 168 | 64.9 | 6.4 | 37 | 12.6 | 4.2 | 57 | 22.5 | 5.8 |
| ≥55 | 157 | 62.2 | 6.7 | 33 | 13.5 | 4.9 | 61 | 24.3 | 5.8 |
| Education | | | | | | | | | |
| ≤High School | 166 | 59.7 | 6.4 | 42 | 12.9 | 4.1 | 79 | 27.4 | 5.9 |
| >High School | 232 | 65.7 | 5.7 | 49 | 13.8 | 4.1 | 64 | 20.5 | 5.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution²Weighted Percent³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Home Smoking

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 173 | 28.5 | 462 | 71.5 | 4.0 |
| Gender | | | | | |
| Male | 64 | 25.5 | 192 | 74.5 | 5.9 |
| Female | 109 | 31.5 | 270 | 68.5 | 5.4 |
| Age | | | | | |
| 18-34 | 40 | 36.1 | 73 | 63.9 | 9.7 |
| 35-54 | 78 | 29.2 | 184 | 70.8 | 6.1 |
| ≥55 | 54 | 23.0 | 198 | 77.0 | 6.1 |
| Education | | | | | |
| ≤High School | 89 | 31.8 | 200 | 68.2 | 6.1 |
| >High School | 83 | 25.7 | 261 | 74.3 | 5.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)

Columbia, Greene

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 66 | 13.0 | 561 | 87.0 | 3.3 |
| Gender | | | | | |
| Male | 45 | 19.7 | 207 | 80.3 | 5.8 |
| Female | 21 | 6.4 | 354 | 93.6 | 3.0 |
| Age | | | | | |
| 18-34 | 26 | 28.3 | 88 | 71.7 | 9.7 |
| 35-54 | 29 | 11.0 | 228 | 89.0 | 4.2 |
| ≥55 | 11 | 4.5 | 237 | 95.5 | 2.7 |
| Education | | | | | |
| ≤High School | 34 | 14.7 | 250 | 85.3 | 5.0 |
| >High School | 31 | 11.6 | 310 | 88.4 | 4.4 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)

Columbia, Greene

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 25 | 4.1 | 603 | 95.9 | 1.8 |
| Gender | | | | | |
| Male | 10 | 4.4 | 242 | 95.6 | 2.9 |
| Female | 15 | 3.8 | 361 | 96.2 | 2.1 |
| Age | | | | | |
| 18-34 | 7 | 6.8 | 106 | 93.2 | 5.3 |
| 35-54 | 7 | 2.3 | 252 | 97.7 | 1.8 |
| ≥55 | 11 | 4.0 | 237 | 96.0 | 2.5 |
| Education | | | | | |
| ≤High School | 8 | 3.2 | 275 | 96.8 | 2.4 |
| >High School | 17 | 4.8 | 326 | 95.2 | 2.6 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Overweight or Obese, BMI(Body Mass Index) 25 or More

Columbia, Greene

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 348 | 58.8 | 261 | 41.2 | 4.4 |
| Gender | | | | | |
| Male | 172 | 66.4 | 85 | 33.6 | 6.5 |
| Female | 176 | 50.5 | 176 | 49.5 | 5.8 |
| Age | | | | | |
| 18-34 | 53 | 46.8 | 54 | 53.2 | 10.4 |
| 35-54 | 134 | 58.0 | 118 | 42.0 | 6.7 |
| ≥55 | 157 | 67.7 | 86 | 32.3 | 6.2 |
| Education | | | | | |
| ≤High School | 175 | 61.3 | 100 | 38.7 | 6.6 |
| >High School | 172 | 56.8 | 160 | 43.2 | 6.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+)

Columbia, Greene

| Demographic Groups | n ¹ | Underweight\ Normal % ² | C.I. ³ | n | Overweight % | C.I. | n | Obese % | C.I. |
|--------------------|----------------|------------------------------------------|-------------------|-----|-----------------|------|-----|------------|------|
| Total | 261 | 41.2 | 4.4 | 215 | 37.5 | 4.4 | 133 | 21.3 | 3.6 |
| Gender | | | | | | | | | |
| Male | 85 | 33.6 | 6.5 | 107 | 42.7 | 6.7 | 65 | 23.8 | 5.6 |
| Female | 176 | 49.5 | 5.8 | 108 | 31.9 | 5.4 | 68 | 18.6 | 4.5 |
| Age | | | | | | | | | |
| 18-34 | 54 | 53.2 | 10.4 | 44 | 39.3 | 10.1 | 9 | 7.5 | 5.1 |
| 35-54 | 118 | 42.0 | 6.7 | 67 | 29.6 | 6.5 | 67 | 28.3 | 6.3 |
| ≥55 | 86 | 32.3 | 6.2 | 101 | 43.7 | 6.8 | 56 | 24.0 | 6.1 |
| Education | | | | | | | | | |
| ≤High School | 100 | 38.7 | 6.6 | 107 | 38.4 | 6.4 | 68 | 23.0 | 5.4 |
| >High School | 160 | 43.2 | 6.0 | 107 | 36.8 | 6.0 | 65 | 20.0 | 4.9 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had Mammography Screening, Women Aged 40 Years and Older

Columbia, Greene

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Female | 240 | 86.0 | 40 | 14.0 | 4.5 |
| Age | | | | | |
| 40-64 | 167 | 87.3 | 27 | 12.7 | 4.9 |
| ≥65 | 73 | 83.4 | 13 | 16.6 | 9.1 |
| Education | | | | | |
| ≤High School | 109 | 79.6 | 27 | 20.4 | 7.5 |
| >High School | 131 | 92.2 | 13 | 7.8 | 4.5 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|----|------|-------------------|
| Female | 188 | 71.2 | 89 | 28.8 | 5.7 |
| Age | | | | | |
| 40-64 | 132 | 72.1 | 61 | 27.9 | 6.7 |
| ≥65 | 56 | 69.4 | 28 | 30.6 | 10.7 |
| Education | | | | | |
| ≤High School | 81 | 61.6 | 53 | 38.4 | 9.0 |
| >High School | 107 | 80.6 | 36 | 19.4 | 6.5 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger

Columbia, Greene

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 65 | 13.5 | 418 | 86.5 | 3.5 |
| Gender | | | | | |
| Male | 16 | 9.2 | 186 | 90.8 | 4.9 |
| Female | 49 | 18.2 | 232 | 81.8 | 5.1 |
| Age | | | | | |
| 18-34 | 30 | 23.0 | 83 | 77.0 | 8.2 |
| 35-54 | 30 | 10.9 | 227 | 89.1 | 4.4 |
| 55-64 | 5 | 4.2 | 108 | 95.8 | 3.8 |
| Education | | | | | |
| ≤High School | 33 | 13.3 | 174 | 86.7 | 4.8 |
| >High School | 32 | 13.6 | 243 | 86.4 | 5.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Sexually Active, Age 18-64 Years Old

Columbia, Greene

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 367 | 83.3 | 90 | 16.7 | 4.1 |
| Gender | | | | | |
| Male | 161 | 82.5 | 33 | 17.5 | 6.6 |
| Female | 206 | 84.3 | 57 | 15.7 | 4.5 |
| Age | | | | | |
| 18-34 | 100 | 85.8 | 9 | 14.2 | 9.0 |
| 35-54 | 200 | 86.9 | 44 | 13.1 | 4.3 |
| 55-64 | 67 | 71.4 | 37 | 28.6 | 9.1 |
| Education | | | | | |
| ≤High School | 156 | 83.3 | 38 | 16.7 | 6.2 |
| >High School | 211 | 83.5 | 51 | 16.5 | 5.4 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 62 | 17.7 | 312 | 82.3 | 4.6 |
| Gender | | | | | |
| Male | 33 | 23.5 | 131 | 76.5 | 7.5 |
| Female | 29 | 11.3 | 181 | 88.7 | 4.4 |
| Age | | | | | |
| 18-34 | 27 | 29.3 | 72 | 70.7 | 10.1 |
| 35-54 | 30 | 13.7 | 173 | 86.3 | 5.2 |
| 55-64 | 5 | 6.7 | 67 | 93.3 | 6.0 |
| Education | | | | | |
| ≤High School | 30 | 18.2 | 132 | 81.8 | 6.7 |
| >High School | 32 | 17.3 | 180 | 82.7 | 6.2 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

Columbia, Greene

| Demographic Groups | Use BC | | | No BC at risk | | | No BC | | |
|--------------------|----------------|----------------|-------------------|---------------|------|------|-------|------|------|
| | n ¹ | % ² | C.I. ³ | n | % | C.I. | n | % | C.I. |
| Total | 199 | 65.6 | 6.2 | 36 | 12.3 | 4.1 | 66 | 22.0 | 5.6 |
| Gender | | | | | | | | | |
| Male | 114 | 62.7 | 8.1 | 21 | 12.0 | 5.2 | 43 | 25.3 | 7.6 |
| Female | 85 | 71.6 | 8.7 | 15 | 13.0 | 6.7 | 23 | 15.4 | 6.7 |
| Age | | | | | | | | | |
| 18-34 | 77 | 65.6 | 10.4 | 13 | 11.8 | 6.4 | 19 | 22.6 | 9.9 |
| 35-59 | 122 | 65.6 | 7.4 | 23 | 12.8 | 5.4 | 47 | 21.6 | 6.2 |
| Education | | | | | | | | | |
| ≤High School | 79 | 67.0 | 9.7 | 16 | 13.9 | 6.9 | 24 | 19.0 | 8.5 |
| >High School | 120 | 64.8 | 8.0 | 20 | 11.4 | 5.1 | 41 | 23.8 | 7.4 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Physician Diagnosed Heart Attack, Angina or Stroke

Columbia, Greene

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 43 | 7.0 | 589 | 93.0 | 2.2 |
| Gender | | | | | |
| Male | 19 | 7.2 | 236 | 92.8 | 3.4 |
| Female | 24 | 6.8 | 353 | 93.2 | 2.8 |
| Age | | | | | |
| 18-64 | 18 | 3.6 | 469 | 96.4 | 1.7 |
| ≥65 | 24 | 19.9 | 113 | 80.1 | 7.7 |
| Education | | | | | |
| ≤High School | 27 | 10.0 | 260 | 90.0 | 4.0 |
| >High School | 16 | 4.6 | 327 | 95.4 | 2.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina)

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 38 | 5.8 | 594 | 94.2 | 1.9 |
| Gender | | | | | |
| Male | 16 | 5.6 | 239 | 94.4 | 2.8 |
| Female | 22 | 6.0 | 355 | 94.0 | 2.6 |
| Age | | | | | |
| 18-64 | 16 | 3.0 | 471 | 97.0 | 1.6 |
| ≥65 | 21 | 15.9 | 116 | 84.1 | 6.6 |
| Education | | | | | |
| ≤High School | 23 | 7.5 | 264 | 92.5 | 3.2 |
| >High School | 15 | 4.4 | 328 | 95.6 | 2.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|----|------|-------------------|
| Male | 112 | 62.6 | 68 | 37.4 | 7.6 |
| Age | | | | | |
| 40-64 | 71 | 53.9 | 61 | 46.1 | 9.2 |
| ≥65 | 41 | 86.1 | 7 | 13.9 | 10.6 |
| Education | | | | | |
| ≤High School | 48 | 64.2 | 27 | 35.8 | 11.7 |
| >High School | 63 | 61.2 | 41 | 38.8 | 10.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|----|------|-------------------|
| Male | 100 | 57.0 | 80 | 43.0 | 7.8 |
| Age | | | | | |
| 40-64 | 64 | 49.4 | 68 | 50.6 | 9.2 |
| ≥65 | 36 | 77.2 | 12 | 22.8 | 12.5 |
| Education | | | | | |
| ≤High School | 42 | 57.3 | 33 | 42.7 | 12.1 |
| >High School | 57 | 56.5 | 47 | 43.5 | 10.2 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had a Digital Rectal Exam, Men Aged 40 and Older

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|----|------|-------------------|
| Male | 162 | 85.7 | 25 | 14.3 | 5.5 |
| Age | | | | | |
| 40-64 | 115 | 84.2 | 21 | 15.8 | 6.6 |
| ≥65 | 47 | 89.5 | 4 | 10.5 | 9.9 |
| Education | | | | | |
| ≤High School | 63 | 80.3 | 16 | 19.7 | 9.1 |
| >High School | 98 | 89.3 | 9 | 10.7 | 6.8 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Used a Blood Stool Test at Home, Aged 50 and Older

Columbia, Greene

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 151 | 47.3 | 176 | 52.7 | 6.0 |
| Gender | | | | | |
| Male | 63 | 50.5 | 61 | 49.5 | 9.4 |
| Female | 88 | 44.5 | 115 | 55.5 | 7.6 |
| Age | | | | | |
| 50-64 | 89 | 46.6 | 105 | 53.4 | 7.8 |
| ≥65 | 62 | 48.2 | 71 | 51.8 | 9.4 |
| Education | | | | | |
| ≤High School | 66 | 43.2 | 92 | 56.8 | 8.5 |
| >High School | 85 | 51.1 | 83 | 48.9 | 8.5 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older

Columbia, Greene

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 61 | 18.9 | 262 | 81.1 | 4.6 |
| Gender | | | | | |
| Male | 26 | 20.5 | 96 | 79.5 | 7.5 |
| Female | 35 | 17.4 | 166 | 82.6 | 5.7 |
| Age | | | | | |
| 50-64 | 37 | 19.6 | 156 | 80.4 | 6.2 |
| ≥65 | 24 | 17.9 | 106 | 82.1 | 7.0 |
| Education | | | | | |
| ≤High School | 26 | 15.0 | 128 | 85.0 | 5.7 |
| >High School | 35 | 22.4 | 133 | 77.6 | 7.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older

Columbia, Greene

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 87 | 26.2 | 236 | 73.8 | 5.2 |
| Gender | | | | | |
| Male | 34 | 25.8 | 88 | 74.2 | 8.1 |
| Female | 53 | 26.6 | 148 | 73.4 | 6.7 |
| Age | | | | | |
| 50-64 | 52 | 25.9 | 141 | 74.1 | 6.7 |
| ≥65 | 35 | 26.7 | 95 | 73.3 | 8.2 |
| Education | | | | | |
| ≤High School | 37 | 22.8 | 117 | 77.2 | 7.0 |
| >High School | 50 | 29.4 | 118 | 70.6 | 7.6 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older

Columbia, Greene

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 172 | 51.9 | 158 | 48.1 | 6.0 |
| Gender | | | | | |
| Male | 65 | 49.2 | 61 | 50.8 | 9.3 |
| Female | 107 | 54.2 | 97 | 45.8 | 7.6 |
| Age | | | | | |
| 50-64 | 89 | 45.3 | 107 | 54.7 | 7.7 |
| ≥65 | 83 | 60.6 | 51 | 39.4 | 9.2 |
| Education | | | | | |
| ≤High School | 79 | 47.4 | 81 | 52.6 | 8.4 |
| >High School | 93 | 56.1 | 76 | 43.9 | 8.4 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older

Columbia, Greene

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 149 | 44.6 | 178 | 55.4 | 6.0 |
| Gender | | | | | |
| Male | 58 | 43.3 | 68 | 56.7 | 9.2 |
| Female | 91 | 45.8 | 110 | 54.2 | 7.7 |
| Age | | | | | |
| 50-64 | 80 | 41.2 | 115 | 58.8 | 7.7 |
| ≥65 | 69 | 49.2 | 63 | 50.8 | 9.4 |
| Education | | | | | |
| ≤High School | 67 | 40.4 | 91 | 59.6 | 8.3 |
| >High School | 82 | 48.6 | 86 | 51.4 | 8.5 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older

Columbia, Greene

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 174 | 52.0 | 156 | 48.0 | 6.0 |
| Gender | | | | | |
| Male | 67 | 50.6 | 59 | 49.4 | 9.4 |
| Female | 107 | 53.2 | 97 | 46.8 | 7.7 |
| Age | | | | | |
| 50-64 | 97 | 49.6 | 99 | 50.4 | 7.8 |
| ≥65 | 77 | 55.2 | 57 | 44.8 | 9.4 |
| Education | | | | | |
| ≤High School | 77 | 46.6 | 82 | 53.4 | 8.4 |
| >High School | 97 | 57.0 | 73 | 43.0 | 8.5 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval