

**Expanded BRFSS, 2003**

**New York Locality**

Table of Contents:

Optional Module 1: Adult Asthma History ..... 1  
Optional Module 5: Diabetes..... 2  
Optional Module 7: Disability/Quality of Life..... 6  
  
Optional Module 10: Health Care Coverage ..... 10  
Optional Module 13: Injury Control - Falls..... 12  
Optional Module 19: Tobacco Consumption ..... 13  
Optional Module 20: Tobacco - Media ..... 14  
Optional Module 21: Tobacco – other products ..... 22  
Optional Module 22: Tobacco – Work site ETS ..... 23  
Optional Module 23: Seat Belts..... 25  
Optional Module 24: Weight Control..... 27

**New York State Department of Health**

Expanded BRFSS, 2003 - New York State Department of Health

## Have had an Asthma Episode or Attack in the Past 12 Months, among Those with Asthma

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	32	64.1	18	35.9	16.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Check Blood for Glucose or Sugar at Least Once per Day among Those with Diabetes

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	15	46.9	18	53.1	20.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Check Feet for Sores or Irritations at Least Once per Day among Those with Diabetes

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	14	37.3	18	62.7	18.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Had Feet Checked by a Medical Professional in Past Twelve Months, among Those with Diabetes

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	21	63.0	10	37.0	21.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Had Dilated Eye Exam in Past 12 Months, among Those with Diabetes

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	30	97.2	2	2.8	4.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Adults with a Disability

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	104	14.3	500	85.7	3.0
Gender					
Male	33	10.8	209	89.2	4.1
Female	71	17.3	291	82.7	4.2
Age					
18-34	15	6.5	164	93.5	3.5
35-54	28	10.4	199	89.6	4.1
≥55	59	29.9	127	70.1	7.6
Education					
≤High School	37	20.1	103	79.9	7.2
>High School	65	11.6	395	88.4	3.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Adults with a Disability Requiring Help

*New York*

Demographic Groups	Disability need help			Disability no help needed			No Disability		
	n <sup>1</sup>	% <sup>2</sup>	C.I. <sup>3</sup>	n	%	C.I.	n	%	C.I.
Total	41	6.3	2.2	63	8.0	2.1	500	85.7	3.0
Gender									
Male	9	4.1	3.0	24	6.7	2.9	209	89.2	4.1
Female	32	8.2	3.2	39	9.1	3.0	291	82.7	4.2
Age									
18-64	20	3.8	1.9	38	6.5	2.2	420	89.8	2.9
≥65	19	19.8	9.1	25	17.4	7.0	70	62.8	10.1
Education									
≤High School	26	15.5	6.7	11	4.6	2.9	103	79.9	7.2
>High School	14	2.5	1.4	51	9.1	2.7	395	88.4	3.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



This page left intentionally blank

This page left intentionally blank

Expanded BRFSS, 2003 - New York State Department of Health

## Main Reason for Not Having Health Care Coverage

*New York*

Demographic Groups	n <sup>1</sup>	Changed or lost job % <sup>2</sup>	C.I. <sup>3</sup>	n	Could not afford %	C.I.	n	Other %	C.I.
Total	9	14.2	10.4	19	23.0	11.1	32	62.8	14.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## No Health Care Coverage for Past Two Years Among Uninsured

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	24	41.6	33	58.4	16.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Had a Fall in Past Three Months

*New York*

Demographic Groups	n <sup>1</sup>	No fall % <sup>2</sup>	C.I. <sup>3</sup>	n	Fall with injury %	C.I.	n	Fall no injury %	C.I.
Total	254	88.1	4.0	17	6.0	3.2	21	5.9	2.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Number of Cigarettes Smoked per day among Everyday Smokers

*New York*

Demographic Groups	n <sup>1</sup>	1-10 cigarettes % <sup>2</sup>	n	≥11 cigarettes %	C.I. <sup>3</sup>
Total	40	57.0	31	43.0	13.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Heard, Read or Seen Information about the Effects of Tobacco from Television

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	331	61.2	240	38.8	4.6
Gender					
Male	135	62.1	95	37.9	7.2
Female	196	60.4	145	39.6	5.9
Age					
18-34	127	75.1	48	24.9	7.0
35-54	118	53.6	101	46.4	7.6
≥55	81	51.3	88	48.7	8.8
Education					
≤High School	92	70.0	42	30.0	8.9
>High School	239	58.0	196	42.0	5.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Heard, Read or Seen Information about the Effects of Tobacco from Radio

*New York*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	114	21.1	457	78.9	4.0
Gender					
Male	47	21.9	179	78.1	6.4
Female	67	20.3	278	79.7	5.0
Age					
18-34	34	20.3	141	79.7	7.2
35-54	34	17.8	182	82.2	6.0
≥55	43	26.2	129	73.8	7.8
Education					
≤High School	43	31.4	91	68.6	9.3
>High School	71	17.1	364	82.9	4.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

**Heard, Read or Seen Information about the Effects of Tobacco from Billboards**

*New York*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	195	36.6	376	63.4	4.7
Gender					
Male	78	36.5	153	63.5	7.4
Female	117	36.7	223	63.3	6.0
Age					
18-34	70	40.0	102	60.0	8.6
35-54	81	38.8	134	61.2	7.4
≥55	40	29.2	134	70.8	8.6
Education					
≤High School	46	38.4	87	61.6	10.2
>High School	149	36.2	286	63.8	5.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Heard, Read or Seen Information about the Effects of Tobacco from Buses

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	153	30.5	413	69.5	4.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Heard, Read or Seen Information about the Effects of Tobacco from Magazines

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	282	49.8	297	50.2	4.8
Gender					
Male	119	49.7	111	50.3	7.6
Female	163	50.0	186	50.0	6.0
Age					
18-34	88	50.2	87	49.8	8.6
35-54	92	44.6	127	55.4	7.6
≥55	95	55.8	80	44.2	8.6
Education					
≤High School	63	47.5	72	52.5	10.1
>High School	219	51.1	222	48.9	5.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Heard, Read or Seen Information about the Effects of Tobacco from a Website

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	60	11.0	523	89.0	3.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Heard, Read or Seen Information about the Effects of Tobacco from Movie Theater

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	97	17.3	481	82.7	3.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Heard or Seen Anti-Tobacco Advertisement that Refers to the NYS Quitline in Past 30 Days

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	221	43.7	329	56.3	4.9
Gender					
Male	83	43.5	140	56.5	7.8
Female	138	43.9	189	56.1	6.2
Age					
18-34	83	52.4	85	47.6	8.8
35-54	81	40.0	127	60.0	7.8
≥55	52	36.1	114	63.9	9.0
Education					
≤High School	57	52.3	72	47.7	10.3
>High School	164	40.6	254	59.4	5.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Used or Tried Any Smokeless Tobacco Products such as Chewing Tobacco or Snuff

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	42	7.2	546	92.8	2.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Work Indoors Most of the Time

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	321	89.9	36	10.1	3.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

**Smoking Policy at Place of Work among Indoor Workers***New York*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Not allowed anywhere %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Allowed in some or all places %</b>	<b>C.I.</b>	<b>n</b>	<b>No official policy %</b>	<b>C.I.</b>
Total	275	86.5	4.4	25	8.3	3.7	19	5.2	2.6
Gender									
Male	117	87.6	6.2	8	6.0	4.9	12	6.4	4.1
Female	158	85.4	6.1	17	10.6	5.5	7	4.0	3.1
Age									
18-34	97	87.3	7.5	12	12.2	7.5	1	0.5	0.9
35-54	124	84.9	6.4	12	7.1	4.6	12	8.0	4.7
≥55	50	88.3	9.7	1	1.5	3.0	6	10.2	9.3
Education									
≤High School	30	87.8	13.6	2	7.2	12.1	2	5.0	7.1
>High School	245	86.3	4.5	23	8.5	3.8	17	5.2	2.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution<sup>2</sup>Weighted Percent<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Use Seatbelt when Ride or Drive in a Car (Always, Nearly Always or Never Ride in a Car)**

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	521	90.0	65	10.0	2.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Use Seatbelt when Ride or Drive in a Car (Always or Never ride in a Car)

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	431	73.8	155	26.2	4.1
Gender					
Male	171	74.0	61	26.0	6.5
Female	260	73.7	94	26.3	5.2
Age					
18-64	337	72.8	131	27.2	4.6
≥65	86	79.0	21	21.0	9.5
Education					
≤High School	100	73.3	35	26.7	8.7
>High School	328	73.9	120	26.1	4.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Trying to Lose Weight

*New York*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	268	44.5	317	55.5	4.7
Gender					
Male	93	38.6	140	61.4	7.3
Female	175	49.6	177	50.4	6.0
Age					
18-34	68	38.1	108	61.9	8.3
35-54	104	45.1	116	54.9	7.5
≥55	93	53.9	85	46.1	8.6
Education					
≤High School	63	47.4	71	52.6	10.1
>High School	205	43.6	243	56.4	5.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Trying to Maintain Current Weight of Those not Trying to Lose Weight

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	205	61.3	113	38.7	6.4
Gender					
Male	83	55.7	57	44.3	9.7
Female	122	67.2	56	32.8	7.9
Age					
18-34	66	51.8	42	48.2	10.9
35-54	77	68.7	40	31.3	9.3
≥55	58	68.1	27	31.9	11.3
Education					
≤High School	40	53.3	32	46.7	14.1
>High School	165	64.9	78	35.1	6.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Eating Fewer Calories or Less Fat to Lose or Maintain Weight

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	325	69.1	140	30.9	5.0
Gender					
Male	116	63.8	56	36.2	8.6
Female	209	73.0	84	27.0	5.7
Age					
18-34	87	65.7	47	34.3	9.4
35-54	120	64.1	57	35.9	8.4
≥55	113	79.7	34	20.3	6.8
Education					
≤High School	71	71.6	27	28.4	10.8
>High School	254	68.2	113	31.8	5.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Using Physical Exercise to Lose or Maintain Weight

*New York*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	337	70.1	136	29.9	4.9
Gender					
Male	134	74.8	42	25.2	7.7
Female	203	66.5	94	33.5	6.3
Age					
18-34	110	78.9	24	21.1	8.9
35-54	136	71.4	45	28.6	7.7
≥55	86	58.3	65	41.7	9.2
Education					
≤High School	48	46.2	55	53.8	11.2
>High School	289	78.7	81	21.3	4.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval