

HHSC Performance Measures

Measure ID	Measure Title	Measure Description
ADD-Cont	Follow-up care for children prescribed ADHD medication (continuation)	Percentage of children ages 6-12 newly prescribed ADHD medication who had appropriate follow up in the continuation and maintenance phase.
ADD-Init	Follow-up care for children prescribed ADHD medication (initiation)	Percentage of children ages 6-12 newly prescribed ADHD medication who had appropriate follow up in the initial 30 days
ADV	Annual Dental Visit	Percentage of children and adolescents, ages 2 to 18 years, who had at least one dental visit within the measurement year
AMB-ED	ED Utilization	Rate of utilization of ambulatory care ED visits per 1,000 member months. Does not include mental health- or chemical dependency-related services
AMM-Acute	Antidepressant medication management (Acute Phase)	Percentage of members age 18 and older with a diagnosis of major depression and were treated with antidepressant medication, who remained on antidepressant medication during the entire 12-week acute treatment phase (84 days).
AMM-Cont	Antidepressant medication management (Continuation Phase)	Percentage of Medicaid enrollees age 18 and older with a diagnosis of major depression and were treated with antidepressant medication, who remained on antidepressant medication for at least 6 months (180 days).
APC	Multiple Concurrent Antipsychotic Use in Children and Adolescents	The percentage of children and adolescents 1–17 years of age who were on two or more concurrent antipsychotic medications
APM	Metabolic Monitoring for Children and Adolescents on antipsychotics	The percentage of children and adolescents 1–17 years of age who had two or more antipsychotic prescriptions and had metabolic testing
APP	Use of First-Line Psychosocial Care for Children and Adolescent on Antipsychotics	The percentage of children and adolescents 1–17 years of age who had a new prescription for an antipsychotic medication and had documentation of psychosocial care as first-line treatment
AWC	Adolescent Well-Care Visits	Percentage of adolescents, ages 12 to 21 years, who had at least one comprehensive well-care visit with a primary care provider during the measurement year
BMS	Adherence to mood stabilizers for individuals with bipolar I disorder	Percentage of patients with bipolar I disorder who received a mood stabilizer medication that had a proportion of days covered (PDC) for mood stabilizer medication ≥ 0.8 during the measurement year.

CBP	Controlling High Blood Pressure (hybrid)	Percentage of Health Home members (sample of MMC members only) ages 18 to 85 who had a diagnosis of hypertension and whose blood pressure was adequately controlled (<140/90) during the measurement year.
CDC-hA1c	Comprehensive diabetes care (HbA1c test)	Percentage of members ages 18 to 75 with diabetes (type 1 and type 2) who had at least one Hemoglobin A1c test within the year.
CHL	Chlamydia Screening in Women	Percentage of Health Home members (women only) ages 21 to 24 who were identified as sexually active and who had at least one test for Chlamydia during the measurement year.
CIS	Childhood Immunization Status	Percentage of members, age 2 years, who were fully immunized. The HEDIS specifications for fully immunized consists of the following vaccines: 4 Diphtheria/Tetanus/Pertussis, 3 Polio, 1 Measles/Mumps/Rubella, 3 H Influenza type B, 3 Hepatitis B, 1 Varicella, and 4 pneumococcal
CWP	Appropriate Testing for Children with Pharyngitis	The percentage of children, ages 2 to 18 years, who were diagnosed with pharyngitis, were prescribed an antibiotic, and who were given a group A streptococcus test
FUA-30 day	Follow-Up After Emergency Department Visit for Alcohol and Other Drug Dependence	Percentage of ED visits for members with a principal diagnosis of alcohol or other drug (AOD) dependence, who had a follow-up visit for AOD within 30 days of the ED visit.
FUA-7 day	Follow-Up After Emergency Department Visit for Alcohol and Other Drug Dependence	Percentage of ED visits for members with a principal diagnosis of alcohol or other drug (AOD) dependence, who had a follow-up visit for AOD within 7 days of the ED visit.
FUH-30 day	Follow Up After Hospitalization for Mental Illness	Percentage of discharges (for treatment of selected mental illness disorders) for members 6 - 64 years of age who were seen on an ambulatory bases or who were in intermediate treatment with a mental health provider within 30 days of discharge.
FUH-7day	Follow Up After Hospitalization for Mental Illness	Percentage of discharges (for treatment of selected mental illness disorders) for members 6 - 64 years of age who were seen on an ambulatory bases or who were in intermediate treatment with a mental health provider within 7 days of discharge.
FUM-30 day	Follow-up After Emergency Department Visit for Mental Illness (FUM)	Percentage of emergency department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness, who had a f/up visit for mental illness within 30 days of the ED visit.
FUM-7 day	Follow-up After Emergency Department Visit for Mental Illness (FUM)	Percentage of emergency department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness, who had a follow-up visit for mental within 7 days of the ED visit.

IET-Engage	Initiation and Engagement of Alcohol and Other Drug Dependence Treatment	Percentage of Health Home enrollees age 13 and older with a new episode of alcohol or other drug (AOD) dependence who initiated treatment and had two or more additional services with a diagnosis of AOD within 30 days of the initiation visit.
IET-Initiation	Initiation and Engagement of Alcohol and Other Drug Dependence Treatment	Percentage of Health Home enrollees age 13 and older with a new episode of alcohol or other drug (AOD) dependence who initiated treatment through an inpatient AOD admission, outpatient visit, intensive outpatient encounter, or partial hospitalization within 14 days of the diagnosis
IMA	Immunizations for Adolescents	Percentage of members, age 13 years, who had one dose of meningococcal vaccine between their 11th and 13th birthdays and one dose of tetanus, diphtheria and pertussis vaccine between their 10th and their 13th birthdays
IPU	Inpatient Utilization – General hospital/Acute Care	Rate of utilization of total acute inpatient stays per 1,000 member months. Does not include mental health- or chemical dependency-related services
LSC	Lead Screening in Children	Percentage of children, age 2 years, who had their blood tested for lead poisoning at least once by their 2nd birthday
MMA-50%	Medication management for people with asthma	Percentage of members age 5 to 64 who were identified as having persistent asthma and filled prescriptions for asthma controller medication during at least 50% of their treatment period
MMA-75%	Medication management for people with asthma	Percentage of members age 5 to 64 who were identified as having persistent asthma and filled prescriptions for asthma controller medication during at least 75% of their treatment period
NFU	Skilled Nursing Home Admission	Number of admissions to a nursing facility from the community that result in a short-term (less than 101 days) during the measurement year per 1,000 member months (18 and older)
PBH	Persistence of beta-blocker treatment after heart attack	Percentage of members who were hospitalized and discharged alive with a diagnosis of AMI and who received persistent beta-blocker treatment for six months after discharge.
PCP	Primary Care	Rate of utilization of primary care visits (PCP) visits per 1,000 member months.
PCR	Plan All-Cause Readmission Rate	Number of acute inpatient stays, for Health Home members age 18 and older, during the measurement year that were followed by an acute readmission for any diagnosis within 30 days
PPR	Potentially Preventable Readmissions	Number of potentially preventable readmission chains per 100,000 enrollees-(at risk admissions followed by one or more clinically related readmissions within 30 days of discharge).

PPV	Potentially Preventable Emergency Room Visits	Number of potentially preventable emergency visits for ambulatory sensitive conditions per 100 enrollees. numerator is the number of potentially preventable visits and the denominator is the number of eligible enrollees
PQI-92	Chronic Condition Hospital Admission Composite— Prevention	Total number of hospital admissions for chronic conditions per 100,000 Health Home members age 18 and older.
URI	Appropriate Treatment for Children with Upper Respiratory Infection	The percentage of children, ages 3 months to 18 years, who were diagnosed with an upper respiratory infection (common cold) and who were NOT given a prescription for an antibiotic
VLS	Viral Load Suppression	Percentage of Medicaid enrollees confirmed HIV-Positive who had a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year
W15	Well-Child Visits in the First 15 Months of Life	Percentage of children who had five or more well-child visits with a primary care provider in their first 15 months of life
W34	Well-Child Visits in the 3rd, 4th, 5th & 6th Year	Percentage of children, ages 3 to 6 years, who had one or more well-child visits with a primary care provider during the measurement year
WCC	Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents	The percentage of Health Home members, ages 3 to 17 years, who had an outpatient visit with a PCP or OB/GYN during the measurement year and who had evidence of the following during the measurement year: BMI percentile documentation, Counseling for nutrition and Counseling for physical activity