

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York State, Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-----------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 11726.6 | 367.1 | 364.0-370.1 | 5580.6 | 410.1 | 405.0-415.2 | 6146.0 | 344.4 | 340.5-348.3 | 3056.2 | 103.3 | 101.6-105.0 | 1537.6 | 127.6 | 124.7-130.7 | 1518.6 | 87.8 | 85.9-89.9 |
| Oral cavity and pharynx | 268.8 | 8.3 | 7.9-8.8 | 183.4 | 13.0 | 12.2-14.0 | 85.4 | 4.8 | 4.3-5.3 | 55.6 | 1.8 | 1.6-2.1 | 41.8 | 3.3 | 2.8-3.8 | 13.8 | 0.8 | 0.6-1.0 |
| Esophagus | 87.6 | 2.9 | 2.6-3.2 | 66.2 | 5.1 | 4.6-5.7 | 21.4 | 1.3 | 1.0-1.5 | 56.8 | 1.9 | 1.7-2.1 | 44.2 | 3.5 | 3.0-4.0 | 12.6 | 0.8 | 0.6-1.0 |
| Stomach | 334.4 | 10.8 | 10.3-11.4 | 178.8 | 13.5 | 12.6-14.5 | 155.6 | 9.0 | 8.4-9.6 | 132.8 | 4.4 | 4.1-4.8 | 76.8 | 6.2 | 5.6-6.9 | 56.0 | 3.2 | 2.9-3.6 |
| Colorectal | 1004.2 | 32.0 | 31.1-33.0 | 534.6 | 39.5 | 37.9-41.1 | 469.6 | 26.7 | 25.6-27.8 | 280.6 | 9.5 | 9.0-10.0 | 151.2 | 12.3 | 11.4-13.2 | 129.4 | 7.5 | 7.0-8.2 |
| Colon excluding rectum | 692.4 | 22.4 | 21.7-23.2 | 351.6 | 26.6 | 25.3-27.9 | 340.8 | 19.5 | 18.6-20.5 | 225.0 | 7.7 | 7.2-8.1 | 117.0 | 9.6 | 8.8-10.4 | 108.0 | 6.3 | 5.8-6.8 |
| Rectum & rectosigmoid | 311.8 | 9.6 | 9.1-10.1 | 183.0 | 12.9 | 12.1-13.8 | 128.8 | 7.2 | 6.6-7.7 | 55.6 | 1.9 | 1.6-2.1 | 34.2 | 2.7 | 2.3-3.1 | 21.4 | 1.3 | 1.0-1.5 |
| Liver / intrahepatic bile duct | 419.4 | 13.2 | 12.6-13.8 | 287.8 | 20.7 | 19.6-21.9 | 131.6 | 7.5 | 6.9-8.1 | 239.4 | 7.8 | 7.4-8.3 | 157.6 | 12.0 | 11.2-13.0 | 81.8 | 4.7 | 4.3-5.2 |
| Pancreas | 361.0 | 11.9 | 11.4-12.5 | 177.4 | 13.7 | 12.8-14.7 | 183.6 | 10.6 | 9.9-11.4 | 232.8 | 8.0 | 7.5-8.4 | 113.8 | 9.3 | 8.5-10.1 | 119.0 | 7.0 | 6.4-7.6 |
| Larynx | 87.4 | 2.7 | 2.5-3.0 | 67.2 | 5.0 | 4.5-5.6 | 20.2 | 1.1 | 0.9-1.3 | 26.4 | 0.9 | 0.7-1.0 | 22.2 | 1.8 | 1.4-2.1 | 4.2 | 0.2 | 0.1-0.3 |
| Lung and bronchus | 966.2 | 32.3 | 31.4-33.2 | 501.8 | 41.0 | 39.3-42.7 | 464.4 | 26.6 | 25.5-27.7 | 465.0 | 15.9 | 15.3-16.6 | 261.8 | 22.1 | 20.9-23.4 | 203.2 | 11.8 | 11.0-12.5 |
| Melanoma of the skin | 92.8 | 3.0 | 2.7-3.3 | 49.4 | 3.7 | 3.3-4.3 | 43.4 | 2.5 | 2.2-2.8 | 12.2 | 0.4 | 0.3-0.5 | 7.8 | 0.6 | 0.4-0.8 | 4.4 | 0.3 | 0.2-0.4 |
| Female breast | | | | | | | 1798.0 | 100.0 | 97.9-102.1 | | | | | | | 234.8 | 13.4 | 12.6-14.2 |
| Cervix uteri | | | | | | | 161.2 | 9.0 | 8.3-9.6 | | | | | | | 37.2 | 2.1 | 1.8-2.4 |
| Corpus uteri and NOS | | | | | | | 437.0 | 23.9 | 22.9-25.0 | | | | | | | 83.0 | 4.7 | 4.3-5.2 |
| Ovary | | | | | | | 164.0 | 9.1 | 8.5-9.8 | | | | | | | 76.2 | 4.3 | 3.9-4.8 |
| Prostate | | | | 1494.0 | 109.3 | 106.7-111.9 | | | | | | | 166.0 | 16.4 | 15.3-17.6 | | | |
| Testis | | | | 83.6 | 4.1 | 3.7-4.5 | | | | | | | 4.2 | 0.2 | 0.1-0.3 | | | |
| Urinary bladder (incl. in situ) | 330.2 | 11.3 | 10.8-11.9 | 236.2 | 20.1 | 18.9-21.3 | 94.0 | 5.5 | 5.0-6.0 | 61.6 | 2.3 | 2.1-2.6 | 40.8 | 4.1 | 3.5-4.7 | 20.8 | 1.3 | 1.0-1.5 |
| Kidney and renal pelvis | 410.2 | 12.6 | 12.0-13.2 | 250.0 | 17.5 | 16.5-18.5 | 160.2 | 8.9 | 8.3-9.5 | 54.0 | 1.8 | 1.6-2.1 | 33.2 | 2.7 | 2.2-3.1 | 20.8 | 1.2 | 1.0-1.5 |
| Brain and other central nervous system | 178.6 | 5.2 | 4.9-5.6 | 100.2 | 6.4 | 5.8-7.0 | 78.4 | 4.3 | 3.9-4.7 | 84.4 | 2.7 | 2.4-2.9 | 45.6 | 3.3 | 2.9-3.8 | 38.8 | 2.2 | 1.9-2.5 |
| Thyroid | 570.2 | 16.1 | 15.5-16.7 | 112.8 | 6.9 | 6.3-7.5 | 457.4 | 24.8 | 23.8-25.9 | 13.0 | 0.5 | 0.4-0.6 | 4.0 | 0.3 | 0.2-0.5 | 9.0 | 0.5 | 0.4-0.7 |
| Hodgkin lymphoma | 94.4 | 2.6 | 2.4-2.9 | 53.2 | 3.1 | 2.8-3.6 | 41.2 | 2.2 | 1.9-2.5 | 8.6 | 0.3 | 0.2-0.4 | 4.4 | 0.3 | 0.2-0.4 | 4.2 | 0.3 | 0.2-0.4 |
| Non-Hodgkin lymphomas | 582.6 | 18.4 | 17.7-19.1 | 300.6 | 21.8 | 20.6-23.0 | 282.0 | 15.9 | 15.1-16.8 | 117.0 | 4.0 | 3.7-4.4 | 64.8 | 5.5 | 4.9-6.1 | 52.2 | 3.1 | 2.7-3.5 |
| Myeloma | 260.4 | 8.5 | 8.0-8.9 | 133.6 | 10.2 | 9.4-11.0 | 126.8 | 7.2 | 6.7-7.8 | 71.8 | 2.5 | 2.2-2.8 | 35.4 | 3.0 | 2.5-3.5 | 36.4 | 2.2 | 1.9-2.5 |
| Leukemias | 391.4 | 11.9 | 11.4-12.5 | 224.2 | 15.5 | 14.5-16.5 | 167.2 | 9.3 | 8.6-9.9 | 133.8 | 4.4 | 4.0-4.7 | 71.0 | 5.4 | 4.9-6.1 | 62.8 | 3.6 | 3.2-4.0 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York State, Non-Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 104317.8 | 503.5 | 502.1-504.9 | 52179.4 | 546.7 | 544.5-548.8 | 52138.4 | 475.6 | 473.7-477.5 | 31042.4 | 143.2 | 142.5-143.9 | 15504.6 | 166.0 | 164.8-167.2 | 15537.8 | 127.6 | 126.7-128.5 |
| Oral cavity and pharynx | 2502.4 | 12.0 | 11.8-12.2 | 1708.2 | 17.7 | 17.3-18.1 | 794.2 | 7.1 | 6.9-7.3 | 480.6 | 2.2 | 2.1-2.3 | 331.4 | 3.4 | 3.3-3.6 | 149.2 | 1.2 | 1.1-1.3 |
| Esophagus | 1012.0 | 4.7 | 4.5-4.8 | 761.4 | 7.8 | 7.6-8.1 | 250.6 | 2.1 | 2.0-2.2 | 776.2 | 3.5 | 3.4-3.6 | 599.2 | 6.2 | 5.9-6.4 | 177.0 | 1.4 | 1.3-1.5 |
| Stomach | 1706.8 | 8.1 | 8.0-8.3 | 1026.6 | 10.9 | 10.6-11.2 | 680.2 | 6.0 | 5.8-6.2 | 668.2 | 3.1 | 3.0-3.3 | 400.0 | 4.3 | 4.1-4.5 | 268.2 | 2.2 | 2.1-2.4 |
| Colorectal | 7969.8 | 38.6 | 38.2-39.0 | 4079.2 | 44.0 | 43.4-44.6 | 3890.6 | 34.2 | 33.7-34.7 | 2659.0 | 12.3 | 12.1-12.5 | 1324.6 | 14.3 | 14.0-14.7 | 1334.4 | 10.7 | 10.5-11.0 |
| Colon excluding rectum | 5643.8 | 27.2 | 26.8-27.5 | 2749.8 | 29.8 | 29.3-30.3 | 2894.0 | 25.1 | 24.6-25.5 | 2129.2 | 9.8 | 9.6-10.0 | 1028.0 | 11.1 | 10.8-11.5 | 1101.2 | 8.8 | 8.6-9.0 |
| Rectum & rectosigmoid | 2326.0 | 11.5 | 11.3-11.7 | 1329.4 | 14.2 | 13.9-14.6 | 996.6 | 9.2 | 8.9-9.4 | 529.8 | 2.5 | 2.4-2.6 | 296.6 | 3.2 | 3.0-3.4 | 233.2 | 1.9 | 1.8-2.0 |
| Liver / intrahepatic bile duct | 1816.8 | 8.3 | 8.1-8.5 | 1294.8 | 12.9 | 12.6-13.3 | 522.0 | 4.4 | 4.2-4.6 | 1198.2 | 5.5 | 5.3-5.6 | 803.6 | 8.2 | 7.9-8.5 | 394.6 | 3.2 | 3.1-3.4 |
| Pancreas | 3190.2 | 14.8 | 14.5-15.0 | 1593.2 | 16.7 | 16.3-17.1 | 1597.0 | 13.2 | 12.9-13.5 | 2443.8 | 11.2 | 11.0-11.4 | 1213.2 | 12.7 | 12.4-13.1 | 1230.6 | 9.9 | 9.6-10.1 |
| Larynx | 656.4 | 3.0 | 2.9-3.1 | 520.2 | 5.3 | 5.1-5.5 | 136.2 | 1.2 | 1.1-1.3 | 182.8 | 0.8 | 0.8-0.9 | 142.6 | 1.5 | 1.4-1.6 | 40.2 | 0.3 | 0.3-0.4 |
| Lung and bronchus | 13268.6 | 61.2 | 60.7-61.6 | 6399.2 | 66.7 | 65.9-67.4 | 6869.4 | 57.3 | 56.7-58.0 | 7316.2 | 33.5 | 33.1-33.8 | 3727.6 | 39.3 | 38.7-39.9 | 3588.6 | 29.2 | 28.8-29.6 |
| Melanoma of the skin | 4168.0 | 20.6 | 20.4-20.9 | 2421.6 | 26.2 | 25.8-26.7 | 1746.4 | 16.7 | 16.3-17.0 | 375.0 | 1.8 | 1.7-1.8 | 235.0 | 2.6 | 2.4-2.7 | 140.0 | 1.2 | 1.1-1.3 |
| Female breast | | | | | | | 15021.0 | 142.2 | 141.1-143.2 | | | | | | | 2270.4 | 19.4 | 19.0-19.7 |
| Cervix uteri | | | | | | | 679.4 | 7.4 | 7.2-7.7 | | | | | | | 202.2 | 1.9 | 1.8-2.1 |
| Corpus uterus and NOS | | | | | | | 3761.0 | 32.9 | 32.4-33.4 | | | | | | | 741.0 | 6.1 | 5.9-6.3 |
| Ovary | | | | | | | 1282.8 | 12.0 | 11.7-12.3 | | | | | | | 803.0 | 6.8 | 6.5-7.0 |
| Prostate | | | | 13748.4 | 133.7 | 132.7-134.8 | | | | | | | 1540.8 | 17.0 | 16.7-17.4 | | | |
| Testis | | | | 496.6 | 6.6 | 6.4-6.9 | | | | | | | 16.6 | 0.2 | 0.2-0.2 | | | |
| Urinary bladder (incl. in situ) | 5116.0 | 23.7 | 23.4-24.0 | 3825.6 | 40.8 | 40.2-41.4 | 1290.4 | 10.6 | 10.4-10.9 | 918.2 | 4.1 | 4.0-4.2 | 642.2 | 7.1 | 6.8-7.3 | 276.0 | 2.1 | 2.0-2.2 |
| Kidney and renal pelvis | 3663.2 | 17.8 | 17.6-18.1 | 2374.0 | 25.1 | 24.6-25.6 | 1289.2 | 11.7 | 11.4-12.0 | 626.4 | 2.9 | 2.8-3.0 | 403.2 | 4.3 | 4.1-4.5 | 223.2 | 1.8 | 1.7-1.9 |
| Brain and other central nervous system | 1250.6 | 6.8 | 6.6-7.0 | 692.2 | 8.0 | 7.7-8.3 | 558.4 | 5.7 | 5.5-6.0 | 804.0 | 4.0 | 3.9-4.1 | 447.6 | 4.8 | 4.6-5.1 | 356.4 | 3.2 | 3.1-3.4 |
| Thyroid | 3456.2 | 20.1 | 19.8-20.4 | 941.6 | 11.0 | 10.7-11.4 | 2514.6 | 28.7 | 28.2-29.2 | 113.8 | 0.5 | 0.5-0.6 | 51.0 | 0.6 | 0.5-0.6 | 62.8 | 0.5 | 0.4-0.6 |
| Hodgkin lymphoma | 538.2 | 3.2 | 3.1-3.4 | 289.4 | 3.6 | 3.4-3.8 | 248.8 | 2.9 | 2.8-3.1 | 55.8 | 0.3 | 0.2-0.3 | 33.2 | 0.4 | 0.3-0.4 | 22.6 | 0.2 | 0.2-0.2 |
| Non-Hodgkin lymphomas | 4484.0 | 21.9 | 21.6-22.2 | 2420.6 | 26.4 | 25.9-26.9 | 2063.4 | 18.4 | 18.1-18.8 | 1078.2 | 5.0 | 4.9-5.1 | 601.8 | 6.6 | 6.4-6.8 | 476.4 | 3.8 | 3.6-4.0 |
| Myeloma | 1746.0 | 8.2 | 8.0-8.4 | 953.8 | 10.1 | 9.8-10.3 | 792.2 | 6.8 | 6.6-7.0 | 629.8 | 2.9 | 2.8-3.0 | 333.8 | 3.6 | 3.4-3.8 | 296.0 | 2.3 | 2.2-2.5 |
| Leukemias | 3359.4 | 16.5 | 16.3-16.8 | 1950.4 | 21.4 | 21.0-21.9 | 1409.0 | 12.7 | 12.4-13.0 | 1240.6 | 5.8 | 5.7-6.0 | 700.8 | 7.8 | 7.5-8.1 | 539.8 | 4.4 | 4.2-4.6 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York State, Non-Hispanic Whites

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 81653.4 | 522.1 | 520.5-523.8 | 41157.2 | 559.6 | 557.1-562.1 | 40496.2 | 499.2 | 496.9-501.5 | 24421.0 | 145.1 | 144.2-145.9 | 12299.4 | 167.2 | 165.8-168.5 | 12121.6 | 129.6 | 128.6-130.7 |
| Oral cavity and pharynx | 2007.6 | 12.7 | 12.4-13.0 | 1393.4 | 18.7 | 18.3-19.2 | 614.2 | 7.3 | 7.1-7.6 | 369.8 | 2.2 | 2.1-2.3 | 253.0 | 3.3 | 3.2-3.5 | 116.8 | 1.2 | 1.1-1.3 |
| Esophagus | 840.6 | 5.1 | 4.9-5.2 | 645.0 | 8.5 | 8.2-8.8 | 195.6 | 2.1 | 2.0-2.3 | 664.2 | 3.9 | 3.8-4.1 | 519.2 | 6.8 | 6.6-7.1 | 145.0 | 1.5 | 1.4-1.6 |
| Stomach | 1076.0 | 6.7 | 6.5-6.9 | 666.4 | 9.1 | 8.8-9.5 | 409.6 | 4.7 | 4.5-5.0 | 404.2 | 2.4 | 2.3-2.6 | 243.6 | 3.3 | 3.2-3.5 | 160.6 | 1.7 | 1.6-1.9 |
| Colorectal | 6020.4 | 38.6 | 38.1-39.0 | 3097.6 | 43.6 | 42.9-44.3 | 2922.8 | 34.3 | 33.7-34.9 | 2027.0 | 12.1 | 11.8-12.3 | 1017.0 | 14.1 | 13.7-14.5 | 1010.0 | 10.5 | 10.2-10.8 |
| Colon excluding rectum | 4294.2 | 27.2 | 26.8-27.6 | 2104.0 | 29.6 | 29.0-30.2 | 2190.2 | 25.1 | 24.6-25.6 | 1611.0 | 9.5 | 9.3-9.8 | 783.0 | 10.8 | 10.5-11.2 | 828.0 | 8.5 | 8.2-8.8 |
| Rectum & rectosigmoid | 1726.2 | 11.4 | 11.1-11.6 | 993.6 | 14.0 | 13.6-14.4 | 732.6 | 9.1 | 8.8-9.4 | 416.0 | 2.6 | 2.4-2.7 | 234.0 | 3.2 | 3.1-3.4 | 182.0 | 2.0 | 1.8-2.1 |
| Liver / intrahepatic bile duct | 1171.4 | 7.0 | 6.8-7.2 | 837.0 | 10.8 | 10.4-11.1 | 334.4 | 3.8 | 3.6-4.0 | 797.4 | 4.7 | 4.5-4.8 | 531.8 | 6.9 | 6.7-7.2 | 265.6 | 2.8 | 2.7-3.0 |
| Pancreas | 2490.2 | 15.0 | 14.7-15.3 | 1270.2 | 17.1 | 16.7-17.5 | 1220.0 | 13.2 | 12.8-13.5 | 1927.2 | 11.3 | 11.1-11.6 | 977.2 | 13.1 | 12.7-13.5 | 950.0 | 9.9 | 9.6-10.2 |
| Larynx | 517.2 | 3.2 | 3.0-3.3 | 411.8 | 5.4 | 5.2-5.7 | 105.4 | 1.3 | 1.2-1.4 | 140.0 | 0.8 | 0.8-0.9 | 109.4 | 1.4 | 1.3-1.6 | 30.6 | 0.3 | 0.3-0.4 |
| Lung and bronchus | 10846.6 | 65.0 | 64.5-65.6 | 5137.4 | 68.4 | 67.6-69.3 | 5709.2 | 63.0 | 62.2-63.7 | 6017.2 | 35.6 | 35.2-36.0 | 3010.4 | 40.4 | 39.7-41.0 | 3006.8 | 32.1 | 31.5-32.6 |
| Melanoma of the skin | 4030.6 | 26.9 | 26.5-27.3 | 2355.8 | 33.1 | 32.5-33.8 | 1674.8 | 22.4 | 21.8-22.9 | 360.6 | 2.2 | 2.1-2.3 | 226.8 | 3.2 | 3.0-3.3 | 133.8 | 1.5 | 1.4-1.6 |
| Female breast | | | | | | | 11468.6 | 148.0 | 146.7-149.3 | | | | | | | 1684.8 | 18.7 | 18.3-19.1 |
| Cervix uteri | | | | | | | 410.4 | 6.6 | 6.3-6.9 | | | | | | | 127.0 | 1.7 | 1.5-1.8 |
| Corpus uterus and NOS | | | | | | | 2792.6 | 33.3 | 32.7-33.8 | | | | | | | 484.4 | 5.2 | 5.0-5.5 |
| Ovary | | | | | | | 984.8 | 12.4 | 12.1-12.8 | | | | | | | 633.2 | 7.0 | 6.8-7.3 |
| Prostate | | | | 9995.8 | 125.4 | 124.3-126.5 | | | | | | | 1125.4 | 15.5 | 15.1-15.9 | | | |
| Testis | | | | 439.6 | 8.6 | 8.2-9.0 | | | | | | | 14.2 | 0.2 | 0.2-0.3 | | | |
| Urinary bladder (incl. in situ) | 4547.2 | 27.3 | 26.9-27.6 | 3421.6 | 46.4 | 45.7-47.1 | 1125.6 | 12.2 | 11.9-12.6 | 801.6 | 4.5 | 4.4-4.7 | 568.0 | 7.8 | 7.5-8.1 | 233.6 | 2.3 | 2.1-2.4 |
| Kidney and renal pelvis | 2896.6 | 18.9 | 18.5-19.2 | 1894.0 | 26.3 | 25.7-26.8 | 1002.6 | 12.3 | 12.0-12.7 | 516.4 | 3.1 | 2.9-3.2 | 330.4 | 4.4 | 4.2-4.7 | 186.0 | 2.0 | 1.8-2.1 |
| Brain and other central nervous system | 1029.8 | 7.6 | 7.4-7.9 | 570.4 | 8.8 | 8.5-9.2 | 459.4 | 6.6 | 6.3-6.9 | 686.4 | 4.5 | 4.4-4.7 | 385.2 | 5.4 | 5.2-5.7 | 301.2 | 3.7 | 3.5-3.9 |
| Thyroid | 2557.8 | 21.1 | 20.7-21.5 | 740.0 | 11.8 | 11.5-12.3 | 1817.8 | 30.2 | 29.6-30.9 | 87.6 | 0.5 | 0.5-0.6 | 40.0 | 0.5 | 0.5-0.6 | 47.6 | 0.5 | 0.4-0.5 |
| Hodgkin lymphoma | 408.0 | 3.5 | 3.4-3.7 | 222.4 | 3.9 | 3.6-4.1 | 185.6 | 3.2 | 3.0-3.4 | 46.4 | 0.3 | 0.3-0.3 | 28.2 | 0.4 | 0.3-0.5 | 18.2 | 0.2 | 0.2-0.3 |
| Non-Hodgkin lymphomas | 3663.6 | 23.5 | 23.2-23.9 | 1985.4 | 28.0 | 27.4-28.6 | 1678.2 | 19.9 | 19.5-20.4 | 888.8 | 5.2 | 5.1-5.4 | 495.6 | 6.9 | 6.6-7.1 | 393.2 | 4.0 | 3.8-4.2 |
| Myeloma | 1170.2 | 7.2 | 7.0-7.4 | 670.0 | 9.1 | 8.7-9.4 | 500.2 | 5.6 | 5.4-5.9 | 446.4 | 2.6 | 2.5-2.7 | 244.4 | 3.3 | 3.1-3.5 | 202.0 | 2.0 | 1.9-2.2 |
| Leukemias | 2808.8 | 18.1 | 17.8-18.5 | 1657.8 | 23.5 | 23.0-24.0 | 1151.0 | 13.8 | 13.4-14.2 | 1021.2 | 6.1 | 5.9-6.3 | 582.0 | 8.2 | 7.9-8.5 | 439.2 | 4.6 | 4.4-4.8 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York State, Non-Hispanic Blacks

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 14984.4 | 464.6 | 461.2-468.0 | 7343.8 | 547.9 | 542.1-553.8 | 7640.6 | 413.9 | 409.7-418.1 | 4741.4 | 150.7 | 148.8-152.7 | 2196.0 | 180.9 | 177.4-184.5 | 2545.4 | 134.1 | 131.7-136.4 |
| Oral cavity and pharynx | 258.2 | 7.9 | 7.5-8.4 | 162.4 | 11.9 | 11.1-12.8 | 95.8 | 5.1 | 4.7-5.6 | 65.4 | 2.0 | 1.8-2.3 | 45.2 | 3.5 | 3.0-4.0 | 20.2 | 1.1 | 0.9-1.3 |
| Esophagus | 115.0 | 3.6 | 3.3-3.9 | 72.4 | 5.7 | 5.1-6.3 | 42.6 | 2.2 | 1.9-2.6 | 75.2 | 2.3 | 2.1-2.6 | 50.2 | 4.1 | 3.6-4.7 | 25.0 | 1.3 | 1.1-1.5 |
| Stomach | 349.2 | 11.3 | 10.7-11.8 | 184.0 | 14.9 | 14.0-16.0 | 165.2 | 9.0 | 8.4-9.6 | 153.4 | 5.0 | 4.6-5.3 | 90.8 | 7.6 | 6.9-8.4 | 62.6 | 3.3 | 2.9-3.7 |
| Colorectal | 1275.0 | 40.2 | 39.2-41.2 | 616.6 | 47.9 | 46.2-49.7 | 658.4 | 35.2 | 34.0-36.5 | 460.8 | 14.8 | 14.2-15.4 | 214.4 | 17.6 | 16.5-18.7 | 246.4 | 12.9 | 12.2-13.7 |
| Colon excluding rectum | 928.6 | 29.4 | 28.5-30.2 | 424.8 | 33.4 | 31.9-34.9 | 503.8 | 26.8 | 25.8-27.9 | 380.4 | 12.2 | 11.6-12.8 | 172.8 | 14.2 | 13.2-15.2 | 207.6 | 10.9 | 10.2-11.6 |
| Rectum & rectosigmoid | 346.4 | 10.8 | 10.3-11.4 | 191.8 | 14.6 | 13.6-15.6 | 154.6 | 8.4 | 7.8-9.0 | 80.4 | 2.6 | 2.3-2.9 | 41.6 | 3.4 | 3.0-3.9 | 38.8 | 2.0 | 1.8-2.4 |
| Liver / intrahepatic bile duct | 355.8 | 10.4 | 9.9-10.9 | 252.2 | 17.7 | 16.7-18.8 | 103.6 | 5.2 | 4.7-5.7 | 237.4 | 7.1 | 6.7-7.6 | 159.6 | 11.7 | 10.9-12.6 | 77.8 | 4.0 | 3.6-4.4 |
| Pancreas | 502.4 | 15.8 | 15.2-16.5 | 223.4 | 17.3 | 16.3-18.4 | 279.0 | 14.7 | 13.9-15.5 | 377.8 | 12.0 | 11.4-12.5 | 169.2 | 13.2 | 12.3-14.2 | 208.6 | 11.0 | 10.3-11.7 |
| Larynx | 105.2 | 3.1 | 2.9-3.4 | 80.2 | 6.0 | 5.4-6.7 | 25.0 | 1.3 | 1.0-1.5 | 36.4 | 1.1 | 1.0-1.3 | 27.2 | 2.2 | 1.8-2.6 | 9.2 | 0.5 | 0.3-0.6 |
| Lung and bronchus | 1559.8 | 48.2 | 47.1-49.3 | 774.2 | 60.0 | 58.1-62.0 | 785.6 | 40.7 | 39.4-42.0 | 896.2 | 28.0 | 27.2-28.9 | 471.6 | 37.9 | 36.3-39.5 | 424.6 | 21.9 | 21.0-22.9 |
| Melanoma of the skin | 29.0 | 0.9 | 0.8-1.1 | 11.2 | 0.9 | 0.7-1.2 | 17.8 | 1.0 | 0.8-1.2 | 6.2 | 0.2 | 0.1-0.3 | 3.4 | 0.3 | 0.1-0.4 | 2.8 | 0.1 | 0.1-0.2 |
| Female breast | | | | | | | 2344.2 | 129.4 | 127.0-131.8 | | | | | | | 462.6 | 24.9 | 23.8-25.9 |
| Cervix uteri | | | | | | | 173.0 | 10.0 | 9.3-10.7 | | | | | | | 57.0 | 3.1 | 2.8-3.5 |
| Corpus uteri and NOS | | | | | | | 692.8 | 35.6 | 34.4-36.8 | | | | | | | 217.2 | 11.1 | 10.5-11.8 |
| Ovary | | | | | | | 173.8 | 9.5 | 8.8-10.1 | | | | | | | 120.8 | 6.3 | 5.8-6.9 |
| Prostate | | | | 2933.6 | 208.2 | 204.7-211.7 | | | | | | | 348.4 | 32.7 | 31.1-34.3 | | | |
| Testis | | | | 27.4 | 1.9 | 1.6-2.3 | | | | | | | 1.4 | 0.1 | 0.1-0.3 | | | |
| Urinary bladder (incl. in situ) | 353.2 | 11.3 | 10.7-11.8 | 236.4 | 19.5 | 18.4-20.7 | 116.8 | 6.2 | 5.7-6.7 | 84.2 | 2.8 | 2.5-3.1 | 49.0 | 4.5 | 3.9-5.1 | 35.2 | 1.8 | 1.6-2.1 |
| Kidney and renal pelvis | 546.4 | 16.8 | 16.1-17.4 | 337.4 | 24.7 | 23.5-26.0 | 209.0 | 11.1 | 10.5-11.9 | 80.2 | 2.5 | 2.3-2.8 | 52.0 | 4.1 | 3.6-4.6 | 28.2 | 1.5 | 1.2-1.7 |
| Brain and other central nervous system | 127.6 | 4.2 | 3.9-4.6 | 69.2 | 5.2 | 4.6-5.7 | 58.4 | 3.5 | 3.1-3.9 | 70.0 | 2.2 | 2.0-2.5 | 36.8 | 2.8 | 2.4-3.2 | 33.2 | 1.9 | 1.6-2.2 |
| Thyroid | 323.8 | 10.6 | 10.0-11.1 | 62.0 | 4.6 | 4.1-5.2 | 261.8 | 15.4 | 14.6-16.3 | 16.8 | 0.5 | 0.4-0.7 | 5.4 | 0.4 | 0.3-0.6 | 11.4 | 0.6 | 0.4-0.8 |
| Hodgkin lymphoma | 93.0 | 3.1 | 2.8-3.4 | 46.2 | 3.3 | 2.9-3.8 | 46.8 | 2.9 | 2.5-3.3 | 7.6 | 0.3 | 0.2-0.4 | 4.2 | 0.3 | 0.2-0.5 | 3.4 | 0.2 | 0.1-0.3 |
| Non-Hodgkin lymphomas | 511.2 | 16.3 | 15.7-17.0 | 258.2 | 19.6 | 18.5-20.8 | 253.0 | 14.0 | 13.3-14.8 | 126.6 | 4.1 | 3.8-4.4 | 67.2 | 5.4 | 4.9-6.1 | 59.4 | 3.2 | 2.8-3.6 |
| Myeloma | 475.4 | 15.0 | 14.4-15.6 | 223.8 | 17.5 | 16.4-18.6 | 251.6 | 13.4 | 12.6-14.1 | 153.6 | 4.9 | 4.6-5.3 | 74.2 | 6.2 | 5.6-6.9 | 79.4 | 4.2 | 3.8-4.6 |
| Leukemias | 337.6 | 10.9 | 10.4-11.5 | 168.0 | 13.2 | 12.3-14.1 | 169.6 | 9.5 | 8.9-10.2 | 148.6 | 4.8 | 4.5-5.2 | 75.2 | 6.1 | 5.5-6.8 | 73.4 | 4.0 | 3.6-4.5 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York City, Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-----------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 8312.4 | 361.6 | 358.1-365.1 | 3961.8 | 409.1 | 403.2-415.0 | 4350.6 | 335.7 | 331.2-340.2 | 2341.0 | 107.2 | 105.2-109.1 | 1172.2 | 134.2 | 130.7-137.8 | 1168.8 | 90.3 | 88.0-92.7 |
| Oral cavity and pharynx | 193.4 | 8.3 | 7.8-8.9 | 130.8 | 13.1 | 12.1-14.2 | 62.6 | 4.8 | 4.3-5.4 | 40.8 | 1.8 | 1.6-2.1 | 30.4 | 3.2 | 2.7-3.8 | 10.4 | 0.8 | 0.6-1.0 |
| Esophagus | 65.4 | 2.9 | 2.6-3.2 | 48.2 | 5.1 | 4.4-5.8 | 17.2 | 1.3 | 1.1-1.6 | 44.6 | 2.0 | 1.8-2.3 | 34.0 | 3.7 | 3.2-4.4 | 10.6 | 0.8 | 0.6-1.1 |
| Stomach | 235.0 | 10.5 | 9.9-11.2 | 125.8 | 13.5 | 12.4-14.6 | 109.2 | 8.6 | 7.9-9.3 | 98.4 | 4.5 | 4.1-4.9 | 57.0 | 6.5 | 5.7-7.3 | 41.4 | 3.2 | 2.8-3.7 |
| Colorectal | 721.2 | 31.9 | 30.9-33.0 | 386.6 | 40.7 | 38.8-42.6 | 334.6 | 25.9 | 24.6-27.2 | 221.0 | 10.2 | 9.6-10.8 | 119.0 | 13.4 | 12.3-14.5 | 102.0 | 7.9 | 7.2-8.6 |
| Colon excluding rectum | 499.0 | 22.3 | 21.4-23.2 | 256.8 | 27.4 | 25.9-29.0 | 242.2 | 18.8 | 17.8-19.9 | 178.6 | 8.2 | 7.7-8.8 | 93.4 | 10.6 | 9.6-11.6 | 85.2 | 6.6 | 6.0-7.3 |
| Rectum & rectosigmoid | 222.2 | 9.6 | 9.1-10.2 | 129.8 | 13.3 | 12.2-14.4 | 92.4 | 7.1 | 6.4-7.8 | 42.4 | 1.9 | 1.7-2.2 | 25.6 | 2.8 | 2.3-3.4 | 16.8 | 1.3 | 1.1-1.6 |
| Liver / intrahepatic bile duct | 316.0 | 13.5 | 12.8-14.2 | 217.6 | 21.7 | 20.4-23.1 | 98.4 | 7.4 | 6.8-8.1 | 181.2 | 8.0 | 7.5-8.6 | 118.8 | 12.5 | 11.5-13.6 | 62.4 | 4.7 | 4.2-5.3 |
| Pancreas | 258.2 | 11.6 | 11.0-12.3 | 127.0 | 13.8 | 12.7-14.9 | 131.2 | 10.2 | 9.4-11.0 | 178.0 | 8.2 | 7.7-8.8 | 85.4 | 9.7 | 8.8-10.7 | 92.6 | 7.2 | 6.5-7.9 |
| Larynx | 65.8 | 2.8 | 2.5-3.1 | 48.2 | 5.0 | 4.4-5.7 | 17.6 | 1.3 | 1.1-1.6 | 21.4 | 0.9 | 0.8-1.1 | 18.4 | 2.0 | 1.6-2.5 | 3.0 | 0.2 | 0.1-0.4 |
| Lung and bronchus | 697.4 | 31.4 | 30.3-32.5 | 364.2 | 40.6 | 38.7-42.6 | 333.2 | 25.4 | 24.2-26.7 | 355.2 | 16.3 | 15.6-17.1 | 196.8 | 22.8 | 21.3-24.3 | 158.4 | 12.2 | 11.3-13.0 |
| Melanoma of the skin | 53.8 | 2.4 | 2.1-2.7 | 25.8 | 2.8 | 2.3-3.3 | 28.0 | 2.2 | 1.8-2.6 | 8.2 | 0.4 | 0.3-0.5 | 5.4 | 0.6 | 0.4-0.9 | 2.8 | 0.2 | 0.1-0.4 |
| Female breast | | | | | | | 1263.2 | 97.5 | 95.0-99.9 | | | | | | | 180.4 | 14.0 | 13.1-14.9 |
| Cervix uteri | | | | | | | 116.8 | 9.3 | 8.5-10.0 | | | | | | | 28.8 | 2.2 | 1.9-2.6 |
| Corpus uterus and NOS | | | | | | | 325.2 | 24.5 | 23.3-25.7 | | | | | | | 65.6 | 5.0 | 4.4-5.5 |
| Ovary | | | | | | | 115.6 | 9.0 | 8.3-9.8 | | | | | | | 56.8 | 4.4 | 3.9-4.9 |
| Prostate | | | | 1088.4 | 110.4 | 107.4-113.5 | | | | | | | 131.2 | 17.3 | 16.0-18.7 | | | |
| Testis | | | | 52.8 | 4.0 | 3.5-4.6 | | | | | | | 2.6 | 0.2 | 0.1-0.4 | | | |
| Urinary bladder (incl. in situ) | 227.8 | 10.5 | 9.9-11.1 | 158.8 | 18.5 | 17.2-19.9 | 69.0 | 5.4 | 4.8-6.0 | 48.2 | 2.4 | 2.1-2.7 | 30.6 | 4.1 | 3.5-4.8 | 17.6 | 1.4 | 1.1-1.7 |
| Kidney and renal pelvis | 288.8 | 12.4 | 11.8-13.1 | 172.8 | 17.3 | 16.1-18.5 | 116.0 | 8.9 | 8.1-9.6 | 41.2 | 1.9 | 1.6-2.2 | 25.6 | 2.8 | 2.4-3.4 | 15.6 | 1.2 | 0.9-1.5 |
| Brain and other central nervous system | 119.2 | 5.1 | 4.7-5.5 | 65.2 | 6.2 | 5.5-6.9 | 54.0 | 4.2 | 3.7-4.7 | 60.2 | 2.7 | 2.4-3.0 | 32.0 | 3.4 | 2.8-3.9 | 28.2 | 2.2 | 1.8-2.6 |
| Thyroid | 371.0 | 15.5 | 14.8-16.2 | 72.8 | 6.7 | 6.0-7.4 | 298.2 | 23.5 | 22.3-24.7 | 10.0 | 0.5 | 0.4-0.6 | 3.0 | 0.4 | 0.2-0.6 | 7.0 | 0.6 | 0.4-0.8 |
| Hodgkin lymphoma | 64.4 | 2.7 | 2.4-3.0 | 36.2 | 3.2 | 2.8-3.7 | 28.2 | 2.2 | 1.8-2.6 | 6.8 | 0.3 | 0.2-0.4 | 3.8 | 0.4 | 0.2-0.6 | 3.0 | 0.2 | 0.1-0.4 |
| Non-Hodgkin lymphomas | 408.8 | 17.9 | 17.2-18.7 | 210.4 | 21.5 | 20.2-22.9 | 198.4 | 15.3 | 14.4-16.3 | 90.8 | 4.2 | 3.8-4.6 | 52.0 | 6.0 | 5.3-6.8 | 38.8 | 3.0 | 2.6-3.5 |
| Myeloma | 191.4 | 8.5 | 7.9-9.1 | 98.8 | 10.6 | 9.6-11.6 | 92.6 | 7.1 | 6.5-7.8 | 56.8 | 2.6 | 2.3-3.0 | 28.8 | 3.3 | 2.8-3.9 | 28.0 | 2.2 | 1.8-2.6 |
| Leukemias | 260.2 | 11.2 | 10.6-11.9 | 150.2 | 14.9 | 13.8-16.1 | 110.0 | 8.5 | 7.8-9.3 | 99.2 | 4.5 | 4.1-4.9 | 50.4 | 5.5 | 4.8-6.2 | 48.8 | 3.8 | 3.3-4.3 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York City, Non-Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 33841.4 | 470.9 | 468.6-473.2 | 16295.8 | 508.0 | 504.4-511.6 | 17545.6 | 449.4 | 446.3-452.4 | 9741.2 | 130.8 | 129.7-132.0 | 4712.0 | 151.0 | 149.1-153.0 | 5029.2 | 117.7 | 116.2-119.2 |
| Oral cavity and pharynx | 753.0 | 10.5 | 10.2-10.9 | 488.8 | 15.2 | 14.6-15.8 | 264.2 | 6.7 | 6.3-7.1 | 156.6 | 2.1 | 2.0-2.3 | 108.0 | 3.4 | 3.1-3.7 | 48.6 | 1.1 | 1.0-1.3 |
| Esophagus | 256.0 | 3.4 | 3.2-3.6 | 181.8 | 5.6 | 5.2-6.0 | 74.2 | 1.7 | 1.6-1.9 | 175.0 | 2.3 | 2.2-2.5 | 126.0 | 3.9 | 3.6-4.2 | 49.0 | 1.1 | 1.0-1.3 |
| Stomach | 755.0 | 10.4 | 10.1-10.7 | 438.0 | 13.8 | 13.2-14.4 | 317.0 | 7.8 | 7.4-8.2 | 327.2 | 4.4 | 4.2-4.7 | 194.6 | 6.3 | 5.9-6.7 | 132.6 | 3.1 | 2.8-3.3 |
| Colorectal | 2792.2 | 38.8 | 38.1-39.4 | 1424.2 | 45.2 | 44.1-46.2 | 1368.0 | 33.9 | 33.0-34.7 | 923.0 | 12.4 | 12.0-12.8 | 446.6 | 14.4 | 13.8-15.0 | 476.4 | 10.9 | 10.5-11.4 |
| Colon excluding rectum | 1964.0 | 27.1 | 26.6-27.7 | 965.4 | 30.7 | 29.8-31.6 | 998.6 | 24.4 | 23.7-25.1 | 755.0 | 10.1 | 9.8-10.4 | 355.2 | 11.4 | 10.9-12.0 | 399.8 | 9.1 | 8.7-9.6 |
| Rectum & rectosigmoid | 828.2 | 11.7 | 11.3-12.0 | 458.8 | 14.4 | 13.8-15.1 | 369.4 | 9.5 | 9.0-9.9 | 168.0 | 2.3 | 2.1-2.5 | 91.4 | 3.0 | 2.7-3.3 | 76.6 | 1.8 | 1.6-2.0 |
| Liver / intrahepatic bile duct | 761.6 | 10.2 | 9.9-10.5 | 537.2 | 16.2 | 15.6-16.8 | 224.4 | 5.4 | 5.0-5.7 | 466.8 | 6.2 | 6.0-6.5 | 310.4 | 9.5 | 9.0-10.0 | 156.4 | 3.6 | 3.4-3.9 |
| Pancreas | 1044.2 | 14.0 | 13.7-14.4 | 499.0 | 15.7 | 15.1-16.3 | 545.2 | 12.7 | 12.2-13.2 | 790.4 | 10.5 | 10.2-10.8 | 375.8 | 11.8 | 11.3-12.4 | 414.6 | 9.4 | 9.0-9.8 |
| Larynx | 197.8 | 2.7 | 2.5-2.8 | 159.6 | 4.9 | 4.6-5.3 | 38.2 | 0.9 | 0.8-1.1 | 57.6 | 0.8 | 0.7-0.9 | 45.0 | 1.4 | 1.2-1.6 | 12.6 | 0.3 | 0.2-0.4 |
| Lung and bronchus | 3732.0 | 50.3 | 49.6-51.0 | 1848.4 | 58.0 | 56.8-59.3 | 1883.6 | 44.8 | 43.9-45.7 | 1962.4 | 26.1 | 25.6-26.7 | 1028.4 | 32.6 | 31.7-33.5 | 934.0 | 21.5 | 20.8-22.1 |
| Melanoma of the skin | 884.4 | 12.4 | 12.1-12.8 | 494.4 | 16.0 | 15.3-16.6 | 390.0 | 10.0 | 9.5-10.5 | 77.4 | 1.1 | 1.0-1.2 | 45.8 | 1.5 | 1.3-1.7 | 31.6 | 0.8 | 0.6-0.9 |
| Female breast | | | | | | | 5201.6 | 138.1 | 136.4-139.9 | | | | | | | 825.4 | 20.0 | 19.4-20.7 |
| Cervix uteri | | | | | | | 315.4 | 9.0 | 8.5-9.4 | | | | | | | 85.0 | 2.2 | 2.0-2.5 |
| Corpus uterus and NOS | | | | | | | 1364.6 | 33.8 | 33.0-34.6 | | | | | | | 320.0 | 7.5 | 7.1-7.9 |
| Ovary | | | | | | | 460.2 | 12.1 | 11.6-12.7 | | | | | | | 281.2 | 6.8 | 6.4-7.2 |
| Prostate | | | | 4463.0 | 131.8 | 130.1-133.6 | | | | | | | 554.4 | 18.3 | 17.6-19.0 | | | |
| Testis | | | | 167.6 | 5.6 | 5.2-6.0 | | | | | | | 4.0 | 0.1 | 0.1-0.2 | | | |
| Urinary bladder (incl. in situ) | 1365.4 | 18.4 | 18.0-18.9 | 1017.4 | 32.7 | 31.8-33.6 | 348.0 | 8.1 | 7.7-8.5 | 262.4 | 3.4 | 3.2-3.6 | 180.2 | 6.0 | 5.6-6.4 | 82.2 | 1.8 | 1.6-1.9 |
| Kidney and renal pelvis | 1115.2 | 15.6 | 15.2-16.0 | 719.6 | 22.4 | 21.7-23.2 | 395.6 | 10.0 | 9.6-10.5 | 182.8 | 2.4 | 2.3-2.6 | 123.0 | 3.9 | 3.6-4.2 | 59.8 | 1.3 | 1.2-1.5 |
| Brain and other central nervous system | 411.2 | 6.2 | 6.0-6.5 | 226.4 | 7.5 | 7.0-7.9 | 184.8 | 5.2 | 4.9-5.6 | 223.6 | 3.2 | 3.0-3.4 | 121.4 | 3.9 | 3.6-4.3 | 102.2 | 2.6 | 2.4-2.9 |
| Thyroid | 1446.8 | 22.3 | 21.8-22.8 | 364.8 | 11.9 | 11.3-12.5 | 1082.0 | 31.7 | 30.9-32.6 | 39.0 | 0.5 | 0.4-0.6 | 15.4 | 0.5 | 0.4-0.6 | 23.6 | 0.5 | 0.4-0.6 |
| Hodgkin lymphoma | 196.2 | 3.1 | 2.9-3.3 | 105.6 | 3.6 | 3.3-3.9 | 90.6 | 2.7 | 2.5-3.0 | 16.8 | 0.2 | 0.2-0.3 | 9.6 | 0.3 | 0.2-0.4 | 7.2 | 0.2 | 0.1-0.3 |
| Non-Hodgkin lymphomas | 1462.0 | 20.7 | 20.2-21.2 | 778.6 | 25.1 | 24.3-25.9 | 683.4 | 17.3 | 16.7-17.9 | 321.0 | 4.3 | 4.1-4.6 | 176.8 | 5.8 | 5.4-6.2 | 144.2 | 3.3 | 3.1-3.6 |
| Myeloma | 668.2 | 9.1 | 8.8-9.4 | 341.2 | 10.7 | 10.2-11.2 | 327.0 | 7.9 | 7.5-8.3 | 217.2 | 2.9 | 2.7-3.1 | 111.4 | 3.5 | 3.2-3.9 | 105.8 | 2.4 | 2.2-2.6 |
| Leukemias | 1009.6 | 14.4 | 14.0-14.8 | 560.6 | 18.3 | 17.6-19.0 | 449.0 | 11.4 | 10.9-11.9 | 377.2 | 5.2 | 5.0-5.4 | 201.4 | 6.7 | 6.3-7.1 | 175.8 | 4.2 | 3.9-4.5 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York City, Non-Hispanic Whites

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 18110.8 | 501.7 | 498.3-505.1 | 8791.2 | 522.4 | 517.4-527.4 | 9319.6 | 495.7 | 490.9-500.5 | 5112.4 | 127.8 | 126.2-129.5 | 2496.8 | 144.4 | 141.9-147.0 | 2615.6 | 116.7 | 114.5-118.8 |
| Oral cavity and pharynx | 400.0 | 11.1 | 10.6-11.6 | 268.6 | 16.0 | 15.2-16.9 | 131.4 | 6.7 | 6.2-7.2 | 77.6 | 2.0 | 1.8-2.2 | 52.6 | 3.1 | 2.7-3.5 | 25.0 | 1.1 | 0.9-1.3 |
| Esophagus | 141.2 | 3.7 | 3.4-4.0 | 103.2 | 6.0 | 5.5-6.6 | 38.0 | 1.7 | 1.5-2.0 | 99.8 | 2.6 | 2.3-2.8 | 73.0 | 4.2 | 3.8-4.7 | 26.8 | 1.1 | 1.0-1.4 |
| Stomach | 284.6 | 7.7 | 7.3-8.1 | 164.0 | 9.8 | 9.1-10.5 | 120.6 | 6.0 | 5.5-6.5 | 128.4 | 3.3 | 3.0-3.6 | 76.4 | 4.5 | 4.0-5.0 | 52.0 | 2.3 | 2.0-2.6 |
| Colorectal | 1419.8 | 38.8 | 37.9-39.8 | 734.4 | 44.3 | 42.9-45.8 | 685.4 | 34.3 | 33.1-35.5 | 475.2 | 11.7 | 11.2-12.2 | 234.6 | 13.6 | 12.8-14.4 | 240.6 | 10.3 | 9.7-10.9 |
| Colon excluding rectum | 1013.8 | 27.2 | 26.4-28.0 | 512.8 | 30.7 | 29.5-32.0 | 501.0 | 24.3 | 23.3-25.4 | 387.8 | 9.4 | 9.0-9.9 | 188.8 | 10.8 | 10.2-11.6 | 199.0 | 8.4 | 7.9-9.0 |
| Rectum & rectosigmoid | 406.0 | 11.6 | 11.1-12.2 | 221.6 | 13.6 | 12.8-14.4 | 184.4 | 9.9 | 9.3-10.6 | 87.4 | 2.3 | 2.1-2.5 | 45.8 | 2.8 | 2.4-3.2 | 41.6 | 1.9 | 1.6-2.2 |
| Liver / intrahepatic bile duct | 293.8 | 7.7 | 7.3-8.1 | 205.2 | 11.8 | 11.1-12.6 | 88.6 | 4.2 | 3.8-4.6 | 190.6 | 4.9 | 4.5-5.2 | 122.8 | 7.1 | 6.5-7.7 | 67.8 | 3.0 | 2.7-3.4 |
| Pancreas | 565.4 | 14.4 | 13.9-15.0 | 279.2 | 16.3 | 15.4-17.2 | 286.2 | 12.8 | 12.1-13.5 | 437.6 | 10.8 | 10.3-11.3 | 219.0 | 12.6 | 11.9-13.4 | 218.6 | 9.3 | 8.7-9.9 |
| Larynx | 102.4 | 2.8 | 2.5-3.0 | 85.0 | 5.0 | 4.5-5.5 | 17.4 | 0.9 | 0.7-1.1 | 29.4 | 0.8 | 0.6-0.9 | 24.0 | 1.4 | 1.2-1.7 | 5.4 | 0.2 | 0.2-0.4 |
| Lung and bronchus | 2059.0 | 53.1 | 52.1-54.2 | 980.6 | 57.0 | 55.4-58.6 | 1078.4 | 50.5 | 49.1-51.9 | 1086.6 | 27.0 | 26.3-27.7 | 543.0 | 31.3 | 30.1-32.5 | 543.6 | 23.7 | 22.8-24.7 |
| Melanoma of the skin | 815.8 | 22.9 | 22.2-23.7 | 463.2 | 28.1 | 26.9-29.3 | 352.6 | 19.1 | 18.2-20.1 | 66.8 | 1.8 | 1.6-2.0 | 39.8 | 2.4 | 2.1-2.7 | 27.0 | 1.3 | 1.1-1.5 |
| Female breast | | | | | | | 2710.4 | 153.9 | 151.1-156.6 | | | | | | | 407.4 | 19.0 | 18.2-20.0 |
| Cervix uteri | | | | | | | 110.2 | 7.1 | 6.5-7.7 | | | | | | | 28.4 | 1.6 | 1.4-2.0 |
| Corpus uteri and NOS | | | | | | | 645.2 | 34.2 | 33.0-35.5 | | | | | | | 123.8 | 5.7 | 5.2-6.2 |
| Ovary | | | | | | | 241.6 | 13.3 | 12.5-14.1 | | | | | | | 155.6 | 7.5 | 6.9-8.0 |
| Prostate | | | | 1971.2 | 111.8 | 109.6-114.1 | | | | | | | 252.0 | 14.2 | 13.4-15.0 | | | |
| Testis | | | | 127.4 | 8.8 | 8.1-9.6 | | | | | | | 2.0 | 0.1 | 0.1-0.2 | | | |
| Urinary bladder (incl. in situ) | 987.2 | 25.3 | 24.6-26.0 | 747.4 | 43.5 | 42.1-45.0 | 239.8 | 11.1 | 10.4-11.8 | 178.8 | 4.1 | 3.8-4.4 | 127.2 | 7.1 | 6.6-7.7 | 51.6 | 2.0 | 1.7-2.3 |
| Kidney and renal pelvis | 614.8 | 17.5 | 16.9-18.1 | 410.2 | 24.9 | 23.8-26.0 | 204.6 | 10.9 | 10.2-11.6 | 106.4 | 2.7 | 2.4-2.9 | 71.4 | 4.1 | 3.7-4.6 | 35.0 | 1.5 | 1.3-1.8 |
| Brain and other central nervous system | 256.4 | 8.1 | 7.6-8.6 | 141.0 | 9.3 | 8.6-10.0 | 115.4 | 7.0 | 6.4-7.7 | 145.2 | 4.1 | 3.8-4.4 | 78.8 | 4.8 | 4.3-5.3 | 66.4 | 3.5 | 3.1-3.9 |
| Thyroid | 780.0 | 25.8 | 25.0-26.7 | 214.8 | 14.2 | 13.3-15.1 | 565.2 | 37.5 | 36.1-39.0 | 21.0 | 0.5 | 0.4-0.6 | 8.2 | 0.5 | 0.3-0.7 | 12.8 | 0.5 | 0.4-0.7 |
| Hodgkin lymphoma | 110.4 | 3.8 | 3.5-4.1 | 61.4 | 4.3 | 3.8-4.9 | 49.0 | 3.2 | 2.8-3.7 | 10.6 | 0.3 | 0.2-0.4 | 7.0 | 0.4 | 0.3-0.6 | 3.6 | 0.2 | 0.1-0.3 |
| Non-Hodgkin lymphomas | 893.4 | 24.8 | 24.1-25.6 | 482.6 | 29.5 | 28.3-30.8 | 410.8 | 20.9 | 19.9-21.9 | 190.6 | 4.7 | 4.4-5.0 | 106.0 | 6.2 | 5.7-6.8 | 84.6 | 3.5 | 3.1-3.9 |
| Myeloma | 275.4 | 7.2 | 6.8-7.6 | 156.0 | 9.1 | 8.5-9.8 | 119.4 | 5.6 | 5.2-6.1 | 91.4 | 2.3 | 2.1-2.5 | 50.6 | 2.9 | 2.6-3.3 | 40.8 | 1.8 | 1.5-2.1 |
| Leukemias | 635.0 | 17.6 | 16.9-18.2 | 365.4 | 22.3 | 21.3-23.4 | 269.6 | 13.6 | 12.8-14.4 | 227.2 | 5.7 | 5.3-6.0 | 121.2 | 7.1 | 6.6-7.7 | 106.0 | 4.7 | 4.2-5.1 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York City, Non-Hispanic Blacks

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 9992.6 | 451.3 | 447.3-455.4 | 4758.4 | 530.0 | 523.0-537.0 | 5234.2 | 405.4 | 400.3-410.4 | 3222.8 | 147.5 | 145.2-149.8 | 1442.6 | 176.0 | 171.8-180.3 | 1780.2 | 132.6 | 129.8-135.5 |
| Oral cavity and pharynx | 172.4 | 7.8 | 7.3-8.3 | 104.0 | 11.6 | 10.6-12.7 | 68.4 | 5.2 | 4.7-5.8 | 44.6 | 2.0 | 1.8-2.3 | 29.6 | 3.4 | 2.9-4.0 | 15.0 | 1.2 | 0.9-1.5 |
| Esophagus | 73.2 | 3.2 | 2.9-3.6 | 45.6 | 5.2 | 4.5-5.9 | 27.6 | 2.1 | 1.7-2.5 | 47.0 | 2.1 | 1.8-2.4 | 29.8 | 3.5 | 3.0-4.2 | 17.2 | 1.2 | 1.0-1.5 |
| Stomach | 233.6 | 10.9 | 10.2-11.5 | 123.0 | 14.9 | 13.7-16.2 | 110.6 | 8.5 | 7.8-9.2 | 106.6 | 5.0 | 4.6-5.4 | 62.0 | 7.8 | 6.9-8.7 | 44.6 | 3.3 | 2.9-3.8 |
| Colorectal | 858.8 | 39.4 | 38.2-40.6 | 405.2 | 47.2 | 45.1-49.4 | 453.6 | 34.5 | 33.1-36.0 | 320.4 | 14.8 | 14.1-15.6 | 142.2 | 17.5 | 16.2-18.9 | 178.2 | 13.2 | 12.3-14.1 |
| Colon excluding rectum | 628.4 | 28.8 | 27.8-29.9 | 279.0 | 32.7 | 31.0-34.5 | 349.4 | 26.4 | 25.2-27.7 | 264.2 | 12.2 | 11.5-12.9 | 112.8 | 13.8 | 12.7-15.1 | 151.4 | 11.2 | 10.4-12.0 |
| Rectum & rectosigmoid | 230.4 | 10.6 | 10.0-11.2 | 126.2 | 14.5 | 13.3-15.7 | 104.2 | 8.1 | 7.4-8.9 | 56.2 | 2.6 | 2.3-2.9 | 29.4 | 3.6 | 3.1-4.3 | 26.8 | 2.0 | 1.7-2.4 |
| Liver / intrahepatic bile duct | 237.4 | 10.2 | 9.6-10.8 | 166.0 | 17.7 | 16.5-19.0 | 71.4 | 5.1 | 4.6-5.7 | 154.6 | 6.8 | 6.3-7.3 | 102.2 | 11.3 | 10.3-12.4 | 52.4 | 3.8 | 3.3-4.3 |
| Pancreas | 332.6 | 15.1 | 14.4-15.9 | 146.6 | 16.9 | 15.7-18.2 | 186.0 | 13.8 | 12.9-14.7 | 251.0 | 11.4 | 10.8-12.0 | 109.2 | 12.6 | 11.5-13.8 | 141.8 | 10.4 | 9.7-11.2 |
| Larynx | 70.0 | 3.0 | 2.7-3.4 | 52.2 | 5.9 | 5.2-6.7 | 17.8 | 1.3 | 1.0-1.6 | 24.8 | 1.1 | 0.9-1.3 | 17.8 | 2.1 | 1.7-2.6 | 7.0 | 0.5 | 0.3-0.7 |
| Lung and bronchus | 975.2 | 43.5 | 42.3-44.8 | 467.6 | 53.3 | 51.1-55.6 | 507.6 | 37.3 | 35.9-38.9 | 564.0 | 25.3 | 24.4-26.3 | 288.4 | 33.9 | 32.1-35.7 | 275.6 | 20.2 | 19.1-21.3 |
| Melanoma of the skin | 19.0 | 0.9 | 0.7-1.1 | 6.8 | 0.8 | 0.6-1.2 | 12.2 | 1.0 | 0.7-1.2 | 4.2 | 0.2 | 0.1-0.3 | 2.2 | 0.2 | 0.1-0.5 | 2.0 | 0.1 | 0.1-0.3 |
| Female breast | | | | | | | 1605.2 | 127.2 | 124.4-130.1 | | | | | | | 330.6 | 25.3 | 24.1-26.6 |
| Cervix uteri | | | | | | | 128.2 | 10.7 | 9.9-11.6 | | | | | | | 42.0 | 3.3 | 2.9-3.8 |
| Corpus uterus and NOS | | | | | | | 515.0 | 37.5 | 36.1-39.0 | | | | | | | 167.0 | 12.0 | 11.2-12.9 |
| Ovary | | | | | | | 126.8 | 9.9 | 9.1-10.7 | | | | | | | 91.2 | 6.7 | 6.1-7.4 |
| Prostate | | | | 1929.4 | 203.6 | 199.4-207.8 | | | | | | | 253.0 | 34.5 | 32.6-36.5 | | | |
| Testis | | | | 18.4 | 2.0 | 1.6-2.5 | | | | | | | 1.4 | 0.2 | 0.1-0.4 | | | |
| Urinary bladder (incl. in situ) | 223.4 | 10.3 | 9.7-10.9 | 149.0 | 18.4 | 17.1-19.8 | 74.4 | 5.5 | 5.0-6.1 | 59.4 | 2.8 | 2.5-3.1 | 34.4 | 4.5 | 3.9-5.3 | 25.0 | 1.8 | 1.5-2.2 |
| Kidney and renal pelvis | 337.4 | 15.1 | 14.3-15.8 | 205.8 | 22.7 | 21.3-24.2 | 131.6 | 9.9 | 9.1-10.7 | 52.8 | 2.4 | 2.1-2.7 | 34.2 | 4.1 | 3.4-4.7 | 18.6 | 1.3 | 1.1-1.6 |
| Brain and other central nervous system | 85.4 | 4.2 | 3.8-4.7 | 46.8 | 5.4 | 4.7-6.1 | 38.6 | 3.4 | 2.9-3.9 | 45.6 | 2.2 | 1.9-2.5 | 25.2 | 2.9 | 2.4-3.5 | 20.4 | 1.7 | 1.4-2.0 |
| Thyroid | 209.2 | 10.1 | 9.5-10.8 | 36.8 | 4.1 | 3.6-4.8 | 172.4 | 14.7 | 13.7-15.8 | 11.2 | 0.5 | 0.4-0.7 | 2.6 | 0.3 | 0.2-0.5 | 8.6 | 0.6 | 0.5-0.9 |
| Hodgkin lymphoma | 60.0 | 3.0 | 2.7-3.4 | 29.4 | 3.3 | 2.8-3.9 | 30.6 | 2.8 | 2.4-3.3 | 5.0 | 0.2 | 0.2-0.4 | 2.0 | 0.2 | 0.1-0.4 | 3.0 | 0.3 | 0.1-0.4 |
| Non-Hodgkin lymphomas | 344.6 | 16.2 | 15.5-17.0 | 167.4 | 19.3 | 17.9-20.7 | 177.2 | 14.2 | 13.3-15.2 | 85.6 | 4.0 | 3.7-4.4 | 42.4 | 5.1 | 4.4-5.9 | 43.2 | 3.3 | 2.9-3.8 |
| Myeloma | 319.4 | 14.6 | 13.8-15.3 | 143.4 | 16.6 | 15.4-17.9 | 176.0 | 13.3 | 12.4-14.2 | 104.8 | 4.8 | 4.4-5.2 | 49.2 | 6.0 | 5.3-6.8 | 55.6 | 4.1 | 3.6-4.6 |
| Leukemias | 223.2 | 10.7 | 10.1-11.4 | 106.8 | 12.7 | 11.6-13.9 | 116.4 | 9.4 | 8.7-10.3 | 99.2 | 4.7 | 4.3-5.1 | 49.6 | 6.0 | 5.2-6.8 | 49.6 | 3.9 | 3.4-4.4 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York State Excl New York City, Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-----------|----------------|------------------|-------------|----------------|------------------|-----------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 3411.4 | 386.9 | 380.8-393.2 | 1617.2 | 418.5 | 408.5-428.7 | 1794.2 | 371.7 | 363.8-379.8 | 715.2 | 93.1 | 89.9-96.4 | 365.4 | 111.0 | 105.5-116.6 | 349.8 | 80.9 | 77.0-84.9 |
| Oral cavity and pharynx | 75.0 | 8.5 | 7.6-9.4 | 52.2 | 13.0 | 11.4-14.9 | 22.8 | 4.7 | 3.9-5.7 | 14.8 | 2.0 | 1.6-2.5 | 11.4 | 3.7 | 2.7-4.8 | 3.4 | 0.8 | 0.5-1.3 |
| Esophagus | 22.2 | 3.0 | 2.4-3.6 | 18.0 | 5.5 | 4.3-6.9 | 4.2 | 1.1 | 0.7-1.7 | 12.2 | 1.5 | 1.1-2.0 | 10.2 | 2.9 | 2.1-3.8 | 2.0 | 0.5 | 0.2-0.9 |
| Stomach | 99.4 | 11.8 | 10.7-13.0 | 53.0 | 13.6 | 11.8-15.5 | 46.4 | 10.4 | 9.1-11.9 | 34.4 | 4.2 | 3.6-5.0 | 19.8 | 5.5 | 4.4-6.8 | 14.6 | 3.3 | 2.6-4.2 |
| Colorectal | 283.0 | 32.6 | 30.9-34.5 | 148.0 | 36.7 | 33.8-39.7 | 135.0 | 29.3 | 27.1-31.7 | 59.6 | 7.7 | 6.8-8.7 | 32.2 | 9.5 | 8.0-11.3 | 27.4 | 6.4 | 5.3-7.6 |
| Colon excluding rectum | 193.4 | 23.1 | 21.6-24.7 | 94.8 | 24.6 | 22.2-27.2 | 98.6 | 22.0 | 20.0-24.1 | 46.4 | 6.1 | 5.3-7.0 | 23.6 | 7.3 | 5.9-8.8 | 22.8 | 5.3 | 4.4-6.4 |
| Rectum & rectosigmoid | 89.6 | 9.5 | 8.6-10.5 | 53.2 | 12.1 | 10.5-13.8 | 36.4 | 7.4 | 6.3-8.6 | 13.2 | 1.6 | 1.2-2.0 | 8.6 | 2.3 | 1.6-3.1 | 4.6 | 1.0 | 0.7-1.6 |
| Liver / intrahepatic bile duct | 103.2 | 12.7 | 11.5-13.9 | 70.0 | 18.6 | 16.5-20.9 | 33.2 | 7.8 | 6.6-9.1 | 58.2 | 7.5 | 6.6-8.4 | 38.8 | 11.0 | 9.3-12.7 | 19.4 | 4.7 | 3.8-5.7 |
| Pancreas | 102.6 | 12.8 | 11.7-14.0 | 50.2 | 13.5 | 11.8-15.5 | 52.4 | 12.1 | 10.7-13.8 | 54.8 | 7.3 | 6.4-8.2 | 28.4 | 8.2 | 6.8-9.8 | 26.4 | 6.4 | 5.4-7.7 |
| Larynx | 21.6 | 2.6 | 2.1-3.2 | 19.0 | 5.3 | 4.2-6.6 | 2.6 | 0.5 | 0.3-0.9 | 5.0 | 0.7 | 0.4-1.0 | 3.8 | 1.2 | 0.7-1.9 | 1.2 | 0.3 | 0.1-0.6 |
| Lung and bronchus | 268.8 | 35.6 | 33.6-37.6 | 137.6 | 42.8 | 39.4-46.4 | 131.2 | 30.6 | 28.3-33.1 | 109.8 | 14.8 | 13.6-16.2 | 65.0 | 20.4 | 18.1-23.0 | 44.8 | 10.6 | 9.2-12.1 |
| Melanoma of the skin | 39.0 | 4.6 | 4.0-5.4 | 23.6 | 6.4 | 5.2-7.8 | 15.4 | 3.4 | 2.6-4.2 | 4.0 | 0.4 | 0.3-0.7 | 2.4 | 0.5 | 0.3-1.0 | 1.6 | 0.3 | 0.1-0.7 |
| Female breast | | | | | | | 534.2 | 107.1 | 103.0-111.4 | | | | | | | 54.4 | 11.8 | 10.4-13.3 |
| Cervix uteri | | | | | | | 44.2 | 8.1 | 7.0-9.3 | | | | | | | 8.4 | 1.7 | 1.2-2.3 |
| Corpus uterus and NOS | | | | | | | 111.8 | 22.7 | 20.8-24.8 | | | | | | | 17.4 | 4.0 | 3.2-4.9 |
| Ovary | | | | | | | 48.4 | 9.4 | 8.2-10.7 | | | | | | | 19.4 | 4.3 | 3.4-5.2 |
| Prostate | | | | 405.4 | 107.5 | 102.5-112.7 | | | | | | | 34.8 | 13.7 | 11.7-16.0 | | | |
| Testis | | | | 30.8 | 4.3 | 3.6-5.0 | | | | | | | 1.6 | 0.2 | 0.1-0.5 | | | |
| Urinary bladder (incl. in situ) | 102.4 | 14.1 | 12.8-15.4 | 77.4 | 24.8 | 22.2-27.6 | 25.0 | 6.1 | 5.0-7.3 | 13.4 | 2.1 | 1.6-2.7 | 10.2 | 4.0 | 3.0-5.3 | 3.2 | 0.8 | 0.5-1.4 |
| Kidney and renal pelvis | 121.4 | 13.2 | 12.1-14.3 | 77.2 | 17.8 | 15.9-19.9 | 44.2 | 9.2 | 8.0-10.6 | 12.8 | 1.7 | 1.3-2.2 | 7.6 | 2.2 | 1.5-3.1 | 5.2 | 1.4 | 0.9-2.0 |
| Brain and other central nervous system | 59.2 | 5.5 | 4.8-6.2 | 34.8 | 6.8 | 5.7-8.0 | 24.4 | 4.4 | 3.6-5.3 | 24.2 | 2.6 | 2.1-3.1 | 13.6 | 3.0 | 2.3-4.0 | 10.6 | 2.1 | 1.6-2.8 |
| Thyroid | 199.2 | 17.5 | 16.4-18.6 | 40.0 | 7.1 | 6.1-8.3 | 159.2 | 28.0 | 26.1-30.1 | 3.0 | 0.4 | 0.2-0.7 | 1.0 | 0.3 | 0.1-0.7 | 2.0 | 0.5 | 0.3-1.0 |
| Hodgkin lymphoma | 29.8 | 2.7 | 2.2-3.2 | 17.0 | 3.1 | 2.4-4.0 | 12.8 | 2.3 | 1.7-3.0 | 1.8 | 0.2 | 0.1-0.4 | 0.6 | 0.1 | 0.0-0.4 | 1.2 | 0.3 | 0.1-0.7 |
| Non-Hodgkin lymphomas | 173.4 | 19.9 | 18.5-21.4 | 90.0 | 23.0 | 20.7-25.5 | 83.4 | 17.8 | 16.0-19.6 | 26.2 | 3.7 | 3.0-4.4 | 12.8 | 4.2 | 3.1-5.4 | 13.4 | 3.3 | 2.5-4.2 |
| Myeloma | 69.0 | 8.5 | 7.5-9.5 | 34.8 | 9.1 | 7.7-10.7 | 34.2 | 7.8 | 6.7-9.1 | 15.0 | 2.1 | 1.6-2.7 | 6.6 | 2.1 | 1.4-3.1 | 8.4 | 2.1 | 1.5-2.8 |
| Leukemias | 131.2 | 14.0 | 12.9-15.2 | 74.0 | 17.4 | 15.4-19.5 | 57.2 | 11.4 | 10.1-12.9 | 34.6 | 4.1 | 3.5-4.8 | 20.6 | 5.3 | 4.2-6.6 | 14.0 | 3.1 | 2.4-3.9 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York State Excl New York City, Non-Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 70457.2 | 520.5 | 518.7-522.3 | 35874.2 | 565.8 | 563.1-568.5 | 34583.0 | 490.1 | 487.6-492.5 | 21301.2 | 149.7 | 148.8-150.6 | 10792.6 | 173.5 | 172.0-175.0 | 10508.6 | 133.0 | 131.9-134.2 |
| Oral cavity and pharynx | 1748.8 | 12.7 | 12.5-13.0 | 1218.8 | 18.9 | 18.4-19.4 | 530.0 | 7.3 | 7.0-7.6 | 324.0 | 2.3 | 2.1-2.4 | 223.4 | 3.5 | 3.3-3.7 | 100.6 | 1.3 | 1.1-1.4 |
| Esophagus | 755.8 | 5.3 | 5.2-5.5 | 579.4 | 8.9 | 8.6-9.3 | 176.4 | 2.3 | 2.1-2.4 | 601.2 | 4.2 | 4.0-4.3 | 473.2 | 7.3 | 7.0-7.6 | 128.0 | 1.6 | 1.5-1.7 |
| Stomach | 951.2 | 6.9 | 6.7-7.1 | 588.2 | 9.4 | 9.0-9.7 | 363.0 | 5.0 | 4.7-5.2 | 341.0 | 2.4 | 2.3-2.6 | 205.4 | 3.3 | 3.1-3.5 | 135.6 | 1.7 | 1.6-1.9 |
| Colorectal | 5177.0 | 38.6 | 38.1-39.1 | 2654.4 | 43.5 | 42.7-44.2 | 2522.6 | 34.5 | 33.9-35.2 | 1736.0 | 12.3 | 12.0-12.6 | 878.0 | 14.3 | 13.9-14.8 | 858.0 | 10.6 | 10.3-11.0 |
| Colon excluding rectum | 3679.6 | 27.3 | 26.9-27.7 | 1784.2 | 29.3 | 28.7-30.0 | 1895.4 | 25.5 | 25.0-26.1 | 1374.2 | 9.7 | 9.5-9.9 | 672.8 | 11.0 | 10.6-11.4 | 701.4 | 8.6 | 8.3-8.9 |
| Rectum & rectosigmoid | 1497.4 | 11.4 | 11.1-11.6 | 870.2 | 14.1 | 13.7-14.6 | 627.2 | 9.0 | 8.6-9.3 | 361.8 | 2.6 | 2.5-2.7 | 205.2 | 3.3 | 3.1-3.5 | 156.6 | 2.0 | 1.9-2.2 |
| Liver / intrahepatic bile duct | 1055.0 | 7.3 | 7.1-7.5 | 757.6 | 11.2 | 10.9-11.6 | 297.4 | 3.9 | 3.7-4.1 | 731.4 | 5.0 | 4.9-5.2 | 493.2 | 7.5 | 7.2-7.8 | 238.2 | 3.0 | 2.9-3.2 |
| Pancreas | 2145.8 | 15.2 | 14.9-15.5 | 1094.0 | 17.2 | 16.7-17.7 | 1051.8 | 13.5 | 13.1-13.9 | 1653.4 | 11.5 | 11.3-11.8 | 837.4 | 13.2 | 12.8-13.6 | 816.0 | 10.1 | 9.8-10.4 |
| Larynx | 458.6 | 3.2 | 3.1-3.4 | 360.6 | 5.5 | 5.2-5.7 | 98.0 | 1.3 | 1.2-1.5 | 125.2 | 0.9 | 0.8-0.9 | 97.6 | 1.5 | 1.4-1.6 | 27.6 | 0.3 | 0.3-0.4 |
| Lung and bronchus | 9535.8 | 66.8 | 66.2-67.4 | 4550.4 | 71.0 | 70.0-71.9 | 4985.4 | 64.1 | 63.3-65.0 | 5353.8 | 37.3 | 36.9-37.8 | 2699.2 | 42.7 | 41.9-43.4 | 2654.6 | 33.4 | 32.8-34.0 |
| Melanoma of the skin | 3283.4 | 25.2 | 24.8-25.6 | 1927.0 | 31.5 | 30.9-32.2 | 1356.4 | 20.8 | 20.2-21.3 | 297.6 | 2.1 | 2.0-2.2 | 189.2 | 3.1 | 2.9-3.3 | 108.4 | 1.4 | 1.3-1.5 |
| Female breast | | | | | | | 9813.8 | 144.1 | 142.7-145.4 | | | | | | | 1445.0 | 19.0 | 18.5-19.5 |
| Cervix uteri | | | | | | | 363.8 | 6.7 | 6.3-7.0 | | | | | | | 117.2 | 1.8 | 1.6-1.9 |
| Corpus uteri and NOS | | | | | | | 2395.8 | 32.5 | 31.9-33.1 | | | | | | | 421.0 | 5.4 | 5.1-5.6 |
| Ovary | | | | | | | 822.6 | 11.9 | 11.5-12.3 | | | | | | | 521.8 | 6.8 | 6.5-7.0 |
| Prostate | | | | 9281.4 | 134.6 | 133.4-135.9 | | | | | | | 986.4 | 16.4 | 16.0-16.9 | | | |
| Testis | | | | 329.0 | 7.4 | 7.0-7.8 | | | | | | | 12.6 | 0.2 | 0.2-0.3 | | | |
| Urinary bladder (incl. in situ) | 3750.2 | 26.4 | 26.0-26.8 | 2807.8 | 44.9 | 44.2-45.7 | 942.4 | 12.0 | 11.6-12.4 | 655.8 | 4.5 | 4.3-4.6 | 462.0 | 7.7 | 7.3-8.0 | 193.8 | 2.3 | 2.1-2.4 |
| Kidney and renal pelvis | 2547.6 | 19.1 | 18.7-19.4 | 1654.0 | 26.4 | 25.9-27.0 | 893.6 | 12.6 | 12.3-13.0 | 443.6 | 3.1 | 3.0-3.2 | 280.2 | 4.4 | 4.2-4.7 | 163.4 | 2.0 | 1.9-2.2 |
| Brain and other central nervous system | 839.2 | 7.1 | 6.9-7.3 | 465.8 | 8.3 | 8.0-8.7 | 373.4 | 6.1 | 5.8-6.4 | 580.4 | 4.4 | 4.2-4.6 | 326.2 | 5.3 | 5.1-5.6 | 254.2 | 3.6 | 3.4-3.8 |
| Thyroid | 2008.2 | 18.8 | 18.4-19.2 | 576.6 | 10.5 | 10.1-11.0 | 1431.6 | 26.9 | 26.2-27.5 | 74.8 | 0.5 | 0.5-0.6 | 35.6 | 0.6 | 0.5-0.7 | 39.2 | 0.5 | 0.4-0.6 |
| Hodgkin lymphoma | 341.8 | 3.3 | 3.1-3.5 | 183.6 | 3.6 | 3.3-3.8 | 158.2 | 3.1 | 2.8-3.3 | 39.0 | 0.3 | 0.3-0.3 | 23.6 | 0.4 | 0.3-0.5 | 15.4 | 0.2 | 0.2-0.3 |
| Non-Hodgkin lymphomas | 3021.6 | 22.5 | 22.2-22.9 | 1641.8 | 26.9 | 26.3-27.5 | 1379.8 | 19.0 | 18.6-19.5 | 757.2 | 5.3 | 5.2-5.5 | 425.0 | 7.0 | 6.7-7.3 | 332.2 | 4.1 | 3.9-4.3 |
| Myeloma | 1077.8 | 7.8 | 7.5-8.0 | 612.6 | 9.7 | 9.4-10.1 | 465.2 | 6.2 | 5.9-6.4 | 412.6 | 2.9 | 2.7-3.0 | 222.4 | 3.6 | 3.4-3.8 | 190.2 | 2.3 | 2.2-2.5 |
| Leukemias | 2349.4 | 17.7 | 17.3-18.0 | 1389.4 | 23.0 | 22.4-23.6 | 960.0 | 13.3 | 12.9-13.7 | 863.4 | 6.2 | 6.0-6.4 | 499.4 | 8.4 | 8.0-8.7 | 364.0 | 4.5 | 4.3-4.8 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York State Excl New York City, Non-Hispanic Whites

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 63539.4 | 528.7 | 526.8-530.7 | 32363.8 | 571.1 | 568.2-574.0 | 31175.6 | 501.0 | 498.4-503.7 | 19308.6 | 150.5 | 149.5-151.5 | 9802.6 | 174.3 | 172.7-175.9 | 9506.0 | 133.8 | 132.5-135.1 |
| Oral cavity and pharynx | 1607.4 | 13.2 | 12.9-13.5 | 1124.6 | 19.5 | 19.0-20.1 | 482.8 | 7.5 | 7.2-7.9 | 292.2 | 2.3 | 2.1-2.4 | 200.4 | 3.4 | 3.2-3.7 | 91.8 | 1.3 | 1.2-1.4 |
| Esophagus | 699.4 | 5.5 | 5.3-5.7 | 541.8 | 9.3 | 8.9-9.6 | 157.6 | 2.3 | 2.1-2.4 | 564.4 | 4.4 | 4.2-4.5 | 446.2 | 7.6 | 7.3-7.9 | 118.2 | 1.6 | 1.5-1.8 |
| Stomach | 791.4 | 6.4 | 6.2-6.6 | 502.4 | 8.9 | 8.6-9.3 | 289.0 | 4.4 | 4.1-4.6 | 275.8 | 2.2 | 2.1-2.3 | 167.2 | 3.0 | 2.8-3.2 | 108.6 | 1.5 | 1.4-1.7 |
| Colorectal | 4600.6 | 38.6 | 38.0-39.1 | 2363.2 | 43.4 | 42.6-44.3 | 2237.4 | 34.4 | 33.7-35.1 | 1551.8 | 12.2 | 11.9-12.5 | 782.4 | 14.2 | 13.8-14.7 | 769.4 | 10.6 | 10.2-10.9 |
| Colon excluding rectum | 3280.4 | 27.3 | 26.8-27.7 | 1591.2 | 29.3 | 28.7-30.0 | 1689.2 | 25.5 | 24.9-26.1 | 1223.2 | 9.6 | 9.3-9.8 | 594.2 | 10.8 | 10.4-11.2 | 629.0 | 8.5 | 8.2-8.9 |
| Rectum & rectosigmoid | 1320.2 | 11.3 | 11.0-11.6 | 772.0 | 14.1 | 13.7-14.6 | 548.2 | 8.9 | 8.5-9.2 | 328.6 | 2.6 | 2.5-2.8 | 188.2 | 3.4 | 3.2-3.6 | 140.4 | 2.0 | 1.9-2.2 |
| Liver / intrahepatic bile duct | 877.6 | 6.8 | 6.6-7.0 | 631.8 | 10.5 | 10.1-10.9 | 245.8 | 3.6 | 3.4-3.8 | 606.8 | 4.7 | 4.5-4.8 | 409.0 | 6.9 | 6.6-7.2 | 197.8 | 2.8 | 2.6-3.0 |
| Pancreas | 1924.6 | 15.1 | 14.8-15.5 | 990.8 | 17.3 | 16.8-17.8 | 933.8 | 13.3 | 12.9-13.7 | 1489.6 | 11.5 | 11.2-11.8 | 758.2 | 13.2 | 12.8-13.7 | 731.4 | 10.0 | 9.7-10.4 |
| Larynx | 414.8 | 3.3 | 3.1-3.4 | 326.8 | 5.6 | 5.3-5.8 | 88.0 | 1.4 | 1.2-1.5 | 110.6 | 0.8 | 0.8-0.9 | 85.4 | 1.5 | 1.3-1.6 | 25.2 | 0.4 | 0.3-0.4 |
| Lung and bronchus | 8787.0 | 68.6 | 67.9-69.2 | 4156.4 | 71.8 | 70.8-72.8 | 4630.6 | 66.7 | 65.8-67.6 | 4930.6 | 38.2 | 37.7-38.7 | 2467.4 | 43.1 | 42.4-43.9 | 2463.2 | 34.5 | 33.9-35.2 |
| Melanoma of the skin | 3214.8 | 28.2 | 27.7-28.6 | 1892.6 | 34.7 | 34.0-35.4 | 1322.2 | 23.5 | 22.9-24.1 | 293.8 | 2.4 | 2.2-2.5 | 187.0 | 3.4 | 3.2-3.6 | 106.8 | 1.6 | 1.4-1.7 |
| Female breast | | | | | | | 8757.8 | 146.3 | 144.8-147.8 | | | | | | | 1277.4 | 18.7 | 18.2-19.2 |
| Cervix uteri | | | | | | | 300.2 | 6.6 | 6.2-6.9 | | | | | | | 98.6 | 1.7 | 1.6-1.9 |
| Corpus uterus and NOS | | | | | | | 2147.2 | 33.0 | 32.4-33.7 | | | | | | | 360.6 | 5.1 | 4.9-5.4 |
| Ovary | | | | | | | 743.2 | 12.2 | 11.8-12.7 | | | | | | | 477.6 | 6.9 | 6.6-7.2 |
| Prostate | | | | 8024.4 | 129.3 | 128.0-130.6 | | | | | | | 873.4 | 15.9 | 15.4-16.4 | | | |
| Testis | | | | 312.2 | 8.6 | 8.1-9.0 | | | | | | | 12.2 | 0.3 | 0.2-0.4 | | | |
| Urinary bladder (incl. in situ) | 3559.6 | 27.9 | 27.5-28.3 | 2673.8 | 47.3 | 46.5-48.1 | 885.8 | 12.6 | 12.2-13.0 | 622.8 | 4.7 | 4.5-4.8 | 440.8 | 8.0 | 7.7-8.3 | 182.0 | 2.3 | 2.2-2.5 |
| Kidney and renal pelvis | 2281.8 | 19.3 | 18.9-19.7 | 1483.8 | 26.7 | 26.1-27.4 | 798.0 | 12.8 | 12.4-13.2 | 410.0 | 3.2 | 3.0-3.3 | 259.0 | 4.5 | 4.3-4.8 | 151.0 | 2.1 | 1.9-2.3 |
| Brain and other central nervous system | 773.4 | 7.6 | 7.3-7.8 | 429.4 | 8.8 | 8.4-9.2 | 344.0 | 6.5 | 6.2-6.8 | 541.2 | 4.7 | 4.5-4.9 | 306.4 | 5.6 | 5.3-5.9 | 234.8 | 3.8 | 3.6-4.0 |
| Thyroid | 1777.4 | 19.6 | 19.2-20.1 | 525.0 | 11.1 | 10.7-11.6 | 1252.4 | 28.0 | 27.3-28.8 | 66.6 | 0.5 | 0.5-0.6 | 31.8 | 0.6 | 0.5-0.7 | 34.8 | 0.5 | 0.4-0.6 |
| Hodgkin lymphoma | 297.4 | 3.4 | 3.3-3.6 | 160.8 | 3.7 | 3.4-4.0 | 136.6 | 3.2 | 2.9-3.5 | 35.8 | 0.3 | 0.3-0.4 | 21.2 | 0.4 | 0.3-0.5 | 14.6 | 0.2 | 0.2-0.3 |
| Non-Hodgkin lymphomas | 2770.2 | 23.1 | 22.7-23.6 | 1502.8 | 27.5 | 26.9-28.2 | 1267.4 | 19.6 | 19.1-20.2 | 698.2 | 5.4 | 5.2-5.6 | 389.6 | 7.1 | 6.7-7.4 | 308.6 | 4.2 | 3.9-4.4 |
| Myeloma | 894.8 | 7.2 | 7.0-7.4 | 514.0 | 9.1 | 8.7-9.4 | 380.8 | 5.6 | 5.4-5.9 | 355.0 | 2.7 | 2.6-2.8 | 193.8 | 3.4 | 3.2-3.7 | 161.2 | 2.1 | 2.0-2.3 |
| Leukemias | 2173.8 | 18.4 | 18.0-18.7 | 1292.4 | 23.9 | 23.3-24.5 | 881.4 | 13.8 | 13.4-14.3 | 794.0 | 6.3 | 6.1-6.5 | 460.8 | 8.5 | 8.2-8.9 | 333.2 | 4.6 | 4.4-4.8 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York State Excl New York City, Non-Hispanic Blacks

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 4990.8 | 496.4 | 490.1-502.8 | 2585.2 | 587.2 | 576.6-597.9 | 2405.6 | 435.2 | 427.3-443.2 | 1518.6 | 159.2 | 155.5-162.9 | 753.4 | 192.5 | 186.1-199.1 | 765.2 | 138.3 | 133.8-142.8 |
| Oral cavity and pharynx | 85.8 | 8.3 | 7.5-9.2 | 58.4 | 12.7 | 11.2-14.3 | 27.4 | 5.0 | 4.2-5.9 | 20.8 | 2.1 | 1.7-2.5 | 15.6 | 3.6 | 2.8-4.6 | 5.2 | 0.9 | 0.6-1.3 |
| Esophagus | 41.8 | 4.3 | 3.7-5.0 | 26.8 | 6.8 | 5.6-8.1 | 15.0 | 2.7 | 2.1-3.3 | 28.2 | 2.9 | 2.5-3.5 | 20.4 | 5.3 | 4.3-6.5 | 7.8 | 1.3 | 0.9-1.8 |
| Stomach | 115.4 | 12.2 | 11.2-13.3 | 60.8 | 14.9 | 13.2-16.8 | 54.6 | 10.3 | 9.1-11.6 | 46.8 | 5.0 | 4.3-5.7 | 28.8 | 7.3 | 6.1-8.7 | 18.0 | 3.3 | 2.6-4.0 |
| Colorectal | 416.2 | 42.0 | 40.2-43.9 | 211.4 | 49.6 | 46.5-52.9 | 204.8 | 36.8 | 34.6-39.2 | 140.4 | 14.7 | 13.7-15.9 | 72.2 | 18.1 | 16.1-20.1 | 68.2 | 12.4 | 11.1-13.8 |
| Colon excluding rectum | 300.2 | 30.7 | 29.1-32.3 | 145.8 | 34.8 | 32.2-37.6 | 154.4 | 27.8 | 25.9-29.9 | 116.2 | 12.2 | 11.2-13.3 | 60.0 | 15.1 | 13.3-17.0 | 56.2 | 10.2 | 9.0-11.5 |
| Rectum & rectosigmoid | 116.0 | 11.4 | 10.5-12.4 | 65.6 | 14.8 | 13.1-16.5 | 50.4 | 9.0 | 7.9-10.2 | 24.2 | 2.5 | 2.1-3.1 | 12.2 | 3.0 | 2.2-3.9 | 12.0 | 2.2 | 1.7-2.8 |
| Liver / intrahepatic bile duct | 118.4 | 10.9 | 10.0-11.8 | 86.2 | 17.9 | 16.2-19.8 | 32.2 | 5.4 | 4.6-6.3 | 82.8 | 8.1 | 7.3-8.9 | 57.4 | 12.7 | 11.2-14.3 | 25.4 | 4.5 | 3.8-5.4 |
| Pancreas | 169.8 | 17.6 | 16.4-18.8 | 76.8 | 18.3 | 16.4-20.3 | 93.0 | 16.9 | 15.4-18.5 | 126.8 | 13.4 | 12.3-14.5 | 60.0 | 14.6 | 12.9-16.5 | 66.8 | 12.3 | 11.0-13.7 |
| Larynx | 35.2 | 3.4 | 2.9-3.9 | 28.0 | 6.2 | 5.2-7.4 | 7.2 | 1.2 | 0.8-1.7 | 11.6 | 1.2 | 0.9-1.5 | 9.4 | 2.3 | 1.7-3.1 | 2.2 | 0.4 | 0.2-0.7 |
| Lung and bronchus | 584.6 | 59.3 | 57.1-61.5 | 306.6 | 74.9 | 71.0-78.9 | 278.0 | 48.9 | 46.3-51.6 | 332.2 | 34.5 | 32.8-36.3 | 183.2 | 46.7 | 43.6-50.0 | 149.0 | 26.4 | 24.5-28.4 |
| Melanoma of the skin | 10.0 | 1.0 | 0.8-1.4 | 4.4 | 1.0 | 0.6-1.6 | 5.6 | 1.0 | 0.7-1.5 | 2.0 | 0.2 | 0.1-0.4 | 1.2 | 0.3 | 0.1-0.7 | 0.8 | 0.1 | 0.0-0.4 |
| Female breast | | | | | | | 738.6 | 134.8 | 130.4-139.3 | | | | | | | 132.0 | 23.8 | 22.0-25.7 |
| Cervix uteri | | | | | | | 44.8 | 8.3 | 7.3-9.5 | | | | | | | 15.0 | 2.8 | 2.2-3.5 |
| Corpus uterus and NOS | | | | | | | 177.8 | 31.0 | 28.9-33.1 | | | | | | | 50.2 | 8.9 | 7.9-10.1 |
| Ovary | | | | | | | 47.0 | 8.5 | 7.4-9.7 | | | | | | | 29.6 | 5.4 | 4.6-6.4 |
| Prostate | | | | 1004.2 | 217.2 | 211.0-223.6 | | | | | | | 95.4 | 28.6 | 26.0-31.4 | | | |
| Testis | | | | 9.0 | 1.8 | 1.3-2.4 | | | | | | | 0.0 | 0.0 | 0.0-0.2 | | | |
| Urinary bladder (incl. in situ) | 129.8 | 13.6 | 12.6-14.7 | 87.4 | 22.0 | 19.9-24.3 | 42.4 | 7.8 | 6.7-8.9 | 24.8 | 2.8 | 2.3-3.4 | 14.6 | 4.4 | 3.4-5.5 | 10.2 | 1.9 | 1.4-2.5 |
| Kidney and renal pelvis | 209.0 | 20.5 | 19.2-21.8 | 131.6 | 28.7 | 26.4-31.1 | 77.4 | 14.0 | 12.6-15.5 | 27.4 | 2.8 | 2.4-3.4 | 17.8 | 4.2 | 3.3-5.2 | 9.6 | 1.8 | 1.3-2.4 |
| Brain and other central nervous system | 42.2 | 4.2 | 3.6-4.8 | 22.4 | 4.8 | 3.9-5.8 | 19.8 | 3.7 | 3.0-4.5 | 24.4 | 2.4 | 2.0-2.9 | 11.6 | 2.5 | 1.9-3.2 | 12.8 | 2.3 | 1.8-2.9 |
| Thyroid | 114.2 | 11.4 | 10.4-12.4 | 25.2 | 5.7 | 4.7-6.8 | 89.0 | 16.8 | 15.2-18.5 | 5.6 | 0.6 | 0.4-0.9 | 2.8 | 0.7 | 0.4-1.2 | 2.8 | 0.5 | 0.3-0.8 |
| Hodgkin lymphoma | 33.0 | 3.2 | 2.7-3.7 | 16.8 | 3.4 | 2.7-4.2 | 16.2 | 3.0 | 2.4-3.8 | 2.6 | 0.3 | 0.1-0.5 | 2.2 | 0.5 | 0.3-1.0 | 0.4 | 0.1 | 0.0-0.3 |
| Non-Hodgkin lymphomas | 166.6 | 16.7 | 15.6-17.9 | 90.8 | 20.7 | 18.7-22.8 | 75.8 | 13.8 | 12.4-15.3 | 41.0 | 4.3 | 3.7-4.9 | 24.8 | 6.2 | 5.1-7.4 | 16.2 | 2.9 | 2.3-3.7 |
| Myeloma | 156.0 | 15.9 | 14.8-17.1 | 80.4 | 19.3 | 17.4-21.4 | 75.6 | 13.5 | 12.1-14.9 | 48.8 | 5.4 | 4.7-6.1 | 25.0 | 6.8 | 5.6-8.1 | 23.8 | 4.4 | 3.7-5.3 |
| Leukemias | 114.4 | 11.6 | 10.6-12.6 | 61.2 | 14.4 | 12.7-16.1 | 53.2 | 9.7 | 8.5-10.9 | 49.4 | 5.2 | 4.6-5.9 | 25.6 | 6.6 | 5.5-8.0 | 23.8 | 4.4 | 3.6-5.2 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Bronx, Hispanic Origin**

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 2772.6 | 385.8 | 379.3-392.4 | 1345.8 | 453.7 | 442.5-465.2 | 1426.8 | 343.7 | 335.7-351.9 | 820.4 | 121.2 | 117.5-125.1 | 417.4 | 157.0 | 150.1-164.2 | 403.0 | 99.2 | 94.9-103.7 |
| Oral cavity and pharynx | 71.0 | 9.7 | 8.7-10.8 | 49.4 | 16.3 | 14.2-18.5 | 21.6 | 5.1 | 4.2-6.2 | 18.0 | 2.6 | 2.1-3.2 | 13.2 | 4.6 | 3.5-5.9 | 4.8 | 1.2 | 0.8-1.8 |
| Esophagus | 24.2 | 3.4 | 2.8-4.1 | 19.4 | 6.6 | 5.3-8.1 | 4.8 | 1.2 | 0.8-1.8 | 16.6 | 2.4 | 1.9-3.0 | 12.8 | 4.6 | 3.5-5.9 | 3.8 | 0.9 | 0.6-1.5 |
| Stomach | 69.2 | 9.9 | 8.9-11.1 | 36.6 | 12.8 | 11.0-14.9 | 32.6 | 8.0 | 6.8-9.4 | 26.0 | 3.9 | 3.3-4.7 | 16.4 | 6.4 | 5.0-7.9 | 9.6 | 2.4 | 1.8-3.3 |
| Colorectal | 230.2 | 32.6 | 30.7-34.6 | 129.6 | 44.1 | 40.6-47.7 | 100.6 | 24.5 | 22.4-26.8 | 75.2 | 11.3 | 10.1-12.5 | 41.0 | 15.3 | 13.2-17.6 | 34.2 | 8.5 | 7.3-9.9 |
| Colon excluding rectum | 158.2 | 22.7 | 21.1-24.4 | 86.6 | 29.9 | 27.1-33.0 | 71.6 | 17.7 | 15.9-19.7 | 61.2 | 9.2 | 8.2-10.3 | 31.0 | 11.6 | 9.8-13.6 | 30.2 | 7.5 | 6.3-8.8 |
| Rectum & rectosigmoid | 72.0 | 9.9 | 8.9-11.0 | 43.0 | 14.2 | 12.3-16.2 | 29.0 | 6.8 | 5.8-8.1 | 14.0 | 2.1 | 1.6-2.7 | 10.0 | 3.7 | 2.7-4.9 | 4.0 | 1.0 | 0.6-1.6 |
| Liver / intrahepatic bile duct | 123.8 | 16.7 | 15.4-18.1 | 88.2 | 28.2 | 25.5-31.0 | 35.6 | 8.5 | 7.3-9.8 | 68.8 | 9.8 | 8.8-10.9 | 48.0 | 16.6 | 14.5-19.0 | 20.8 | 5.0 | 4.1-6.1 |
| Pancreas | 84.0 | 12.2 | 11.0-13.4 | 42.2 | 15.1 | 13.1-17.4 | 41.8 | 10.2 | 8.9-11.7 | 52.6 | 7.9 | 7.0-8.9 | 26.4 | 10.1 | 8.4-12.0 | 26.2 | 6.5 | 5.4-7.8 |
| Larynx | 28.0 | 3.9 | 3.2-4.6 | 20.2 | 6.9 | 5.6-8.5 | 7.8 | 1.8 | 1.3-2.5 | 9.2 | 1.3 | 0.9-1.7 | 7.6 | 2.7 | 1.9-3.7 | 1.6 | 0.4 | 0.2-0.7 |
| Lung and bronchus | 256.6 | 37.2 | 35.2-39.4 | 133.8 | 49.0 | 45.2-53.0 | 122.8 | 29.7 | 27.4-32.2 | 135.8 | 20.2 | 18.7-21.8 | 77.0 | 29.2 | 26.3-32.4 | 58.8 | 14.4 | 12.8-16.2 |
| Melanoma of the skin | 10.4 | 1.5 | 1.1-2.0 | 4.6 | 1.6 | 1.0-2.4 | 5.8 | 1.4 | 0.9-2.0 | 1.8 | 0.2 | 0.1-0.5 | 1.4 | 0.5 | 0.2-1.0 | 0.4 | 0.1 | 0.0-0.4 |
| Female breast | | | | | | | 412.6 | 99.1 | 94.9-103.6 | | | | | | | 63.8 | 15.7 | 14.0-17.5 |
| Cervix uteri | | | | | | | 37.0 | 9.1 | 7.8-10.5 | | | | | | | 10.4 | 2.5 | 1.9-3.3 |
| Corpus uterus and NOS | | | | | | | 103.8 | 24.4 | 22.4-26.7 | | | | | | | 22.6 | 5.3 | 4.4-6.4 |
| Ovary | | | | | | | 37.8 | 9.2 | 7.9-10.6 | | | | | | | 17.6 | 4.4 | 3.5-5.4 |
| Prostate | | | | 375.4 | 124.3 | 118.6-130.3 | | | | | | | 48.2 | 21.1 | 18.5-24.0 | | | |
| Testis | | | | 12.4 | 3.1 | 2.4-4.0 | | | | | | | 0.8 | 0.2 | 0.1-0.6 | | | |
| Urinary bladder (incl. in situ) | 76.0 | 11.3 | 10.1-12.5 | 50.8 | 19.0 | 16.7-21.6 | 25.2 | 6.3 | 5.2-7.5 | 15.6 | 2.5 | 2.0-3.1 | 9.6 | 4.2 | 3.1-5.6 | 6.0 | 1.6 | 1.0-2.2 |
| Kidney and renal pelvis | 100.2 | 13.7 | 12.5-15.0 | 59.8 | 19.7 | 17.4-22.1 | 40.4 | 9.6 | 8.3-11.0 | 16.2 | 2.4 | 1.9-3.0 | 10.8 | 3.9 | 2.9-5.1 | 5.4 | 1.3 | 0.9-1.9 |
| Brain and other central nervous system | 39.6 | 5.3 | 4.6-6.1 | 20.6 | 6.3 | 5.1-7.7 | 19.0 | 4.5 | 3.7-5.5 | 20.4 | 2.9 | 2.4-3.5 | 9.2 | 3.2 | 2.3-4.3 | 11.2 | 2.7 | 2.1-3.6 |
| Thyroid | 100.8 | 13.4 | 12.2-14.6 | 18.0 | 5.7 | 4.5-7.0 | 82.8 | 19.8 | 18.0-21.9 | 3.4 | 0.5 | 0.3-0.8 | 1.2 | 0.4 | 0.1-0.9 | 2.2 | 0.6 | 0.3-1.0 |
| Hodgkin lymphoma | 23.2 | 3.0 | 2.4-3.6 | 11.8 | 3.3 | 2.5-4.3 | 11.4 | 2.7 | 2.0-3.5 | 2.0 | 0.3 | 0.1-0.5 | 1.4 | 0.5 | 0.2-0.9 | 0.6 | 0.2 | 0.0-0.4 |
| Non-Hodgkin lymphomas | 132.2 | 18.6 | 17.1-20.1 | 64.6 | 21.9 | 19.5-24.6 | 67.6 | 16.3 | 14.6-18.2 | 30.8 | 4.6 | 3.9-5.4 | 16.2 | 6.1 | 4.8-7.6 | 14.6 | 3.7 | 2.9-4.6 |
| Myeloma | 72.6 | 10.2 | 9.2-11.3 | 39.2 | 13.6 | 11.7-15.7 | 33.4 | 8.0 | 6.8-9.3 | 22.0 | 3.3 | 2.7-3.9 | 12.6 | 4.7 | 3.5-6.0 | 9.4 | 2.3 | 1.7-3.1 |
| Leukemias | 86.0 | 11.7 | 10.6-12.9 | 45.0 | 14.5 | 12.6-16.6 | 41.0 | 9.7 | 8.4-11.2 | 34.0 | 4.9 | 4.1-5.7 | 16.6 | 5.7 | 4.5-7.1 | 17.4 | 4.3 | 3.4-5.3 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Bronx, Non-Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 3708.8 | 493.4 | 486.2-500.8 | 1803.2 | 551.0 | 539.5-562.8 | 1905.6 | 456.9 | 447.5-466.5 | 1220.2 | 157.2 | 153.2-161.3 | 584.8 | 185.5 | 178.7-192.5 | 635.4 | 139.5 | 134.5-144.6 |
| Oral cavity and pharynx | 78.4 | 10.5 | 9.5-11.7 | 51.0 | 15.6 | 13.7-17.7 | 27.4 | 6.5 | 5.4-7.7 | 21.2 | 2.8 | 2.3-3.4 | 13.8 | 4.3 | 3.3-5.5 | 7.4 | 1.7 | 1.2-2.4 |
| Esophagus | 26.0 | 3.3 | 2.8-4.0 | 17.4 | 5.4 | 4.3-6.6 | 8.6 | 2.0 | 1.4-2.7 | 19.4 | 2.4 | 2.0-3.0 | 11.8 | 3.6 | 2.7-4.6 | 7.6 | 1.6 | 1.1-2.3 |
| Stomach | 74.8 | 10.1 | 9.1-11.2 | 39.2 | 12.7 | 10.9-14.6 | 35.6 | 8.3 | 7.1-9.6 | 32.6 | 4.2 | 3.6-5.0 | 18.2 | 5.8 | 4.7-7.2 | 14.4 | 3.1 | 2.4-4.0 |
| Colorectal | 303.4 | 40.6 | 38.6-42.8 | 143.6 | 45.2 | 41.9-48.7 | 159.8 | 37.3 | 34.7-40.1 | 113.6 | 14.6 | 13.4-15.9 | 52.0 | 16.6 | 14.6-18.7 | 61.6 | 13.2 | 11.7-14.9 |
| Colon excluding rectum | 215.2 | 28.6 | 26.9-30.4 | 99.4 | 31.5 | 28.7-34.5 | 115.8 | 26.5 | 24.3-28.9 | 93.8 | 12.0 | 10.9-13.2 | 40.8 | 12.9 | 11.2-14.9 | 53.0 | 11.4 | 10.0-12.9 |
| Rectum & rectosigmoid | 88.2 | 12.0 | 10.9-13.2 | 44.2 | 13.7 | 11.9-15.7 | 44.0 | 10.8 | 9.3-12.3 | 19.8 | 2.6 | 2.1-3.2 | 11.2 | 3.6 | 2.7-4.7 | 8.6 | 1.8 | 1.3-2.5 |
| Liver / intrahepatic bile duct | 108.4 | 13.8 | 12.6-15.0 | 75.6 | 22.3 | 20.1-24.8 | 32.8 | 7.2 | 6.1-8.4 | 65.0 | 8.3 | 7.4-9.3 | 43.0 | 12.9 | 11.2-14.8 | 22.0 | 4.7 | 3.8-5.7 |
| Pancreas | 118.0 | 15.1 | 13.9-16.4 | 59.4 | 18.3 | 16.2-20.6 | 58.6 | 12.6 | 11.2-14.2 | 93.8 | 12.0 | 10.9-13.2 | 46.4 | 14.5 | 12.6-16.5 | 47.4 | 10.2 | 8.9-11.6 |
| Larynx | 31.4 | 4.1 | 3.5-4.8 | 22.4 | 6.8 | 5.6-8.2 | 9.0 | 2.1 | 1.5-2.9 | 9.0 | 1.2 | 0.8-1.6 | 6.6 | 2.1 | 1.5-3.0 | 2.4 | 0.5 | 0.3-1.0 |
| Lung and bronchus | 410.0 | 52.8 | 50.5-55.1 | 202.4 | 62.2 | 58.3-66.2 | 207.6 | 46.0 | 43.2-49.0 | 254.2 | 32.3 | 30.5-34.2 | 133.0 | 41.1 | 38.0-44.5 | 121.2 | 26.0 | 24.0-28.3 |
| Melanoma of the skin | 48.2 | 6.6 | 5.8-7.5 | 26.2 | 8.8 | 7.3-10.5 | 22.0 | 5.1 | 4.2-6.3 | 5.8 | 0.7 | 0.5-1.1 | 3.6 | 1.2 | 0.7-1.9 | 2.2 | 0.4 | 0.2-0.8 |
| Female breast | | | | | | | 534.0 | 131.9 | 126.8-137.2 | | | | | | | 106.4 | 24.2 | 22.1-26.4 |
| Cervix uteri | | | | | | | 39.2 | 11.0 | 9.4-12.7 | | | | | | | 10.8 | 2.9 | 2.1-3.8 |
| Corpus uterus and NOS | | | | | | | 162.2 | 37.2 | 34.6-39.9 | | | | | | | 50.2 | 11.2 | 9.8-12.7 |
| Ovary | | | | | | | 49.2 | 12.1 | 10.6-13.7 | | | | | | | 27.0 | 6.1 | 5.1-7.3 |
| Prostate | | | | 581.0 | 167.3 | 161.2-173.7 | | | | | | | 78.0 | 25.9 | 23.3-28.6 | | | |
| Testis | | | | 10.4 | 3.6 | 2.7-4.7 | | | | | | | 1.6 | 0.6 | 0.3-1.2 | | | |
| Urinary bladder (incl. in situ) | 123.8 | 16.0 | 14.8-17.4 | 88.2 | 28.4 | 25.8-31.3 | 35.6 | 7.8 | 6.6-9.0 | 30.4 | 3.8 | 3.2-4.4 | 18.8 | 6.2 | 5.0-7.6 | 11.6 | 2.2 | 1.6-2.9 |
| Kidney and renal pelvis | 136.6 | 18.0 | 16.7-19.5 | 83.4 | 25.4 | 23.0-28.1 | 53.2 | 12.5 | 11.0-14.1 | 26.2 | 3.4 | 2.9-4.1 | 16.4 | 5.3 | 4.2-6.5 | 9.8 | 2.1 | 1.5-2.8 |
| Brain and other central nervous system | 37.4 | 5.5 | 4.7-6.4 | 19.6 | 6.5 | 5.2-8.0 | 17.8 | 4.8 | 3.8-5.9 | 23.0 | 3.1 | 2.5-3.7 | 14.0 | 4.4 | 3.4-5.6 | 9.0 | 2.1 | 1.5-2.9 |
| Thyroid | 115.8 | 17.5 | 16.0-19.0 | 23.0 | 7.4 | 6.1-9.0 | 92.8 | 26.1 | 23.7-28.7 | 4.2 | 0.6 | 0.3-0.9 | 2.2 | 0.7 | 0.4-1.3 | 2.0 | 0.4 | 0.2-0.8 |
| Hodgkin lymphoma | 22.0 | 3.5 | 2.8-4.2 | 12.8 | 4.3 | 3.3-5.5 | 9.2 | 2.7 | 2.0-3.7 | 2.6 | 0.4 | 0.2-0.6 | 1.4 | 0.4 | 0.2-0.9 | 1.2 | 0.3 | 0.1-0.7 |
| Non-Hodgkin lymphomas | 148.8 | 20.7 | 19.2-22.3 | 78.0 | 24.9 | 22.5-27.6 | 70.8 | 17.5 | 15.7-19.5 | 40.2 | 5.3 | 4.6-6.1 | 23.2 | 7.4 | 6.1-9.0 | 17.0 | 3.8 | 3.0-4.8 |
| Myeloma | 98.4 | 13.0 | 11.8-14.2 | 47.2 | 14.4 | 12.6-16.4 | 51.2 | 12.0 | 10.6-13.7 | 29.8 | 3.7 | 3.2-4.4 | 15.2 | 4.7 | 3.7-5.9 | 14.6 | 3.0 | 2.4-3.9 |
| Leukemias | 100.2 | 13.7 | 12.5-15.0 | 51.4 | 16.5 | 14.5-18.7 | 48.8 | 11.7 | 10.2-13.4 | 41.6 | 5.4 | 4.7-6.2 | 21.6 | 7.1 | 5.8-8.6 | 20.0 | 4.3 | 3.5-5.4 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Bronx, Non-Hispanic Whites

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 1190.0 | 546.5 | 531.2-562.1 | 593.0 | 576.3 | 554.6-598.8 | 597.0 | 532.5 | 510.5-555.3 | 409.4 | 155.3 | 148.0-162.9 | 206.8 | 183.0 | 171.5-195.2 | 202.6 | 135.2 | 125.7-145.4 |
| Oral cavity and pharynx | 28.6 | 14.2 | 11.8-17.1 | 19.2 | 20.1 | 16.1-25.0 | 9.4 | 8.5 | 6.0-11.9 | 8.0 | 3.3 | 2.3-4.7 | 4.6 | 4.3 | 2.7-6.7 | 3.4 | 2.6 | 1.3-4.6 |
| Esophagus | 10.0 | 4.2 | 3.1-5.8 | 6.8 | 6.3 | 4.3-9.0 | 3.2 | 2.5 | 1.4-4.5 | 9.0 | 3.9 | 2.8-5.4 | 6.0 | 5.7 | 3.8-8.4 | 3.0 | 2.2 | 1.2-4.1 |
| Stomach | 19.2 | 8.8 | 7.0-11.0 | 11.4 | 11.8 | 8.8-15.6 | 7.8 | 6.1 | 4.1-8.8 | 7.0 | 2.7 | 1.8-4.0 | 3.4 | 3.0 | 1.7-5.1 | 3.6 | 2.5 | 1.3-4.6 |
| Colorectal | 100.0 | 44.7 | 40.4-49.3 | 49.2 | 50.3 | 43.8-57.6 | 50.8 | 39.6 | 34.1-45.9 | 39.0 | 14.4 | 12.3-16.9 | 18.8 | 16.8 | 13.4-20.9 | 20.2 | 12.4 | 9.7-15.7 |
| Colon excluding rectum | 73.6 | 31.7 | 28.2-35.6 | 35.6 | 36.0 | 30.5-42.2 | 38.0 | 27.9 | 23.5-33.0 | 32.8 | 11.9 | 10.0-14.2 | 15.8 | 14.0 | 10.9-17.7 | 17.0 | 10.2 | 7.9-13.3 |
| Rectum & rectosigmoid | 26.4 | 13.0 | 10.6-15.7 | 13.6 | 14.3 | 10.9-18.5 | 12.8 | 11.7 | 8.6-15.7 | 6.2 | 2.5 | 1.6-3.8 | 3.0 | 2.9 | 1.6-5.0 | 3.2 | 2.1 | 1.0-4.1 |
| Liver / intrahepatic bile duct | 27.8 | 12.3 | 10.2-14.8 | 20.6 | 19.4 | 15.7-23.8 | 7.2 | 6.2 | 4.1-9.0 | 17.6 | 7.5 | 5.9-9.4 | 11.6 | 11.0 | 8.3-14.5 | 6.0 | 4.4 | 2.7-6.8 |
| Pancreas | 36.2 | 13.7 | 11.6-16.2 | 20.2 | 18.4 | 14.8-22.6 | 16.0 | 9.7 | 7.3-12.8 | 29.2 | 10.3 | 8.6-12.3 | 15.0 | 12.7 | 10.0-16.2 | 14.2 | 8.5 | 6.4-11.3 |
| Larynx | 12.2 | 5.8 | 4.3-7.6 | 9.0 | 9.2 | 6.6-12.5 | 3.2 | 2.7 | 1.5-4.8 | 2.8 | 1.1 | 0.6-2.0 | 2.4 | 2.1 | 1.1-3.9 | 0.4 | 0.3 | 0.0-1.6 |
| Lung and bronchus | 147.0 | 58.8 | 54.4-63.5 | 74.0 | 66.6 | 59.8-74.1 | 73.0 | 52.6 | 47.0-59.0 | 93.0 | 35.3 | 32.0-38.9 | 49.4 | 43.6 | 38.2-49.7 | 43.6 | 28.3 | 24.3-32.9 |
| Melanoma of the skin | 42.2 | 20.5 | 17.5-23.9 | 24.2 | 25.6 | 20.9-31.1 | 18.0 | 16.3 | 12.6-20.9 | 5.0 | 2.1 | 1.3-3.3 | 3.2 | 2.9 | 1.6-5.0 | 1.8 | 1.5 | 0.6-3.4 |
| Female breast | | | | | | | 158.4 | 154.5 | 142.3-167.5 | | | | | | | 30.6 | 20.2 | 16.5-24.5 |
| Cervix uteri | | | | | | | 5.4 | 7.7 | 4.9-11.5 | | | | | | | 2.2 | 2.8 | 1.3-5.3 |
| Corpus uterus and NOS | | | | | | | 44.4 | 41.3 | 35.4-47.9 | | | | | | | 8.4 | 6.9 | 4.7-9.9 |
| Ovary | | | | | | | 19.4 | 18.8 | 14.8-23.8 | | | | | | | 11.0 | 9.1 | 6.6-12.4 |
| Prostate | | | | 129.0 | 117.2 | 108.1-127.0 | | | | | | | 21.4 | 16.6 | 13.5-20.4 | | | |
| Testis | | | | 5.0 | 7.6 | 4.9-11.4 | | | | | | | 0.6 | 0.7 | 0.1-2.3 | | | |
| Urinary bladder (incl. in situ) | 66.6 | 26.0 | 23.1-29.3 | 48.4 | 43.4 | 37.9-49.6 | 18.2 | 12.2 | 9.6-15.6 | 16.6 | 5.1 | 4.0-6.5 | 12.0 | 9.5 | 7.2-12.5 | 4.6 | 1.8 | 1.1-3.2 |
| Kidney and renal pelvis | 43.6 | 19.9 | 17.1-23.0 | 27.8 | 27.0 | 22.5-32.3 | 15.8 | 13.7 | 10.5-17.7 | 10.4 | 4.4 | 3.1-6.0 | 6.4 | 6.5 | 4.3-9.6 | 4.0 | 2.4 | 1.4-4.2 |
| Brain and other central nervous system | 14.4 | 7.5 | 5.7-9.8 | 8.2 | 9.4 | 6.6-13.2 | 6.2 | 5.7 | 3.6-8.6 | 11.8 | 5.4 | 4.0-7.3 | 7.0 | 7.5 | 5.0-10.8 | 4.8 | 3.4 | 2.1-5.6 |
| Thyroid | 40.0 | 28.5 | 24.4-33.1 | 11.4 | 15.5 | 11.5-20.4 | 28.6 | 42.2 | 35.0-50.4 | 1.4 | 0.5 | 0.2-1.2 | 0.6 | 0.6 | 0.1-2.0 | 0.8 | 0.4 | 0.1-1.7 |
| Hodgkin lymphoma | 5.8 | 4.0 | 2.6-5.9 | 3.4 | 4.7 | 2.6-7.6 | 2.4 | 3.4 | 1.6-6.2 | 0.6 | 0.4 | 0.1-1.2 | 0.4 | 0.4 | 0.0-1.9 | 0.2 | 0.3 | 0.0-1.8 |
| Non-Hodgkin lymphomas | 53.6 | 25.3 | 22.0-29.0 | 29.2 | 30.1 | 25.1-35.8 | 24.4 | 20.9 | 16.6-25.9 | 15.4 | 5.8 | 4.4-7.5 | 8.4 | 7.5 | 5.3-10.5 | 7.0 | 4.3 | 2.8-6.6 |
| Myeloma | 24.2 | 10.0 | 8.2-12.2 | 14.0 | 12.7 | 9.9-16.3 | 10.2 | 7.7 | 5.4-10.7 | 6.2 | 2.1 | 1.3-3.1 | 3.0 | 2.3 | 1.2-4.0 | 3.2 | 2.1 | 1.1-4.0 |
| Leukemias | 42.0 | 19.2 | 16.3-22.4 | 23.4 | 23.7 | 19.3-28.9 | 18.6 | 15.3 | 11.8-19.6 | 19.4 | 7.8 | 6.1-9.8 | 10.6 | 10.0 | 7.3-13.5 | 8.8 | 6.2 | 4.0-9.1 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Bronx, Non-Hispanic Blacks

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 2237.6 | 482.0 | 472.9-491.2 | 1072.6 | 555.4 | 539.9-571.3 | 1165.0 | 437.7 | 426.3-449.3 | 720.4 | 159.3 | 154.1-164.8 | 326.6 | 186.3 | 176.9-196.1 | 393.8 | 144.6 | 138.2-151.3 |
| Oral cavity and pharynx | 43.0 | 9.2 | 8.0-10.5 | 27.2 | 13.9 | 11.6-16.5 | 15.8 | 5.8 | 4.6-7.3 | 11.4 | 2.5 | 1.9-3.2 | 7.4 | 3.8 | 2.6-5.3 | 4.0 | 1.6 | 1.0-2.5 |
| Esophagus | 13.4 | 2.9 | 2.3-3.7 | 8.6 | 5.1 | 3.6-6.9 | 4.8 | 1.8 | 1.1-2.7 | 9.2 | 2.0 | 1.4-2.7 | 5.2 | 2.9 | 1.9-4.4 | 4.0 | 1.5 | 0.9-2.3 |
| Stomach | 48.8 | 11.1 | 9.7-12.6 | 23.8 | 14.1 | 11.6-17.1 | 25.0 | 9.4 | 7.8-11.2 | 22.0 | 5.0 | 4.1-6.1 | 12.2 | 7.1 | 5.4-9.2 | 9.8 | 3.6 | 2.7-4.8 |
| Colorectal | 177.6 | 39.0 | 36.4-41.7 | 82.0 | 43.8 | 39.4-48.5 | 95.6 | 35.9 | 32.7-39.3 | 65.0 | 14.7 | 13.1-16.4 | 28.6 | 16.4 | 13.7-19.5 | 36.4 | 13.4 | 11.5-15.6 |
| Colon excluding rectum | 125.0 | 27.6 | 25.4-29.9 | 54.6 | 29.4 | 25.9-33.4 | 70.4 | 26.4 | 23.7-29.4 | 54.0 | 12.2 | 10.7-13.8 | 22.2 | 12.7 | 10.3-15.4 | 31.8 | 11.7 | 9.9-13.7 |
| Rectum & rectosigmoid | 52.6 | 11.4 | 10.0-12.9 | 27.4 | 14.3 | 11.9-17.1 | 25.2 | 9.5 | 7.9-11.3 | 11.0 | 2.5 | 1.9-3.3 | 6.4 | 3.8 | 2.5-5.4 | 4.6 | 1.7 | 1.1-2.6 |
| Liver / intrahepatic bile duct | 68.6 | 14.0 | 12.5-15.6 | 47.8 | 23.7 | 20.7-27.1 | 20.8 | 7.1 | 5.8-8.7 | 41.0 | 8.3 | 7.2-9.6 | 27.4 | 13.4 | 11.2-16.0 | 13.6 | 4.7 | 3.6-6.0 |
| Pancreas | 75.4 | 16.4 | 14.8-18.2 | 35.8 | 18.9 | 16.1-22.0 | 39.6 | 14.5 | 12.5-16.7 | 56.8 | 12.5 | 11.1-14.1 | 26.6 | 14.0 | 11.6-16.8 | 30.2 | 11.2 | 9.5-13.2 |
| Larynx | 18.2 | 3.8 | 3.0-4.6 | 12.6 | 6.5 | 4.9-8.4 | 5.6 | 2.0 | 1.3-2.9 | 5.8 | 1.3 | 0.8-1.8 | 3.8 | 2.3 | 1.4-3.7 | 2.0 | 0.7 | 0.3-1.3 |
| Lung and bronchus | 240.0 | 51.7 | 48.7-54.8 | 114.4 | 61.2 | 56.0-66.7 | 125.6 | 45.3 | 41.8-49.1 | 142.0 | 30.8 | 28.6-33.2 | 71.4 | 39.5 | 35.3-44.0 | 70.6 | 25.4 | 22.7-28.2 |
| Melanoma of the skin | 4.4 | 1.1 | 0.7-1.6 | 1.2 | 0.9 | 0.3-1.8 | 3.2 | 1.3 | 0.7-2.1 | 0.4 | 0.1 | 0.0-0.3 | 0.2 | 0.1 | 0.0-0.6 | 0.2 | 0.1 | 0.0-0.5 |
| Female breast | | | | | | | 332.2 | 125.7 | 119.6-132.0 | | | | | | | 70.8 | 26.5 | 23.8-29.5 |
| Cervix uteri | | | | | | | 29.0 | 11.9 | 10.0-14.0 | | | | | | | 7.8 | 3.1 | 2.2-4.3 |
| Corpus uterus and NOS | | | | | | | 108.0 | 38.5 | 35.3-41.9 | | | | | | | 39.0 | 14.0 | 12.1-16.2 |
| Ovary | | | | | | | 25.2 | 9.4 | 7.8-11.3 | | | | | | | 14.6 | 5.3 | 4.2-6.7 |
| Prostate | | | | 411.6 | 201.4 | 192.4-210.7 | | | | | | | 51.8 | 34.7 | 30.4-39.3 | | | |
| Testis | | | | 3.6 | 1.9 | 1.1-3.0 | | | | | | | 0.8 | 0.6 | 0.2-1.4 | | | |
| Urinary bladder (incl. in situ) | 48.8 | 11.1 | 9.8-12.7 | 33.0 | 19.8 | 16.7-23.2 | 15.8 | 6.0 | 4.7-7.5 | 12.6 | 2.9 | 2.2-3.7 | 6.2 | 3.7 | 2.4-5.3 | 6.4 | 2.3 | 1.6-3.3 |
| Kidney and renal pelvis | 84.4 | 17.9 | 16.2-19.8 | 50.0 | 25.9 | 22.6-29.5 | 34.4 | 12.6 | 10.7-14.6 | 13.8 | 3.0 | 2.3-3.9 | 8.4 | 4.6 | 3.3-6.3 | 5.4 | 2.0 | 1.3-2.9 |
| Brain and other central nervous system | 19.8 | 4.6 | 3.8-5.7 | 9.6 | 5.0 | 3.6-6.7 | 10.2 | 4.3 | 3.2-5.7 | 9.8 | 2.1 | 1.6-2.8 | 6.0 | 3.1 | 2.0-4.5 | 3.8 | 1.5 | 0.9-2.4 |
| Thyroid | 57.4 | 12.7 | 11.3-14.3 | 7.6 | 3.5 | 2.5-4.9 | 49.8 | 19.9 | 17.4-22.6 | 2.2 | 0.5 | 0.2-0.9 | 1.0 | 0.5 | 0.2-1.2 | 1.2 | 0.5 | 0.2-1.1 |
| Hodgkin lymphoma | 13.4 | 3.1 | 2.4-4.0 | 7.4 | 3.8 | 2.6-5.2 | 6.0 | 2.7 | 1.8-3.8 | 1.8 | 0.4 | 0.2-0.8 | 0.8 | 0.5 | 0.1-1.2 | 1.0 | 0.4 | 0.1-1.0 |
| Non-Hodgkin lymphomas | 84.4 | 19.0 | 17.2-21.0 | 43.0 | 23.3 | 20.1-26.7 | 41.4 | 16.3 | 14.1-18.7 | 21.4 | 4.8 | 3.9-5.9 | 12.6 | 7.3 | 5.5-9.5 | 8.8 | 3.4 | 2.4-4.5 |
| Myeloma | 68.2 | 15.0 | 13.4-16.7 | 30.0 | 16.0 | 13.4-18.9 | 38.2 | 14.5 | 12.5-16.7 | 20.6 | 4.6 | 3.7-5.6 | 10.4 | 5.8 | 4.2-7.7 | 10.2 | 3.8 | 2.8-5.0 |
| Leukemias | 48.8 | 10.9 | 9.6-12.4 | 22.8 | 12.0 | 9.8-14.5 | 26.0 | 10.1 | 8.4-12.1 | 18.8 | 4.1 | 3.3-5.0 | 8.6 | 4.6 | 3.3-6.3 | 10.2 | 3.7 | 2.8-5.0 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Kings (Brooklyn), Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 1607.4 | 361.1 | 353.1-369.2 | 751.4 | 399.4 | 386.2-413.0 | 856.0 | 341.8 | 331.5-352.3 | 479.8 | 115.7 | 111.0-120.5 | 239.0 | 141.6 | 133.3-150.2 | 240.8 | 99.0 | 93.4-104.8 |
| Oral cavity and pharynx | 38.0 | 8.5 | 7.3-9.8 | 24.2 | 12.6 | 10.3-15.1 | 13.8 | 5.5 | 4.3-7.0 | 9.6 | 2.2 | 1.6-2.9 | 7.2 | 3.9 | 2.7-5.4 | 2.4 | 1.0 | 0.5-1.7 |
| Esophagus | 15.2 | 3.4 | 2.7-4.2 | 11.2 | 5.9 | 4.4-7.8 | 4.0 | 1.6 | 1.0-2.5 | 9.6 | 2.3 | 1.7-3.0 | 7.8 | 4.4 | 3.1-6.1 | 1.8 | 0.7 | 0.3-1.4 |
| Stomach | 49.8 | 11.5 | 10.1-13.0 | 30.2 | 16.5 | 13.8-19.4 | 19.6 | 7.8 | 6.3-9.6 | 25.0 | 6.1 | 5.0-7.3 | 14.4 | 8.4 | 6.5-10.7 | 10.6 | 4.4 | 3.3-5.7 |
| Colorectal | 135.4 | 31.7 | 29.4-34.3 | 74.2 | 41.5 | 37.2-46.1 | 61.2 | 25.0 | 22.3-28.0 | 45.8 | 11.2 | 9.8-12.8 | 23.4 | 13.7 | 11.2-16.6 | 22.4 | 9.5 | 7.8-11.4 |
| Colon excluding rectum | 94.0 | 22.2 | 20.2-24.3 | 49.6 | 27.8 | 24.3-31.7 | 44.4 | 18.2 | 15.9-20.8 | 38.0 | 9.4 | 8.1-10.9 | 19.0 | 11.5 | 9.2-14.1 | 19.0 | 8.0 | 6.5-9.8 |
| Rectum & rectosigmoid | 41.4 | 9.6 | 8.3-11.0 | 24.6 | 13.7 | 11.3-16.4 | 16.8 | 6.8 | 5.4-8.4 | 7.8 | 1.8 | 1.3-2.5 | 4.4 | 2.3 | 1.4-3.5 | 3.4 | 1.5 | 0.8-2.3 |
| Liver / intrahepatic bile duct | 61.0 | 13.4 | 11.9-15.0 | 42.0 | 21.3 | 18.4-24.5 | 19.0 | 7.4 | 6.0-9.1 | 36.8 | 8.4 | 7.2-9.7 | 25.4 | 13.2 | 10.9-15.7 | 11.4 | 4.7 | 3.5-6.1 |
| Pancreas | 50.4 | 11.7 | 10.3-13.3 | 24.2 | 13.4 | 11.0-16.1 | 26.2 | 10.6 | 8.8-12.6 | 36.8 | 8.8 | 7.6-10.2 | 17.8 | 10.4 | 8.3-12.9 | 19.0 | 7.8 | 6.3-9.5 |
| Larynx | 13.6 | 2.9 | 2.2-3.7 | 9.2 | 4.7 | 3.4-6.3 | 4.4 | 1.6 | 1.0-2.5 | 5.4 | 1.2 | 0.8-1.8 | 5.2 | 2.9 | 1.8-4.2 | 0.2 | 0.1 | 0.0-0.4 |
| Lung and bronchus | 142.4 | 33.4 | 30.9-36.0 | 70.4 | 40.7 | 36.4-45.4 | 72.0 | 28.7 | 25.8-31.9 | 71.2 | 17.3 | 15.5-19.2 | 38.4 | 23.2 | 19.9-26.9 | 32.8 | 13.4 | 11.4-15.6 |
| Melanoma of the skin | 9.4 | 2.3 | 1.7-3.0 | 4.4 | 2.6 | 1.6-3.9 | 5.0 | 2.1 | 1.3-3.1 | 1.4 | 0.4 | 0.1-0.7 | 0.6 | 0.4 | 0.1-1.1 | 0.8 | 0.3 | 0.1-0.9 |
| Female breast | | | | | | | 239.4 | 95.3 | 89.9-100.9 | | | | | | | 36.8 | 15.0 | 12.9-17.4 |
| Cervix uteri | | | | | | | 26.8 | 10.8 | 9.0-12.8 | | | | | | | 7.8 | 3.2 | 2.3-4.4 |
| Corpus uterus and NOS | | | | | | | 69.8 | 27.2 | 24.4-30.3 | | | | | | | 16.4 | 6.6 | 5.2-8.2 |
| Ovary | | | | | | | 23.0 | 9.2 | 7.6-11.0 | | | | | | | 11.2 | 4.4 | 3.3-5.7 |
| Prostate | | | | 189.4 | 98.1 | 91.8-104.8 | | | | | | | 26.6 | 18.5 | 15.4-22.0 | | | |
| Testis | | | | 11.0 | 4.1 | 3.1-5.4 | | | | | | | 0.2 | 0.1 | 0.0-0.5 | | | |
| Urinary bladder (incl. in situ) | 47.8 | 11.5 | 10.1-13.1 | 32.4 | 20.3 | 17.2-23.8 | 15.4 | 6.2 | 4.9-7.7 | 10.2 | 2.7 | 2.0-3.5 | 6.6 | 4.6 | 3.1-6.5 | 3.6 | 1.5 | 0.9-2.4 |
| Kidney and renal pelvis | 55.6 | 12.2 | 10.8-13.8 | 32.2 | 16.6 | 14.1-19.5 | 23.4 | 9.2 | 7.6-11.0 | 7.0 | 1.7 | 1.2-2.4 | 4.0 | 2.4 | 1.4-3.7 | 3.0 | 1.3 | 0.7-2.1 |
| Brain and other central nervous system | 24.4 | 5.2 | 4.3-6.3 | 13.2 | 6.4 | 4.9-8.2 | 11.2 | 4.4 | 3.3-5.7 | 11.2 | 2.5 | 1.9-3.2 | 6.4 | 3.2 | 2.2-4.6 | 4.8 | 1.9 | 1.2-2.9 |
| Thyroid | 71.8 | 15.2 | 13.7-16.9 | 13.2 | 6.1 | 4.7-7.8 | 58.6 | 23.6 | 20.9-26.5 | 0.8 | 0.2 | 0.1-0.5 | 0.2 | 0.1 | 0.0-0.5 | 0.6 | 0.2 | 0.0-0.7 |
| Hodgkin lymphoma | 15.0 | 3.2 | 2.5-4.0 | 8.4 | 3.8 | 2.7-5.2 | 6.6 | 2.6 | 1.8-3.7 | 2.0 | 0.5 | 0.2-0.9 | 1.0 | 0.5 | 0.2-1.2 | 1.0 | 0.4 | 0.1-1.0 |
| Non-Hodgkin lymphomas | 75.8 | 17.0 | 15.3-18.8 | 41.2 | 20.8 | 17.9-23.9 | 34.6 | 13.9 | 11.9-16.2 | 16.0 | 3.9 | 3.1-4.9 | 9.2 | 5.4 | 3.9-7.2 | 6.8 | 2.8 | 1.9-3.9 |
| Myeloma | 33.6 | 7.7 | 6.6-9.0 | 15.8 | 9.0 | 7.1-11.3 | 17.8 | 7.0 | 5.6-8.6 | 12.6 | 3.2 | 2.4-4.0 | 5.2 | 3.3 | 2.1-4.9 | 7.4 | 3.1 | 2.2-4.2 |
| Leukemias | 50.0 | 10.9 | 9.6-12.4 | 29.8 | 14.8 | 12.4-17.5 | 20.2 | 8.0 | 6.5-9.7 | 16.8 | 3.9 | 3.1-4.9 | 9.2 | 5.2 | 3.7-7.0 | 7.6 | 3.1 | 2.2-4.3 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Kings (Brooklyn), Non-Hispanic Origin**

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 10650.8 | 459.6 | 455.6-463.6 | 5096.4 | 504.5 | 498.2-510.8 | 5554.4 | 432.5 | 427.2-437.7 | 3075.6 | 130.5 | 128.4-132.7 | 1476.4 | 152.3 | 148.8-155.9 | 1599.2 | 116.6 | 114.0-119.3 |
| Oral cavity and pharynx | 220.0 | 9.6 | 9.0-10.2 | 143.0 | 14.2 | 13.2-15.3 | 77.0 | 5.9 | 5.3-6.5 | 42.4 | 1.9 | 1.6-2.1 | 30.2 | 3.1 | 2.6-3.7 | 12.2 | 0.9 | 0.7-1.1 |
| Esophagus | 72.4 | 3.0 | 2.7-3.4 | 50.4 | 5.0 | 4.4-5.6 | 22.0 | 1.6 | 1.3-1.9 | 46.6 | 1.9 | 1.7-2.2 | 36.0 | 3.6 | 3.0-4.1 | 10.6 | 0.7 | 0.6-1.0 |
| Stomach | 265.4 | 11.4 | 10.8-12.1 | 153.8 | 15.5 | 14.4-16.6 | 111.6 | 8.4 | 7.7-9.2 | 118.2 | 5.1 | 4.7-5.6 | 68.8 | 7.2 | 6.5-8.1 | 49.4 | 3.6 | 3.1-4.1 |
| Colorectal | 944.2 | 40.7 | 39.5-41.9 | 479.2 | 48.2 | 46.2-50.2 | 465.0 | 35.2 | 33.7-36.7 | 304.4 | 12.9 | 12.2-13.5 | 147.8 | 15.3 | 14.2-16.5 | 156.6 | 11.1 | 10.4-12.0 |
| Colon excluding rectum | 661.6 | 28.4 | 27.4-29.4 | 322.6 | 32.5 | 30.9-34.2 | 339.0 | 25.4 | 24.2-26.7 | 251.0 | 10.5 | 10.0-11.2 | 120.0 | 12.4 | 11.4-13.5 | 131.0 | 9.3 | 8.5-10.0 |
| Rectum & rectosigmoid | 282.6 | 12.3 | 11.6-13.0 | 156.6 | 15.6 | 14.5-16.8 | 126.0 | 9.7 | 9.0-10.5 | 53.4 | 2.3 | 2.0-2.6 | 27.8 | 2.9 | 2.4-3.5 | 25.6 | 1.9 | 1.6-2.3 |
| Liver / intrahepatic bile duct | 225.0 | 9.4 | 8.9-10.0 | 158.8 | 15.3 | 14.3-16.5 | 66.2 | 4.9 | 4.3-5.4 | 142.8 | 6.0 | 5.5-6.5 | 93.0 | 9.1 | 8.3-10.0 | 49.8 | 3.6 | 3.2-4.1 |
| Pancreas | 336.0 | 14.2 | 13.5-14.9 | 157.8 | 15.7 | 14.6-16.9 | 178.2 | 12.9 | 12.1-13.9 | 244.8 | 10.3 | 9.7-10.9 | 115.4 | 11.6 | 10.7-12.7 | 129.4 | 9.2 | 8.5-10.0 |
| Larynx | 61.2 | 2.6 | 2.3-2.9 | 49.8 | 5.0 | 4.3-5.6 | 11.4 | 0.8 | 0.6-1.1 | 17.0 | 0.7 | 0.6-0.9 | 14.2 | 1.5 | 1.2-1.9 | 2.8 | 0.2 | 0.1-0.4 |
| Lung and bronchus | 1105.0 | 46.8 | 45.6-48.1 | 588.6 | 59.4 | 57.3-61.7 | 516.4 | 38.1 | 36.6-39.6 | 588.8 | 24.9 | 24.0-25.8 | 331.8 | 33.9 | 32.3-35.6 | 257.0 | 18.6 | 17.5-19.6 |
| Melanoma of the skin | 225.4 | 9.9 | 9.3-10.5 | 119.2 | 12.3 | 11.3-13.3 | 106.2 | 8.3 | 7.6-9.1 | 20.0 | 0.9 | 0.7-1.1 | 11.2 | 1.2 | 0.9-1.6 | 8.8 | 0.6 | 0.5-0.9 |
| Female breast | | | | | | | 1641.8 | 132.1 | 129.2-135.1 | | | | | | | 287.2 | 21.6 | 20.5-22.8 |
| Cervix uteri | | | | | | | 112.2 | 9.4 | 8.6-10.2 | | | | | | | 30.6 | 2.4 | 2.1-2.9 |
| Corpus uterus and NOS | | | | | | | 456.0 | 34.4 | 32.9-35.8 | | | | | | | 111.6 | 8.0 | 7.4-8.8 |
| Ovary | | | | | | | 148.2 | 11.6 | 10.8-12.5 | | | | | | | 98.6 | 7.3 | 6.7-8.0 |
| Prostate | | | | 1425.4 | 135.3 | 132.1-138.5 | | | | | | | 178.6 | 19.2 | 18.0-20.5 | | | |
| Testis | | | | 57.0 | 5.4 | 4.8-6.1 | | | | | | | 1.2 | 0.1 | 0.0-0.3 | | | |
| Urinary bladder (incl. in situ) | 393.2 | 16.7 | 16.0-17.5 | 300.2 | 30.9 | 29.3-32.5 | 93.0 | 6.9 | 6.2-7.5 | 73.8 | 3.1 | 2.8-3.4 | 51.0 | 5.5 | 4.9-6.3 | 22.8 | 1.5 | 1.3-1.9 |
| Kidney and renal pelvis | 356.6 | 15.3 | 14.6-16.1 | 230.4 | 22.6 | 21.2-23.9 | 126.2 | 9.7 | 8.9-10.5 | 51.8 | 2.2 | 1.9-2.5 | 37.2 | 3.7 | 3.2-4.3 | 14.6 | 1.0 | 0.8-1.3 |
| Brain and other central nervous system | 138.2 | 6.2 | 5.8-6.7 | 77.4 | 7.7 | 6.9-8.5 | 60.8 | 5.1 | 4.5-5.7 | 71.4 | 3.2 | 2.8-3.5 | 39.0 | 3.9 | 3.4-4.5 | 32.4 | 2.6 | 2.2-3.0 |
| Thyroid | 502.6 | 22.8 | 21.9-23.8 | 123.0 | 12.2 | 11.2-13.2 | 379.6 | 32.0 | 30.6-33.5 | 14.6 | 0.6 | 0.5-0.8 | 3.8 | 0.4 | 0.3-0.7 | 10.8 | 0.7 | 0.6-1.0 |
| Hodgkin lymphoma | 71.6 | 3.3 | 3.0-3.7 | 37.6 | 3.7 | 3.2-4.3 | 34.0 | 2.9 | 2.5-3.4 | 4.4 | 0.2 | 0.1-0.3 | 1.8 | 0.2 | 0.1-0.4 | 2.6 | 0.2 | 0.1-0.4 |
| Non-Hodgkin lymphomas | 436.8 | 19.2 | 18.4-20.0 | 231.6 | 23.4 | 22.1-24.9 | 205.2 | 15.8 | 14.9-16.9 | 95.2 | 4.1 | 3.7-4.5 | 48.0 | 5.0 | 4.4-5.7 | 47.2 | 3.4 | 3.0-3.9 |
| Myeloma | 219.8 | 9.4 | 8.8-10.0 | 111.6 | 11.2 | 10.3-12.2 | 108.2 | 8.1 | 7.4-8.8 | 70.8 | 3.0 | 2.6-3.3 | 36.4 | 3.7 | 3.2-4.3 | 34.4 | 2.4 | 2.1-2.8 |
| Leukemias | 306.0 | 13.4 | 12.8-14.1 | 162.4 | 16.6 | 15.5-17.8 | 143.6 | 11.1 | 10.3-12.0 | 116.2 | 5.0 | 4.6-5.5 | 59.6 | 6.2 | 5.5-7.0 | 56.6 | 4.2 | 3.7-4.8 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Kings (Brooklyn), Non-Hispanic Whites

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 5154.2 | 487.2 | 481.1-493.4 | 2456.0 | 505.2 | 496.1-514.4 | 2698.2 | 484.1 | 475.6-492.7 | 1432.0 | 125.7 | 122.7-128.7 | 695.8 | 142.4 | 137.6-147.3 | 736.2 | 114.5 | 110.7-118.5 |
| Oral cavity and pharynx | 98.2 | 9.4 | 8.6-10.3 | 64.6 | 13.6 | 12.1-15.2 | 33.6 | 5.5 | 4.7-6.5 | 15.4 | 1.5 | 1.1-1.8 | 10.4 | 2.3 | 1.7-3.0 | 5.0 | 0.8 | 0.5-1.1 |
| Esophagus | 32.4 | 2.9 | 2.5-3.4 | 24.6 | 5.1 | 4.2-6.1 | 7.8 | 1.1 | 0.8-1.6 | 21.6 | 1.9 | 1.6-2.3 | 16.8 | 3.4 | 2.7-4.2 | 4.8 | 0.7 | 0.4-1.1 |
| Stomach | 101.6 | 9.5 | 8.7-10.4 | 59.0 | 12.3 | 10.9-13.8 | 42.6 | 7.3 | 6.3-8.4 | 47.6 | 4.3 | 3.8-4.9 | 27.6 | 5.8 | 4.9-6.9 | 20.0 | 3.1 | 2.5-3.8 |
| Colorectal | 452.2 | 42.1 | 40.3-43.9 | 235.4 | 48.5 | 45.7-51.4 | 216.8 | 37.0 | 34.7-39.4 | 146.0 | 12.5 | 11.6-13.5 | 69.8 | 14.3 | 12.8-15.9 | 76.2 | 11.1 | 10.0-12.4 |
| Colon excluding rectum | 318.0 | 29.1 | 27.7-30.6 | 163.8 | 33.6 | 31.3-36.0 | 154.2 | 25.7 | 23.8-27.7 | 119.6 | 10.1 | 9.3-11.0 | 56.0 | 11.3 | 10.0-12.7 | 63.6 | 9.2 | 8.1-10.3 |
| Rectum & rectosigmoid | 134.2 | 12.9 | 12.0-14.0 | 71.6 | 14.9 | 13.4-16.6 | 62.6 | 11.3 | 10.0-12.7 | 26.4 | 2.4 | 2.0-2.9 | 13.8 | 3.0 | 2.3-3.8 | 12.6 | 2.0 | 1.5-2.6 |
| Liver / intrahepatic bile duct | 79.2 | 7.3 | 6.6-8.0 | 53.2 | 10.7 | 9.4-12.1 | 26.0 | 4.3 | 3.6-5.2 | 51.4 | 4.5 | 4.0-5.1 | 30.0 | 6.0 | 5.1-7.0 | 21.4 | 3.4 | 2.7-4.1 |
| Pancreas | 170.8 | 15.3 | 14.3-16.4 | 82.6 | 16.9 | 15.3-18.6 | 88.2 | 13.8 | 12.5-15.3 | 125.0 | 10.8 | 10.0-11.8 | 63.6 | 13.0 | 11.6-14.6 | 61.4 | 9.0 | 7.9-10.1 |
| Larynx | 28.6 | 2.7 | 2.2-3.1 | 24.8 | 5.1 | 4.2-6.1 | 3.8 | 0.7 | 0.4-1.1 | 7.4 | 0.7 | 0.5-1.0 | 6.4 | 1.4 | 1.0-2.0 | 1.0 | 0.2 | 0.1-0.5 |
| Lung and bronchus | 544.8 | 49.4 | 47.5-51.3 | 288.6 | 59.7 | 56.6-62.9 | 256.2 | 41.8 | 39.5-44.2 | 298.2 | 26.0 | 24.7-27.4 | 169.0 | 34.6 | 32.2-37.0 | 129.2 | 19.6 | 18.0-21.2 |
| Melanoma of the skin | 204.2 | 19.6 | 18.4-20.8 | 109.0 | 22.8 | 20.9-24.9 | 95.2 | 17.3 | 15.7-19.0 | 16.6 | 1.5 | 1.2-1.9 | 9.0 | 2.0 | 1.4-2.6 | 7.6 | 1.1 | 0.8-1.6 |
| Female breast | | | | | | | 781.6 | 148.8 | 144.0-153.7 | | | | | | | 128.8 | 21.3 | 19.6-23.2 |
| Cervix uteri | | | | | | | 38.0 | 7.8 | 6.7-9.0 | | | | | | | 8.6 | 1.7 | 1.2-2.3 |
| Corpus uterus and NOS | | | | | | | 185.0 | 33.7 | 31.5-36.1 | | | | | | | 34.2 | 5.5 | 4.6-6.4 |
| Ovary | | | | | | | 71.0 | 12.8 | 11.4-14.3 | | | | | | | 47.0 | 7.8 | 6.8-8.9 |
| Prostate | | | | 499.2 | 100.5 | 96.6-104.6 | | | | | | | 57.2 | 11.5 | 10.1-12.9 | | | |
| Testis | | | | 43.0 | 8.5 | 7.4-9.8 | | | | | | | 0.8 | 0.2 | 0.0-0.4 | | | |
| Urinary bladder (incl. in situ) | 271.2 | 24.5 | 23.2-25.9 | 211.0 | 43.5 | 40.9-46.3 | 60.2 | 9.9 | 8.7-11.1 | 46.2 | 3.7 | 3.2-4.2 | 33.2 | 6.6 | 5.6-7.7 | 13.0 | 1.8 | 1.4-2.3 |
| Kidney and renal pelvis | 194.2 | 18.7 | 17.5-19.9 | 130.6 | 27.0 | 24.9-29.2 | 63.6 | 11.5 | 10.2-12.9 | 28.2 | 2.5 | 2.1-2.9 | 20.0 | 4.0 | 3.3-4.9 | 8.2 | 1.2 | 0.9-1.7 |
| Brain and other central nervous system | 82.8 | 8.3 | 7.5-9.1 | 47.2 | 9.9 | 8.6-11.3 | 35.6 | 6.8 | 5.8-8.0 | 43.8 | 4.2 | 3.7-4.8 | 24.0 | 5.0 | 4.1-6.0 | 19.8 | 3.6 | 2.9-4.4 |
| Thyroid | 271.8 | 27.9 | 26.4-29.5 | 66.4 | 14.0 | 12.5-15.6 | 205.4 | 41.5 | 38.9-44.3 | 7.8 | 0.7 | 0.5-0.9 | 2.6 | 0.6 | 0.3-1.0 | 5.2 | 0.7 | 0.5-1.1 |
| Hodgkin lymphoma | 39.8 | 4.0 | 3.4-4.6 | 22.0 | 4.5 | 3.7-5.5 | 17.8 | 3.5 | 2.8-4.3 | 2.0 | 0.2 | 0.1-0.3 | 1.2 | 0.2 | 0.1-0.5 | 0.8 | 0.2 | 0.0-0.4 |
| Non-Hodgkin lymphomas | 251.8 | 24.0 | 22.6-25.4 | 136.0 | 28.6 | 26.4-30.8 | 115.8 | 20.1 | 18.4-21.9 | 50.2 | 4.3 | 3.7-4.9 | 26.6 | 5.4 | 4.5-6.5 | 23.6 | 3.4 | 2.8-4.1 |
| Myeloma | 66.4 | 6.0 | 5.4-6.7 | 37.8 | 7.6 | 6.5-8.8 | 28.6 | 4.8 | 4.0-5.7 | 22.4 | 2.0 | 1.6-2.4 | 13.2 | 2.7 | 2.1-3.5 | 9.2 | 1.4 | 1.0-1.9 |
| Leukemias | 180.4 | 16.8 | 15.7-18.0 | 97.0 | 20.1 | 18.3-22.0 | 83.4 | 14.1 | 12.7-15.6 | 62.6 | 5.5 | 4.9-6.2 | 31.2 | 6.4 | 5.4-7.5 | 31.4 | 4.9 | 4.1-5.8 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Kings (Brooklyn), Non-Hispanic Blacks

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 4001.8 | 429.5 | 423.4-435.6 | 1883.6 | 509.9 | 499.3-520.6 | 2118.2 | 383.3 | 375.8-390.8 | 1285.8 | 139.7 | 136.3-143.3 | 573.0 | 168.4 | 162.1-174.9 | 712.8 | 124.5 | 120.4-128.8 |
| Oral cavity and pharynx | 64.0 | 6.8 | 6.1-7.7 | 38.4 | 10.6 | 9.1-12.2 | 25.6 | 4.5 | 3.7-5.3 | 16.2 | 1.8 | 1.4-2.2 | 11.6 | 3.3 | 2.5-4.3 | 4.6 | 0.8 | 0.5-1.2 |
| Esophagus | 29.4 | 3.1 | 2.6-3.6 | 17.2 | 4.6 | 3.7-5.7 | 12.2 | 2.1 | 1.6-2.7 | 18.0 | 1.8 | 1.5-2.3 | 12.4 | 3.5 | 2.6-4.5 | 5.6 | 0.9 | 0.6-1.3 |
| Stomach | 98.8 | 10.9 | 9.9-11.9 | 53.0 | 15.5 | 13.6-17.5 | 45.8 | 8.3 | 7.2-9.4 | 47.0 | 5.2 | 4.6-6.0 | 27.0 | 8.2 | 6.8-9.7 | 20.0 | 3.5 | 2.8-4.3 |
| Colorectal | 354.2 | 38.4 | 36.6-40.3 | 164.6 | 46.5 | 43.3-49.9 | 189.6 | 33.5 | 31.3-35.8 | 125.6 | 13.7 | 12.6-14.9 | 58.8 | 17.3 | 15.3-19.4 | 66.8 | 11.6 | 10.4-13.0 |
| Colon excluding rectum | 259.8 | 28.1 | 26.6-29.7 | 112.4 | 31.7 | 29.1-34.6 | 147.4 | 25.9 | 24.1-27.9 | 104.0 | 11.3 | 10.3-12.4 | 48.6 | 14.3 | 12.5-16.2 | 55.4 | 9.6 | 8.5-10.8 |
| Rectum & rectosigmoid | 94.4 | 10.3 | 9.4-11.3 | 52.2 | 14.8 | 13.0-16.8 | 42.2 | 7.5 | 6.5-8.7 | 21.6 | 2.4 | 2.0-2.9 | 10.2 | 3.0 | 2.2-4.0 | 11.4 | 2.0 | 1.5-2.7 |
| Liver / intrahepatic bile duct | 77.4 | 7.9 | 7.1-8.7 | 52.2 | 13.5 | 11.9-15.3 | 25.2 | 4.2 | 3.5-5.1 | 57.2 | 6.0 | 5.3-6.8 | 35.6 | 9.6 | 8.2-11.2 | 21.6 | 3.8 | 3.1-4.6 |
| Pancreas | 131.2 | 14.1 | 13.1-15.3 | 56.6 | 15.7 | 13.9-17.7 | 74.6 | 13.0 | 11.7-14.4 | 95.8 | 10.3 | 9.4-11.3 | 40.8 | 11.3 | 9.7-13.0 | 55.0 | 9.6 | 8.5-10.8 |
| Larynx | 26.2 | 2.7 | 2.3-3.2 | 19.2 | 5.4 | 4.3-6.6 | 7.0 | 1.1 | 0.8-1.6 | 8.8 | 1.0 | 0.7-1.3 | 7.0 | 2.1 | 1.4-2.9 | 1.8 | 0.3 | 0.1-0.6 |
| Lung and bronchus | 356.0 | 37.6 | 35.9-39.5 | 175.8 | 48.5 | 45.3-51.9 | 180.2 | 31.1 | 29.1-33.2 | 204.0 | 21.7 | 20.4-23.1 | 107.2 | 30.1 | 27.6-32.9 | 96.8 | 16.8 | 15.3-18.4 |
| Melanoma of the skin | 7.4 | 0.8 | 0.6-1.1 | 3.2 | 0.9 | 0.5-1.5 | 4.2 | 0.8 | 0.5-1.2 | 2.2 | 0.2 | 0.1-0.4 | 1.6 | 0.5 | 0.2-0.9 | 0.6 | 0.1 | 0.0-0.3 |
| Female breast | | | | | | | 662.2 | 123.0 | 118.8-127.5 | | | | | | | 139.4 | 25.0 | 23.1-27.0 |
| Cervix uteri | | | | | | | 54.8 | 10.4 | 9.2-11.8 | | | | | | | 19.4 | 3.5 | 2.8-4.3 |
| Corpus uterus and NOS | | | | | | | 223.6 | 37.9 | 35.6-40.2 | | | | | | | 71.4 | 12.0 | 10.8-13.3 |
| Ovary | | | | | | | 56.4 | 10.2 | 9.0-11.5 | | | | | | | 42.6 | 7.4 | 6.4-8.4 |
| Prostate | | | | 797.2 | 205.2 | 198.7-211.9 | | | | | | | 110.2 | 35.7 | 32.7-38.9 | | | |
| Testis | | | | 6.0 | 1.7 | 1.1-2.4 | | | | | | | 0.2 | 0.0 | 0.0-0.3 | | | |
| Urinary bladder (incl. in situ) | 83.8 | 9.1 | 8.2-10.0 | 57.4 | 16.9 | 14.9-19.1 | 26.4 | 4.6 | 3.8-5.5 | 21.6 | 2.5 | 2.0-3.0 | 13.0 | 4.3 | 3.3-5.5 | 8.6 | 1.5 | 1.0-2.0 |
| Kidney and renal pelvis | 121.8 | 12.9 | 11.8-14.0 | 73.0 | 19.6 | 17.6-21.8 | 48.8 | 8.4 | 7.4-9.6 | 19.2 | 2.0 | 1.6-2.5 | 13.8 | 3.9 | 3.0-4.9 | 5.4 | 0.8 | 0.6-1.3 |
| Brain and other central nervous system | 37.4 | 4.4 | 3.8-5.1 | 20.8 | 5.8 | 4.7-7.1 | 16.6 | 3.5 | 2.7-4.3 | 18.0 | 2.1 | 1.7-2.6 | 10.0 | 2.9 | 2.1-3.8 | 8.0 | 1.5 | 1.1-2.1 |
| Thyroid | 80.8 | 9.4 | 8.5-10.4 | 15.6 | 4.5 | 3.6-5.7 | 65.2 | 13.0 | 11.6-14.5 | 5.2 | 0.6 | 0.4-0.8 | 0.2 | 0.1 | 0.0-0.4 | 5.0 | 0.8 | 0.5-1.3 |
| Hodgkin lymphoma | 24.0 | 2.9 | 2.4-3.5 | 10.6 | 3.0 | 2.2-3.9 | 13.4 | 2.9 | 2.2-3.7 | 2.2 | 0.3 | 0.1-0.5 | 0.6 | 0.2 | 0.0-0.5 | 1.6 | 0.3 | 0.1-0.6 |
| Non-Hodgkin lymphomas | 130.4 | 14.5 | 13.4-15.7 | 62.0 | 17.2 | 15.3-19.3 | 68.4 | 12.7 | 11.3-14.2 | 34.8 | 3.9 | 3.3-4.5 | 15.4 | 4.4 | 3.5-5.6 | 19.4 | 3.5 | 2.8-4.3 |
| Myeloma | 136.2 | 14.7 | 13.6-15.9 | 63.4 | 18.0 | 16.1-20.2 | 72.8 | 12.8 | 11.5-14.2 | 43.6 | 4.7 | 4.1-5.4 | 20.4 | 6.1 | 4.9-7.4 | 23.2 | 3.9 | 3.2-4.7 |
| Leukemias | 90.8 | 10.4 | 9.5-11.5 | 44.0 | 12.8 | 11.1-14.6 | 46.8 | 8.9 | 7.8-10.2 | 42.4 | 4.8 | 4.2-5.5 | 21.8 | 6.3 | 5.1-7.7 | 20.6 | 3.8 | 3.1-4.7 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York (Manhattan), Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|------------|----------------|------------------|-------------|----------------|------------------|-----------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 1649.2 | 349.4 | 341.8-357.2 | 799.2 | 411.4 | 398.6-424.5 | 850.0 | 315.6 | 305.9-325.5 | 494.2 | 102.4 | 98.4-106.6 | 240.2 | 129.3 | 122.0-136.9 | 254.0 | 86.0 | 81.2-91.0 |
| Oral cavity and pharynx | 40.8 | 8.7 | 7.5-10.0 | 28.4 | 14.5 | 12.2-17.1 | 12.4 | 4.6 | 3.5-6.0 | 7.2 | 1.5 | 1.0-2.1 | 5.4 | 2.9 | 1.9-4.2 | 1.8 | 0.6 | 0.2-1.1 |
| Esophagus | 11.8 | 2.4 | 1.8-3.1 | 7.2 | 3.8 | 2.7-5.3 | 4.6 | 1.5 | 0.9-2.2 | 10.2 | 2.1 | 1.6-2.8 | 7.2 | 3.8 | 2.7-5.3 | 3.0 | 1.0 | 0.5-1.6 |
| Stomach | 42.8 | 9.2 | 8.0-10.6 | 22.0 | 11.9 | 9.7-14.3 | 20.8 | 7.7 | 6.2-9.4 | 17.0 | 3.5 | 2.8-4.4 | 10.8 | 5.8 | 4.4-7.6 | 6.2 | 2.0 | 1.4-2.9 |
| Colorectal | 154.2 | 32.4 | 30.1-34.8 | 80.6 | 41.7 | 37.7-46.1 | 73.6 | 25.9 | 23.3-28.8 | 48.2 | 10.0 | 8.8-11.4 | 24.6 | 13.1 | 10.9-15.7 | 23.6 | 8.0 | 6.6-9.6 |
| Colon excluding rectum | 105.8 | 22.3 | 20.4-24.3 | 52.2 | 27.4 | 24.1-30.9 | 53.6 | 18.9 | 16.6-21.3 | 36.8 | 7.6 | 6.5-8.8 | 18.6 | 9.8 | 7.9-12.0 | 18.2 | 6.1 | 4.8-7.5 |
| Rectum & rectosigmoid | 48.4 | 10.2 | 8.9-11.5 | 28.4 | 14.4 | 12.1-17.0 | 20.0 | 7.1 | 5.7-8.7 | 11.4 | 2.5 | 1.9-3.2 | 6.0 | 3.3 | 2.2-4.7 | 5.4 | 1.9 | 1.2-2.8 |
| Liver / intrahepatic bile duct | 64.6 | 13.2 | 11.8-14.7 | 41.8 | 20.9 | 18.1-23.9 | 22.8 | 7.7 | 6.3-9.3 | 37.2 | 7.6 | 6.6-8.8 | 23.2 | 12.1 | 10.0-14.6 | 14.0 | 4.7 | 3.6-6.0 |
| Pancreas | 51.4 | 10.6 | 9.4-12.0 | 25.2 | 13.2 | 11.0-15.7 | 26.2 | 8.8 | 7.4-10.6 | 41.2 | 8.4 | 7.3-9.7 | 17.8 | 9.4 | 7.5-11.6 | 23.4 | 7.7 | 6.4-9.3 |
| Larynx | 13.2 | 2.7 | 2.1-3.5 | 10.0 | 5.1 | 3.7-6.7 | 3.2 | 1.1 | 0.7-1.9 | 4.0 | 0.8 | 0.5-1.2 | 3.2 | 1.6 | 0.9-2.7 | 0.8 | 0.2 | 0.1-0.7 |
| Lung and bronchus | 143.4 | 29.7 | 27.5-32.0 | 73.4 | 38.7 | 34.8-43.0 | 70.0 | 23.9 | 21.5-26.6 | 74.8 | 15.4 | 13.9-17.1 | 38.8 | 20.6 | 17.8-23.8 | 36.0 | 12.1 | 10.3-14.0 |
| Melanoma of the skin | 14.4 | 3.1 | 2.4-4.0 | 8.8 | 4.6 | 3.3-6.2 | 5.6 | 2.1 | 1.4-3.1 | 2.6 | 0.5 | 0.3-0.9 | 2.0 | 1.1 | 0.5-2.0 | 0.6 | 0.2 | 0.0-0.6 |
| Female breast | | | | | | | 255.8 | 97.7 | 92.3-103.4 | | | | | | | 37.6 | 13.5 | 11.6-15.6 |
| Cervix uteri | | | | | | | 17.2 | 7.0 | 5.6-8.7 | | | | | | | 3.8 | 1.5 | 0.9-2.3 |
| Corpus uterus and NOS | | | | | | | 67.2 | 25.1 | 22.4-28.0 | | | | | | | 11.6 | 4.0 | 3.0-5.2 |
| Ovary | | | | | | | 22.0 | 8.7 | 7.1-10.5 | | | | | | | 14.0 | 4.9 | 3.8-6.2 |
| Prostate | | | | 228.6 | 114.5 | 107.9-121.4 | | | | | | | 28.2 | 15.9 | 13.4-18.8 | | | |
| Testis | | | | 9.8 | 4.1 | 3.0-5.5 | | | | | | | 0.4 | 0.2 | 0.0-0.7 | | | |
| Urinary bladder (incl. in situ) | 51.2 | 10.5 | 9.3-11.9 | 38.2 | 20.6 | 17.8-23.8 | 13.0 | 4.3 | 3.3-5.6 | 11.8 | 2.4 | 1.8-3.1 | 7.4 | 4.2 | 3.0-5.8 | 4.4 | 1.4 | 0.9-2.1 |
| Kidney and renal pelvis | 48.2 | 10.5 | 9.2-11.9 | 28.8 | 14.9 | 12.5-17.6 | 19.4 | 7.5 | 6.1-9.2 | 8.8 | 1.8 | 1.3-2.5 | 4.6 | 2.5 | 1.6-3.8 | 4.2 | 1.4 | 0.8-2.2 |
| Brain and other central nervous system | 23.2 | 5.3 | 4.4-6.4 | 13.2 | 6.8 | 5.2-8.6 | 10.0 | 4.1 | 3.0-5.5 | 12.2 | 2.6 | 2.0-3.4 | 6.8 | 3.6 | 2.5-5.0 | 5.4 | 1.8 | 1.2-2.7 |
| Thyroid | 57.0 | 12.9 | 11.4-14.5 | 14.0 | 7.0 | 5.4-8.9 | 43.0 | 18.4 | 15.9-21.1 | 2.4 | 0.5 | 0.3-0.9 | 0.4 | 0.2 | 0.0-0.9 | 2.0 | 0.7 | 0.3-1.3 |
| Hodgkin lymphoma | 8.6 | 2.0 | 1.4-2.7 | 5.0 | 2.5 | 1.6-3.7 | 3.6 | 1.6 | 0.9-2.5 | 1.2 | 0.3 | 0.1-0.6 | 0.6 | 0.3 | 0.1-1.0 | 0.6 | 0.2 | 0.0-0.7 |
| Non-Hodgkin lymphomas | 80.4 | 17.4 | 15.7-19.2 | 41.4 | 21.7 | 18.8-24.9 | 39.0 | 14.4 | 12.4-16.6 | 18.8 | 4.0 | 3.2-4.9 | 10.8 | 6.0 | 4.5-7.8 | 8.0 | 2.8 | 2.0-3.8 |
| Myeloma | 36.2 | 7.6 | 6.6-8.9 | 19.4 | 10.2 | 8.2-12.4 | 16.8 | 5.9 | 4.7-7.3 | 11.2 | 2.3 | 1.7-3.0 | 6.2 | 3.3 | 2.2-4.7 | 5.0 | 1.6 | 1.0-2.4 |
| Leukemias | 45.6 | 10.0 | 8.7-11.4 | 27.6 | 14.7 | 12.3-17.4 | 18.0 | 6.8 | 5.4-8.4 | 21.6 | 4.5 | 3.7-5.5 | 10.4 | 5.6 | 4.1-7.3 | 11.2 | 3.7 | 2.8-4.9 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York (Manhattan), Non-Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 7113.6 | 480.8 | 475.7-486.0 | 3402.6 | 507.9 | 500.1-515.7 | 3711.0 | 468.4 | 461.4-475.5 | 1892.4 | 121.7 | 119.2-124.2 | 909.6 | 137.1 | 133.1-141.2 | 982.8 | 111.5 | 108.3-114.8 |
| Oral cavity and pharynx | 185.2 | 12.5 | 11.6-13.3 | 123.4 | 18.3 | 16.9-19.8 | 61.8 | 7.6 | 6.8-8.5 | 34.6 | 2.2 | 1.9-2.6 | 23.4 | 3.5 | 2.9-4.2 | 11.2 | 1.2 | 0.9-1.6 |
| Esophagus | 51.2 | 3.3 | 2.9-3.8 | 33.6 | 5.0 | 4.2-5.8 | 17.6 | 2.0 | 1.6-2.5 | 36.8 | 2.3 | 2.0-2.7 | 24.8 | 3.7 | 3.1-4.4 | 12.0 | 1.3 | 1.0-1.7 |
| Stomach | 106.6 | 7.1 | 6.5-7.7 | 62.0 | 9.3 | 8.3-10.4 | 44.6 | 5.3 | 4.6-6.1 | 51.4 | 3.3 | 2.9-3.7 | 32.0 | 4.7 | 4.0-5.5 | 19.4 | 2.2 | 1.8-2.8 |
| Colorectal | 489.4 | 33.2 | 31.9-34.6 | 253.4 | 38.5 | 36.4-40.8 | 236.0 | 29.0 | 27.3-30.8 | 156.0 | 10.1 | 9.4-10.8 | 78.6 | 12.0 | 10.8-13.3 | 77.4 | 8.6 | 7.8-9.6 |
| Colon excluding rectum | 337.2 | 22.6 | 21.5-23.8 | 168.0 | 25.7 | 23.9-27.5 | 169.2 | 20.3 | 18.9-21.8 | 120.2 | 7.7 | 7.1-8.4 | 59.2 | 9.0 | 8.0-10.1 | 61.0 | 6.8 | 6.0-7.6 |
| Rectum & rectosigmoid | 152.2 | 10.6 | 9.8-11.4 | 85.4 | 12.9 | 11.7-14.2 | 66.8 | 8.7 | 7.8-9.7 | 35.8 | 2.4 | 2.0-2.8 | 19.4 | 3.0 | 2.4-3.7 | 16.4 | 1.8 | 1.4-2.3 |
| Liver / intrahepatic bile duct | 143.6 | 9.4 | 8.7-10.1 | 97.8 | 14.3 | 13.0-15.6 | 45.8 | 5.3 | 4.6-6.0 | 86.6 | 5.5 | 5.0-6.1 | 57.2 | 8.4 | 7.5-9.5 | 29.4 | 3.2 | 2.7-3.8 |
| Pancreas | 212.4 | 13.8 | 13.0-14.7 | 105.4 | 15.9 | 14.5-17.3 | 107.0 | 12.2 | 11.2-13.4 | 165.2 | 10.5 | 9.8-11.3 | 81.4 | 12.2 | 11.1-13.5 | 83.8 | 9.2 | 8.4-10.2 |
| Larynx | 33.4 | 2.2 | 1.9-2.6 | 26.8 | 3.9 | 3.3-4.6 | 6.6 | 0.8 | 0.6-1.2 | 10.4 | 0.6 | 0.5-0.8 | 8.0 | 1.1 | 0.8-1.6 | 2.4 | 0.3 | 0.1-0.5 |
| Lung and bronchus | 763.6 | 49.5 | 47.9-51.1 | 331.2 | 49.3 | 47.0-51.8 | 432.4 | 49.7 | 47.6-51.9 | 381.8 | 24.3 | 23.2-25.4 | 175.2 | 26.3 | 24.5-28.1 | 206.6 | 22.8 | 21.4-24.3 |
| Melanoma of the skin | 337.8 | 22.6 | 21.6-23.8 | 195.0 | 29.4 | 27.5-31.3 | 142.8 | 17.6 | 16.3-19.0 | 21.2 | 1.3 | 1.1-1.6 | 13.6 | 2.0 | 1.6-2.6 | 7.6 | 0.8 | 0.6-1.2 |
| Female breast | | | | | | | 1165.8 | 155.4 | 151.3-159.6 | | | | | | | 152.8 | 18.1 | 16.8-19.5 |
| Cervix uteri | | | | | | | 51.0 | 7.1 | 6.2-8.1 | | | | | | | 13.4 | 1.7 | 1.3-2.2 |
| Corpus uterus and NOS | | | | | | | 251.0 | 31.4 | 29.6-33.3 | | | | | | | 51.0 | 5.8 | 5.1-6.6 |
| Ovary | | | | | | | 90.2 | 11.6 | 10.5-12.7 | | | | | | | 60.2 | 7.1 | 6.3-8.0 |
| Prostate | | | | 904.4 | 130.1 | 126.3-133.9 | | | | | | | 114.6 | 17.3 | 15.9-18.8 | | | |
| Testis | | | | 55.8 | 8.0 | 7.0-9.1 | | | | | | | 0.4 | 0.0 | 0.0-0.2 | | | |
| Urinary bladder (incl. in situ) | 305.8 | 19.8 | 18.8-20.8 | 221.8 | 33.6 | 31.6-35.6 | 84.0 | 9.6 | 8.7-10.6 | 56.8 | 3.4 | 3.0-3.9 | 38.6 | 5.8 | 5.0-6.7 | 18.2 | 1.8 | 1.5-2.3 |
| Kidney and renal pelvis | 191.4 | 13.2 | 12.3-14.1 | 125.2 | 18.8 | 17.4-20.4 | 66.2 | 8.4 | 7.5-9.5 | 30.8 | 1.9 | 1.6-2.3 | 20.2 | 3.0 | 2.4-3.7 | 10.6 | 1.1 | 0.8-1.5 |
| Brain and other central nervous system | 97.6 | 7.4 | 6.7-8.1 | 55.4 | 9.2 | 8.1-10.4 | 42.2 | 5.8 | 5.0-6.7 | 49.8 | 3.5 | 3.0-3.9 | 28.2 | 4.3 | 3.6-5.1 | 21.6 | 2.8 | 2.3-3.4 |
| Thyroid | 282.2 | 20.4 | 19.3-21.5 | 77.8 | 11.7 | 10.6-13.0 | 204.4 | 28.4 | 26.6-30.3 | 5.0 | 0.3 | 0.2-0.5 | 3.0 | 0.4 | 0.2-0.8 | 2.0 | 0.2 | 0.1-0.4 |
| Hodgkin lymphoma | 41.2 | 3.0 | 2.6-3.5 | 19.8 | 3.1 | 2.5-3.8 | 21.4 | 2.9 | 2.4-3.6 | 3.2 | 0.2 | 0.1-0.3 | 2.0 | 0.3 | 0.1-0.5 | 1.2 | 0.1 | 0.0-0.3 |
| Non-Hodgkin lymphomas | 356.0 | 24.1 | 23.0-25.3 | 188.2 | 28.8 | 26.9-30.7 | 167.8 | 20.4 | 19.0-21.9 | 67.2 | 4.3 | 3.9-4.8 | 36.4 | 5.5 | 4.7-6.4 | 30.8 | 3.4 | 2.8-4.0 |
| Myeloma | 122.6 | 8.0 | 7.4-8.7 | 65.2 | 9.7 | 8.7-10.9 | 57.4 | 6.7 | 6.0-7.6 | 38.6 | 2.5 | 2.1-2.9 | 20.0 | 3.0 | 2.5-3.7 | 18.6 | 2.1 | 1.7-2.6 |
| Leukemias | 227.8 | 15.7 | 14.8-16.7 | 131.6 | 20.5 | 18.9-22.2 | 96.2 | 11.9 | 10.8-13.1 | 79.8 | 5.3 | 4.8-5.9 | 41.8 | 6.7 | 5.8-7.7 | 38.0 | 4.3 | 3.7-5.0 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York (Manhattan), Non-Hispanic Whites

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 4832.2 | 482.6 | 476.3-489.0 | 2342.0 | 497.6 | 488.4-506.9 | 2490.2 | 480.8 | 471.9-489.8 | 1151.8 | 107.3 | 104.5-110.1 | 557.2 | 117.4 | 113.0-121.9 | 594.6 | 100.4 | 96.7-104.3 |
| Oral cavity and pharynx | 129.4 | 12.7 | 11.7-13.7 | 89.0 | 18.6 | 16.9-20.5 | 40.4 | 7.4 | 6.4-8.6 | 21.6 | 2.0 | 1.6-2.4 | 15.0 | 3.1 | 2.4-3.9 | 6.6 | 1.0 | 0.7-1.5 |
| Esophagus | 33.4 | 3.2 | 2.7-3.8 | 22.0 | 4.7 | 3.8-5.7 | 11.4 | 1.9 | 1.5-2.6 | 23.8 | 2.2 | 1.8-2.6 | 16.2 | 3.3 | 2.6-4.2 | 7.6 | 1.2 | 0.9-1.7 |
| Stomach | 46.6 | 4.6 | 4.0-5.2 | 24.4 | 5.2 | 4.3-6.3 | 22.2 | 4.1 | 3.3-5.0 | 24.4 | 2.3 | 1.9-2.8 | 15.6 | 3.2 | 2.5-4.0 | 8.8 | 1.6 | 1.1-2.2 |
| Colorectal | 301.6 | 30.4 | 28.9-32.1 | 156.8 | 34.1 | 31.7-36.7 | 144.8 | 27.2 | 25.2-29.4 | 86.4 | 8.1 | 7.4-9.0 | 43.8 | 9.3 | 8.1-10.7 | 42.6 | 7.2 | 6.2-8.3 |
| Colon excluding rectum | 211.4 | 21.0 | 19.7-22.3 | 107.2 | 23.3 | 21.3-25.5 | 104.2 | 19.0 | 17.3-20.8 | 65.4 | 6.1 | 5.4-6.8 | 33.8 | 7.1 | 6.1-8.3 | 31.6 | 5.3 | 4.5-6.3 |
| Rectum & rectosigmoid | 90.2 | 9.5 | 8.6-10.4 | 49.6 | 10.8 | 9.5-12.3 | 40.6 | 8.2 | 7.1-9.5 | 21.0 | 2.1 | 1.7-2.5 | 10.0 | 2.2 | 1.6-2.9 | 11.0 | 1.9 | 1.4-2.5 |
| Liver / intrahepatic bile duct | 64.6 | 6.3 | 5.6-7.0 | 43.0 | 9.0 | 7.8-10.4 | 21.6 | 3.7 | 3.0-4.6 | 36.8 | 3.4 | 2.9-4.0 | 24.4 | 5.1 | 4.2-6.1 | 12.4 | 2.0 | 1.5-2.6 |
| Pancreas | 137.0 | 13.1 | 12.1-14.1 | 72.4 | 15.4 | 13.8-17.1 | 64.6 | 11.2 | 9.9-12.6 | 107.0 | 9.9 | 9.1-10.8 | 58.4 | 12.4 | 11.0-13.9 | 48.6 | 8.0 | 7.0-9.2 |
| Larynx | 18.6 | 1.8 | 1.5-2.3 | 15.6 | 3.2 | 2.6-4.1 | 3.0 | 0.6 | 0.3-1.0 | 5.4 | 0.5 | 0.3-0.7 | 4.2 | 0.9 | 0.5-1.4 | 1.2 | 0.2 | 0.1-0.5 |
| Lung and bronchus | 477.2 | 44.8 | 43.0-46.6 | 193.0 | 40.3 | 37.8-43.0 | 284.2 | 48.7 | 46.2-51.4 | 223.0 | 20.5 | 19.3-21.8 | 91.8 | 19.4 | 17.6-21.3 | 131.2 | 21.4 | 19.7-23.1 |
| Melanoma of the skin | 315.6 | 31.5 | 29.9-33.2 | 185.0 | 39.7 | 37.1-42.4 | 130.6 | 25.0 | 23.0-27.1 | 19.0 | 1.8 | 1.4-2.2 | 12.8 | 2.7 | 2.1-3.5 | 6.2 | 1.0 | 0.7-1.5 |
| Female breast | | | | | | | 810.2 | 166.5 | 161.2-172.0 | | | | | | | 89.4 | 15.7 | 14.2-17.3 |
| Cervix uteri | | | | | | | 25.6 | 5.7 | 4.8-6.9 | | | | | | | 5.6 | 1.0 | 0.7-1.5 |
| Corpus uteri and NOS | | | | | | | 159.2 | 30.3 | 28.1-32.6 | | | | | | | 29.4 | 5.1 | 4.3-6.0 |
| Ovary | | | | | | | 62.0 | 12.0 | 10.6-13.5 | | | | | | | 43.0 | 7.7 | 6.6-8.8 |
| Prostate | | | | 603.6 | 122.6 | 118.2-127.1 | | | | | | | 71.0 | 14.9 | 13.3-16.5 | | | |
| Testis | | | | 46.4 | 10.0 | 8.6-11.4 | | | | | | | 0.0 | 0.0 | 0.0-0.3 | | | |
| Urinary bladder (incl. in situ) | 241.6 | 22.8 | 21.5-24.1 | 179.0 | 37.9 | 35.4-40.5 | 62.6 | 10.9 | 9.7-12.2 | 39.0 | 3.4 | 2.9-3.9 | 27.6 | 5.7 | 4.8-6.8 | 11.4 | 1.7 | 1.3-2.3 |
| Kidney and renal pelvis | 119.6 | 12.2 | 11.2-13.3 | 83.6 | 18.1 | 16.4-20.0 | 36.0 | 6.9 | 5.8-8.1 | 18.8 | 1.7 | 1.4-2.2 | 13.8 | 2.9 | 2.2-3.7 | 5.0 | 0.8 | 0.5-1.2 |
| Brain and other central nervous system | 76.4 | 8.8 | 7.9-9.9 | 44.0 | 10.7 | 9.2-12.4 | 32.4 | 7.2 | 6.0-8.6 | 38.8 | 4.0 | 3.4-4.7 | 22.4 | 4.9 | 4.0-5.9 | 16.4 | 3.3 | 2.6-4.2 |
| Thyroid | 182.2 | 20.0 | 18.7-21.4 | 54.0 | 11.8 | 10.4-13.4 | 128.2 | 28.2 | 25.9-30.7 | 3.0 | 0.3 | 0.1-0.5 | 1.4 | 0.3 | 0.1-0.6 | 1.6 | 0.3 | 0.1-0.6 |
| Hodgkin lymphoma | 27.8 | 3.2 | 2.6-3.8 | 13.2 | 3.1 | 2.3-4.1 | 14.6 | 3.2 | 2.4-4.1 | 2.6 | 0.2 | 0.1-0.4 | 1.8 | 0.3 | 0.2-0.7 | 0.8 | 0.1 | 0.0-0.4 |
| Non-Hodgkin lymphomas | 263.6 | 26.1 | 24.7-27.7 | 141.4 | 30.6 | 28.4-33.1 | 122.2 | 22.3 | 20.4-24.2 | 47.8 | 4.4 | 3.9-5.0 | 27.0 | 5.7 | 4.8-6.8 | 20.8 | 3.4 | 2.7-4.1 |
| Myeloma | 75.8 | 7.2 | 6.5-8.0 | 44.0 | 9.2 | 8.0-10.5 | 31.8 | 5.5 | 4.6-6.4 | 23.6 | 2.2 | 1.8-2.7 | 13.4 | 2.8 | 2.2-3.6 | 10.2 | 1.7 | 1.3-2.4 |
| Leukemias | 171.6 | 17.4 | 16.2-18.7 | 102.8 | 23.0 | 20.9-25.1 | 68.8 | 12.7 | 11.3-14.3 | 55.2 | 5.2 | 4.6-5.9 | 29.8 | 6.5 | 5.5-7.7 | 25.4 | 4.1 | 3.4-5.0 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
New York (Manhattan), Non-Hispanic Blacks

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 1331.4 | 512.6 | 500.1-525.3 | 634.6 | 597.6 | 576.4-619.4 | 696.8 | 466.7 | 450.9-483.0 | 487.0 | 185.2 | 177.8-192.9 | 220.6 | 225.0 | 211.6-239.1 | 266.4 | 165.1 | 156.2-174.5 |
| Oral cavity and pharynx | 25.8 | 10.0 | 8.3-11.9 | 15.6 | 14.6 | 11.5-18.4 | 10.2 | 6.9 | 5.1-9.2 | 7.8 | 2.9 | 2.1-4.0 | 4.4 | 4.2 | 2.6-6.5 | 3.4 | 2.1 | 1.2-3.5 |
| Esophagus | 10.6 | 3.9 | 2.9-5.2 | 5.8 | 5.5 | 3.6-8.0 | 4.8 | 3.0 | 1.9-4.5 | 7.4 | 2.8 | 2.0-3.9 | 4.0 | 4.3 | 2.6-6.8 | 3.4 | 2.1 | 1.2-3.4 |
| Stomach | 25.8 | 10.0 | 8.3-11.9 | 14.6 | 14.2 | 11.1-18.0 | 11.2 | 7.2 | 5.4-9.5 | 12.2 | 4.5 | 3.5-5.9 | 6.8 | 6.4 | 4.4-9.1 | 5.4 | 3.1 | 2.1-4.7 |
| Colorectal | 108.6 | 42.1 | 38.6-45.9 | 57.2 | 56.4 | 49.8-63.5 | 51.4 | 33.3 | 29.3-37.9 | 46.4 | 17.7 | 15.5-20.2 | 23.2 | 24.3 | 20.0-29.3 | 23.2 | 13.7 | 11.3-16.5 |
| Colon excluding rectum | 77.6 | 29.9 | 26.9-33.1 | 39.6 | 39.4 | 34.0-45.5 | 38.0 | 24.2 | 20.8-28.0 | 36.0 | 13.8 | 11.8-16.0 | 16.6 | 17.3 | 13.7-21.6 | 19.4 | 11.5 | 9.3-14.1 |
| Rectum & rectosigmoid | 31.0 | 12.2 | 10.3-14.4 | 17.6 | 16.9 | 13.5-21.0 | 13.4 | 9.2 | 7.1-11.8 | 10.4 | 3.9 | 2.9-5.2 | 6.6 | 7.0 | 4.8-9.9 | 3.8 | 2.2 | 1.3-3.6 |
| Liver / intrahepatic bile duct | 42.6 | 15.5 | 13.5-17.8 | 30.4 | 27.6 | 23.3-32.5 | 12.2 | 7.5 | 5.7-9.7 | 26.8 | 9.9 | 8.3-11.8 | 17.6 | 17.1 | 13.6-21.2 | 9.2 | 5.6 | 4.1-7.6 |
| Pancreas | 50.8 | 19.3 | 17.0-21.9 | 23.2 | 22.7 | 18.6-27.3 | 27.6 | 17.0 | 14.2-20.2 | 41.4 | 15.6 | 13.5-17.9 | 16.2 | 16.1 | 12.7-20.1 | 25.2 | 15.2 | 12.7-18.2 |
| Larynx | 10.4 | 3.8 | 2.9-5.1 | 7.8 | 7.1 | 5.0-9.9 | 2.6 | 1.7 | 0.9-3.1 | 4.4 | 1.6 | 1.0-2.4 | 3.2 | 3.0 | 1.7-4.9 | 1.2 | 0.7 | 0.3-1.7 |
| Lung and bronchus | 167.4 | 63.1 | 58.9-67.6 | 75.2 | 72.6 | 65.3-80.6 | 92.2 | 57.1 | 52.0-62.7 | 101.4 | 38.1 | 34.9-41.7 | 49.2 | 48.7 | 42.7-55.4 | 52.2 | 31.9 | 28.1-36.1 |
| Melanoma of the skin | 3.4 | 1.2 | 0.7-2.0 | 1.0 | 0.9 | 0.3-2.3 | 2.4 | 1.6 | 0.8-2.9 | 0.6 | 0.2 | 0.0-0.7 | 0.0 | 0.0 | 0.0-0.8 | 0.6 | 0.3 | 0.1-1.2 |
| Female breast | | | | | | | 205.2 | 144.1 | 135.2-153.6 | | | | | | | 47.2 | 30.7 | 26.7-35.1 |
| Cervix uteri | | | | | | | 16.2 | 11.5 | 9.1-14.4 | | | | | | | 5.8 | 4.0 | 2.7-5.9 |
| Corpus uterus and NOS | | | | | | | 59.2 | 38.2 | 33.9-43.0 | | | | | | | 18.4 | 11.2 | 9.0-13.8 |
| Ovary | | | | | | | 14.4 | 10.0 | 7.8-12.7 | | | | | | | 11.0 | 6.8 | 5.1-9.0 |
| Prostate | | | | 221.4 | 199.5 | 187.6-212.0 | | | | | | | 35.4 | 38.8 | 33.2-45.0 | | | |
| Testis | | | | 4.2 | 3.7 | 2.2-5.8 | | | | | | | 0.2 | 0.2 | 0.0-1.3 | | | |
| Urinary bladder (incl. in situ) | 36.0 | 13.8 | 11.8-16.0 | 22.0 | 22.8 | 18.7-27.6 | 14.0 | 8.7 | 6.8-11.2 | 12.4 | 4.6 | 3.5-6.0 | 7.2 | 8.0 | 5.6-11.1 | 5.2 | 2.8 | 1.8-4.3 |
| Kidney and renal pelvis | 48.8 | 18.8 | 16.5-21.4 | 29.0 | 26.1 | 21.9-30.9 | 19.8 | 13.2 | 10.7-16.2 | 7.4 | 2.8 | 2.0-3.9 | 3.2 | 3.2 | 1.8-5.3 | 4.2 | 2.5 | 1.5-3.9 |
| Brain and other central nervous system | 9.4 | 3.9 | 2.9-5.3 | 4.6 | 4.6 | 2.8-6.9 | 4.8 | 3.4 | 2.1-5.2 | 5.4 | 2.1 | 1.4-3.1 | 3.2 | 3.0 | 1.7-4.9 | 2.2 | 1.4 | 0.7-2.7 |
| Thyroid | 24.2 | 10.0 | 8.3-12.0 | 4.2 | 3.5 | 2.2-5.5 | 20.0 | 15.8 | 12.7-19.3 | 0.8 | 0.3 | 0.1-0.9 | 0.6 | 0.6 | 0.1-1.8 | 0.2 | 0.1 | 0.0-0.9 |
| Hodgkin lymphoma | 9.4 | 4.0 | 2.9-5.4 | 4.8 | 4.7 | 2.9-7.0 | 4.6 | 3.6 | 2.2-5.4 | 0.4 | 0.2 | 0.0-0.6 | 0.2 | 0.2 | 0.0-1.1 | 0.2 | 0.1 | 0.0-0.9 |
| Non-Hodgkin lymphomas | 51.4 | 20.8 | 18.2-23.6 | 25.4 | 24.2 | 20.1-29.0 | 26.0 | 18.2 | 15.1-21.8 | 10.8 | 4.4 | 3.3-5.8 | 5.4 | 5.7 | 3.7-8.3 | 5.4 | 3.5 | 2.3-5.3 |
| Myeloma | 36.8 | 14.3 | 12.3-16.6 | 15.8 | 15.5 | 12.2-19.5 | 21.0 | 13.5 | 10.9-16.4 | 12.4 | 4.7 | 3.6-6.1 | 5.4 | 5.6 | 3.6-8.2 | 7.0 | 4.2 | 2.9-5.9 |
| Leukemias | 28.4 | 11.4 | 9.6-13.5 | 13.6 | 13.6 | 10.5-17.4 | 14.8 | 10.0 | 7.7-12.8 | 13.8 | 5.6 | 4.3-7.2 | 6.0 | 6.5 | 4.3-9.3 | 7.8 | 5.2 | 3.7-7.3 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Queens, Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-----------|----------------|------------------|-------------|----------------|------------------|-----------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 2004.2 | 338.1 | 331.3-344.9 | 935.6 | 361.5 | 350.6-372.6 | 1068.6 | 331.6 | 322.7-340.8 | 485.2 | 88.6 | 85.0-92.3 | 244.2 | 108.3 | 102.0-114.9 | 241.0 | 76.5 | 72.2-81.0 |
| Oral cavity and pharynx | 37.4 | 6.2 | 5.3-7.1 | 24.4 | 8.9 | 7.3-10.8 | 13.0 | 4.0 | 3.1-5.1 | 4.6 | 0.8 | 0.5-1.2 | 3.8 | 1.5 | 0.9-2.4 | 0.8 | 0.2 | 0.1-0.6 |
| Esophagus | 11.4 | 2.0 | 1.5-2.6 | 8.2 | 3.2 | 2.2-4.4 | 3.2 | 1.0 | 0.6-1.7 | 6.4 | 1.2 | 0.8-1.6 | 4.8 | 2.0 | 1.3-3.1 | 1.6 | 0.5 | 0.2-1.0 |
| Stomach | 67.6 | 11.8 | 10.6-13.2 | 34.4 | 13.6 | 11.5-15.9 | 33.2 | 10.7 | 9.1-12.5 | 27.8 | 4.9 | 4.1-5.8 | 14.2 | 5.7 | 4.4-7.3 | 13.6 | 4.3 | 3.4-5.5 |
| Colorectal | 178.0 | 30.7 | 28.6-32.8 | 90.4 | 35.8 | 32.4-39.4 | 87.6 | 27.4 | 24.9-30.2 | 44.2 | 8.0 | 7.0-9.2 | 25.4 | 10.5 | 8.7-12.7 | 18.8 | 6.1 | 4.9-7.5 |
| Colon excluding rectum | 123.6 | 21.5 | 19.8-23.3 | 59.6 | 24.0 | 21.2-27.1 | 64.0 | 20.1 | 17.9-22.5 | 36.2 | 6.6 | 5.7-7.7 | 21.0 | 8.8 | 7.1-10.8 | 15.2 | 4.9 | 3.9-6.2 |
| Rectum & rectosigmoid | 54.4 | 9.1 | 8.1-10.3 | 30.8 | 11.8 | 9.9-13.9 | 23.6 | 7.3 | 6.1-8.8 | 8.0 | 1.4 | 1.0-1.9 | 4.4 | 1.7 | 1.0-2.7 | 3.6 | 1.1 | 0.7-1.8 |
| Liver / intrahepatic bile duct | 56.8 | 9.9 | 8.7-11.1 | 38.6 | 15.3 | 13.1-17.8 | 18.2 | 5.8 | 4.6-7.1 | 33.4 | 5.9 | 5.0-6.9 | 19.4 | 7.9 | 6.3-9.7 | 14.0 | 4.4 | 3.4-5.6 |
| Pancreas | 61.8 | 11.5 | 10.2-12.9 | 29.4 | 12.5 | 10.4-14.8 | 32.4 | 10.6 | 9.0-12.4 | 40.6 | 7.7 | 6.6-8.8 | 19.0 | 8.3 | 6.7-10.3 | 21.6 | 7.1 | 5.8-8.6 |
| Larynx | 9.6 | 1.7 | 1.3-2.3 | 8.0 | 3.3 | 2.3-4.6 | 1.6 | 0.6 | 0.2-1.1 | 2.4 | 0.4 | 0.2-0.8 | 2.2 | 1.0 | 0.5-1.7 | 0.2 | 0.1 | 0.0-0.4 |
| Lung and bronchus | 131.6 | 23.6 | 21.7-25.5 | 73.2 | 31.6 | 28.3-35.2 | 58.4 | 18.3 | 16.2-20.5 | 65.6 | 12.2 | 10.9-13.7 | 38.2 | 17.6 | 15.0-20.4 | 27.4 | 8.6 | 7.2-10.2 |
| Melanoma of the skin | 16.2 | 2.9 | 2.3-3.6 | 6.8 | 2.7 | 1.8-3.8 | 9.4 | 3.0 | 2.2-4.0 | 2.0 | 0.4 | 0.2-0.7 | 1.4 | 0.6 | 0.2-1.2 | 0.6 | 0.2 | 0.0-0.6 |
| Female breast | | | | | | | 309.2 | 94.4 | 89.7-99.3 | | | | | | | 36.6 | 11.3 | 9.7-13.1 |
| Cervix uteri | | | | | | | 32.8 | 10.3 | 8.8-12.1 | | | | | | | 6.6 | 2.0 | 1.4-2.9 |
| Corpus uterus and NOS | | | | | | | 74.8 | 22.2 | 20.0-24.6 | | | | | | | 13.2 | 4.1 | 3.1-5.2 |
| Ovary | | | | | | | 29.0 | 9.1 | 7.7-10.8 | | | | | | | 11.6 | 3.7 | 2.8-4.8 |
| Prostate | | | | 265.2 | 100.9 | 95.3-106.8 | | | | | | | 26.4 | 14.1 | 11.7-16.7 | | | |
| Testis | | | | 17.0 | 4.8 | 3.8-5.9 | | | | | | | 1.0 | 0.3 | 0.1-0.7 | | | |
| Urinary bladder (incl. in situ) | 45.8 | 8.5 | 7.4-9.7 | 31.2 | 14.3 | 12.1-16.9 | 14.6 | 4.7 | 3.7-5.9 | 9.4 | 2.0 | 1.5-2.7 | 6.2 | 3.6 | 2.4-5.1 | 3.2 | 1.1 | 0.6-1.8 |
| Kidney and renal pelvis | 75.8 | 12.4 | 11.2-13.7 | 46.6 | 16.9 | 14.7-19.4 | 29.2 | 8.8 | 7.4-10.4 | 8.0 | 1.4 | 1.0-2.0 | 5.2 | 2.2 | 1.4-3.2 | 2.8 | 0.9 | 0.5-1.5 |
| Brain and other central nervous system | 28.4 | 4.8 | 4.0-5.7 | 16.2 | 6.0 | 4.7-7.5 | 12.2 | 3.8 | 2.9-4.9 | 15.4 | 2.7 | 2.2-3.5 | 9.2 | 3.9 | 2.8-5.3 | 6.2 | 2.0 | 1.3-2.8 |
| Thyroid | 123.0 | 18.9 | 17.4-20.5 | 24.2 | 7.7 | 6.3-9.2 | 98.8 | 30.0 | 27.4-32.8 | 3.2 | 0.6 | 0.3-1.0 | 1.0 | 0.5 | 0.1-1.1 | 2.2 | 0.7 | 0.4-1.3 |
| Hodgkin lymphoma | 15.2 | 2.4 | 1.9-3.0 | 9.4 | 3.2 | 2.3-4.3 | 5.8 | 1.8 | 1.2-2.6 | 1.6 | 0.3 | 0.1-0.5 | 0.8 | 0.2 | 0.1-0.6 | 0.8 | 0.3 | 0.1-0.7 |
| Non-Hodgkin lymphomas | 107.6 | 18.5 | 16.9-20.2 | 57.4 | 22.1 | 19.5-25.0 | 50.2 | 15.8 | 13.9-17.9 | 23.6 | 4.3 | 3.6-5.2 | 14.8 | 6.6 | 5.1-8.4 | 8.8 | 2.7 | 2.0-3.7 |
| Myeloma | 44.6 | 7.8 | 6.8-8.9 | 22.4 | 8.8 | 7.2-10.7 | 22.2 | 7.0 | 5.7-8.4 | 10.0 | 1.9 | 1.4-2.5 | 4.2 | 1.8 | 1.1-2.8 | 5.8 | 1.9 | 1.2-2.7 |
| Leukemias | 67.4 | 11.3 | 10.1-12.6 | 40.8 | 14.9 | 12.8-17.2 | 26.6 | 8.4 | 7.1-10.0 | 24.4 | 4.3 | 3.5-5.1 | 13.4 | 5.6 | 4.3-7.2 | 11.0 | 3.4 | 2.6-4.5 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Queens, Non-Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 9585.8 | 450.3 | 446.2-454.5 | 4631.8 | 478.8 | 472.5-485.2 | 4954.0 | 434.9 | 429.3-440.6 | 2759.8 | 123.7 | 121.6-125.8 | 1351.8 | 142.8 | 139.3-146.3 | 1408.0 | 111.1 | 108.5-113.8 |
| Oral cavity and pharynx | 215.8 | 10.2 | 9.6-10.8 | 134.6 | 13.8 | 12.7-14.9 | 81.2 | 7.2 | 6.5-7.9 | 47.4 | 2.1 | 1.9-2.4 | 32.2 | 3.3 | 2.8-3.8 | 15.2 | 1.2 | 0.9-1.5 |
| Esophagus | 81.0 | 3.6 | 3.2-4.0 | 60.6 | 6.1 | 5.4-6.9 | 20.4 | 1.6 | 1.3-1.9 | 55.4 | 2.4 | 2.2-2.8 | 41.0 | 4.2 | 3.6-4.8 | 14.4 | 1.1 | 0.8-1.4 |
| Stomach | 263.6 | 12.2 | 11.5-12.9 | 155.2 | 16.2 | 15.1-17.4 | 108.4 | 9.1 | 8.3-9.9 | 105.8 | 4.8 | 4.4-5.3 | 65.6 | 7.1 | 6.3-7.9 | 40.2 | 3.1 | 2.7-3.6 |
| Colorectal | 847.2 | 39.6 | 38.4-40.9 | 439.6 | 46.1 | 44.2-48.1 | 407.6 | 34.4 | 32.9-36.0 | 275.6 | 12.3 | 11.7-13.0 | 132.2 | 14.0 | 13.0-15.2 | 143.4 | 11.1 | 10.3-12.0 |
| Colon excluding rectum | 597.0 | 27.8 | 26.8-28.8 | 298.6 | 31.5 | 29.8-33.1 | 298.4 | 24.9 | 23.6-26.2 | 229.4 | 10.3 | 9.7-10.9 | 106.6 | 11.4 | 10.4-12.4 | 122.8 | 9.5 | 8.7-10.3 |
| Rectum & rectosigmoid | 250.2 | 11.8 | 11.2-12.5 | 141.0 | 14.7 | 13.6-15.8 | 109.2 | 9.5 | 8.7-10.4 | 46.2 | 2.1 | 1.8-2.4 | 25.6 | 2.7 | 2.2-3.2 | 20.6 | 1.6 | 1.3-2.0 |
| Liver / intrahepatic bile duct | 228.8 | 10.4 | 9.8-11.0 | 165.2 | 16.5 | 15.4-17.7 | 63.6 | 5.3 | 4.7-6.0 | 136.8 | 6.1 | 5.7-6.6 | 92.4 | 9.4 | 8.5-10.3 | 44.4 | 3.6 | 3.1-4.1 |
| Pancreas | 296.0 | 13.2 | 12.5-13.9 | 137.0 | 14.3 | 13.2-15.4 | 159.0 | 12.3 | 11.4-13.2 | 223.0 | 9.8 | 9.2-10.4 | 103.4 | 10.6 | 9.7-11.6 | 119.6 | 9.0 | 8.3-9.8 |
| Larynx | 53.2 | 2.4 | 2.1-2.7 | 45.2 | 4.6 | 4.0-5.3 | 8.0 | 0.6 | 0.4-0.9 | 16.8 | 0.7 | 0.6-0.9 | 13.4 | 1.3 | 1.0-1.7 | 3.4 | 0.3 | 0.1-0.4 |
| Lung and bronchus | 1088.6 | 49.0 | 47.7-50.4 | 556.0 | 57.6 | 55.4-59.8 | 532.6 | 42.7 | 41.1-44.4 | 537.6 | 23.8 | 22.9-24.7 | 288.4 | 30.1 | 28.5-31.7 | 249.2 | 19.2 | 18.1-20.4 |
| Melanoma of the skin | 190.6 | 8.9 | 8.4-9.5 | 106.6 | 11.4 | 10.4-12.4 | 84.0 | 7.3 | 6.6-8.0 | 22.4 | 1.1 | 0.9-1.3 | 12.8 | 1.4 | 1.1-1.8 | 9.6 | 0.8 | 0.6-1.1 |
| Female breast | | | | | | | 1482.8 | 134.5 | 131.3-137.7 | | | | | | | 226.0 | 18.7 | 17.6-19.9 |
| Cervix uteri | | | | | | | 94.2 | 9.3 | 8.5-10.2 | | | | | | | 24.8 | 2.2 | 1.8-2.6 |
| Corpus uterus and NOS | | | | | | | 383.8 | 32.6 | 31.1-34.1 | | | | | | | 87.2 | 6.9 | 6.3-7.6 |
| Ovary | | | | | | | 133.8 | 12.5 | 11.5-13.5 | | | | | | | 74.8 | 6.2 | 5.5-6.8 |
| Prostate | | | | 1238.0 | 119.8 | 116.8-122.9 | | | | | | | 147.6 | 16.1 | 14.9-17.3 | | | |
| Testis | | | | 33.4 | 4.1 | 3.5-4.8 | | | | | | | 0.6 | 0.1 | 0.0-0.2 | | | |
| Urinary bladder (incl. in situ) | 397.2 | 17.9 | 17.1-18.7 | 298.2 | 31.6 | 30.0-33.3 | 99.0 | 7.7 | 7.0-8.5 | 73.8 | 3.2 | 2.9-3.6 | 51.8 | 5.6 | 4.9-6.3 | 22.0 | 1.6 | 1.3-2.0 |
| Kidney and renal pelvis | 312.4 | 14.8 | 14.1-15.6 | 199.4 | 20.9 | 19.6-22.2 | 113.0 | 9.8 | 9.0-10.7 | 54.0 | 2.4 | 2.1-2.7 | 34.4 | 3.7 | 3.1-4.3 | 19.6 | 1.5 | 1.2-1.8 |
| Brain and other central nervous system | 104.6 | 5.5 | 5.0-6.0 | 57.2 | 6.5 | 5.7-7.3 | 47.4 | 4.6 | 4.0-5.3 | 59.2 | 2.9 | 2.6-3.3 | 29.2 | 3.2 | 2.7-3.8 | 30.0 | 2.6 | 2.2-3.1 |
| Thyroid | 390.8 | 21.7 | 20.8-22.8 | 94.4 | 10.7 | 9.7-11.7 | 296.4 | 32.0 | 30.3-33.7 | 11.6 | 0.5 | 0.4-0.7 | 4.8 | 0.5 | 0.3-0.8 | 6.8 | 0.5 | 0.3-0.7 |
| Hodgkin lymphoma | 46.8 | 2.8 | 2.4-3.2 | 26.2 | 3.2 | 2.7-3.9 | 20.6 | 2.3 | 1.9-2.9 | 4.0 | 0.2 | 0.1-0.3 | 2.6 | 0.3 | 0.1-0.5 | 1.4 | 0.1 | 0.0-0.3 |
| Non-Hodgkin lymphomas | 391.0 | 18.5 | 17.7-19.4 | 212.4 | 22.7 | 21.3-24.1 | 178.6 | 15.3 | 14.3-16.4 | 96.2 | 4.3 | 3.9-4.7 | 56.0 | 6.0 | 5.3-6.8 | 40.2 | 3.0 | 2.6-3.5 |
| Myeloma | 179.2 | 8.2 | 7.6-8.7 | 90.2 | 9.3 | 8.4-10.2 | 89.0 | 7.2 | 6.6-8.0 | 63.4 | 2.8 | 2.5-3.1 | 31.6 | 3.3 | 2.8-3.8 | 31.8 | 2.4 | 2.1-2.9 |
| Leukemias | 283.2 | 13.7 | 13.0-14.5 | 159.4 | 17.5 | 16.2-18.8 | 123.8 | 10.9 | 10.0-11.8 | 106.8 | 4.9 | 4.4-5.3 | 62.0 | 6.7 | 6.0-7.5 | 44.8 | 3.6 | 3.1-4.1 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Queens, Non-Hispanic Whites

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 4580.0 | 502.1 | 495.2-509.1 | 2257.2 | 527.0 | 517.0-537.2 | 2322.8 | 493.3 | 483.4-503.3 | 1442.0 | 136.3 | 133.0-139.7 | 706.0 | 154.7 | 149.5-160.0 | 736.0 | 124.0 | 119.7-128.5 |
| Oral cavity and pharynx | 100.0 | 11.0 | 10.0-12.1 | 65.6 | 15.3 | 13.6-17.1 | 34.4 | 7.2 | 6.1-8.6 | 23.4 | 2.4 | 1.9-2.9 | 15.4 | 3.4 | 2.7-4.4 | 8.0 | 1.5 | 1.1-2.1 |
| Esophagus | 43.6 | 4.4 | 3.8-5.1 | 32.2 | 7.3 | 6.2-8.6 | 11.4 | 1.9 | 1.4-2.6 | 30.8 | 3.1 | 2.6-3.6 | 22.8 | 5.2 | 4.3-6.3 | 8.0 | 1.2 | 0.9-1.8 |
| Stomach | 82.2 | 8.7 | 7.9-9.7 | 47.8 | 11.3 | 9.9-12.9 | 34.4 | 6.5 | 5.4-7.7 | 34.2 | 3.3 | 2.8-3.9 | 21.2 | 4.9 | 4.0-6.0 | 13.0 | 1.9 | 1.4-2.5 |
| Colorectal | 393.2 | 42.3 | 40.3-44.3 | 203.6 | 48.4 | 45.4-51.6 | 189.6 | 37.2 | 34.6-39.9 | 141.8 | 13.0 | 12.0-14.0 | 72.0 | 15.6 | 14.0-17.4 | 69.8 | 11.1 | 9.8-12.4 |
| Colon excluding rectum | 281.8 | 29.5 | 27.9-31.2 | 140.8 | 33.1 | 30.7-35.8 | 141.0 | 26.6 | 24.5-28.9 | 118.0 | 10.7 | 9.8-11.6 | 58.2 | 12.6 | 11.1-14.2 | 59.8 | 9.4 | 8.3-10.6 |
| Rectum & rectosigmoid | 111.4 | 12.7 | 11.6-13.9 | 62.8 | 15.3 | 13.6-17.1 | 48.6 | 10.6 | 9.2-12.1 | 23.8 | 2.3 | 1.9-2.8 | 13.8 | 3.1 | 2.4-3.9 | 10.0 | 1.7 | 1.2-2.3 |
| Liver / intrahepatic bile duct | 81.6 | 8.2 | 7.4-9.1 | 59.6 | 13.3 | 11.8-14.9 | 22.0 | 3.9 | 3.2-4.8 | 56.2 | 5.5 | 4.9-6.3 | 37.4 | 8.4 | 7.2-9.7 | 18.8 | 3.1 | 2.5-3.9 |
| Pancreas | 150.0 | 14.3 | 13.3-15.5 | 70.2 | 15.7 | 14.1-17.6 | 79.8 | 13.0 | 11.6-14.5 | 120.4 | 11.0 | 10.1-11.9 | 56.4 | 12.2 | 10.8-13.8 | 64.0 | 9.8 | 8.7-11.1 |
| Larynx | 27.2 | 2.9 | 2.4-3.4 | 23.0 | 5.3 | 4.4-6.4 | 4.2 | 0.8 | 0.5-1.3 | 10.0 | 1.0 | 0.8-1.4 | 8.8 | 2.0 | 1.5-2.7 | 1.2 | 0.2 | 0.1-0.6 |
| Lung and bronchus | 578.4 | 57.6 | 55.4-59.8 | 282.4 | 63.2 | 59.9-66.7 | 296.0 | 53.1 | 50.3-56.1 | 303.0 | 28.7 | 27.3-30.3 | 149.6 | 32.7 | 30.4-35.2 | 153.4 | 25.7 | 23.8-27.7 |
| Melanoma of the skin | 174.2 | 19.3 | 18.0-20.8 | 99.2 | 23.3 | 21.2-25.5 | 75.0 | 16.6 | 14.8-18.6 | 18.2 | 2.0 | 1.6-2.5 | 10.2 | 2.4 | 1.8-3.2 | 8.0 | 1.7 | 1.1-2.4 |
| Female breast | | | | | | | 643.4 | 147.9 | 142.4-153.7 | | | | | | | 112.4 | 20.7 | 18.8-22.7 |
| Cervix uteri | | | | | | | 26.8 | 7.3 | 6.0-8.8 | | | | | | | 8.2 | 2.1 | 1.4-2.9 |
| Corpus uterus and NOS | | | | | | | 163.0 | 35.1 | 32.5-37.8 | | | | | | | 36.2 | 6.2 | 5.3-7.3 |
| Ovary | | | | | | | 55.2 | 12.5 | 10.9-14.2 | | | | | | | 36.8 | 6.7 | 5.8-7.9 |
| Prostate | | | | 495.0 | 109.6 | 105.2-114.1 | | | | | | | 72.4 | 15.1 | 13.5-16.8 | | | |
| Testis | | | | 22.8 | 7.9 | 6.5-9.6 | | | | | | | 0.4 | 0.1 | 0.0-0.4 | | | |
| Urinary bladder (incl. in situ) | 270.2 | 26.5 | 25.0-28.0 | 205.4 | 45.8 | 43.0-48.7 | 64.8 | 11.3 | 10.0-12.7 | 52.6 | 4.5 | 3.9-5.1 | 36.6 | 7.6 | 6.6-8.9 | 16.0 | 2.3 | 1.8-3.0 |
| Kidney and renal pelvis | 152.0 | 17.6 | 16.3-19.0 | 95.2 | 23.3 | 21.2-25.6 | 56.8 | 12.3 | 10.7-14.0 | 31.0 | 2.9 | 2.4-3.5 | 18.0 | 3.9 | 3.1-4.9 | 13.0 | 2.1 | 1.6-2.8 |
| Brain and other central nervous system | 55.0 | 7.3 | 6.4-8.3 | 27.6 | 7.6 | 6.3-9.0 | 27.4 | 7.1 | 5.8-8.6 | 33.8 | 3.6 | 3.1-4.3 | 16.2 | 3.9 | 3.1-4.9 | 17.6 | 3.5 | 2.7-4.4 |
| Thyroid | 154.6 | 23.9 | 22.1-25.7 | 43.4 | 12.6 | 10.9-14.5 | 111.2 | 35.7 | 32.6-39.1 | 6.2 | 0.6 | 0.4-0.9 | 2.6 | 0.6 | 0.3-1.1 | 3.6 | 0.5 | 0.3-1.0 |
| Hodgkin lymphoma | 25.4 | 4.4 | 3.6-5.3 | 15.4 | 5.5 | 4.2-6.9 | 10.0 | 3.3 | 2.4-4.4 | 3.0 | 0.3 | 0.2-0.6 | 2.0 | 0.5 | 0.2-1.0 | 1.0 | 0.2 | 0.1-0.6 |
| Non-Hodgkin lymphomas | 213.2 | 23.3 | 21.8-24.9 | 116.8 | 27.9 | 25.6-30.4 | 96.4 | 19.5 | 17.6-21.5 | 57.2 | 5.2 | 4.6-5.8 | 32.4 | 7.1 | 6.0-8.4 | 24.8 | 3.6 | 3.0-4.4 |
| Myeloma | 72.0 | 7.3 | 6.6-8.2 | 39.0 | 8.9 | 7.7-10.3 | 33.0 | 6.0 | 5.0-7.1 | 27.2 | 2.5 | 2.1-2.9 | 14.4 | 3.1 | 2.4-3.9 | 12.8 | 2.0 | 1.5-2.6 |
| Leukemias | 157.2 | 17.2 | 15.9-18.5 | 92.0 | 22.1 | 20.0-24.4 | 65.2 | 13.1 | 11.5-14.8 | 61.0 | 5.6 | 4.9-6.3 | 35.2 | 7.6 | 6.5-8.9 | 25.8 | 4.2 | 3.4-5.1 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Queens, Non-Hispanic Blacks**

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 2195.0 | 427.0 | 418.9-435.3 | 1052.0 | 500.0 | 486.1-514.2 | 1143.0 | 384.0 | 373.8-394.5 | 666.2 | 131.3 | 126.8-135.9 | 293.0 | 155.3 | 147.2-163.8 | 373.2 | 119.8 | 114.3-125.5 |
| Oral cavity and pharynx | 36.4 | 7.3 | 6.2-8.4 | 20.8 | 10.0 | 8.1-12.2 | 15.6 | 5.4 | 4.3-6.9 | 8.8 | 1.7 | 1.2-2.3 | 6.0 | 2.9 | 1.9-4.2 | 2.8 | 0.9 | 0.5-1.5 |
| Esophagus | 17.8 | 3.4 | 2.7-4.2 | 13.0 | 6.2 | 4.8-8.0 | 4.8 | 1.6 | 1.0-2.4 | 11.6 | 2.2 | 1.6-2.8 | 7.8 | 3.8 | 2.7-5.3 | 3.8 | 1.2 | 0.7-1.9 |
| Stomach | 56.6 | 11.2 | 9.9-12.7 | 29.6 | 15.4 | 12.9-18.2 | 27.0 | 8.9 | 7.5-10.7 | 23.0 | 4.6 | 3.8-5.6 | 15.0 | 8.4 | 6.6-10.6 | 8.0 | 2.6 | 1.8-3.6 |
| Colorectal | 200.0 | 39.7 | 37.2-42.3 | 92.0 | 45.7 | 41.4-50.2 | 108.0 | 35.8 | 32.8-39.0 | 76.4 | 15.3 | 13.8-16.9 | 28.4 | 14.9 | 12.5-17.7 | 48.0 | 15.5 | 13.5-17.6 |
| Colon excluding rectum | 153.0 | 30.4 | 28.2-32.6 | 66.8 | 33.5 | 29.8-37.4 | 86.2 | 28.3 | 25.6-31.2 | 64.8 | 13.0 | 11.6-14.6 | 23.6 | 12.6 | 10.4-15.2 | 41.2 | 13.3 | 11.5-15.3 |
| Rectum & rectosigmoid | 47.0 | 9.4 | 8.2-10.7 | 25.2 | 12.2 | 10.1-14.7 | 21.8 | 7.5 | 6.1-9.1 | 11.6 | 2.3 | 1.7-3.0 | 4.8 | 2.3 | 1.5-3.5 | 6.8 | 2.2 | 1.5-3.1 |
| Liver / intrahepatic bile duct | 41.4 | 7.9 | 6.9-9.1 | 29.6 | 14.0 | 11.8-16.6 | 11.8 | 3.8 | 2.9-5.0 | 25.6 | 4.9 | 4.0-5.8 | 18.6 | 9.0 | 7.2-11.2 | 7.0 | 2.2 | 1.5-3.2 |
| Pancreas | 68.6 | 13.4 | 12.0-15.0 | 27.4 | 14.1 | 11.8-16.8 | 41.2 | 13.1 | 11.4-15.1 | 53.6 | 10.4 | 9.2-11.7 | 24.0 | 12.2 | 10.1-14.7 | 29.6 | 9.1 | 7.7-10.8 |
| Larynx | 13.8 | 2.5 | 2.0-3.2 | 11.2 | 5.5 | 4.1-7.2 | 2.6 | 0.8 | 0.4-1.4 | 5.4 | 1.0 | 0.6-1.4 | 3.4 | 1.6 | 0.9-2.6 | 2.0 | 0.6 | 0.3-1.1 |
| Lung and bronchus | 185.6 | 35.9 | 33.6-38.3 | 90.6 | 44.8 | 40.6-49.3 | 95.0 | 30.2 | 27.5-33.2 | 101.8 | 19.8 | 18.1-21.6 | 53.0 | 27.5 | 24.2-31.2 | 48.8 | 15.2 | 13.4-17.3 |
| Melanoma of the skin | 3.6 | 0.7 | 0.4-1.2 | 1.4 | 0.7 | 0.3-1.5 | 2.2 | 0.7 | 0.4-1.4 | 1.0 | 0.2 | 0.1-0.4 | 0.4 | 0.2 | 0.0-0.7 | 0.6 | 0.2 | 0.0-0.6 |
| Female breast | | | | | | | 375.2 | 128.8 | 122.8-135.0 | | | | | | | 69.6 | 23.1 | 20.7-25.7 |
| Cervix uteri | | | | | | | 26.0 | 9.8 | 8.1-11.7 | | | | | | | 8.2 | 2.7 | 1.9-3.8 |
| Corpus uterus and NOS | | | | | | | 113.4 | 35.8 | 32.8-39.0 | | | | | | | 34.6 | 10.7 | 9.2-12.5 |
| Ovary | | | | | | | 28.0 | 9.5 | 7.9-11.3 | | | | | | | 21.6 | 6.8 | 5.6-8.3 |
| Prostate | | | | 454.2 | 201.9 | 193.4-210.6 | | | | | | | 52.0 | 30.6 | 26.9-34.6 | | | |
| Testis | | | | 4.2 | 2.0 | 1.2-3.1 | | | | | | | 0.2 | 0.1 | 0.0-0.7 | | | |
| Urinary bladder (incl. in situ) | 51.4 | 10.0 | 8.8-11.4 | 35.2 | 18.5 | 15.8-21.6 | 16.2 | 5.0 | 4.0-6.3 | 11.2 | 2.2 | 1.7-2.9 | 7.4 | 3.9 | 2.7-5.4 | 3.8 | 1.3 | 0.8-2.0 |
| Kidney and renal pelvis | 75.2 | 14.7 | 13.2-16.3 | 49.6 | 23.9 | 20.9-27.2 | 25.6 | 8.5 | 7.1-10.2 | 11.2 | 2.3 | 1.7-3.0 | 7.8 | 4.2 | 3.0-5.8 | 3.4 | 1.1 | 0.6-1.8 |
| Brain and other central nervous system | 16.2 | 3.6 | 2.8-4.5 | 10.6 | 5.3 | 3.9-7.0 | 5.6 | 2.1 | 1.4-3.2 | 11.2 | 2.6 | 1.9-3.4 | 5.2 | 2.9 | 1.9-4.3 | 6.0 | 2.3 | 1.5-3.3 |
| Thyroid | 40.8 | 8.7 | 7.5-10.0 | 7.6 | 3.8 | 2.7-5.3 | 33.2 | 12.6 | 10.7-14.7 | 2.4 | 0.5 | 0.2-0.8 | 0.6 | 0.3 | 0.1-0.9 | 1.8 | 0.6 | 0.3-1.1 |
| Hodgkin lymphoma | 11.0 | 2.5 | 1.9-3.3 | 5.2 | 2.5 | 1.6-3.8 | 5.8 | 2.5 | 1.7-3.7 | 0.4 | 0.1 | 0.0-0.3 | 0.2 | 0.1 | 0.0-0.5 | 0.2 | 0.1 | 0.0-0.5 |
| Non-Hodgkin lymphomas | 68.8 | 13.8 | 12.4-15.4 | 32.8 | 16.1 | 13.7-18.9 | 36.0 | 12.5 | 10.7-14.6 | 17.4 | 3.5 | 2.8-4.3 | 8.4 | 4.3 | 3.0-5.8 | 9.0 | 2.9 | 2.1-3.9 |
| Myeloma | 69.6 | 13.7 | 12.2-15.2 | 29.6 | 14.4 | 12.1-17.0 | 40.0 | 13.1 | 11.3-15.1 | 26.0 | 5.1 | 4.2-6.1 | 11.8 | 6.1 | 4.6-8.0 | 14.2 | 4.5 | 3.5-5.7 |
| Leukemias | 50.4 | 10.5 | 9.2-11.9 | 23.2 | 12.0 | 9.8-14.5 | 27.2 | 9.7 | 8.1-11.6 | 22.2 | 4.4 | 3.6-5.4 | 12.4 | 6.3 | 4.8-8.1 | 9.8 | 3.4 | 2.5-4.5 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Richmond (Staten Island), Non-Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 2782.4 | 542.7 | 533.4-552.1 | 1361.8 | 578.9 | 564.8-593.3 | 1420.6 | 520.8 | 508.2-533.7 | 793.2 | 149.4 | 144.7-154.3 | 389.4 | 171.8 | 164.1-179.9 | 403.8 | 134.1 | 128.2-140.3 |
| Oral cavity and pharynx | 53.6 | 10.2 | 8.9-11.5 | 36.8 | 15.3 | 13.0-17.7 | 16.8 | 5.7 | 4.5-7.2 | 11.0 | 2.0 | 1.5-2.6 | 8.4 | 3.4 | 2.4-4.6 | 2.6 | 0.8 | 0.4-1.4 |
| Esophagus | 25.4 | 4.6 | 3.8-5.5 | 19.8 | 7.9 | 6.4-9.7 | 5.6 | 1.9 | 1.3-2.9 | 16.8 | 3.1 | 2.4-3.8 | 12.4 | 4.9 | 3.8-6.4 | 4.4 | 1.5 | 0.9-2.4 |
| Stomach | 44.6 | 8.5 | 7.4-9.8 | 27.8 | 12.1 | 10.1-14.4 | 16.8 | 5.8 | 4.6-7.3 | 19.2 | 3.6 | 2.9-4.5 | 10.0 | 4.3 | 3.2-5.8 | 9.2 | 3.2 | 2.3-4.3 |
| Colorectal | 208.0 | 40.4 | 37.9-43.0 | 108.4 | 47.2 | 43.2-51.5 | 99.6 | 34.5 | 31.5-37.9 | 73.4 | 13.9 | 12.5-15.5 | 36.0 | 16.2 | 13.8-18.8 | 37.4 | 12.0 | 10.3-14.0 |
| Colon excluding rectum | 153.0 | 29.5 | 27.4-31.7 | 76.8 | 33.4 | 30.0-37.1 | 76.2 | 26.1 | 23.5-29.0 | 60.6 | 11.5 | 10.2-12.9 | 28.6 | 12.9 | 10.8-15.3 | 32.0 | 10.4 | 8.8-12.2 |
| Rectum & rectosigmoid | 55.0 | 10.9 | 9.6-12.3 | 31.6 | 13.8 | 11.7-16.3 | 23.4 | 8.4 | 6.9-10.2 | 12.8 | 2.4 | 1.8-3.1 | 7.4 | 3.3 | 2.3-4.6 | 5.4 | 1.7 | 1.1-2.5 |
| Liver / intrahepatic bile duct | 55.8 | 10.0 | 8.9-11.3 | 39.8 | 15.5 | 13.4-17.9 | 16.0 | 5.4 | 4.3-6.8 | 35.6 | 6.5 | 5.5-7.5 | 24.8 | 10.0 | 8.2-12.0 | 10.8 | 3.6 | 2.7-4.7 |
| Pancreas | 81.8 | 15.3 | 13.8-16.9 | 39.4 | 16.6 | 14.3-19.2 | 42.4 | 14.1 | 12.2-16.2 | 63.6 | 11.7 | 10.4-13.1 | 29.2 | 12.3 | 10.3-14.5 | 34.4 | 11.3 | 9.7-13.2 |
| Larynx | 18.6 | 3.5 | 2.8-4.3 | 15.4 | 6.5 | 5.1-8.3 | 3.2 | 1.2 | 0.7-2.0 | 4.4 | 0.8 | 0.5-1.2 | 2.8 | 1.1 | 0.6-1.9 | 1.6 | 0.5 | 0.2-1.1 |
| Lung and bronchus | 364.8 | 68.4 | 65.2-71.7 | 170.2 | 72.9 | 68.0-78.2 | 194.6 | 65.8 | 61.7-70.2 | 200.0 | 37.5 | 35.2-40.0 | 100.0 | 44.0 | 40.1-48.1 | 100.0 | 33.0 | 30.1-36.1 |
| Melanoma of the skin | 82.4 | 16.3 | 14.8-18.1 | 47.4 | 20.7 | 18.0-23.6 | 35.0 | 13.2 | 11.2-15.4 | 8.0 | 1.6 | 1.1-2.2 | 4.6 | 2.3 | 1.5-3.5 | 3.4 | 1.2 | 0.7-1.9 |
| Female breast | | | | | | | 377.2 | 141.5 | 134.9-148.4 | | | | | | | 53.0 | 17.2 | 15.2-19.6 |
| Cervix uteri | | | | | | | 18.8 | 8.2 | 6.6-10.2 | | | | | | | 5.4 | 2.0 | 1.3-3.0 |
| Corpus uterus and NOS | | | | | | | 111.6 | 38.3 | 35.1-41.8 | | | | | | | 20.0 | 6.6 | 5.3-8.1 |
| Ovary | | | | | | | 38.8 | 14.6 | 12.6-17.0 | | | | | | | 20.6 | 6.8 | 5.5-8.3 |
| Prostate | | | | 314.2 | 122.4 | 116.3-128.8 | | | | | | | 35.6 | 17.2 | 14.7-19.9 | | | |
| Testis | | | | 11.0 | 6.4 | 4.8-8.4 | | | | | | | 0.2 | 0.1 | 0.0-0.6 | | | |
| Urinary bladder (incl. in situ) | 145.4 | 27.1 | 25.2-29.2 | 109.0 | 47.0 | 43.0-51.2 | 36.4 | 12.0 | 10.3-14.0 | 27.6 | 5.2 | 4.4-6.2 | 20.0 | 9.4 | 7.6-11.5 | 7.6 | 2.5 | 1.7-3.5 |
| Kidney and renal pelvis | 118.2 | 22.8 | 20.9-24.8 | 81.2 | 34.3 | 30.9-37.9 | 37.0 | 13.0 | 11.1-15.1 | 20.0 | 3.7 | 3.0-4.6 | 14.8 | 6.3 | 4.9-8.0 | 5.2 | 1.7 | 1.1-2.6 |
| Brain and other central nervous system | 33.4 | 7.3 | 6.2-8.6 | 16.8 | 7.9 | 6.2-9.9 | 16.6 | 6.7 | 5.3-8.5 | 20.2 | 3.9 | 3.1-4.7 | 11.0 | 4.6 | 3.4-6.1 | 9.2 | 3.2 | 2.3-4.3 |
| Thyroid | 155.4 | 36.0 | 33.4-38.7 | 46.6 | 22.1 | 19.3-25.3 | 108.8 | 48.9 | 44.7-53.4 | 3.6 | 0.7 | 0.4-1.1 | 1.6 | 0.6 | 0.3-1.3 | 2.0 | 0.7 | 0.3-1.3 |
| Hodgkin lymphoma | 14.6 | 3.7 | 2.9-4.7 | 9.2 | 4.8 | 3.5-6.5 | 5.4 | 2.7 | 1.7-4.0 | 2.6 | 0.5 | 0.3-0.9 | 1.8 | 0.8 | 0.3-1.5 | 0.8 | 0.3 | 0.1-0.8 |
| Non-Hodgkin lymphomas | 129.4 | 25.8 | 23.7-27.9 | 68.4 | 30.4 | 27.2-33.9 | 61.0 | 22.0 | 19.5-24.7 | 22.2 | 4.4 | 3.6-5.4 | 13.2 | 6.2 | 4.7-7.9 | 9.0 | 3.1 | 2.2-4.2 |
| Myeloma | 48.2 | 8.9 | 7.8-10.2 | 27.0 | 11.5 | 9.6-13.7 | 21.2 | 6.9 | 5.7-8.5 | 14.6 | 2.7 | 2.1-3.5 | 8.2 | 3.5 | 2.5-4.8 | 6.4 | 2.1 | 1.5-3.1 |
| Leukemias | 92.4 | 18.6 | 16.9-20.4 | 55.8 | 24.7 | 21.8-27.8 | 36.6 | 13.8 | 11.8-16.1 | 32.8 | 6.5 | 5.5-7.7 | 16.4 | 7.9 | 6.2-9.8 | 16.4 | 5.7 | 4.5-7.2 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Richmond (Staten Island), Non-Hispanic Whites

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 2354.4 | 568.4 | 557.7-579.2 | 1143.0 | 596.0 | 580.2-612.3 | 1211.4 | 555.5 | 540.7-570.5 | 677.2 | 153.4 | 148.2-158.8 | 331.0 | 175.8 | 167.3-184.8 | 346.2 | 138.1 | 131.4-145.1 |
| Oral cavity and pharynx | 43.8 | 10.2 | 8.9-11.8 | 30.2 | 15.2 | 12.8-18.0 | 13.6 | 5.8 | 4.5-7.5 | 9.2 | 1.9 | 1.4-2.6 | 7.2 | 3.4 | 2.3-4.8 | 2.0 | 0.7 | 0.3-1.5 |
| Esophagus | 21.8 | 4.9 | 4.0-5.9 | 17.6 | 8.6 | 6.9-10.7 | 4.2 | 1.8 | 1.1-2.9 | 14.6 | 3.2 | 2.5-4.1 | 11.2 | 5.5 | 4.1-7.2 | 3.4 | 1.4 | 0.8-2.4 |
| Stomach | 35.0 | 8.2 | 7.0-9.6 | 21.4 | 11.6 | 9.5-14.2 | 13.6 | 5.7 | 4.4-7.4 | 15.2 | 3.3 | 2.6-4.2 | 8.6 | 4.4 | 3.2-6.0 | 6.6 | 2.5 | 1.7-3.7 |
| Colorectal | 172.8 | 41.2 | 38.4-44.2 | 89.4 | 47.8 | 43.3-52.6 | 83.4 | 35.6 | 32.1-39.4 | 62.0 | 14.1 | 12.5-15.9 | 30.2 | 16.4 | 13.8-19.4 | 31.8 | 12.2 | 10.3-14.4 |
| Colon excluding rectum | 129.0 | 30.4 | 28.0-32.9 | 65.4 | 34.7 | 30.9-38.8 | 63.6 | 26.8 | 23.8-30.1 | 52.0 | 11.9 | 10.4-13.5 | 25.0 | 13.7 | 11.4-16.5 | 27.0 | 10.4 | 8.7-12.4 |
| Rectum & rectosigmoid | 43.8 | 10.8 | 9.4-12.4 | 24.0 | 13.1 | 10.8-15.8 | 19.8 | 8.8 | 7.1-10.9 | 10.0 | 2.2 | 1.6-3.0 | 5.2 | 2.7 | 1.7-4.1 | 4.8 | 1.8 | 1.1-2.8 |
| Liver / intrahepatic bile duct | 40.6 | 9.0 | 7.8-10.4 | 28.8 | 13.9 | 11.7-16.5 | 11.8 | 4.9 | 3.7-6.5 | 28.6 | 6.4 | 5.4-7.6 | 19.4 | 9.7 | 7.8-11.9 | 9.2 | 3.7 | 2.7-5.1 |
| Pancreas | 71.4 | 16.3 | 14.6-18.1 | 33.8 | 17.5 | 14.9-20.5 | 37.6 | 15.1 | 12.9-17.5 | 56.0 | 12.5 | 11.1-14.1 | 25.6 | 13.0 | 10.8-15.6 | 30.4 | 12.1 | 10.2-14.3 |
| Larynx | 15.8 | 3.7 | 2.9-4.6 | 12.6 | 6.6 | 5.0-8.5 | 3.2 | 1.4 | 0.8-2.4 | 3.8 | 0.8 | 0.5-1.3 | 2.2 | 1.0 | 0.5-2.0 | 1.6 | 0.6 | 0.3-1.4 |
| Lung and bronchus | 311.6 | 70.7 | 67.1-74.4 | 142.6 | 73.4 | 68.0-79.1 | 169.0 | 69.6 | 64.9-74.7 | 169.4 | 38.1 | 35.5-40.8 | 83.2 | 43.9 | 39.7-48.5 | 86.2 | 34.0 | 30.8-37.5 |
| Melanoma of the skin | 79.6 | 19.4 | 17.5-21.6 | 45.8 | 24.2 | 21.1-27.7 | 33.8 | 16.0 | 13.5-18.8 | 8.0 | 1.9 | 1.3-2.6 | 4.6 | 2.7 | 1.7-4.1 | 3.4 | 1.4 | 0.8-2.4 |
| Female breast | | | | | | | 316.8 | 150.2 | 142.4-158.3 | | | | | | | 46.2 | 17.9 | 15.6-20.6 |
| Cervix uteri | | | | | | | 14.4 | 8.2 | 6.3-10.6 | | | | | | | 3.8 | 1.8 | 1.0-2.9 |
| Corpus uterus and NOS | | | | | | | 93.6 | 40.4 | 36.7-44.5 | | | | | | | 15.6 | 6.3 | 5.0-8.0 |
| Ovary | | | | | | | 34.0 | 16.2 | 13.7-19.1 | | | | | | | 17.8 | 7.1 | 5.7-8.9 |
| Prostate | | | | 244.4 | 116.5 | 109.9-123.4 | | | | | | | 30.0 | 16.8 | 14.2-19.8 | | | |
| Testis | | | | 10.2 | 8.0 | 5.9-10.6 | | | | | | | 0.2 | 0.1 | 0.0-0.7 | | | |
| Urinary bladder (incl. in situ) | 137.6 | 31.1 | 28.8-33.6 | 103.6 | 53.8 | 49.1-58.8 | 34.0 | 13.6 | 11.6-15.9 | 24.4 | 5.5 | 4.5-6.6 | 17.8 | 9.9 | 7.9-12.3 | 6.6 | 2.5 | 1.7-3.7 |
| Kidney and renal pelvis | 105.4 | 25.4 | 23.2-27.8 | 73.0 | 38.1 | 34.1-42.4 | 32.4 | 14.3 | 12.1-16.9 | 18.0 | 4.1 | 3.3-5.1 | 13.2 | 6.9 | 5.3-8.9 | 4.8 | 2.0 | 1.2-3.0 |
| Brain and other central nervous system | 27.8 | 7.8 | 6.5-9.3 | 14.0 | 8.4 | 6.4-10.7 | 13.8 | 7.3 | 5.5-9.4 | 17.0 | 4.1 | 3.2-5.2 | 9.2 | 4.9 | 3.5-6.7 | 7.8 | 3.4 | 2.4-4.8 |
| Thyroid | 131.4 | 39.7 | 36.6-43.1 | 39.6 | 24.1 | 20.7-27.8 | 91.8 | 54.8 | 49.5-60.4 | 2.6 | 0.5 | 0.3-0.9 | 1.0 | 0.4 | 0.1-1.1 | 1.6 | 0.5 | 0.2-1.2 |
| Hodgkin lymphoma | 11.6 | 4.0 | 3.0-5.2 | 7.4 | 5.2 | 3.6-7.3 | 4.2 | 2.8 | 1.7-4.4 | 2.4 | 0.5 | 0.3-1.0 | 1.6 | 0.8 | 0.3-1.7 | 0.8 | 0.4 | 0.1-1.1 |
| Non-Hodgkin lymphomas | 111.2 | 27.1 | 24.8-29.6 | 59.2 | 32.1 | 28.4-36.2 | 52.0 | 23.0 | 20.1-26.2 | 20.0 | 4.8 | 3.9-5.9 | 11.6 | 6.5 | 4.9-8.5 | 8.4 | 3.4 | 2.4-4.7 |
| Myeloma | 37.0 | 8.3 | 7.2-9.7 | 21.2 | 10.9 | 8.9-13.3 | 15.8 | 6.2 | 4.9-7.9 | 12.0 | 2.7 | 2.1-3.6 | 6.6 | 3.4 | 2.3-4.9 | 5.4 | 2.3 | 1.5-3.4 |
| Leukemias | 83.8 | 20.8 | 18.7-23.0 | 50.2 | 26.7 | 23.4-30.4 | 33.6 | 16.1 | 13.6-18.9 | 29.0 | 6.8 | 5.7-8.1 | 14.4 | 8.1 | 6.3-10.3 | 14.6 | 6.0 | 4.6-7.7 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Nassau, Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-----------|----------------|------------------|-----------|----------------|------------------|-----------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 670.8 | 379.2 | 365.8-393.0 | 302.4 | 397.4 | 375.7-419.9 | 368.4 | 376.6 | 359.1-394.7 | 127.8 | 80.3 | 73.9-87.1 | 59.4 | 84.3 | 74.1-95.4 | 68.4 | 76.8 | 68.7-85.6 |
| Oral cavity and pharynx | 14.8 | 8.5 | 6.5-10.7 | 9.2 | 12.2 | 8.7-16.6 | 5.6 | 5.7 | 3.7-8.3 | 1.8 | 1.2 | 0.5-2.3 | 1.6 | 2.8 | 1.1-5.4 | 0.2 | 0.2 | 0.0-1.0 |
| Esophagus | 4.8 | 3.1 | 1.9-4.6 | 4.2 | 6.5 | 3.8-10.0 | 0.6 | 0.7 | 0.1-1.9 | 1.6 | 0.9 | 0.4-1.7 | 1.6 | 1.9 | 0.8-3.9 | 0.0 | 0.0 | 0.0-0.7 |
| Stomach | 24.6 | 13.9 | 11.5-16.8 | 11.4 | 12.9 | 9.5-17.1 | 13.2 | 14.3 | 11.0-18.3 | 7.8 | 4.1 | 2.8-5.6 | 4.8 | 5.1 | 3.2-7.8 | 3.0 | 3.1 | 1.7-5.1 |
| Colorectal | 54.2 | 31.7 | 27.8-35.9 | 27.6 | 36.0 | 29.7-43.2 | 26.6 | 28.5 | 23.8-34.0 | 10.0 | 6.3 | 4.6-8.4 | 5.8 | 8.2 | 5.3-12.1 | 4.2 | 4.9 | 3.0-7.5 |
| Colon excluding rectum | 37.2 | 22.4 | 19.1-26.0 | 18.0 | 24.6 | 19.4-30.8 | 19.2 | 20.9 | 16.8-25.7 | 7.8 | 5.1 | 3.6-7.0 | 4.0 | 6.0 | 3.5-9.5 | 3.8 | 4.4 | 2.6-6.9 |
| Rectum & rectosigmoid | 17.0 | 9.3 | 7.3-11.6 | 9.6 | 11.4 | 8.1-15.5 | 7.4 | 7.6 | 5.3-10.6 | 2.2 | 1.3 | 0.6-2.3 | 1.8 | 2.2 | 0.9-4.4 | 0.4 | 0.5 | 0.1-1.7 |
| Liver / intrahepatic bile duct | 17.6 | 11.6 | 9.2-14.4 | 10.6 | 15.4 | 11.2-20.5 | 7.0 | 8.6 | 5.9-11.9 | 8.6 | 5.5 | 3.9-7.5 | 4.6 | 7.1 | 4.3-10.8 | 4.0 | 4.4 | 2.7-6.8 |
| Pancreas | 19.6 | 12.5 | 10.0-15.4 | 10.0 | 14.5 | 10.4-19.4 | 9.6 | 11.0 | 8.0-14.6 | 12.2 | 8.0 | 6.0-10.4 | 6.2 | 9.3 | 6.1-13.3 | 6.0 | 6.9 | 4.6-9.9 |
| Larynx | 4.6 | 2.8 | 1.7-4.3 | 4.0 | 5.6 | 3.2-8.9 | 0.6 | 0.7 | 0.1-2.0 | 0.2 | 0.2 | 0.0-0.8 | 0.2 | 0.5 | 0.0-2.1 | 0.0 | 0.0 | 0.0-0.7 |
| Lung and bronchus | 44.6 | 28.7 | 24.9-32.8 | 21.2 | 33.8 | 27.3-41.3 | 23.4 | 25.9 | 21.3-31.2 | 15.8 | 9.7 | 7.6-12.2 | 8.8 | 12.4 | 8.7-17.0 | 7.0 | 7.7 | 5.3-10.8 |
| Melanoma of the skin | 8.0 | 4.3 | 3.0-5.9 | 4.4 | 5.0 | 3.0-7.8 | 3.6 | 3.7 | 2.2-5.9 | 1.0 | 0.7 | 0.2-1.5 | 0.4 | 0.6 | 0.1-2.2 | 0.6 | 0.7 | 0.1-2.0 |
| Female breast | | | | | | | 110.6 | 109.4 | 100.2-119.1 | | | | | | | 10.6 | 11.1 | 8.2-14.6 |
| Cervix uteri | | | | | | | 9.2 | 8.6 | 6.3-11.5 | | | | | | | 1.6 | 1.9 | 0.8-3.8 |
| Corpus uteri and NOS | | | | | | | 19.2 | 18.3 | 14.8-22.5 | | | | | | | 3.4 | 4.1 | 2.3-6.5 |
| Ovary | | | | | | | 9.2 | 8.6 | 6.2-11.5 | | | | | | | 5.2 | 5.5 | 3.5-8.0 |
| Prostate | | | | 74.6 | 100.6 | 89.8-112.2 | | | | | | | 6.6 | 12.1 | 8.2-17.1 | | | |
| Testis | | | | 4.6 | 3.7 | 2.4-5.7 | | | | | | | 0.0 | 0.0 | 0.0-0.9 | | | |
| Urinary bladder (incl. in situ) | 19.2 | 12.8 | 10.2-15.7 | 13.6 | 21.7 | 16.5-27.8 | 5.6 | 6.5 | 4.3-9.4 | 2.0 | 1.5 | 0.7-2.8 | 1.2 | 2.3 | 0.8-4.9 | 0.8 | 1.0 | 0.3-2.5 |
| Kidney and renal pelvis | 23.4 | 13.0 | 10.6-15.7 | 13.6 | 16.3 | 12.3-21.1 | 9.8 | 10.2 | 7.5-13.6 | 1.4 | 1.1 | 0.4-2.2 | 0.6 | 1.3 | 0.3-3.4 | 0.8 | 1.0 | 0.3-2.5 |
| Brain and other central nervous system | 14.6 | 7.2 | 5.6-9.2 | 8.2 | 8.3 | 5.7-11.6 | 6.4 | 6.1 | 4.1-8.7 | 5.6 | 2.7 | 1.8-4.0 | 3.2 | 3.0 | 1.6-5.0 | 2.4 | 2.3 | 1.2-4.1 |
| Thyroid | 37.8 | 17.4 | 14.9-20.1 | 8.4 | 8.0 | 5.7-11.0 | 29.4 | 26.8 | 22.6-31.6 | 1.2 | 0.9 | 0.3-1.9 | 0.2 | 0.3 | 0.0-1.7 | 1.0 | 1.3 | 0.4-2.9 |
| Hodgkin lymphoma | 6.4 | 3.1 | 2.1-4.5 | 3.0 | 3.2 | 1.7-5.6 | 3.4 | 3.2 | 1.8-5.2 | 0.0 | 0.0 | 0.0-0.4 | 0.0 | 0.0 | 0.0-0.9 | 0.0 | 0.0 | 0.0-0.7 |
| Non-Hodgkin lymphomas | 33.4 | 18.2 | 15.4-21.4 | 17.6 | 21.1 | 16.5-26.4 | 15.8 | 16.2 | 12.7-20.2 | 5.2 | 3.4 | 2.2-5.1 | 3.0 | 4.6 | 2.4-7.8 | 2.2 | 2.5 | 1.2-4.5 |
| Myeloma | 15.0 | 9.2 | 7.1-11.6 | 6.4 | 9.0 | 5.9-13.0 | 8.6 | 9.4 | 6.8-12.7 | 2.2 | 1.3 | 0.6-2.4 | 0.4 | 0.4 | 0.0-1.5 | 1.8 | 2.0 | 0.9-3.9 |
| Leukemias | 26.8 | 14.6 | 12.1-17.5 | 16.2 | 19.9 | 15.4-25.2 | 10.6 | 10.9 | 8.1-14.3 | 5.4 | 3.0 | 1.9-4.5 | 3.0 | 2.9 | 1.5-4.9 | 2.4 | 2.8 | 1.4-5.0 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Nassau, Non-Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 8389.2 | 527.9 | 522.6-533.2 | 4175.2 | 567.5 | 559.5-575.5 | 4214.0 | 503.0 | 495.7-510.3 | 2252.8 | 130.4 | 127.9-132.9 | 1090.2 | 147.7 | 143.7-151.8 | 1162.6 | 119.1 | 115.9-122.4 |
| Oral cavity and pharynx | 180.2 | 11.3 | 10.6-12.2 | 118.6 | 16.2 | 14.9-17.7 | 61.6 | 7.2 | 6.4-8.2 | 28.0 | 1.6 | 1.3-1.9 | 14.0 | 1.9 | 1.5-2.4 | 14.0 | 1.4 | 1.0-1.8 |
| Esophagus | 71.6 | 4.3 | 3.8-4.8 | 51.8 | 7.0 | 6.2-8.0 | 19.8 | 2.0 | 1.6-2.5 | 51.0 | 3.1 | 2.7-3.5 | 39.0 | 5.4 | 4.6-6.2 | 12.0 | 1.2 | 0.9-1.6 |
| Stomach | 123.8 | 7.6 | 7.0-8.2 | 73.2 | 10.0 | 8.9-11.1 | 50.6 | 5.7 | 5.0-6.5 | 41.4 | 2.4 | 2.1-2.8 | 22.4 | 3.1 | 2.5-3.7 | 19.0 | 2.0 | 1.6-2.4 |
| Colorectal | 614.4 | 38.4 | 37.0-39.9 | 296.2 | 41.7 | 39.5-44.0 | 318.2 | 35.6 | 33.7-37.5 | 196.4 | 11.4 | 10.7-12.2 | 96.0 | 13.4 | 12.2-14.7 | 100.4 | 9.8 | 8.9-10.8 |
| Colon excluding rectum | 446.4 | 27.4 | 26.3-28.7 | 206.6 | 28.9 | 27.1-30.8 | 239.8 | 26.2 | 24.6-27.8 | 163.2 | 9.4 | 8.7-10.1 | 75.0 | 10.4 | 9.3-11.5 | 88.2 | 8.6 | 7.8-9.5 |
| Rectum & rectosigmoid | 168.0 | 11.0 | 10.2-11.8 | 89.6 | 12.8 | 11.6-14.1 | 78.4 | 9.4 | 8.4-10.4 | 33.2 | 2.0 | 1.7-2.4 | 21.0 | 3.0 | 2.4-3.6 | 12.2 | 1.2 | 0.9-1.6 |
| Liver / intrahepatic bile duct | 109.6 | 6.5 | 5.9-7.1 | 78.0 | 9.9 | 8.9-11.0 | 31.6 | 3.5 | 3.0-4.2 | 78.2 | 4.6 | 4.1-5.1 | 53.4 | 6.9 | 6.1-7.9 | 24.8 | 2.6 | 2.2-3.2 |
| Pancreas | 274.4 | 16.1 | 15.2-17.0 | 137.2 | 18.3 | 16.9-19.8 | 137.2 | 14.3 | 13.2-15.5 | 207.4 | 11.9 | 11.2-12.7 | 103.6 | 13.8 | 12.6-15.1 | 103.8 | 10.3 | 9.4-11.3 |
| Larynx | 42.0 | 2.5 | 2.2-2.9 | 34.4 | 4.5 | 3.9-5.3 | 7.6 | 0.8 | 0.6-1.2 | 12.8 | 0.7 | 0.6-0.9 | 10.4 | 1.3 | 1.0-1.8 | 2.4 | 0.2 | 0.1-0.4 |
| Lung and bronchus | 929.6 | 55.1 | 53.5-56.7 | 408.4 | 55.3 | 52.9-57.8 | 521.2 | 55.6 | 53.4-57.9 | 463.2 | 26.6 | 25.5-27.8 | 220.0 | 29.6 | 27.8-31.4 | 243.2 | 24.7 | 23.3-26.2 |
| Melanoma of the skin | 390.8 | 24.9 | 23.8-26.1 | 232.8 | 32.3 | 30.4-34.3 | 158.0 | 19.8 | 18.3-21.3 | 32.4 | 1.9 | 1.6-2.2 | 21.2 | 2.9 | 2.4-3.6 | 11.2 | 1.1 | 0.8-1.5 |
| Female breast | | | | | | | 1234.0 | 155.8 | 151.7-160.0 | | | | | | | 174.4 | 18.7 | 17.4-20.1 |
| Cervix uteri | | | | | | | 41.0 | 6.0 | 5.1-6.9 | | | | | | | 12.8 | 1.6 | 1.2-2.0 |
| Corpus uterus and NOS | | | | | | | 303.4 | 33.8 | 32.1-35.7 | | | | | | | 60.0 | 6.3 | 5.6-7.1 |
| Ovary | | | | | | | 101.6 | 12.6 | 11.5-13.8 | | | | | | | 55.6 | 6.0 | 5.3-6.8 |
| Prostate | | | | 1163.8 | 146.2 | 142.4-150.1 | | | | | | | 101.8 | 13.7 | 12.5-14.9 | | | |
| Testis | | | | 30.2 | 6.2 | 5.3-7.4 | | | | | | | 1.6 | 0.3 | 0.1-0.5 | | | |
| Urinary bladder (incl. in situ) | 440.4 | 25.8 | 24.7-26.9 | 333.4 | 45.1 | 43.0-47.4 | 107.0 | 11.2 | 10.2-12.3 | 76.8 | 4.2 | 3.8-4.7 | 53.2 | 7.2 | 6.4-8.2 | 23.6 | 2.3 | 1.9-2.8 |
| Kidney and renal pelvis | 295.4 | 18.6 | 17.7-19.7 | 190.6 | 26.1 | 24.4-27.9 | 104.8 | 12.2 | 11.1-13.4 | 48.2 | 2.7 | 2.4-3.1 | 31.6 | 4.1 | 3.5-4.9 | 16.6 | 1.6 | 1.2-2.0 |
| Brain and other central nervous system | 98.8 | 7.2 | 6.5-7.9 | 52.8 | 8.1 | 7.1-9.2 | 46.0 | 6.4 | 5.5-7.3 | 65.0 | 4.1 | 3.7-4.6 | 36.6 | 5.0 | 4.2-5.8 | 28.4 | 3.4 | 2.8-4.1 |
| Thyroid | 309.8 | 25.0 | 23.7-26.3 | 95.8 | 15.3 | 13.9-16.8 | 214.0 | 34.1 | 32.0-36.4 | 11.6 | 0.7 | 0.5-0.9 | 5.0 | 0.7 | 0.5-1.1 | 6.6 | 0.6 | 0.4-0.9 |
| Hodgkin lymphoma | 45.2 | 3.8 | 3.3-4.4 | 23.6 | 4.1 | 3.3-4.9 | 21.6 | 3.7 | 3.0-4.5 | 5.0 | 0.3 | 0.2-0.5 | 2.6 | 0.4 | 0.2-0.6 | 2.4 | 0.3 | 0.1-0.5 |
| Non-Hodgkin lymphomas | 370.8 | 23.5 | 22.4-24.7 | 197.6 | 28.0 | 26.2-29.9 | 173.2 | 19.9 | 18.5-21.4 | 93.8 | 5.4 | 4.9-5.9 | 50.6 | 6.8 | 6.0-7.7 | 43.2 | 4.3 | 3.7-5.0 |
| Myeloma | 140.8 | 8.4 | 7.8-9.0 | 78.8 | 10.6 | 9.6-11.8 | 62.0 | 6.6 | 5.9-7.4 | 50.0 | 2.8 | 2.5-3.2 | 27.2 | 3.6 | 3.0-4.3 | 22.8 | 2.2 | 1.8-2.7 |
| Leukemias | 277.6 | 17.7 | 16.7-18.7 | 165.8 | 23.5 | 21.8-25.2 | 111.8 | 13.3 | 12.1-14.5 | 103.4 | 6.0 | 5.4-6.5 | 62.6 | 8.8 | 7.8-9.8 | 40.8 | 4.0 | 3.4-4.6 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Nassau, Non-Hispanic Whites

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 6997.8 | 554.3 | 548.0-560.5 | 3516.0 | 592.5 | 583.4-601.8 | 3481.8 | 531.6 | 522.9-540.4 | 1918.4 | 134.6 | 131.8-137.5 | 939.2 | 153.4 | 148.9-158.0 | 979.2 | 122.2 | 118.5-126.0 |
| Oral cavity and pharynx | 153.0 | 12.0 | 11.1-13.0 | 102.2 | 17.3 | 15.7-18.9 | 50.8 | 7.5 | 6.5-8.6 | 23.2 | 1.6 | 1.3-2.0 | 11.0 | 1.8 | 1.4-2.4 | 12.2 | 1.4 | 1.1-1.9 |
| Esophagus | 61.4 | 4.5 | 4.0-5.0 | 45.4 | 7.5 | 6.5-8.6 | 16.0 | 1.9 | 1.5-2.5 | 45.2 | 3.3 | 2.9-3.8 | 35.4 | 6.0 | 5.1-7.0 | 9.8 | 1.1 | 0.8-1.5 |
| Stomach | 88.6 | 6.6 | 6.0-7.3 | 56.0 | 9.3 | 8.2-10.5 | 32.6 | 4.6 | 3.8-5.4 | 27.6 | 1.9 | 1.6-2.3 | 15.8 | 2.6 | 2.0-3.3 | 11.8 | 1.4 | 1.0-1.9 |
| Colorectal | 498.8 | 38.8 | 37.1-40.4 | 239.4 | 42.0 | 39.5-44.6 | 259.4 | 35.9 | 33.7-38.1 | 163.2 | 11.6 | 10.8-12.5 | 78.4 | 13.5 | 12.1-14.9 | 84.8 | 10.0 | 9.0-11.1 |
| Colon excluding rectum | 365.0 | 27.6 | 26.3-29.0 | 168.0 | 28.9 | 26.9-31.1 | 197.0 | 26.5 | 24.7-28.4 | 135.4 | 9.4 | 8.7-10.2 | 61.2 | 10.4 | 9.2-11.7 | 74.2 | 8.6 | 7.7-9.7 |
| Rectum & rectosigmoid | 133.8 | 11.1 | 10.2-12.1 | 71.4 | 13.1 | 11.7-14.6 | 62.4 | 9.4 | 8.3-10.6 | 27.8 | 2.1 | 1.8-2.6 | 17.2 | 3.1 | 2.4-3.9 | 10.6 | 1.4 | 1.0-1.9 |
| Liver / intrahepatic bile duct | 83.6 | 6.1 | 5.5-6.8 | 62.0 | 9.7 | 8.6-10.9 | 21.6 | 3.0 | 2.4-3.8 | 60.6 | 4.3 | 3.8-4.8 | 42.6 | 6.7 | 5.8-7.7 | 18.0 | 2.3 | 1.8-2.9 |
| Pancreas | 232.4 | 16.8 | 15.8-17.8 | 117.6 | 19.3 | 17.7-21.0 | 114.8 | 14.8 | 13.5-16.2 | 177.6 | 12.4 | 11.5-13.3 | 89.8 | 14.6 | 13.2-16.1 | 87.8 | 10.5 | 9.5-11.6 |
| Larynx | 36.6 | 2.8 | 2.4-3.2 | 30.2 | 5.0 | 4.2-5.9 | 6.4 | 0.9 | 0.6-1.3 | 10.4 | 0.7 | 0.5-1.0 | 8.2 | 1.3 | 0.9-1.8 | 2.2 | 0.3 | 0.1-0.5 |
| Lung and bronchus | 815.4 | 59.4 | 57.5-61.3 | 351.0 | 57.7 | 55.0-60.6 | 464.4 | 61.5 | 59.0-64.2 | 403.2 | 28.1 | 26.9-29.4 | 189.4 | 30.6 | 28.7-32.7 | 213.8 | 26.6 | 24.9-28.3 |
| Melanoma of the skin | 383.0 | 31.4 | 29.9-33.0 | 228.8 | 39.5 | 37.2-42.1 | 154.2 | 25.7 | 23.7-27.8 | 32.0 | 2.3 | 2.0-2.8 | 21.0 | 3.6 | 2.9-4.4 | 11.0 | 1.4 | 1.0-1.9 |
| Female breast | | | | | | | 995.2 | 163.4 | 158.4-168.5 | | | | | | | 142.6 | 18.8 | 17.4-20.4 |
| Cervix uteri | | | | | | | 27.0 | 5.6 | 4.6-6.7 | | | | | | | 8.8 | 1.4 | 1.0-1.9 |
| Corpus uterus and NOS | | | | | | | 243.2 | 34.7 | 32.7-36.8 | | | | | | | 45.8 | 6.0 | 5.2-6.9 |
| Ovary | | | | | | | 81.0 | 12.9 | 11.5-14.4 | | | | | | | 47.8 | 6.5 | 5.6-7.4 |
| Prostate | | | | 905.0 | 140.1 | 136.0-144.4 | | | | | | | 85.2 | 13.4 | 12.1-14.8 | | | |
| Testis | | | | 27.2 | 8.0 | 6.6-9.5 | | | | | | | 1.4 | 0.3 | 0.1-0.6 | | | |
| Urinary bladder (incl. in situ) | 403.6 | 29.0 | 27.7-30.3 | 310.2 | 50.9 | 48.4-53.6 | 93.4 | 12.0 | 10.9-13.3 | 71.8 | 4.7 | 4.2-5.3 | 51.0 | 8.2 | 7.2-9.3 | 20.8 | 2.5 | 2.0-3.1 |
| Kidney and renal pelvis | 253.8 | 20.4 | 19.2-21.7 | 164.8 | 28.4 | 26.4-30.6 | 89.0 | 13.3 | 12.0-14.8 | 41.8 | 2.8 | 2.5-3.3 | 27.4 | 4.3 | 3.6-5.1 | 14.4 | 1.7 | 1.3-2.1 |
| Brain and other central nervous system | 83.8 | 7.9 | 7.1-8.8 | 44.8 | 8.8 | 7.6-10.2 | 39.0 | 7.1 | 6.0-8.3 | 58.8 | 4.7 | 4.1-5.3 | 33.2 | 5.6 | 4.7-6.6 | 25.6 | 3.9 | 3.2-4.8 |
| Thyroid | 248.2 | 27.5 | 25.8-29.2 | 81.0 | 17.2 | 15.5-19.2 | 167.2 | 37.5 | 34.8-40.4 | 9.4 | 0.6 | 0.5-0.9 | 4.0 | 0.7 | 0.4-1.1 | 5.4 | 0.6 | 0.4-0.9 |
| Hodgkin lymphoma | 35.8 | 4.1 | 3.5-4.8 | 19.2 | 4.4 | 3.5-5.4 | 16.6 | 3.9 | 3.1-5.0 | 4.2 | 0.3 | 0.2-0.5 | 2.2 | 0.4 | 0.2-0.7 | 2.0 | 0.2 | 0.1-0.5 |
| Non-Hodgkin lymphomas | 317.2 | 24.9 | 23.6-26.2 | 167.8 | 29.3 | 27.2-31.5 | 149.4 | 21.3 | 19.7-23.1 | 80.4 | 5.5 | 4.9-6.1 | 43.0 | 6.9 | 6.0-8.0 | 37.4 | 4.4 | 3.8-5.2 |
| Myeloma | 104.8 | 7.8 | 7.1-8.5 | 61.6 | 10.2 | 9.1-11.5 | 43.2 | 5.8 | 5.0-6.7 | 40.8 | 2.7 | 2.4-3.2 | 23.0 | 3.7 | 3.0-4.5 | 17.8 | 2.0 | 1.6-2.5 |
| Leukemias | 239.8 | 19.1 | 18.0-20.3 | 145.2 | 25.2 | 23.3-27.2 | 94.6 | 14.4 | 13.0-15.9 | 92.6 | 6.5 | 5.9-7.1 | 56.4 | 9.5 | 8.4-10.7 | 36.2 | 4.4 | 3.7-5.2 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Nassau, Non-Hispanic Blacks**

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 819.6 | 458.8 | 444.6-473.4 | 398.0 | 533.7 | 509.6-558.6 | 421.6 | 412.9 | 395.1-431.3 | 214.4 | 124.3 | 116.9-132.1 | 90.0 | 132.8 | 120.4-146.1 | 124.4 | 119.8 | 110.5-129.8 |
| Oral cavity and pharynx | 11.6 | 6.4 | 4.8-8.3 | 6.8 | 8.5 | 5.8-12.0 | 4.8 | 4.8 | 3.0-7.2 | 3.0 | 1.6 | 0.9-2.6 | 1.8 | 2.2 | 1.0-4.2 | 1.2 | 1.1 | 0.4-2.5 |
| Esophagus | 6.6 | 3.8 | 2.6-5.5 | 4.0 | 5.9 | 3.5-9.2 | 2.6 | 2.5 | 1.3-4.3 | 3.2 | 1.8 | 1.0-3.0 | 2.2 | 3.1 | 1.5-5.6 | 1.0 | 0.9 | 0.3-2.3 |
| Stomach | 21.2 | 12.3 | 10.0-14.9 | 10.4 | 14.9 | 11.0-19.8 | 10.8 | 10.5 | 7.9-13.8 | 7.8 | 4.5 | 3.2-6.2 | 3.4 | 5.3 | 3.0-8.5 | 4.4 | 4.2 | 2.6-6.4 |
| Colorectal | 73.4 | 41.6 | 37.3-46.2 | 36.2 | 49.7 | 42.4-57.9 | 37.2 | 36.1 | 31.0-41.8 | 21.8 | 12.8 | 10.4-15.5 | 10.8 | 15.8 | 11.7-20.7 | 11.0 | 10.6 | 8.0-13.9 |
| Colon excluding rectum | 55.2 | 31.3 | 27.7-35.3 | 25.8 | 35.8 | 29.7-42.8 | 29.4 | 28.2 | 23.8-33.3 | 18.4 | 10.7 | 8.6-13.2 | 8.8 | 12.6 | 9.0-17.0 | 9.6 | 9.3 | 6.8-12.4 |
| Rectum & rectosigmoid | 18.2 | 10.3 | 8.2-12.7 | 10.4 | 13.9 | 10.2-18.5 | 7.8 | 7.9 | 5.5-10.9 | 3.4 | 2.0 | 1.2-3.3 | 2.0 | 3.2 | 1.5-5.9 | 1.4 | 1.3 | 0.5-2.8 |
| Liver / intrahepatic bile duct | 10.2 | 5.5 | 4.1-7.3 | 6.4 | 8.3 | 5.6-11.8 | 3.8 | 3.5 | 2.1-5.5 | 8.4 | 4.8 | 3.4-6.5 | 5.4 | 7.0 | 4.6-10.3 | 3.0 | 3.0 | 1.7-5.1 |
| Pancreas | 26.0 | 14.6 | 12.2-17.4 | 10.4 | 14.2 | 10.5-18.8 | 15.6 | 14.8 | 11.7-18.6 | 19.2 | 11.3 | 9.1-13.9 | 8.0 | 11.6 | 8.1-15.9 | 11.2 | 11.0 | 8.3-14.4 |
| Larynx | 3.8 | 2.0 | 1.2-3.2 | 3.0 | 3.7 | 2.0-6.2 | 0.8 | 0.7 | 0.2-2.0 | 1.6 | 0.9 | 0.4-1.8 | 1.4 | 1.9 | 0.7-4.0 | 0.2 | 0.2 | 0.0-1.2 |
| Lung and bronchus | 69.0 | 39.5 | 35.4-44.0 | 34.0 | 50.5 | 42.9-58.9 | 35.0 | 33.2 | 28.4-38.6 | 37.4 | 21.6 | 18.6-25.0 | 18.4 | 27.9 | 22.3-34.4 | 19.0 | 17.9 | 14.5-22.0 |
| Melanoma of the skin | 0.6 | 0.3 | 0.1-1.0 | 0.2 | 0.4 | 0.0-2.0 | 0.4 | 0.3 | 0.0-1.4 | 0.2 | 0.1 | 0.0-0.8 | 0.2 | 0.4 | 0.0-2.0 | 0.0 | 0.0 | 0.0-0.9 |
| Female breast | | | | | | | 135.6 | 134.7 | 124.5-145.5 | | | | | | | 22.2 | 20.9 | 17.2-25.4 |
| Cervix uteri | | | | | | | 8.6 | 8.4 | 6.1-11.5 | | | | | | | 3.4 | 3.5 | 2.0-5.7 |
| Corpus uterus and NOS | | | | | | | 39.8 | 36.5 | 31.6-42.1 | | | | | | | 11.0 | 10.2 | 7.6-13.3 |
| Ovary | | | | | | | 10.4 | 10.3 | 7.6-13.6 | | | | | | | 4.8 | 4.9 | 3.1-7.3 |
| Prostate | | | | 182.4 | 232.3 | 216.9-248.5 | | | | | | | 12.6 | 20.8 | 15.8-26.7 | | | |
| Testis | | | | 0.6 | 0.9 | 0.2-2.6 | | | | | | | 0.0 | 0.0 | 0.0-1.1 | | | |
| Urinary bladder (incl. in situ) | 18.6 | 10.9 | 8.7-13.4 | 10.8 | 15.8 | 11.7-20.8 | 7.8 | 7.8 | 5.5-10.7 | 3.6 | 2.2 | 1.3-3.5 | 1.4 | 2.4 | 0.9-4.9 | 2.2 | 2.2 | 1.1-4.0 |
| Kidney and renal pelvis | 28.8 | 16.2 | 13.6-19.1 | 17.2 | 22.7 | 18.0-28.3 | 11.6 | 11.5 | 8.7-15.0 | 4.6 | 2.6 | 1.7-4.0 | 3.2 | 4.7 | 2.6-7.7 | 1.4 | 1.4 | 0.5-3.0 |
| Brain and other central nervous system | 6.4 | 4.0 | 2.7-5.7 | 3.0 | 4.0 | 2.2-6.8 | 3.4 | 4.1 | 2.3-6.6 | 4.0 | 2.3 | 1.4-3.6 | 1.8 | 2.3 | 1.0-4.5 | 2.2 | 2.2 | 1.1-4.0 |
| Thyroid | 19.4 | 11.1 | 9.0-13.6 | 4.6 | 6.1 | 3.8-9.3 | 14.8 | 15.3 | 11.9-19.3 | 0.8 | 0.5 | 0.1-1.2 | 0.6 | 0.9 | 0.2-2.6 | 0.2 | 0.2 | 0.0-1.2 |
| Hodgkin lymphoma | 4.4 | 2.7 | 1.7-4.1 | 1.8 | 2.4 | 1.1-4.6 | 2.6 | 2.8 | 1.5-4.9 | 0.4 | 0.3 | 0.0-1.1 | 0.2 | 0.3 | 0.0-1.8 | 0.2 | 0.3 | 0.0-1.5 |
| Non-Hodgkin lymphomas | 25.8 | 15.3 | 12.7-18.2 | 13.8 | 19.9 | 15.3-25.3 | 12.0 | 12.2 | 9.3-15.8 | 7.2 | 4.1 | 2.9-5.7 | 3.2 | 4.3 | 2.4-7.1 | 4.0 | 4.0 | 2.4-6.2 |
| Myeloma | 27.4 | 15.0 | 12.6-17.8 | 12.0 | 16.3 | 12.3-21.3 | 15.4 | 14.2 | 11.2-17.8 | 7.4 | 4.3 | 3.0-6.0 | 3.4 | 5.0 | 2.8-8.1 | 4.0 | 3.9 | 2.4-6.1 |
| Leukemias | 19.4 | 11.5 | 9.3-14.1 | 10.4 | 15.5 | 11.5-20.4 | 9.0 | 9.0 | 6.5-12.2 | 5.8 | 3.4 | 2.2-4.9 | 2.4 | 3.6 | 1.8-6.4 | 3.4 | 3.1 | 1.8-5.1 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Suffolk, Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-----------|----------------|------------------|------------|----------------|------------------|-----------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 760.0 | 390.2 | 376.8-403.8 | 341.0 | 401.7 | 380.3-423.8 | 419.0 | 395.4 | 377.9-413.5 | 144.2 | 88.9 | 82.1-96.1 | 69.6 | 100.5 | 89.1-112.8 | 74.6 | 81.9 | 73.4-90.9 |
| Oral cavity and pharynx | 14.6 | 6.9 | 5.3-8.8 | 9.6 | 9.7 | 6.8-13.3 | 5.0 | 4.5 | 2.8-6.7 | 3.0 | 1.8 | 1.0-3.0 | 2.0 | 2.6 | 1.1-5.0 | 1.0 | 1.1 | 0.3-2.5 |
| Esophagus | 4.2 | 2.7 | 1.6-4.1 | 3.2 | 4.5 | 2.4-7.6 | 1.0 | 1.3 | 0.4-2.9 | 2.6 | 1.5 | 0.8-2.7 | 1.8 | 2.7 | 1.1-5.1 | 0.8 | 0.7 | 0.2-2.0 |
| Stomach | 22.0 | 12.2 | 9.9-14.9 | 12.0 | 15.4 | 11.3-20.3 | 10.0 | 10.1 | 7.4-13.5 | 7.2 | 4.5 | 3.0-6.3 | 4.6 | 7.1 | 4.3-10.8 | 2.6 | 2.7 | 1.4-4.6 |
| Colorectal | 63.8 | 33.3 | 29.4-37.4 | 33.2 | 35.7 | 29.8-42.4 | 30.6 | 31.0 | 26.1-36.6 | 10.2 | 6.8 | 4.9-9.0 | 5.6 | 8.3 | 5.2-12.4 | 4.6 | 5.5 | 3.5-8.3 |
| Colon excluding rectum | 45.0 | 24.7 | 21.4-28.4 | 21.4 | 24.3 | 19.3-30.0 | 23.6 | 24.9 | 20.4-30.0 | 8.2 | 5.5 | 3.9-7.5 | 4.4 | 6.8 | 4.0-10.6 | 3.8 | 4.5 | 2.7-7.0 |
| Rectum & rectosigmoid | 18.8 | 8.6 | 6.8-10.6 | 11.8 | 11.4 | 8.4-15.2 | 7.0 | 6.2 | 4.2-8.6 | 2.0 | 1.3 | 0.6-2.3 | 1.2 | 1.5 | 0.4-3.4 | 0.8 | 1.0 | 0.3-2.5 |
| Liver / intrahepatic bile duct | 23.2 | 13.4 | 10.9-16.2 | 15.0 | 18.9 | 14.5-24.2 | 8.2 | 8.8 | 6.2-12.0 | 11.8 | 7.1 | 5.3-9.3 | 7.8 | 10.1 | 6.9-14.2 | 4.0 | 4.6 | 2.8-7.2 |
| Pancreas | 24.4 | 13.1 | 10.7-15.8 | 11.8 | 13.2 | 9.7-17.5 | 12.6 | 12.9 | 9.8-16.6 | 11.6 | 7.0 | 5.2-9.2 | 6.0 | 8.1 | 5.2-11.8 | 5.6 | 6.3 | 4.1-9.1 |
| Larynx | 2.4 | 1.3 | 0.6-2.4 | 2.0 | 2.4 | 1.0-4.5 | 0.4 | 0.4 | 0.0-1.5 | 0.8 | 0.6 | 0.1-1.4 | 0.4 | 0.4 | 0.0-1.7 | 0.4 | 0.5 | 0.1-1.7 |
| Lung and bronchus | 54.8 | 35.2 | 30.9-39.8 | 24.2 | 36.6 | 29.8-44.2 | 30.6 | 34.6 | 29.2-40.8 | 22.0 | 14.2 | 11.5-17.2 | 10.2 | 14.9 | 10.7-20.0 | 11.8 | 13.6 | 10.2-17.6 |
| Melanoma of the skin | 9.8 | 5.9 | 4.2-7.9 | 6.2 | 9.5 | 6.2-13.9 | 3.6 | 3.6 | 2.0-5.8 | 0.8 | 0.4 | 0.1-1.0 | 0.6 | 0.7 | 0.1-2.0 | 0.2 | 0.1 | 0.0-0.9 |
| Female breast | | | | | | | 113.8 | 104.6 | 95.9-114.0 | | | | | | | 9.2 | 10.1 | 7.2-13.5 |
| Cervix uteri | | | | | | | 10.8 | 8.9 | 6.6-11.8 | | | | | | | 2.0 | 1.7 | 0.8-3.1 |
| Corpus uterus and NOS | | | | | | | 25.0 | 23.2 | 19.2-27.8 | | | | | | | 4.2 | 4.3 | 2.6-6.6 |
| Ovary | | | | | | | 12.4 | 11.3 | 8.5-14.7 | | | | | | | 4.4 | 4.6 | 2.8-7.0 |
| Prostate | | | | 85.0 | 101.0 | 90.7-112.0 | | | | | | | 6.2 | 12.9 | 8.6-18.3 | | | |
| Testis | | | | 8.2 | 5.2 | 3.7-7.2 | | | | | | | 0.4 | 0.3 | 0.0-1.2 | | | |
| Urinary bladder (incl. in situ) | 19.4 | 13.1 | 10.5-16.1 | 14.6 | 22.9 | 17.5-29.2 | 4.8 | 6.0 | 3.8-8.9 | 3.0 | 2.1 | 1.1-3.4 | 2.4 | 4.3 | 2.1-7.5 | 0.6 | 0.6 | 0.1-1.7 |
| Kidney and renal pelvis | 27.6 | 13.5 | 11.2-16.2 | 16.8 | 17.0 | 13.1-21.6 | 10.8 | 10.4 | 7.7-13.7 | 3.0 | 1.8 | 1.0-3.0 | 1.4 | 1.4 | 0.5-3.2 | 1.6 | 2.0 | 0.9-4.0 |
| Brain and other central nervous system | 14.8 | 6.1 | 4.7-7.8 | 8.0 | 8.5 | 5.6-12.1 | 6.8 | 4.7 | 3.2-6.6 | 5.0 | 2.2 | 1.4-3.4 | 3.0 | 3.3 | 1.6-5.9 | 2.0 | 1.4 | 0.7-2.7 |
| Thyroid | 60.2 | 22.8 | 20.2-25.6 | 9.2 | 7.0 | 5.0-9.6 | 51.0 | 39.6 | 34.8-44.9 | 0.4 | 0.2 | 0.0-0.8 | 0.2 | 0.1 | 0.0-1.0 | 0.2 | 0.3 | 0.0-1.2 |
| Hodgkin lymphoma | 5.4 | 2.7 | 1.7-4.0 | 2.8 | 2.6 | 1.3-4.6 | 2.6 | 2.7 | 1.4-4.7 | 0.4 | 0.2 | 0.0-0.9 | 0.0 | 0.0 | 0.0-0.8 | 0.4 | 0.4 | 0.0-1.6 |
| Non-Hodgkin lymphomas | 37.6 | 19.5 | 16.6-22.8 | 19.4 | 22.6 | 17.7-28.3 | 18.2 | 17.5 | 13.9-21.7 | 5.8 | 3.9 | 2.6-5.7 | 2.2 | 3.4 | 1.6-6.1 | 3.6 | 4.4 | 2.6-7.0 |
| Myeloma | 16.0 | 8.8 | 6.8-11.0 | 8.2 | 9.0 | 6.2-12.6 | 7.8 | 8.3 | 5.8-11.4 | 5.0 | 3.2 | 2.0-4.8 | 2.2 | 3.0 | 1.4-5.5 | 2.8 | 3.2 | 1.7-5.4 |
| Leukemias | 31.6 | 14.5 | 12.1-17.2 | 17.0 | 17.2 | 13.2-21.9 | 14.6 | 12.8 | 9.9-16.2 | 7.8 | 4.6 | 3.1-6.4 | 4.4 | 6.1 | 3.6-9.5 | 3.4 | 3.5 | 1.9-5.6 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Suffolk, Non-Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 9291.2 | 549.8 | 544.6-555.1 | 4751.0 | 606.6 | 598.7-614.7 | 4540.2 | 510.7 | 503.7-517.8 | 2548.2 | 143.8 | 141.3-146.4 | 1285.8 | 167.5 | 163.3-171.7 | 1262.4 | 127.7 | 124.5-131.0 |
| Oral cavity and pharynx | 217.0 | 12.5 | 11.8-13.3 | 150.6 | 18.7 | 17.4-20.2 | 66.4 | 7.2 | 6.4-8.1 | 35.0 | 2.0 | 1.7-2.4 | 24.8 | 3.2 | 2.7-3.9 | 10.2 | 1.1 | 0.8-1.4 |
| Esophagus | 89.2 | 5.0 | 4.5-5.5 | 69.2 | 8.4 | 7.5-9.4 | 20.0 | 2.0 | 1.6-2.5 | 70.6 | 3.8 | 3.5-4.3 | 57.2 | 7.0 | 6.2-7.9 | 13.4 | 1.3 | 1.0-1.7 |
| Stomach | 123.8 | 7.1 | 6.5-7.7 | 74.6 | 9.4 | 8.4-10.4 | 49.2 | 5.3 | 4.6-6.0 | 47.6 | 2.7 | 2.4-3.1 | 26.6 | 3.4 | 2.8-4.0 | 21.0 | 2.2 | 1.8-2.7 |
| Colorectal | 702.4 | 41.7 | 40.2-43.1 | 369.8 | 48.2 | 45.9-50.5 | 332.6 | 36.4 | 34.6-38.3 | 215.4 | 12.3 | 11.5-13.1 | 111.2 | 14.7 | 13.5-16.0 | 104.2 | 10.5 | 9.6-11.5 |
| Colon excluding rectum | 498.2 | 29.5 | 28.4-30.8 | 246.8 | 32.4 | 30.6-34.3 | 251.4 | 27.2 | 25.7-28.9 | 172.6 | 9.9 | 9.2-10.6 | 86.6 | 11.5 | 10.4-12.7 | 86.0 | 8.6 | 7.8-9.5 |
| Rectum & rectosigmoid | 204.2 | 12.1 | 11.4-12.9 | 123.0 | 15.8 | 14.5-17.1 | 81.2 | 9.1 | 8.2-10.1 | 42.8 | 2.4 | 2.1-2.8 | 24.6 | 3.2 | 2.6-3.8 | 18.2 | 1.9 | 1.5-2.3 |
| Liver / intrahepatic bile duct | 130.4 | 7.3 | 6.7-7.9 | 95.4 | 11.5 | 10.5-12.7 | 35.0 | 3.6 | 3.1-4.2 | 91.8 | 5.1 | 4.6-5.6 | 59.8 | 7.5 | 6.6-8.4 | 32.0 | 3.2 | 2.7-3.7 |
| Pancreas | 286.8 | 16.3 | 15.5-17.2 | 150.0 | 19.0 | 17.6-20.4 | 136.8 | 14.1 | 13.0-15.2 | 204.0 | 11.5 | 10.8-12.3 | 105.6 | 13.4 | 12.3-14.6 | 98.4 | 9.9 | 9.1-10.9 |
| Larynx | 57.2 | 3.2 | 2.8-3.6 | 46.0 | 5.7 | 4.9-6.5 | 11.2 | 1.1 | 0.8-1.5 | 14.2 | 0.8 | 0.6-1.0 | 12.0 | 1.5 | 1.1-2.0 | 2.2 | 0.2 | 0.1-0.4 |
| Lung and bronchus | 1208.8 | 68.0 | 66.3-69.8 | 557.4 | 71.1 | 68.4-73.9 | 651.4 | 66.3 | 64.0-68.7 | 598.6 | 33.5 | 32.3-34.8 | 292.0 | 37.8 | 35.8-39.8 | 306.6 | 30.5 | 29.0-32.1 |
| Melanoma of the skin | 472.4 | 28.4 | 27.2-29.7 | 287.0 | 37.3 | 35.3-39.4 | 185.4 | 21.8 | 20.4-23.4 | 37.8 | 2.1 | 1.8-2.4 | 25.2 | 3.2 | 2.7-3.8 | 12.6 | 1.3 | 1.0-1.7 |
| Female breast | | | | | | | 1249.8 | 144.9 | 141.1-148.8 | | | | | | | 177.6 | 18.4 | 17.1-19.7 |
| Cervix uteri | | | | | | | 41.4 | 5.9 | 5.0-6.8 | | | | | | | 13.2 | 1.5 | 1.2-2.0 |
| Corpus uterus and NOS | | | | | | | 311.4 | 33.0 | 31.3-34.8 | | | | | | | 55.0 | 5.7 | 5.0-6.4 |
| Ovary | | | | | | | 112.2 | 12.8 | 11.7-13.9 | | | | | | | 70.6 | 7.2 | 6.4-8.0 |
| Prostate | | | | 1186.8 | 139.8 | 136.2-143.5 | | | | | | | 120.6 | 16.3 | 15.0-17.6 | | | |
| Testis | | | | 41.0 | 7.8 | 6.7-9.0 | | | | | | | 1.4 | 0.2 | 0.1-0.4 | | | |
| Urinary bladder (incl. in situ) | 511.0 | 29.2 | 28.0-30.3 | 378.6 | 49.5 | 47.2-51.8 | 132.4 | 13.5 | 12.5-14.6 | 89.2 | 5.0 | 4.5-5.4 | 63.2 | 8.6 | 7.7-9.6 | 26.0 | 2.5 | 2.1-3.0 |
| Kidney and renal pelvis | 347.8 | 20.6 | 19.6-21.7 | 223.0 | 28.5 | 26.7-30.3 | 124.8 | 13.9 | 12.7-15.1 | 54.0 | 3.0 | 2.6-3.4 | 35.6 | 4.5 | 3.9-5.3 | 18.4 | 1.8 | 1.5-2.3 |
| Brain and other central nervous system | 110.0 | 7.4 | 6.8-8.1 | 60.0 | 8.6 | 7.6-9.7 | 50.0 | 6.2 | 5.4-7.2 | 75.8 | 4.4 | 4.0-4.9 | 40.2 | 5.1 | 4.4-5.9 | 35.6 | 3.9 | 3.3-4.5 |
| Thyroid | 305.2 | 23.1 | 21.9-24.4 | 93.8 | 13.7 | 12.4-15.1 | 211.4 | 32.2 | 30.1-34.3 | 10.4 | 0.6 | 0.4-0.8 | 4.4 | 0.6 | 0.3-0.9 | 6.0 | 0.6 | 0.4-0.9 |
| Hodgkin lymphoma | 45.0 | 3.5 | 3.1-4.0 | 23.4 | 3.6 | 2.9-4.3 | 21.6 | 3.5 | 2.8-4.2 | 4.2 | 0.3 | 0.2-0.5 | 2.0 | 0.3 | 0.1-0.6 | 2.2 | 0.3 | 0.1-0.5 |
| Non-Hodgkin lymphomas | 399.4 | 23.9 | 22.8-25.0 | 219.8 | 29.0 | 27.3-30.8 | 179.6 | 19.8 | 18.5-21.3 | 84.6 | 4.8 | 4.3-5.3 | 49.6 | 6.6 | 5.8-7.5 | 35.0 | 3.5 | 3.0-4.0 |
| Myeloma | 153.0 | 8.8 | 8.2-9.5 | 88.6 | 11.4 | 10.4-12.6 | 64.4 | 6.7 | 5.9-7.5 | 49.6 | 2.7 | 2.4-3.1 | 28.6 | 3.7 | 3.1-4.3 | 21.0 | 2.0 | 1.6-2.5 |
| Leukemias | 342.0 | 20.6 | 19.6-21.6 | 203.6 | 27.1 | 25.4-28.9 | 138.4 | 15.4 | 14.2-16.6 | 100.6 | 5.9 | 5.4-6.4 | 60.4 | 8.3 | 7.4-9.3 | 40.2 | 4.1 | 3.5-4.8 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Suffolk, Non-Hispanic Whites

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 8459.8 | 559.3 | 553.7-564.9 | 4326.6 | 614.1 | 605.6-622.6 | 4133.2 | 522.4 | 514.8-530.1 | 2340.6 | 145.3 | 142.6-148.1 | 1179.8 | 168.6 | 164.2-173.0 | 1160.8 | 129.5 | 126.1-133.1 |
| Oral cavity and pharynx | 202.8 | 13.0 | 12.2-13.9 | 141.8 | 19.6 | 18.1-21.1 | 61.0 | 7.4 | 6.5-8.3 | 32.4 | 2.1 | 1.8-2.4 | 22.6 | 3.2 | 2.6-3.9 | 9.8 | 1.2 | 0.8-1.6 |
| Esophagus | 83.4 | 5.2 | 4.7-5.7 | 65.0 | 8.8 | 7.8-9.8 | 18.4 | 2.1 | 1.7-2.6 | 66.4 | 4.0 | 3.6-4.5 | 53.6 | 7.2 | 6.3-8.1 | 12.8 | 1.4 | 1.1-1.8 |
| Stomach | 110.0 | 6.9 | 6.4-7.6 | 66.6 | 9.3 | 8.3-10.4 | 43.4 | 5.1 | 4.4-5.8 | 40.8 | 2.6 | 2.2-3.0 | 22.2 | 3.1 | 2.5-3.8 | 18.6 | 2.2 | 1.7-2.8 |
| Colorectal | 631.4 | 41.8 | 40.3-43.4 | 334.4 | 48.4 | 46.0-50.9 | 297.0 | 36.4 | 34.5-38.5 | 197.0 | 12.4 | 11.6-13.3 | 102.8 | 15.0 | 13.7-16.4 | 94.2 | 10.4 | 9.5-11.5 |
| Colon excluding rectum | 451.8 | 29.9 | 28.6-31.3 | 226.4 | 33.1 | 31.1-35.2 | 225.4 | 27.3 | 25.6-29.1 | 157.6 | 9.9 | 9.2-10.7 | 80.0 | 11.8 | 10.6-13.0 | 77.6 | 8.5 | 7.7-9.5 |
| Rectum & rectosigmoid | 179.6 | 11.9 | 11.1-12.8 | 108.0 | 15.3 | 14.0-16.8 | 71.6 | 9.1 | 8.1-10.2 | 39.4 | 2.5 | 2.1-2.9 | 22.8 | 3.3 | 2.7-4.0 | 16.6 | 1.9 | 1.5-2.4 |
| Liver / intrahepatic bile duct | 112.4 | 6.9 | 6.4-7.6 | 81.0 | 10.9 | 9.8-12.0 | 31.4 | 3.6 | 3.0-4.3 | 78.8 | 4.8 | 4.3-5.3 | 51.4 | 7.0 | 6.2-8.0 | 27.4 | 2.9 | 2.5-3.5 |
| Pancreas | 261.0 | 16.3 | 15.4-17.2 | 137.4 | 19.0 | 17.6-20.6 | 123.6 | 14.0 | 12.9-15.2 | 188.2 | 11.7 | 10.9-12.5 | 97.0 | 13.5 | 12.3-14.9 | 91.2 | 10.1 | 9.2-11.1 |
| Larynx | 52.4 | 3.2 | 2.8-3.7 | 41.8 | 5.8 | 5.0-6.6 | 10.6 | 1.2 | 0.9-1.6 | 13.0 | 0.8 | 0.6-1.0 | 10.8 | 1.5 | 1.1-2.0 | 2.2 | 0.2 | 0.1-0.5 |
| Lung and bronchus | 1133.6 | 70.3 | 68.4-72.2 | 517.0 | 72.4 | 69.6-75.4 | 616.6 | 69.5 | 67.0-72.0 | 559.2 | 34.4 | 33.2-35.8 | 269.4 | 38.3 | 36.2-40.4 | 289.8 | 31.9 | 30.2-33.6 |
| Melanoma of the skin | 461.2 | 31.3 | 29.9-32.7 | 281.2 | 40.6 | 38.5-42.9 | 180.0 | 24.3 | 22.6-26.1 | 37.0 | 2.2 | 1.9-2.6 | 24.6 | 3.4 | 2.8-4.1 | 12.4 | 1.4 | 1.1-1.9 |
| Female breast | | | | | | | 1126.4 | 147.3 | 143.2-151.5 | | | | | | | 160.4 | 18.3 | 17.0-19.7 |
| Cervix uteri | | | | | | | 35.0 | 5.8 | 4.9-6.8 | | | | | | | 11.8 | 1.6 | 1.2-2.1 |
| Corpus uterus and NOS | | | | | | | 277.0 | 32.8 | 31.0-34.7 | | | | | | | 48.6 | 5.5 | 4.8-6.3 |
| Ovary | | | | | | | 103.2 | 13.2 | 12.0-14.5 | | | | | | | 66.2 | 7.4 | 6.6-8.3 |
| Prostate | | | | 1033.0 | 134.7 | 131.0-138.5 | | | | | | | 106.6 | 15.6 | 14.3-17.0 | | | |
| Testis | | | | 39.0 | 8.9 | 7.7-10.3 | | | | | | | 1.4 | 0.2 | 0.1-0.5 | | | |
| Urinary bladder (incl. in situ) | 485.8 | 30.6 | 29.3-31.9 | 360.6 | 51.8 | 49.3-54.3 | 125.2 | 14.2 | 13.0-15.4 | 84.4 | 5.1 | 4.6-5.6 | 60.2 | 8.9 | 7.9-10.0 | 24.2 | 2.5 | 2.1-3.0 |
| Kidney and renal pelvis | 312.8 | 20.8 | 19.7-21.9 | 199.6 | 28.5 | 26.6-30.4 | 113.2 | 14.1 | 12.9-15.4 | 49.2 | 3.0 | 2.6-3.4 | 32.8 | 4.6 | 3.9-5.3 | 16.4 | 1.8 | 1.4-2.3 |
| Brain and other central nervous system | 101.2 | 7.7 | 7.0-8.5 | 55.6 | 9.0 | 7.9-10.2 | 45.6 | 6.5 | 5.6-7.6 | 71.0 | 4.7 | 4.2-5.2 | 38.8 | 5.5 | 4.7-6.4 | 32.2 | 3.9 | 3.3-4.6 |
| Thyroid | 274.2 | 24.3 | 23.0-25.8 | 87.8 | 14.7 | 13.3-16.3 | 186.4 | 33.7 | 31.4-36.1 | 9.6 | 0.6 | 0.4-0.8 | 4.2 | 0.6 | 0.4-1.0 | 5.4 | 0.6 | 0.4-0.9 |
| Hodgkin lymphoma | 40.8 | 3.9 | 3.3-4.5 | 21.0 | 3.8 | 3.1-4.7 | 19.8 | 3.9 | 3.1-4.8 | 4.0 | 0.3 | 0.2-0.5 | 1.8 | 0.3 | 0.1-0.6 | 2.2 | 0.3 | 0.1-0.6 |
| Non-Hodgkin lymphomas | 366.0 | 24.3 | 23.2-25.5 | 201.4 | 29.4 | 27.6-31.4 | 164.6 | 20.3 | 18.8-21.8 | 78.6 | 4.9 | 4.4-5.4 | 45.8 | 6.6 | 5.8-7.5 | 32.8 | 3.6 | 3.0-4.2 |
| Myeloma | 131.4 | 8.3 | 7.7-9.0 | 76.2 | 10.8 | 9.7-12.0 | 55.2 | 6.4 | 5.6-7.2 | 42.2 | 2.5 | 2.2-2.9 | 25.2 | 3.5 | 2.9-4.2 | 17.0 | 1.8 | 1.4-2.2 |
| Leukemias | 319.8 | 21.6 | 20.5-22.8 | 191.6 | 28.6 | 26.7-30.5 | 128.2 | 15.9 | 14.6-17.4 | 94.0 | 6.0 | 5.5-6.6 | 56.0 | 8.4 | 7.4-9.5 | 38.0 | 4.3 | 3.7-5.0 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Suffolk, Non-Hispanic Blacks

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 577.6 | 516.1 | 496.8-535.9 | 293.8 | 604.6 | 572.1-638.4 | 283.8 | 455.5 | 431.7-480.4 | 149.4 | 143.6 | 133.2-154.6 | 76.6 | 184.8 | 165.7-205.5 | 72.8 | 117.8 | 105.8-130.8 |
| Oral cavity and pharynx | 8.6 | 7.4 | 5.3-10.0 | 5.2 | 9.8 | 6.3-14.7 | 3.4 | 5.3 | 3.0-8.6 | 1.6 | 1.6 | 0.7-3.1 | 1.4 | 3.2 | 1.3-6.7 | 0.2 | 0.3 | 0.0-1.8 |
| Esophagus | 4.8 | 4.0 | 2.5-6.1 | 3.4 | 7.1 | 3.9-11.8 | 1.4 | 1.9 | 0.7-4.0 | 3.6 | 3.4 | 2.0-5.5 | 3.0 | 7.5 | 4.0-12.6 | 0.6 | 0.8 | 0.2-2.6 |
| Stomach | 10.0 | 8.9 | 6.5-11.8 | 5.6 | 12.2 | 7.9-18.0 | 4.4 | 6.9 | 4.2-10.5 | 3.8 | 3.6 | 2.1-5.6 | 2.8 | 6.7 | 3.5-11.4 | 1.0 | 1.5 | 0.5-3.7 |
| Colorectal | 46.4 | 42.1 | 36.7-48.1 | 21.2 | 47.4 | 38.2-58.1 | 25.2 | 39.4 | 32.7-47.1 | 12.6 | 12.1 | 9.2-15.6 | 6.0 | 15.5 | 10.2-22.4 | 6.6 | 10.4 | 7.1-14.8 |
| Colon excluding rectum | 32.4 | 30.3 | 25.7-35.5 | 13.2 | 31.4 | 23.8-40.5 | 19.2 | 30.4 | 24.5-37.3 | 10.6 | 10.2 | 7.6-13.4 | 4.4 | 11.6 | 7.1-17.8 | 6.2 | 9.7 | 6.5-13.8 |
| Rectum & rectosigmoid | 14.0 | 11.8 | 9.2-15.1 | 8.0 | 16.0 | 11.2-22.2 | 6.0 | 9.0 | 6.0-13.0 | 2.0 | 1.9 | 0.9-3.6 | 1.6 | 3.9 | 1.5-7.9 | 0.4 | 0.8 | 0.1-2.7 |
| Liver / intrahepatic bile duct | 9.4 | 7.6 | 5.5-10.2 | 8.0 | 14.6 | 10.3-20.2 | 1.4 | 2.0 | 0.8-4.2 | 6.2 | 5.1 | 3.4-7.4 | 4.0 | 7.6 | 4.5-12.1 | 2.2 | 3.2 | 1.6-6.0 |
| Pancreas | 18.2 | 17.5 | 14.0-21.6 | 9.2 | 21.8 | 15.6-29.5 | 9.0 | 14.6 | 10.6-19.6 | 11.0 | 10.4 | 7.7-13.6 | 6.4 | 14.7 | 9.8-21.1 | 4.6 | 7.4 | 4.6-11.3 |
| Larynx | 3.6 | 3.0 | 1.8-4.9 | 3.0 | 6.0 | 3.3-10.1 | 0.6 | 0.9 | 0.2-2.7 | 0.8 | 0.6 | 0.1-1.5 | 0.8 | 1.2 | 0.3-3.4 | 0.0 | 0.0 | 0.0-1.3 |
| Lung and bronchus | 56.2 | 52.5 | 46.4-59.3 | 28.4 | 63.8 | 53.1-75.8 | 27.8 | 44.9 | 37.6-53.2 | 29.6 | 27.9 | 23.4-32.9 | 17.8 | 41.3 | 32.7-51.3 | 11.8 | 18.5 | 14.0-24.1 |
| Melanoma of the skin | 1.6 | 1.4 | 0.6-2.8 | 0.8 | 1.5 | 0.4-4.0 | 0.8 | 1.3 | 0.3-3.5 | 0.6 | 0.6 | 0.1-1.8 | 0.4 | 0.9 | 0.1-3.2 | 0.2 | 0.3 | 0.0-1.9 |
| Female breast | | | | | | | 86.0 | 137.5 | 124.5-151.4 | | | | | | | 12.4 | 18.9 | 14.4-24.4 |
| Cervix uteri | | | | | | | 5.0 | 8.4 | 5.4-12.5 | | | | | | | 1.0 | 1.7 | 0.5-4.1 |
| Corpus uterus and NOS | | | | | | | 23.8 | 37.0 | 30.5-44.5 | | | | | | | 5.4 | 9.0 | 5.9-13.3 |
| Ovary | | | | | | | 5.6 | 9.2 | 6.0-13.3 | | | | | | | 3.2 | 5.7 | 3.3-9.3 |
| Prostate | | | | 121.4 | 237.7 | 218.1-258.6 | | | | | | | 11.0 | 30.1 | 22.4-39.5 | | | |
| Testis | | | | 1.4 | 2.7 | 1.1-5.7 | | | | | | | 0.0 | 0.0 | 0.0-1.6 | | | |
| Urinary bladder (incl. in situ) | 17.8 | 16.9 | 13.5-20.9 | 12.4 | 28.2 | 21.2-36.8 | 5.4 | 8.9 | 5.8-13.1 | 3.8 | 4.1 | 2.5-6.5 | 2.0 | 6.3 | 2.9-11.4 | 1.8 | 3.1 | 1.4-6.0 |
| Kidney and renal pelvis | 25.0 | 21.6 | 17.9-25.9 | 15.4 | 30.4 | 23.6-38.5 | 9.6 | 15.2 | 11.1-20.3 | 3.2 | 3.4 | 1.9-5.5 | 2.0 | 4.6 | 2.1-8.6 | 1.2 | 2.3 | 0.8-5.0 |
| Brain and other central nervous system | 6.0 | 5.3 | 3.6-7.7 | 3.0 | 5.8 | 3.2-9.7 | 3.0 | 4.7 | 2.6-7.8 | 3.6 | 3.1 | 1.8-4.9 | 1.2 | 2.0 | 0.7-4.7 | 2.4 | 3.7 | 1.9-6.6 |
| Thyroid | 15.2 | 13.6 | 10.7-17.1 | 2.8 | 5.7 | 3.1-9.7 | 12.4 | 20.2 | 15.4-26.1 | 0.6 | 0.5 | 0.1-1.6 | 0.2 | 0.3 | 0.0-2.1 | 0.4 | 0.6 | 0.1-2.3 |
| Hodgkin lymphoma | 3.2 | 2.6 | 1.5-4.4 | 1.8 | 3.1 | 1.4-6.1 | 1.4 | 2.1 | 0.8-4.5 | 0.2 | 0.1 | 0.0-0.9 | 0.2 | 0.3 | 0.0-2.0 | 0.0 | 0.0 | 0.0-1.3 |
| Non-Hodgkin lymphomas | 22.0 | 19.4 | 15.8-23.5 | 12.2 | 25.6 | 19.2-33.3 | 9.8 | 15.2 | 11.1-20.2 | 4.2 | 4.1 | 2.5-6.3 | 3.0 | 7.5 | 4.0-12.4 | 1.2 | 1.8 | 0.6-4.1 |
| Myeloma | 16.6 | 15.8 | 12.4-19.7 | 9.2 | 19.3 | 13.8-26.2 | 7.4 | 12.5 | 8.8-17.3 | 5.4 | 5.5 | 3.6-8.1 | 2.6 | 7.2 | 3.7-12.3 | 2.8 | 4.7 | 2.6-8.0 |
| Leukemias | 13.0 | 12.1 | 9.2-15.5 | 6.6 | 13.1 | 8.8-18.7 | 6.4 | 11.2 | 7.6-15.8 | 4.8 | 4.9 | 3.1-7.4 | 3.0 | 7.6 | 4.1-12.7 | 1.8 | 3.1 | 1.4-5.9 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Westchester, Hispanic Origin**

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-----------|----------------|------------------|------------|----------------|------------------|-----------|----------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | |
| All Invasive Malignant Tumors | 688.0 | 380.2 | 366.9-393.9 | 317.8 | 410.2 | 388.4-432.8 | 370.2 | 366.7 | 349.6-384.5 | 136.8 | 86.3 | 79.7-93.4 | 69.6 | 107.8 | 95.9-120.6 | 67.2 | 73.1 | 65.3-81.7 | |
| Oral cavity and pharynx | 12.6 | 7.1 | 5.3-9.2 | 8.0 | 9.7 | 6.7-13.5 | 4.6 | 4.9 | 3.0-7.4 | 3.0 | 1.9 | 1.0-3.2 | 2.2 | 3.3 | 1.5-5.9 | 0.8 | 0.9 | 0.2-2.3 | |
| Esophagus | 3.2 | 2.0 | 1.1-3.3 | 2.6 | 3.5 | 1.7-6.1 | 0.6 | 0.9 | 0.2-2.4 | 0.8 | 0.5 | 0.1-1.2 | 0.8 | 1.1 | 0.3-2.8 | 0.0 | 0.0 | 0.0-0.7 | |
| Stomach | 23.0 | 13.7 | 11.2-16.6 | 11.2 | 14.9 | 11.0-19.7 | 11.8 | 12.9 | 9.7-16.7 | 7.4 | 4.5 | 3.1-6.2 | 3.8 | 5.5 | 3.2-8.8 | 3.6 | 3.9 | 2.3-6.2 | |
| Colorectal | 60.4 | 33.7 | 29.7-37.9 | 31.8 | 40.5 | 33.8-47.9 | 28.6 | 28.7 | 24.0-33.9 | 11.6 | 7.0 | 5.2-9.1 | 6.0 | 8.7 | 5.6-12.6 | 5.6 | 6.0 | 3.9-8.7 | |
| Colon excluding rectum | 41.6 | 23.8 | 20.5-27.5 | 20.2 | 26.2 | 20.9-32.4 | 21.4 | 22.1 | 18.0-26.8 | 8.6 | 5.2 | 3.7-7.1 | 3.6 | 5.5 | 3.1-8.8 | 5.0 | 5.3 | 3.4-7.8 | |
| Rectum & rectosigmoid | 18.8 | 9.8 | 7.8-12.2 | 11.6 | 14.2 | 10.5-18.8 | 7.2 | 6.6 | 4.6-9.2 | 3.0 | 1.8 | 1.0-3.0 | 2.4 | 3.2 | 1.5-5.7 | 0.6 | 0.7 | 0.1-2.0 | |
| Liver / intrahepatic bile duct | 19.6 | 11.9 | 9.5-14.6 | 12.6 | 17.9 | 13.4-23.2 | 7.0 | 7.4 | 5.1-10.4 | 10.4 | 6.4 | 4.7-8.5 | 7.2 | 10.0 | 6.8-14.1 | 3.2 | 3.8 | 2.1-6.2 | |
| Pancreas | 22.4 | 13.9 | 11.4-16.9 | 10.4 | 13.7 | 10.0-18.4 | 12.0 | 13.9 | 10.5-18.0 | 11.2 | 7.4 | 5.5-9.7 | 5.4 | 7.5 | 4.7-11.1 | 5.8 | 7.1 | 4.7-10.2 | |
| Larynx | 4.2 | 2.6 | 1.6-4.0 | 3.6 | 5.5 | 3.1-8.8 | 0.6 | 0.5 | 0.1-1.5 | 1.0 | 0.7 | 0.2-1.5 | 0.8 | 1.4 | 0.4-3.4 | 0.2 | 0.2 | 0.0-1.0 | |
| Lung and bronchus | 48.2 | 30.7 | 26.7-35.0 | 23.2 | 36.4 | 29.6-44.1 | 25.0 | 27.1 | 22.4-32.5 | 19.4 | 13.0 | 10.4-16.0 | 13.0 | 21.3 | 16.1-27.4 | 6.4 | 7.2 | 4.9-10.3 | |
| Melanoma of the skin | 5.8 | 3.4 | 2.2-5.0 | 3.8 | 5.2 | 2.9-8.4 | 2.0 | 2.2 | 1.0-4.0 | 1.4 | 0.6 | 0.2-1.4 | 1.0 | 0.9 | 0.3-2.1 | 0.4 | 0.4 | 0.0-1.4 | |
| Female breast | | | | | | | 115.2 | 109.2 | 100.2-118.8 | | | | | | | | 12.8 | 12.9 | 9.8-16.5 |
| Cervix uteri | | | | | | | 8.4 | 7.5 | 5.3-10.1 | | | | | | | | 0.6 | 0.5 | 0.1-1.6 |
| Corpus uterus and NOS | | | | | | | 23.8 | 23.6 | 19.4-28.4 | | | | | | | | 4.0 | 3.9 | 2.4-6.1 |
| Ovary | | | | | | | 10.0 | 9.9 | 7.3-13.2 | | | | | | | | 3.0 | 3.1 | 1.7-5.1 |
| Prostate | | | | 88.8 | 114.8 | 103.5-126.9 | | | | | | | 6.6 | 12.8 | 8.7-17.9 | | | | |
| Testis | | | | 5.8 | 4.4 | 3.0-6.5 | | | | | | | 0.4 | 0.3 | 0.0-1.4 | | | | |
| Urinary bladder (incl. in situ) | 17.6 | 11.8 | 9.4-14.6 | 12.6 | 20.0 | 15.1-25.9 | 5.0 | 5.8 | 3.7-8.6 | 3.0 | 2.2 | 1.2-3.6 | 2.0 | 3.8 | 1.7-6.8 | 1.0 | 1.3 | 0.4-2.9 | |
| Kidney and renal pelvis | 22.6 | 11.8 | 9.6-14.4 | 15.2 | 17.6 | 13.6-22.5 | 7.4 | 7.1 | 4.9-9.9 | 2.4 | 1.6 | 0.8-2.9 | 2.0 | 3.2 | 1.4-5.9 | 0.4 | 0.5 | 0.1-1.8 | |
| Brain and other central nervous system | 9.2 | 4.1 | 3.0-5.6 | 5.8 | 5.2 | 3.4-7.6 | 3.4 | 3.1 | 1.8-5.0 | 3.2 | 1.7 | 0.9-2.7 | 1.8 | 2.4 | 1.0-4.7 | 1.4 | 1.2 | 0.5-2.5 | |
| Thyroid | 42.8 | 19.2 | 16.6-22.1 | 9.0 | 7.7 | 5.6-10.5 | 33.8 | 30.0 | 25.5-35.0 | 0.4 | 0.3 | 0.0-1.0 | 0.2 | 0.4 | 0.0-1.9 | 0.2 | 0.3 | 0.0-1.3 | |
| Hodgkin lymphoma | 5.8 | 2.7 | 1.7-3.9 | 3.2 | 3.6 | 1.9-6.0 | 2.6 | 2.1 | 1.1-3.7 | 0.6 | 0.4 | 0.1-1.2 | 0.2 | 0.2 | 0.0-1.3 | 0.4 | 0.6 | 0.1-1.9 | |
| Non-Hodgkin lymphomas | 35.4 | 19.5 | 16.6-22.8 | 20.0 | 24.7 | 19.6-30.5 | 15.4 | 15.5 | 12.1-19.6 | 8.0 | 5.6 | 3.9-7.6 | 3.4 | 6.0 | 3.4-9.5 | 4.6 | 5.4 | 3.4-8.1 | |
| Myeloma | 15.0 | 8.9 | 6.9-11.2 | 6.6 | 9.0 | 5.9-12.9 | 8.4 | 8.8 | 6.2-11.9 | 2.8 | 2.1 | 1.1-3.5 | 1.2 | 2.6 | 1.0-5.4 | 1.6 | 1.8 | 0.8-3.6 | |
| Leukemias | 26.6 | 14.6 | 12.1-17.5 | 14.6 | 18.5 | 14.1-23.8 | 12.0 | 11.7 | 8.8-15.1 | 6.6 | 4.0 | 2.7-5.7 | 4.2 | 6.3 | 3.7-9.8 | 2.4 | 2.4 | 1.2-4.2 | |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Westchester, Non-Hispanic Origin

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 5156.0 | 494.1 | 487.8-500.4 | 2536.0 | 534.9 | 525.4-544.6 | 2620.0 | 469.4 | 460.8-478.0 | 1390.4 | 120.9 | 118.0-123.9 | 669.8 | 138.7 | 133.9-143.6 | 720.6 | 109.2 | 105.4-113.0 |
| Oral cavity and pharynx | 105.0 | 10.0 | 9.2-11.0 | 72.6 | 15.1 | 13.6-16.9 | 32.4 | 5.8 | 4.9-6.9 | 18.0 | 1.6 | 1.3-2.0 | 12.2 | 2.5 | 1.9-3.3 | 5.8 | 0.9 | 0.6-1.4 |
| Esophagus | 43.8 | 4.0 | 3.5-4.6 | 30.8 | 6.5 | 5.5-7.6 | 13.0 | 2.0 | 1.6-2.7 | 32.8 | 2.9 | 2.5-3.5 | 23.8 | 5.0 | 4.2-6.1 | 9.0 | 1.4 | 1.0-1.9 |
| Stomach | 80.2 | 7.5 | 6.7-8.3 | 47.2 | 10.2 | 8.9-11.6 | 33.0 | 5.4 | 4.6-6.4 | 32.8 | 2.9 | 2.5-3.5 | 19.6 | 4.1 | 3.3-5.0 | 13.2 | 2.1 | 1.6-2.8 |
| Colorectal | 357.0 | 33.9 | 32.3-35.6 | 175.8 | 38.1 | 35.5-40.8 | 181.2 | 30.7 | 28.6-32.9 | 121.8 | 10.5 | 9.6-11.4 | 57.4 | 12.0 | 10.6-13.5 | 64.4 | 9.2 | 8.2-10.4 |
| Colon excluding rectum | 255.0 | 24.1 | 22.7-25.5 | 122.2 | 26.5 | 24.4-28.8 | 132.8 | 22.2 | 20.4-24.1 | 97.4 | 8.3 | 7.6-9.1 | 46.6 | 9.7 | 8.5-11.1 | 50.8 | 7.2 | 6.3-8.2 |
| Rectum & rectosigmoid | 102.0 | 9.9 | 9.0-10.8 | 53.6 | 11.5 | 10.2-13.1 | 48.4 | 8.5 | 7.4-9.8 | 24.4 | 2.2 | 1.8-2.6 | 10.8 | 2.3 | 1.7-3.1 | 13.6 | 2.0 | 1.5-2.6 |
| Liver / intrahepatic bile duct | 78.2 | 7.0 | 6.3-7.8 | 53.4 | 10.6 | 9.4-12.1 | 24.8 | 4.0 | 3.3-4.9 | 50.0 | 4.4 | 3.8-5.0 | 34.2 | 6.7 | 5.7-7.8 | 15.8 | 2.6 | 2.0-3.3 |
| Pancreas | 162.2 | 14.5 | 13.5-15.6 | 82.6 | 17.4 | 15.8-19.3 | 79.6 | 12.2 | 11.0-13.5 | 115.8 | 10.1 | 9.2-11.0 | 56.6 | 11.7 | 10.3-13.2 | 59.2 | 8.8 | 7.8-9.9 |
| Larynx | 27.0 | 2.4 | 2.0-2.9 | 21.8 | 4.3 | 3.5-5.3 | 5.2 | 0.9 | 0.6-1.3 | 8.2 | 0.7 | 0.5-1.0 | 7.4 | 1.5 | 1.0-2.1 | 0.8 | 0.1 | 0.0-0.4 |
| Lung and bronchus | 549.2 | 48.9 | 47.0-50.8 | 243.8 | 50.4 | 47.5-53.4 | 305.4 | 48.2 | 45.7-50.8 | 286.4 | 24.7 | 23.4-26.1 | 130.0 | 26.7 | 24.7-28.9 | 156.4 | 23.1 | 21.4-24.9 |
| Melanoma of the skin | 224.4 | 21.4 | 20.1-22.8 | 129.2 | 27.7 | 25.5-30.0 | 95.2 | 16.9 | 15.3-18.6 | 17.4 | 1.5 | 1.2-1.8 | 10.8 | 2.2 | 1.7-3.0 | 6.6 | 0.9 | 0.6-1.3 |
| Female breast | | | | | | | 808.4 | 153.5 | 148.5-158.6 | | | | | | | 107.8 | 17.7 | 16.1-19.4 |
| Cervix uteri | | | | | | | 24.6 | 5.5 | 4.5-6.7 | | | | | | | 7.0 | 1.3 | 0.9-1.9 |
| Corpus uterus and NOS | | | | | | | 170.4 | 29.2 | 27.2-31.3 | | | | | | | 37.0 | 5.8 | 5.0-6.7 |
| Ovary | | | | | | | 75.0 | 14.3 | 12.7-15.9 | | | | | | | 44.0 | 7.0 | 6.1-8.1 |
| Prostate | | | | 713.6 | 141.5 | 136.8-146.3 | | | | | | | 79.6 | 16.4 | 14.8-18.1 | | | |
| Testis | | | | 19.4 | 6.3 | 5.1-7.7 | | | | | | | 0.4 | 0.1 | 0.0-0.4 | | | |
| Urinary bladder (incl. in situ) | 260.6 | 23.1 | 21.8-24.4 | 189.4 | 39.2 | 36.7-41.9 | 71.2 | 11.2 | 10.0-12.5 | 41.4 | 3.4 | 2.9-3.9 | 28.8 | 5.9 | 5.0-7.0 | 12.6 | 1.7 | 1.3-2.2 |
| Kidney and renal pelvis | 167.0 | 16.3 | 15.1-17.5 | 110.6 | 23.7 | 21.7-25.9 | 56.4 | 10.2 | 9.0-11.5 | 23.0 | 2.1 | 1.7-2.5 | 16.0 | 3.4 | 2.7-4.2 | 7.0 | 1.1 | 0.8-1.6 |
| Brain and other central nervous system | 60.2 | 6.8 | 6.0-7.6 | 31.8 | 7.7 | 6.5-9.1 | 28.4 | 5.9 | 4.9-7.1 | 39.2 | 3.9 | 3.4-4.5 | 21.6 | 4.9 | 4.0-6.0 | 17.6 | 3.0 | 2.4-3.8 |
| Thyroid | 173.2 | 21.4 | 19.9-23.0 | 52.2 | 13.1 | 11.4-14.9 | 121.0 | 29.2 | 26.8-31.8 | 8.2 | 0.7 | 0.5-1.0 | 5.0 | 1.0 | 0.7-1.5 | 3.2 | 0.5 | 0.3-0.8 |
| Hodgkin lymphoma | 25.6 | 3.3 | 2.8-4.0 | 11.8 | 3.2 | 2.4-4.2 | 13.8 | 3.5 | 2.7-4.5 | 2.4 | 0.2 | 0.1-0.4 | 1.0 | 0.2 | 0.1-0.5 | 1.4 | 0.2 | 0.1-0.5 |
| Non-Hodgkin lymphomas | 251.0 | 24.1 | 22.7-25.6 | 137.8 | 30.3 | 28.0-32.7 | 113.2 | 19.3 | 17.7-21.1 | 48.0 | 4.0 | 3.5-4.6 | 25.0 | 5.0 | 4.2-6.0 | 23.0 | 3.3 | 2.7-4.0 |
| Myeloma | 97.2 | 9.1 | 8.3-10.0 | 53.0 | 11.2 | 9.8-12.7 | 44.2 | 7.5 | 6.5-8.6 | 36.4 | 3.1 | 2.7-3.6 | 19.2 | 4.0 | 3.2-4.9 | 17.2 | 2.5 | 2.0-3.1 |
| Leukemias | 199.8 | 19.1 | 17.8-20.4 | 116.4 | 25.3 | 23.2-27.5 | 83.4 | 14.2 | 12.7-15.7 | 63.4 | 5.4 | 4.8-6.1 | 35.0 | 7.2 | 6.2-8.4 | 28.4 | 4.1 | 3.5-5.0 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Westchester, Non-Hispanic Whites

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 4153.0 | 512.5 | 505.0-520.0 | 2053.8 | 545.2 | 534.2-556.4 | 2099.2 | 495.3 | 484.9-505.8 | 1118.6 | 120.1 | 116.8-123.4 | 539.8 | 136.2 | 131.0-141.6 | 578.8 | 109.6 | 105.3-114.0 |
| Oral cavity and pharynx | 86.8 | 10.6 | 9.6-11.7 | 59.4 | 15.6 | 13.8-17.6 | 27.4 | 6.3 | 5.2-7.6 | 14.6 | 1.6 | 1.3-2.1 | 10.0 | 2.5 | 1.9-3.4 | 4.6 | 0.9 | 0.5-1.4 |
| Esophagus | 35.0 | 4.1 | 3.5-4.8 | 24.8 | 6.6 | 5.4-7.9 | 10.2 | 2.0 | 1.5-2.8 | 27.0 | 3.0 | 2.5-3.6 | 19.0 | 4.9 | 4.0-6.1 | 8.0 | 1.5 | 1.1-2.2 |
| Stomach | 56.6 | 6.6 | 5.8-7.5 | 37.2 | 9.9 | 8.5-11.5 | 19.4 | 4.1 | 3.2-5.1 | 21.4 | 2.3 | 1.9-2.9 | 14.0 | 3.5 | 2.7-4.4 | 7.4 | 1.5 | 1.0-2.3 |
| Colorectal | 276.2 | 33.7 | 31.8-35.6 | 137.6 | 37.8 | 34.9-40.9 | 138.6 | 30.4 | 27.9-33.0 | 95.6 | 10.1 | 9.1-11.1 | 44.2 | 11.2 | 9.7-12.9 | 51.4 | 9.1 | 7.9-10.5 |
| Colon excluding rectum | 198.2 | 23.9 | 22.4-25.6 | 96.6 | 26.5 | 24.1-29.1 | 101.6 | 21.9 | 19.8-24.1 | 76.2 | 7.8 | 7.0-8.7 | 35.0 | 8.8 | 7.5-10.2 | 41.2 | 7.1 | 6.1-8.2 |
| Rectum & rectosigmoid | 78.0 | 9.7 | 8.7-10.8 | 41.0 | 11.3 | 9.7-13.1 | 37.0 | 8.5 | 7.2-9.9 | 19.4 | 2.2 | 1.8-2.8 | 9.2 | 2.5 | 1.8-3.3 | 10.2 | 2.0 | 1.5-2.8 |
| Liver / intrahepatic bile duct | 56.4 | 6.3 | 5.6-7.2 | 39.2 | 9.8 | 8.4-11.3 | 17.2 | 3.4 | 2.7-4.3 | 34.2 | 3.7 | 3.1-4.3 | 24.4 | 5.9 | 4.9-7.1 | 9.8 | 2.0 | 1.4-2.7 |
| Pancreas | 132.4 | 14.9 | 13.7-16.2 | 70.2 | 18.5 | 16.5-20.6 | 62.2 | 12.0 | 10.6-13.6 | 94.4 | 10.2 | 9.3-11.3 | 47.2 | 12.1 | 10.6-13.8 | 47.2 | 8.7 | 7.6-10.0 |
| Larynx | 22.6 | 2.6 | 2.1-3.1 | 18.6 | 4.6 | 3.7-5.7 | 4.0 | 0.9 | 0.5-1.5 | 6.2 | 0.6 | 0.4-1.0 | 5.6 | 1.4 | 0.9-2.0 | 0.6 | 0.1 | 0.0-0.5 |
| Lung and bronchus | 457.4 | 50.8 | 48.7-53.0 | 196.6 | 50.0 | 46.8-53.3 | 260.8 | 52.2 | 49.3-55.3 | 238.6 | 25.5 | 24.0-27.0 | 104.4 | 26.2 | 24.0-28.6 | 134.2 | 24.8 | 22.9-26.9 |
| Melanoma of the skin | 217.8 | 27.1 | 25.4-28.9 | 125.8 | 34.0 | 31.3-36.9 | 92.0 | 22.0 | 19.8-24.4 | 17.2 | 1.8 | 1.4-2.3 | 10.6 | 2.7 | 2.0-3.6 | 6.6 | 1.1 | 0.7-1.6 |
| Female breast | | | | | | | 639.6 | 162.9 | 156.8-169.2 | | | | | | | 84.2 | 17.6 | 15.8-19.6 |
| Cervix uteri | | | | | | | 15.0 | 4.9 | 3.7-6.3 | | | | | | | 5.2 | 1.3 | 0.8-2.0 |
| Corpus uterus and NOS | | | | | | | 130.8 | 29.1 | 26.9-31.6 | | | | | | | 26.4 | 5.2 | 4.3-6.2 |
| Ovary | | | | | | | 61.2 | 15.4 | 13.5-17.4 | | | | | | | 36.2 | 7.4 | 6.3-8.6 |
| Prostate | | | | 527.4 | 130.0 | 125.0-135.2 | | | | | | | 64.0 | 15.6 | 13.9-17.5 | | | |
| Testis | | | | 17.6 | 8.3 | 6.6-10.3 | | | | | | | 0.4 | 0.2 | 0.0-0.6 | | | |
| Urinary bladder (incl. in situ) | 232.6 | 26.0 | 24.5-27.7 | 170.2 | 43.5 | 40.6-46.7 | 62.4 | 12.6 | 11.1-14.2 | 35.0 | 3.4 | 2.9-4.0 | 24.4 | 6.0 | 5.0-7.2 | 10.6 | 1.7 | 1.2-2.3 |
| Kidney and renal pelvis | 133.6 | 16.9 | 15.6-18.4 | 90.4 | 24.8 | 22.5-27.4 | 43.2 | 10.3 | 8.8-11.9 | 18.8 | 2.1 | 1.7-2.6 | 13.0 | 3.4 | 2.6-4.4 | 5.8 | 1.2 | 0.7-1.9 |
| Brain and other central nervous system | 49.6 | 7.4 | 6.4-8.5 | 25.8 | 8.2 | 6.7-9.9 | 23.8 | 6.6 | 5.3-8.1 | 32.4 | 4.1 | 3.5-4.9 | 18.0 | 5.2 | 4.1-6.5 | 14.4 | 3.1 | 2.4-4.1 |
| Thyroid | 138.2 | 24.0 | 22.0-26.0 | 44.2 | 15.0 | 12.9-17.3 | 94.0 | 32.8 | 29.6-36.2 | 6.2 | 0.7 | 0.4-1.0 | 3.8 | 1.0 | 0.6-1.6 | 2.4 | 0.4 | 0.2-0.9 |
| Hodgkin lymphoma | 20.8 | 3.9 | 3.1-4.7 | 8.6 | 3.2 | 2.3-4.4 | 12.2 | 4.5 | 3.4-5.9 | 2.2 | 0.2 | 0.1-0.4 | 0.8 | 0.2 | 0.0-0.6 | 1.4 | 0.2 | 0.1-0.6 |
| Non-Hodgkin lymphomas | 209.8 | 25.7 | 24.0-27.4 | 118.2 | 32.6 | 29.9-35.5 | 91.6 | 20.1 | 18.1-22.2 | 39.8 | 4.1 | 3.5-4.7 | 20.8 | 5.1 | 4.2-6.2 | 19.0 | 3.2 | 2.6-4.1 |
| Myeloma | 66.4 | 7.9 | 7.0-8.8 | 38.0 | 10.0 | 8.6-11.7 | 28.4 | 6.1 | 5.1-7.3 | 25.0 | 2.6 | 2.1-3.1 | 14.2 | 3.5 | 2.7-4.5 | 10.8 | 1.9 | 1.4-2.5 |
| Leukemias | 173.8 | 21.2 | 19.8-22.8 | 103.2 | 28.2 | 25.7-30.9 | 70.6 | 15.5 | 13.8-17.4 | 53.0 | 5.6 | 5.0-6.4 | 29.8 | 7.5 | 6.3-8.9 | 23.2 | 4.2 | 3.4-5.2 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Gender and Hispanic Ethnicity, 2015-2019
Westchester, Non-Hispanic Blacks

| Site of cancer | Incidence | | | | | | | | | Mortality | | | | | | | | |
|--|-------------------|------------------|-------------|---------------|------------------|-------------|---------------|------------------|-------------|-------------------|------------------|-------------|----------------|------------------|-------------|----------------|------------------|-------------|
| | Males and Females | | | Males | | | Females | | | Males and Females | | | Males | | | Females | | |
| | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Cases | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI | Avg Ann Deaths | Rate per 100,000 | 95% CI |
| All Invasive Malignant Tumors | 749.6 | 467.5 | 452.4-483.1 | 357.0 | 545.3 | 519.5-572.1 | 392.6 | 422.6 | 403.7-442.3 | 216.0 | 136.3 | 128.2-144.8 | 100.4 | 167.0 | 152.2-182.7 | 115.6 | 119.1 | 109.4-129.5 |
| Oral cavity and pharynx | 11.4 | 7.5 | 5.7-9.8 | 8.2 | 13.0 | 9.2-17.8 | 3.2 | 4.0 | 2.2-6.6 | 2.2 | 1.5 | 0.8-2.8 | 1.6 | 2.7 | 1.1-5.4 | 0.6 | 0.8 | 0.2-2.3 |
| Esophagus | 7.0 | 4.2 | 2.9-5.9 | 4.6 | 7.2 | 4.5-10.9 | 2.4 | 2.3 | 1.2-4.2 | 4.8 | 3.1 | 1.9-4.6 | 4.0 | 7.0 | 4.2-10.8 | 0.8 | 0.7 | 0.2-2.1 |
| Stomach | 18.0 | 11.9 | 9.5-14.7 | 6.8 | 12.1 | 8.3-16.9 | 11.2 | 11.8 | 8.9-15.5 | 8.2 | 5.4 | 3.8-7.3 | 3.8 | 6.4 | 3.8-10.0 | 4.4 | 4.6 | 2.8-7.2 |
| Colorectal | 59.6 | 37.3 | 33.1-41.8 | 26.6 | 41.9 | 34.9-50.0 | 33.0 | 34.7 | 29.5-40.6 | 21.2 | 13.4 | 10.9-16.2 | 10.0 | 16.4 | 12.0-21.8 | 11.2 | 11.3 | 8.5-14.8 |
| Colon excluding rectum | 43.4 | 27.4 | 23.8-31.4 | 18.4 | 30.0 | 24.0-37.1 | 25.0 | 26.1 | 21.6-31.3 | 17.0 | 10.8 | 8.6-13.4 | 8.6 | 14.3 | 10.2-19.5 | 8.4 | 8.8 | 6.3-12.0 |
| Rectum & rectosigmoid | 16.2 | 9.9 | 7.8-12.4 | 8.2 | 11.9 | 8.5-16.3 | 8.0 | 8.6 | 6.1-11.8 | 4.2 | 2.6 | 1.6-4.0 | 1.4 | 2.1 | 0.8-4.4 | 2.8 | 2.5 | 1.4-4.4 |
| Liver / intrahepatic bile duct | 16.0 | 9.3 | 7.3-11.6 | 10.4 | 14.4 | 10.7-19.1 | 5.6 | 5.8 | 3.8-8.6 | 11.2 | 6.7 | 5.0-8.7 | 6.8 | 9.5 | 6.5-13.5 | 4.4 | 4.7 | 2.9-7.3 |
| Pancreas | 23.8 | 15.1 | 12.4-18.1 | 9.6 | 15.5 | 11.3-20.7 | 14.2 | 14.7 | 11.4-18.7 | 16.6 | 10.5 | 8.3-13.1 | 7.0 | 11.3 | 7.8-15.9 | 9.6 | 10.0 | 7.3-13.4 |
| Larynx | 4.0 | 2.3 | 1.4-3.6 | 3.2 | 4.4 | 2.5-7.4 | 0.8 | 0.8 | 0.2-2.2 | 1.8 | 1.1 | 0.5-2.2 | 1.6 | 2.7 | 1.1-5.4 | 0.2 | 0.2 | 0.0-1.3 |
| Lung and bronchus | 71.6 | 44.9 | 40.3-49.9 | 35.2 | 56.6 | 48.3-65.9 | 36.4 | 37.5 | 32.2-43.6 | 38.6 | 24.4 | 21.1-28.2 | 20.0 | 33.8 | 27.3-41.3 | 18.6 | 18.8 | 15.1-23.1 |
| Melanoma of the skin | 0.8 | 0.4 | 0.1-1.2 | 0.4 | 0.5 | 0.1-1.9 | 0.4 | 0.4 | 0.0-1.6 | 0.0 | 0.0 | 0.0-0.5 | 0.0 | 0.0 | 0.0-1.2 | 0.0 | 0.0 | 0.0-1.0 |
| Female breast | | | | | | | 121.8 | 133.6 | 122.9-145.0 | | | | | | | 19.6 | 21.2 | 17.1-26.0 |
| Cervix uteri | | | | | | | 7.0 | 7.9 | 5.4-11.1 | | | | | | | 1.4 | 1.5 | 0.6-3.2 |
| Corpus uterus and NOS | | | | | | | 30.2 | 30.9 | 26.1-36.5 | | | | | | | 9.0 | 9.2 | 6.7-12.5 |
| Ovary | | | | | | | 8.6 | 9.1 | 6.5-12.5 | | | | | | | 6.0 | 6.1 | 4.1-8.9 |
| Prostate | | | | 147.4 | 216.5 | 200.7-233.2 | | | | | | | 14.0 | 26.4 | 20.5-33.4 | | | |
| Testis | | | | 1.0 | 1.4 | 0.4-3.4 | | | | | | | 0.0 | 0.0 | 0.0-1.2 | | | |
| Urinary bladder (incl. in situ) | 20.0 | 12.1 | 9.8-14.8 | 12.8 | 20.6 | 15.7-26.5 | 7.2 | 7.3 | 5.1-10.3 | 4.6 | 3.0 | 1.9-4.5 | 3.0 | 5.5 | 3.1-9.2 | 1.6 | 1.6 | 0.7-3.3 |
| Kidney and renal pelvis | 27.4 | 16.7 | 14.0-19.9 | 16.0 | 22.9 | 18.0-28.7 | 11.4 | 12.3 | 9.3-16.1 | 3.6 | 2.3 | 1.3-3.6 | 2.4 | 3.6 | 1.8-6.4 | 1.2 | 1.3 | 0.5-2.9 |
| Brain and other central nervous system | 7.2 | 5.0 | 3.4-6.9 | 3.4 | 5.5 | 3.2-8.9 | 3.8 | 4.9 | 2.9-7.7 | 3.8 | 2.5 | 1.5-3.9 | 1.6 | 2.6 | 1.1-5.1 | 2.2 | 2.5 | 1.2-4.6 |
| Thyroid | 22.8 | 15.3 | 12.5-18.4 | 4.8 | 7.5 | 4.8-11.4 | 18.0 | 21.9 | 17.4-27.1 | 1.6 | 1.0 | 0.4-1.9 | 0.8 | 1.1 | 0.3-3.1 | 0.8 | 0.7 | 0.2-2.1 |
| Hodgkin lymphoma | 4.2 | 2.9 | 1.8-4.5 | 2.8 | 4.4 | 2.4-7.5 | 1.4 | 1.7 | 0.7-3.5 | 0.2 | 0.1 | 0.0-0.8 | 0.2 | 0.3 | 0.0-1.9 | 0.0 | 0.0 | 0.0-1.0 |
| Non-Hodgkin lymphomas | 29.2 | 18.2 | 15.3-21.4 | 13.4 | 20.2 | 15.6-25.9 | 15.8 | 16.8 | 13.2-21.2 | 6.8 | 4.2 | 2.9-5.9 | 4.0 | 6.2 | 3.8-9.8 | 2.8 | 2.9 | 1.6-5.0 |
| Myeloma | 28.2 | 17.7 | 14.8-20.9 | 13.4 | 21.3 | 16.3-27.2 | 14.8 | 15.5 | 12.1-19.6 | 10.6 | 6.8 | 5.1-9.0 | 4.8 | 7.5 | 4.7-11.3 | 5.8 | 6.0 | 4.0-8.8 |
| Leukemias | 18.6 | 11.8 | 9.5-14.5 | 9.0 | 14.2 | 10.3-19.2 | 9.6 | 10.4 | 7.6-14.0 | 8.8 | 5.3 | 3.8-7.2 | 4.0 | 6.5 | 3.9-10.3 | 4.8 | 4.8 | 3.1-7.4 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution. NOS=Not otherwise specified.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.