

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Colorectal, All Races**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	5680	34.2	33.3-35.1	2764	40.4	38.9-42.1	2916	30.1	29.0-31.2
1977	5930	35.5	34.6-36.5	2871	42.4	40.8-44.0	3059	31.3	30.2-32.4
1978	5948	35.5	34.5-36.4	2914	42.7	41.1-44.4	3034	30.9	29.8-32.0
1979	6315	37.6	36.6-38.5	3005	43.9	42.3-45.6	3310	33.5	32.3-34.6
1980	6434	37.9	37.0-38.9	3122	45.5	43.9-47.2	3312	33.1	31.9-34.2
1981	6574	38.5	37.5-39.4	3259	47.6	45.9-49.3	3315	32.8	31.6-33.9
1982	6673	38.8	37.9-39.8	3263	47.2	45.5-48.9	3410	33.4	32.3-34.6
1983	6592	38.0	37.0-38.9	3254	46.6	45.0-48.3	3338	32.5	31.4-33.6
1984	6491	37.5	36.6-38.4	3146	45.3	43.7-47.0	3345	32.5	31.3-33.6
1985	6540	37.4	36.5-38.3	3240	46.4	44.7-48.0	3300	31.8	30.7-32.9
1986	6389	36.4	35.5-37.3	3102	44.0	42.4-45.6	3287	31.6	30.5-32.7
1987	6145	34.7	33.8-35.6	2986	42.0	40.5-43.6	3159	29.9	28.9-31.0
1988	5974	33.7	32.8-34.5	2870	40.3	38.8-41.9	3104	29.3	28.3-30.4
1989	5664	31.9	31.0-32.7	2793	39.0	37.5-40.5	2871	27.2	26.2-28.2
1990	5532	30.8	30.0-31.7	2732	37.7	36.2-39.2	2800	26.2	25.2-27.2
1991	5605	31.2	30.4-32.0	2768	38.2	36.8-39.7	2837	26.5	25.5-27.5
1992	5258	29.1	28.3-29.9	2551	34.8	33.4-36.2	2707	25.3	24.4-26.3
1993	5250	28.8	28.0-29.6	2481	33.3	31.9-34.6	2769	25.6	24.7-26.6
1994	5707	31.1	30.3-32.0	2751	37.1	35.7-38.5	2956	27.3	26.3-28.3
1995	5602	30.4	29.6-31.2	2724	36.5	35.1-37.9	2878	26.4	25.4-27.4
1996	5693	30.7	29.9-31.5	2779	36.7	35.3-38.1	2914	26.6	25.6-27.6
1997	5982	32.0	31.2-32.9	2892	37.6	36.2-39.0	3090	28.0	27.0-29.0
1998	6129	32.5	31.7-33.3	2933	37.6	36.2-39.0	3196	28.9	27.9-30.0
1999	6131	32.0	31.2-32.8	2888	36.5	35.1-37.9	3243	28.9	27.9-29.9
2000	6127	31.6	30.8-32.4	2977	36.9	35.6-38.3	3150	27.7	26.7-28.7
2001	5874	29.9	29.2-30.7	2849	35.1	33.8-36.4	3025	26.4	25.4-27.3
2002	5998	30.2	29.4-31.0	2853	34.3	33.1-35.6	3145	27.1	26.2-28.1
2003	5939	29.6	28.8-30.3	2888	34.3	33.1-35.6	3051	26.2	25.2-27.1
2004	5606	27.6	26.9-28.4	2655	30.9	29.8-32.1	2951	25.1	24.2-26.1
2005	5419	26.5	25.8-27.2	2632	30.6	29.4-31.8	2787	23.6	22.7-24.5
2006	5193	25.2	24.5-25.9	2501	28.7	27.6-29.8	2692	22.6	21.8-23.5
2007	5216	25.1	24.4-25.8	2552	28.7	27.5-29.8	2664	22.4	21.5-23.3
2008	5143	24.4	23.7-25.1	2506	27.7	26.6-28.8	2637	21.9	21.1-22.8
2009	4970	23.2	22.5-23.9	2463	26.6	25.6-27.7	2507	20.5	19.6-21.3
2010	5058	23.4	22.8-24.1	2493	26.5	25.5-27.6	2565	21.0	20.1-21.8
2011	4947	22.6	21.9-23.2	2483	26.0	25.0-27.1	2464	19.8	19.1-20.7
2012	4815	21.6	21.0-22.2	2430	24.8	23.8-25.8	2385	19.0	18.2-19.8
2013	4956	21.9	21.3-22.5	2499	25.1	24.1-26.1	2457	19.3	18.6-20.1
2014	5036	21.9	21.3-22.6	2553	25.0	24.1-26.1	2483	19.4	18.6-20.2
2015	4857	21.0	20.4-21.6	2460	23.8	22.8-24.8	2397	18.6	17.8-19.4
2016	5242	22.3	21.7-23.0	2732	26.1	25.1-27.2	2510	19.3	18.5-20.1
2017	5211	22.1	21.5-22.7	2726	25.6	24.7-26.6	2485	19.2	18.4-20.0
2018	5152	21.4	20.9-22.1	2626	24.5	23.5-25.4	2526	18.9	18.1-19.7
2019	5215	21.6	21.0-22.2	2771	25.5	24.5-26.5	2444	18.5	17.7-19.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Colorectal, Whites**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	5256	34.7	33.7-35.7	2577	41.3	39.6-43.0	2679	30.3	29.2-31.5
1977	5398	35.6	34.6-36.5	2653	42.8	41.2-44.6	2745	31.0	29.8-32.2
1978	5386	35.3	34.4-36.3	2674	43.1	41.4-44.8	2712	30.4	29.2-31.6
1979	5713	37.5	36.5-38.5	2732	44.0	42.3-45.7	2981	33.3	32.1-34.6
1980	5842	38.2	37.2-39.2	2860	46.1	44.4-47.9	2982	33.0	31.9-34.3
1981	5982	39.0	38.0-40.0	2984	48.3	46.5-50.1	2998	33.1	31.9-34.3
1982	6052	39.3	38.3-40.3	2970	47.9	46.1-49.7	3082	33.7	32.5-34.9
1983	5944	38.4	37.4-39.4	2973	47.6	45.9-49.5	2971	32.4	31.3-33.6
1984	5892	38.3	37.3-39.3	2872	46.4	44.6-48.2	3020	33.0	31.8-34.2
1985	5893	38.1	37.1-39.1	2963	47.6	45.8-49.4	2930	31.9	30.8-33.1
1986	5707	36.9	35.9-37.9	2789	44.7	43.0-46.4	2918	31.8	30.6-33.0
1987	5531	35.6	34.6-36.5	2713	43.1	41.5-44.9	2818	30.5	29.4-31.7
1988	5329	34.3	33.4-35.2	2583	41.2	39.6-42.9	2746	29.7	28.6-30.9
1989	5006	32.3	31.4-33.2	2474	39.5	38.0-41.2	2532	27.4	26.4-28.5
1990	4892	31.4	30.5-32.3	2428	38.2	36.7-39.8	2464	26.6	25.5-27.7
1991	4934	31.7	30.8-32.6	2480	39.3	37.7-40.9	2454	26.5	25.5-27.6
1992	4600	29.5	28.7-30.4	2271	35.7	34.2-37.2	2329	25.3	24.2-26.3
1993	4577	29.3	28.4-30.1	2177	33.8	32.3-35.2	2400	26.0	25.0-27.1
1994	4931	31.5	30.6-32.3	2380	37.2	35.7-38.8	2551	27.6	26.6-28.8
1995	4784	30.5	29.7-31.4	2353	36.7	35.2-38.3	2431	26.3	25.2-27.4
1996	4823	30.7	29.8-31.6	2369	36.7	35.2-38.2	2454	26.5	25.4-27.6
1997	5077	32.2	31.3-33.1	2456	37.6	36.1-39.1	2621	28.1	27.0-29.2
1998	5193	32.8	31.9-33.7	2526	38.2	36.7-39.8	2667	28.8	27.7-30.0
1999	5129	32.0	31.1-32.9	2469	37.1	35.6-38.6	2660	28.4	27.3-29.5
2000	5198	32.2	31.4-33.1	2561	37.9	36.5-39.4	2637	27.9	26.8-29.0
2001	4892	30.1	29.3-31.0	2386	35.2	33.8-36.6	2506	26.5	25.4-27.6
2002	4989	30.5	29.6-31.4	2407	34.9	33.5-36.3	2582	27.0	26.0-28.1
2003	4927	29.9	29.1-30.8	2419	34.8	33.5-36.3	2508	26.2	25.2-27.3
2004	4561	27.6	26.8-28.4	2171	30.8	29.6-32.2	2390	25.0	24.0-26.1
2005	4341	26.1	25.3-26.9	2117	30.0	28.8-31.4	2224	23.3	22.3-24.3
2006	4156	25.0	24.2-25.8	2041	28.8	27.5-30.1	2115	22.1	21.1-23.1
2007	4220	25.2	24.5-26.0	2074	28.8	27.5-30.1	2146	22.6	21.6-23.6
2008	4091	24.3	23.5-25.1	2018	27.7	26.5-29.0	2073	21.6	20.7-22.6
2009	3960	23.2	22.5-24.0	1970	26.6	25.4-27.8	1990	20.5	19.6-21.4
2010	4028	23.5	22.8-24.3	1973	26.3	25.2-27.5	2055	21.2	20.3-22.2
2011	3912	22.6	21.9-23.4	1965	25.9	24.8-27.1	1947	20.0	19.1-21.0
2012	3792	21.8	21.1-22.5	1906	24.7	23.6-25.8	1886	19.4	18.5-20.3
2013	3908	22.3	21.5-23.0	1984	25.4	24.3-26.6	1924	19.7	18.8-20.6
2014	3843	21.6	20.9-22.3	1981	24.9	23.7-26.0	1862	18.9	18.0-19.8
2015	3750	21.1	20.4-21.8	1884	23.6	22.5-24.7	1866	19.0	18.1-19.9
2016	3994	22.3	21.6-23.0	2066	25.6	24.5-26.7	1928	19.5	18.6-20.5
2017	3960	22.0	21.3-22.8	2054	25.0	23.9-26.2	1906	19.5	18.6-20.5
2018	3894	21.5	20.8-22.2	1999	24.3	23.2-25.4	1895	19.0	18.1-19.9
2019	3894	21.6	20.9-22.3	2101	25.5	24.4-26.7	1793	18.2	17.3-19.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Colorectal, Blacks**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	355	26.2	23.3-29.4	152	27.4	22.6-32.7	203	25.3	21.6-29.3
1977	472	35.1	31.6-38.7	184	35.9	30.2-42.3	288	34.6	30.4-39.2
1978	496	36.4	32.9-40.0	198	36.2	30.6-42.4	298	36.4	32.1-41.1
1979	531	36.8	33.4-40.3	228	40.0	34.3-46.3	303	34.7	30.7-39.1
1980	534	35.0	31.9-38.4	227	38.2	32.8-44.3	307	33.3	29.4-37.4
1981	525	34.1	31.0-37.4	236	40.9	35.2-47.2	289	30.4	26.9-34.3
1982	547	33.2	30.3-36.3	253	40.2	34.9-46.1	294	29.1	25.8-32.8
1983	567	33.4	30.5-36.4	238	37.7	32.4-43.4	329	31.4	28.0-35.1
1984	532	31.3	28.5-34.2	236	35.5	30.7-40.8	296	28.6	25.4-32.2
1985	575	32.6	29.9-35.6	247	39.0	33.7-44.8	328	29.7	26.4-33.2
1986	614	33.9	31.1-36.9	269	38.4	33.5-43.8	345	31.2	27.9-34.8
1987	541	28.7	26.2-31.3	234	33.8	29.2-39.0	307	26.0	23.1-29.2
1988	573	30.2	27.6-32.9	244	33.8	29.3-38.8	329	28.1	25.0-31.3
1989	569	28.7	26.3-31.3	272	37.4	32.6-42.6	297	24.2	21.5-27.2
1990	557	28.2	25.8-30.7	249	34.0	29.5-38.9	308	24.9	22.1-27.9
1991	568	28.3	26.0-30.9	244	31.4	27.3-36.0	324	26.3	23.4-29.3
1992	559	26.7	24.5-29.1	226	28.8	24.8-33.2	333	25.8	23.1-28.8
1993	556	26.5	24.3-28.9	246	31.2	27.0-35.8	310	24.0	21.3-26.8
1994	609	28.6	26.3-31.0	275	34.4	30.1-39.1	334	25.3	22.6-28.2
1995	643	28.9	26.7-31.3	291	35.4	31.1-40.2	352	25.5	22.9-28.4
1996	722	32.2	29.8-34.7	325	38.1	33.7-42.9	397	28.8	26.0-31.8
1997	748	32.4	30.1-34.9	354	40.9	36.4-45.8	394	27.7	25.0-30.6
1998	775	32.9	30.6-35.4	321	35.4	31.3-39.8	454	31.4	28.6-34.5
1999	825	33.5	31.2-35.9	336	35.2	31.2-39.4	489	32.5	29.6-35.5
2000	718	28.4	26.3-30.6	314	32.9	29.1-37.0	404	25.9	23.5-28.6
2001	777	30.2	28.1-32.4	348	36.5	32.5-40.8	429	27.0	24.5-29.7
2002	797	30.3	28.2-32.5	331	33.0	29.3-37.0	466	28.9	26.3-31.7
2003	788	28.9	26.9-31.1	347	33.4	29.7-37.4	441	26.6	24.2-29.3
2004	809	29.2	27.2-31.3	364	33.9	30.3-37.8	445	26.4	24.0-29.0
2005	820	29.0	27.0-31.1	374	34.0	30.4-37.9	446	26.1	23.7-28.7
2006	794	27.6	25.7-29.7	343	30.1	26.8-33.7	451	26.0	23.6-28.5
2007	758	25.9	24.1-27.9	366	31.1	27.8-34.7	392	22.4	20.2-24.8
2008	810	27.1	25.2-29.1	374	31.4	28.1-35.0	436	24.3	22.0-26.7
2009	754	24.7	22.9-26.6	362	29.2	26.1-32.6	392	21.5	19.4-23.8
2010	756	24.1	22.4-26.0	361	29.3	26.2-32.7	395	21.1	19.1-23.3
2011	762	23.7	22.0-25.4	370	28.5	25.5-31.7	392	20.4	18.4-22.5
2012	749	22.6	20.9-24.3	377	28.2	25.3-31.3	372	18.9	17.0-20.9
2013	751	22.1	20.5-23.7	357	25.5	22.8-28.4	394	19.5	17.6-21.6
2014	844	24.4	22.8-26.2	376	26.4	23.7-29.3	468	23.0	21.0-25.3
2015	778	21.6	20.1-23.2	397	26.9	24.2-29.8	381	18.0	16.2-19.9
2016	896	24.5	22.9-26.2	473	31.9	29.0-35.0	423	19.6	17.8-21.6
2017	827	22.5	21.0-24.1	439	29.4	26.6-32.4	388	18.0	16.2-20.0
2018	831	21.8	20.4-23.4	403	25.8	23.2-28.5	428	19.1	17.3-21.0
2019	874	22.7	21.2-24.3	434	27.7	25.0-30.5	440	19.5	17.7-21.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Colorectal, All Races**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	2453	33.2	31.8-34.6	1175	39.4	37.0-41.8	1278	29.3	27.6-31.0
1977	2542	34.8	33.5-36.3	1195	41.3	38.9-43.8	1347	30.9	29.2-32.6
1978	2565	35.1	33.7-36.5	1242	42.8	40.3-45.3	1323	30.5	28.8-32.2
1979	2716	37.5	36.1-39.0	1289	45.0	42.5-47.7	1427	32.9	31.2-34.7
1980	2785	38.5	37.1-40.0	1339	47.0	44.5-49.7	1446	33.4	31.6-35.2
1981	2705	37.5	36.1-39.0	1296	45.8	43.3-48.5	1409	32.3	30.6-34.1
1982	2754	38.2	36.7-39.6	1328	47.4	44.8-50.1	1426	32.6	30.9-34.4
1983	2698	37.1	35.7-38.5	1297	45.8	43.3-48.5	1401	32.0	30.3-33.8
1984	2580	35.7	34.3-37.2	1233	44.1	41.6-46.7	1347	30.8	29.1-32.5
1985	2597	35.7	34.3-37.1	1294	45.9	43.4-48.6	1303	29.7	28.1-31.4
1986	2652	36.5	35.1-37.9	1313	46.3	43.7-48.9	1339	30.5	28.8-32.2
1987	2390	32.7	31.4-34.1	1127	39.8	37.4-42.3	1263	28.4	26.8-30.0
1988	2366	32.7	31.4-34.1	1105	39.5	37.2-42.0	1261	28.5	27.0-30.2
1989	2199	30.5	29.3-31.8	1053	37.3	35.1-39.7	1146	26.3	24.8-27.9
1990	2152	29.9	28.6-31.2	1024	36.3	34.0-38.6	1128	25.7	24.2-27.2
1991	2197	30.8	29.6-32.2	1070	38.8	36.5-41.3	1127	25.9	24.4-27.5
1992	2022	28.5	27.2-29.8	902	32.5	30.4-34.7	1120	26.1	24.6-27.7
1993	1973	27.7	26.5-28.9	905	32.0	29.9-34.2	1068	24.8	23.3-26.3
1994	2272	31.8	30.5-33.1	1049	37.3	35.1-39.7	1223	28.3	26.7-30.0
1995	2267	31.7	30.4-33.0	1061	37.8	35.5-40.2	1206	27.8	26.3-29.5
1996	2299	32.0	30.7-33.4	1097	38.3	36.0-40.7	1202	27.8	26.2-29.4
1997	2415	33.5	32.1-34.8	1105	38.5	36.2-40.9	1310	30.0	28.4-31.7
1998	2498	34.2	32.9-35.6	1134	39.0	36.7-41.3	1364	31.2	29.6-32.9
1999	2451	32.9	31.6-34.3	1112	37.3	35.1-39.5	1339	30.2	28.6-31.8
2000	2449	32.4	31.1-33.7	1174	38.4	36.2-40.7	1275	28.2	26.7-29.8
2001	2364	30.9	29.7-32.2	1122	36.7	34.5-38.9	1242	27.2	25.7-28.8
2002	2397	30.9	29.7-32.2	1114	35.3	33.2-37.4	1283	27.8	26.3-29.4
2003	2380	30.5	29.3-31.7	1142	36.2	34.1-38.4	1238	26.7	25.2-28.2
2004	2232	28.3	27.2-29.5	1041	32.4	30.4-34.5	1191	25.5	24.1-27.0
2005	2227	28.1	26.9-29.3	1056	32.8	30.8-34.9	1171	25.0	23.6-26.5
2006	2113	26.6	25.4-27.7	985	30.2	28.3-32.2	1128	24.1	22.7-25.5
2007	2053	25.5	24.4-26.6	999	29.9	28.0-31.8	1054	22.3	21.0-23.7
2008	2016	24.9	23.8-26.0	972	28.6	26.8-30.5	1044	22.1	20.8-23.5
2009	2003	24.3	23.3-25.4	983	28.5	26.7-30.4	1020	21.2	19.9-22.6
2010	1950	23.4	22.3-24.5	972	27.6	25.8-29.4	978	20.3	19.0-21.6
2011	1877	22.0	21.1-23.1	945	26.3	24.6-28.1	932	18.9	17.7-20.1
2012	1848	21.2	20.2-22.2	937	25.2	23.6-26.9	911	18.1	17.0-19.4
2013	1976	22.4	21.4-23.4	969	25.7	24.1-27.4	1007	19.8	18.5-21.1
2014	1977	21.9	20.9-22.9	1002	25.5	23.9-27.2	975	19.1	17.9-20.4
2015	1872	20.4	19.5-21.4	952	24.0	22.5-25.6	920	17.6	16.5-18.8
2016	2054	22.1	21.1-23.1	1094	27.2	25.6-28.9	960	18.2	17.1-19.4
2017	2047	21.7	20.7-22.6	1060	25.5	24.0-27.2	987	18.7	17.5-19.9
2018	1987	20.6	19.7-21.6	1019	24.4	22.9-26.0	968	17.7	16.5-18.8
2019	2098	21.8	20.8-22.7	1111	26.1	24.6-27.8	987	18.3	17.1-19.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Colorectal, Whites**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	2141	34.2	32.7-35.7	1030	40.5	37.9-43.1	1111	30.1	28.3-32.0
1977	2138	34.8	33.3-36.4	1031	42.0	39.3-44.7	1107	30.4	28.5-32.3
1978	2147	35.1	33.6-36.6	1071	43.9	41.2-46.8	1076	29.6	27.8-31.5
1979	2261	37.6	36.1-39.3	1089	45.6	42.9-48.5	1172	32.7	30.8-34.7
1980	2342	39.5	37.9-41.2	1147	49.0	46.1-52.0	1195	33.7	31.8-35.8
1981	2261	38.6	37.0-40.2	1099	47.6	44.7-50.6	1162	32.8	30.8-34.8
1982	2290	39.4	37.7-41.1	1105	48.7	45.8-51.8	1185	33.7	31.8-35.8
1983	2203	37.7	36.1-39.4	1083	47.5	44.6-50.5	1120	31.9	30.0-33.9
1984	2122	37.1	35.6-38.8	1024	46.1	43.2-49.1	1098	31.9	30.0-33.9
1985	2092	36.6	35.0-38.2	1076	48.1	45.2-51.2	1016	29.5	27.7-31.4
1986	2112	37.4	35.8-39.1	1068	48.2	45.3-51.3	1044	30.5	28.7-32.5
1987	1911	34.0	32.4-35.5	919	41.7	39.0-44.5	992	29.0	27.2-31.0
1988	1888	34.3	32.7-35.9	897	41.8	39.1-44.7	991	29.6	27.7-31.6
1989	1720	31.8	30.3-33.4	822	38.8	36.1-41.6	898	27.2	25.4-29.2
1990	1659	30.8	29.3-32.3	795	37.3	34.8-40.1	864	26.2	24.5-28.1
1991	1677	31.7	30.2-33.2	849	41.2	38.4-44.1	828	25.6	23.8-27.5
1992	1539	29.7	28.2-31.2	710	34.7	32.1-37.3	829	26.5	24.7-28.4
1993	1475	28.8	27.3-30.3	684	33.1	30.6-35.7	791	25.7	23.9-27.6
1994	1716	33.5	31.9-35.1	795	39.0	36.3-41.8	921	29.9	28.0-32.0
1995	1655	32.8	31.2-34.4	782	39.0	36.3-41.9	873	28.6	26.7-30.7
1996	1648	32.9	31.3-34.6	797	39.5	36.8-42.3	851	28.3	26.4-30.3
1997	1714	34.2	32.6-35.9	773	38.5	35.8-41.3	941	31.1	29.1-33.2
1998	1790	35.8	34.2-37.5	833	41.2	38.5-44.1	957	32.3	30.3-34.5
1999	1695	33.7	32.1-35.3	808	39.5	36.8-42.3	887	29.8	27.8-31.9
2000	1749	34.6	33.0-36.3	867	41.9	39.2-44.8	882	29.1	27.2-31.2
2001	1630	32.1	30.5-33.7	794	38.4	35.8-41.2	836	27.8	25.9-29.8
2002	1660	32.6	31.0-34.2	787	37.5	34.9-40.2	873	28.7	26.8-30.7
2003	1633	32.1	30.6-33.7	808	38.8	36.2-41.6	825	27.3	25.5-29.3
2004	1461	28.7	27.3-30.2	713	34.1	31.6-36.7	748	24.9	23.1-26.8
2005	1435	28.3	26.9-29.9	691	33.1	30.7-35.7	744	25.1	23.3-27.0
2006	1340	26.8	25.4-28.3	658	31.6	29.2-34.1	682	23.3	21.5-25.2
2007	1312	26.2	24.8-27.7	645	30.6	28.3-33.1	667	22.9	21.2-24.8
2008	1252	25.2	23.8-26.7	626	29.6	27.3-32.1	626	21.9	20.2-23.7
2009	1264	25.2	23.8-26.7	628	29.5	27.2-31.9	636	21.9	20.2-23.8
2010	1201	24.0	22.6-25.4	604	28.2	25.9-30.5	597	20.7	19.0-22.5
2011	1124	22.1	20.8-23.5	575	26.4	24.3-28.7	549	18.8	17.2-20.5
2012	1088	21.2	20.0-22.6	545	24.4	22.4-26.6	543	18.6	17.0-20.2
2013	1202	23.3	22.0-24.7	598	26.8	24.7-29.1	604	20.5	18.9-22.3
2014	1128	21.6	20.3-22.9	601	26.0	24.0-28.2	527	18.1	16.5-19.7
2015	1076	20.5	19.3-21.8	537	23.4	21.4-25.5	539	18.2	16.7-19.9
2016	1126	21.3	20.1-22.7	602	25.9	23.8-28.1	524	17.9	16.4-19.6
2017	1155	21.7	20.4-23.0	587	24.4	22.5-26.5	568	19.4	17.8-21.1
2018	1102	20.4	19.2-21.7	576	24.0	22.1-26.1	526	17.5	16.0-19.2
2019	1153	21.7	20.5-23.1	636	26.6	24.5-28.8	517	17.6	16.0-19.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Colorectal, Blacks**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	277	26.4	23.0-30.0	125	30.7	24.9-37.3	152	23.7	19.8-28.1
1977	371	36.2	32.2-40.5	147	38.4	31.6-46.2	224	34.9	30.1-40.2
1978	380	36.9	32.9-41.2	147	36.7	30.2-44.2	233	37.0	32.1-42.4
1979	405	36.5	32.7-40.5	165	39.2	32.7-46.5	240	35.0	30.4-40.0
1980	405	34.2	30.7-37.9	168	35.9	30.1-42.5	237	33.0	28.7-37.8
1981	391	32.5	29.1-36.1	169	38.0	31.7-45.0	222	29.7	25.7-34.0
1982	421	32.8	29.5-36.2	198	41.1	35.0-47.9	223	27.9	24.2-32.0
1983	447	33.6	30.4-37.1	189	39.6	33.5-46.5	258	30.9	27.1-35.1
1984	407	30.9	27.8-34.2	180	35.3	29.9-41.4	227	28.1	24.4-32.1
1985	452	32.8	29.6-36.1	195	40.0	33.9-46.7	257	29.4	25.8-33.3
1986	484	34.1	31.0-37.4	208	38.6	33.0-44.8	276	31.5	27.8-35.6
1987	415	28.4	25.6-31.4	173	33.3	28.1-39.2	242	26.0	22.7-29.5
1988	426	29.2	26.3-32.2	179	33.3	28.1-39.1	247	26.9	23.5-30.5
1989	413	27.0	24.4-29.9	194	35.1	29.9-40.9	219	22.9	19.9-26.2
1990	430	28.3	25.6-31.2	189	34.6	29.4-40.4	241	25.0	21.8-28.4
1991	437	28.6	25.9-31.5	186	32.6	27.6-38.0	251	26.2	23.0-29.7
1992	412	26.0	23.4-28.7	155	27.0	22.5-32.1	257	25.7	22.6-29.1
1993	417	26.5	23.9-29.2	182	32.5	27.4-38.0	235	23.5	20.6-26.8
1994	446	27.5	24.9-30.2	191	32.5	27.7-37.8	255	24.9	21.9-28.2
1995	482	28.6	26.1-31.4	217	35.9	30.9-41.4	265	24.9	22.0-28.2
1996	536	31.7	29.0-34.6	231	37.1	32.1-42.6	305	28.8	25.6-32.2
1997	572	33.3	30.5-36.2	261	42.0	36.6-47.9	311	28.8	25.6-32.2
1998	581	33.0	30.4-35.9	233	35.5	30.8-40.6	348	31.6	28.3-35.1
1999	614	33.2	30.6-36.0	238	34.6	30.1-39.5	376	32.5	29.3-35.9
2000	523	27.5	25.2-30.0	221	31.9	27.6-36.7	302	25.3	22.5-28.3
2001	574	29.8	27.4-32.4	241	34.4	30.0-39.4	333	27.6	24.7-30.7
2002	577	29.3	26.9-31.8	242	32.8	28.6-37.5	335	27.4	24.5-30.5
2003	574	28.4	26.1-30.8	237	31.5	27.4-36.0	337	26.8	24.0-29.9
2004	578	28.2	25.9-30.6	234	30.5	26.5-34.9	344	26.9	24.1-29.9
2005	593	28.4	26.1-30.8	257	32.8	28.7-37.3	336	26.1	23.3-29.0
2006	574	27.3	25.0-29.6	231	29.2	25.4-33.5	343	26.2	23.5-29.1
2007	547	25.6	23.5-27.9	262	31.7	27.8-36.0	285	21.8	19.3-24.5
2008	580	26.6	24.4-28.9	260	30.4	26.6-34.5	320	24.0	21.4-26.8
2009	539	24.5	22.4-26.7	253	29.2	25.6-33.2	286	21.3	18.9-24.0
2010	545	24.1	22.1-26.2	249	28.5	24.9-32.5	296	21.6	19.2-24.3
2011	541	23.1	21.2-25.2	253	27.6	24.1-31.4	288	20.2	17.9-22.7
2012	543	22.8	20.9-24.8	271	29.3	25.8-33.2	272	18.7	16.5-21.1
2013	548	22.3	20.5-24.3	253	25.7	22.5-29.2	295	20.0	17.7-22.4
2014	598	24.1	22.1-26.1	264	26.5	23.2-30.0	334	22.6	20.2-25.2
2015	541	21.2	19.4-23.1	275	27.1	23.9-30.7	266	17.3	15.2-19.5
2016	654	25.0	23.1-27.1	336	32.3	28.9-36.1	318	20.2	18.0-22.7
2017	567	21.6	19.9-23.5	289	27.9	24.7-31.4	278	17.6	15.6-19.9
2018	567	21.0	19.2-22.8	272	24.9	21.9-28.1	295	18.3	16.2-20.5
2019	604	22.2	20.5-24.1	295	27.3	24.2-30.7	309	18.9	16.8-21.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Colorectal, All Races**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	3227	34.9	33.7-36.2	1589	41.2	39.1-43.4	1638	30.7	29.2-32.3
1977	3388	36.1	34.9-37.4	1676	43.2	41.1-45.5	1712	31.5	30.1-33.1
1978	3383	35.8	34.5-37.0	1672	42.7	40.5-44.9	1711	31.2	29.7-32.7
1979	3598	37.6	36.4-38.9	1716	43.1	41.0-45.3	1882	33.8	32.3-35.4
1980	3649	37.5	36.3-38.7	1783	44.3	42.2-46.5	1866	32.9	31.4-34.4
1981	3868	39.2	38.0-40.5	1962	48.9	46.6-51.2	1906	33.1	31.6-34.6
1982	3919	39.4	38.1-40.6	1935	47.0	44.8-49.2	1984	34.0	32.6-35.6
1983	3892	38.6	37.4-39.9	1955	47.1	44.9-49.3	1937	32.9	31.4-34.4
1984	3910	38.7	37.5-40.0	1913	46.1	43.9-48.3	1997	33.7	32.2-35.2
1985	3936	38.6	37.4-39.8	1946	46.6	44.4-48.8	1990	33.3	31.8-34.8
1986	3737	36.4	35.3-37.6	1789	42.4	40.4-44.5	1948	32.4	31.0-33.9
1987	3752	36.0	34.9-37.2	1858	43.4	41.4-45.6	1894	31.1	29.7-32.5
1988	3608	34.3	33.2-35.4	1765	40.7	38.8-42.8	1843	29.9	28.6-31.3
1989	3465	32.8	31.7-33.9	1740	40.2	38.2-42.2	1725	27.8	26.5-29.2
1990	3379	31.4	30.4-32.5	1707	38.6	36.7-40.5	1672	26.6	25.3-27.9
1991	3408	31.4	30.3-32.5	1698	37.7	35.9-39.6	1710	26.9	25.6-28.2
1992	3236	29.5	28.5-30.5	1649	36.2	34.4-38.1	1587	24.8	23.6-26.0
1993	3276	29.5	28.5-30.5	1576	34.1	32.4-35.8	1700	26.2	25.0-27.5
1994	3435	30.8	29.7-31.8	1702	37.1	35.3-38.9	1733	26.6	25.3-27.9
1995	3335	29.6	28.6-30.6	1663	35.7	33.9-37.5	1672	25.5	24.2-26.7
1996	3394	29.9	28.9-31.0	1682	35.8	34.1-37.6	1712	25.8	24.6-27.1
1997	3567	31.1	30.1-32.2	1787	37.0	35.3-38.8	1780	26.7	25.4-28.0
1998	3631	31.4	30.4-32.4	1799	36.7	35.0-38.5	1832	27.4	26.1-28.7
1999	3680	31.4	30.4-32.4	1776	36.1	34.4-37.8	1904	28.0	26.7-29.3
2000	3678	31.0	30.0-32.0	1803	36.0	34.3-37.7	1875	27.3	26.1-28.6
2001	3509	29.3	28.3-30.3	1726	34.1	32.5-35.8	1783	25.8	24.6-27.1
2002	3601	29.7	28.7-30.7	1739	33.7	32.1-35.3	1862	26.6	25.4-27.9
2003	3559	29.0	28.0-30.0	1746	33.2	31.7-34.8	1813	25.8	24.6-27.0
2004	3374	27.2	26.3-28.1	1614	30.1	28.6-31.6	1760	24.8	23.7-26.0
2005	3192	25.4	24.5-26.3	1576	29.2	27.8-30.7	1616	22.6	21.5-23.8
2006	3080	24.4	23.5-25.2	1516	27.8	26.4-29.2	1564	21.6	20.6-22.8
2007	3163	24.8	23.9-25.7	1553	27.9	26.5-29.4	1610	22.4	21.3-23.5
2008	3127	24.1	23.2-24.9	1534	27.2	25.8-28.6	1593	21.7	20.7-22.9
2009	2967	22.5	21.7-23.4	1480	25.6	24.3-27.0	1487	20.0	18.9-21.0
2010	3108	23.4	22.6-24.2	1521	25.9	24.6-27.3	1587	21.3	20.3-22.4
2011	3070	22.9	22.1-23.7	1538	25.8	24.5-27.2	1532	20.5	19.5-21.6
2012	2967	21.8	21.0-22.6	1493	24.5	23.2-25.8	1474	19.5	18.5-20.6
2013	2980	21.6	20.8-22.4	1530	24.7	23.4-26.0	1450	19.1	18.1-20.1
2014	3059	22.0	21.2-22.8	1551	24.8	23.6-26.1	1508	19.6	18.6-20.6
2015	2985	21.3	20.6-22.1	1508	23.7	22.5-24.9	1477	19.3	18.3-20.3
2016	3188	22.5	21.7-23.4	1638	25.5	24.3-26.8	1550	19.9	18.9-21.0
2017	3164	22.4	21.6-23.2	1666	25.7	24.5-27.0	1498	19.5	18.5-20.6
2018	3165	22.1	21.3-22.9	1607	24.6	23.3-25.8	1558	19.8	18.8-20.9
2019	3117	21.6	20.8-22.4	1660	25.1	23.9-26.4	1457	18.6	17.6-19.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Colorectal, Whites**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	3115	35.0	33.7-36.3	1547	41.7	39.5-44.0	1568	30.5	29.0-32.0
1977	3260	36.1	34.8-37.3	1622	43.5	41.3-45.8	1638	31.3	29.8-32.9
1978	3239	35.6	34.3-36.8	1603	42.5	40.4-44.8	1636	30.9	29.4-32.5
1979	3451	37.5	36.2-38.8	1643	43.0	40.8-45.2	1808	33.7	32.2-35.4
1980	3500	37.4	36.1-38.7	1713	44.2	42.1-46.5	1787	32.7	31.2-34.3
1981	3720	39.2	38.0-40.5	1884	48.8	46.5-51.2	1836	33.2	31.7-34.7
1982	3762	39.4	38.1-40.7	1865	47.2	45.0-49.5	1897	33.8	32.3-35.4
1983	3739	38.8	37.5-40.0	1888	47.5	45.3-49.8	1851	32.8	31.3-34.3
1984	3769	39.0	37.7-40.2	1848	46.5	44.3-48.7	1921	33.7	32.2-35.3
1985	3797	38.9	37.7-40.2	1887	47.1	44.9-49.4	1910	33.4	31.9-35.0
1986	3595	36.7	35.5-37.9	1721	42.7	40.6-44.8	1874	32.6	31.2-34.2
1987	3617	36.4	35.2-37.7	1793	43.9	41.8-46.1	1824	31.4	30.0-32.9
1988	3441	34.3	33.1-35.5	1686	40.9	38.9-43.0	1755	29.8	28.4-31.3
1989	3286	32.7	31.5-33.8	1652	40.1	38.1-42.1	1634	27.6	26.2-29.0
1990	3232	31.6	30.5-32.8	1632	38.8	36.8-40.8	1600	26.7	25.4-28.1
1991	3257	31.6	30.5-32.7	1631	38.2	36.3-40.1	1626	26.9	25.6-28.3
1992	3061	29.4	28.4-30.5	1561	36.2	34.4-38.1	1500	24.7	23.4-26.0
1993	3101	29.5	28.5-30.6	1493	34.2	32.4-36.0	1608	26.2	24.9-27.5
1994	3215	30.5	29.4-31.6	1585	36.5	34.7-38.4	1630	26.5	25.2-27.9
1995	3129	29.5	28.4-30.5	1571	35.7	33.9-37.5	1558	25.1	23.9-26.4
1996	3175	29.7	28.7-30.8	1572	35.6	33.8-37.4	1603	25.7	24.4-27.0
1997	3363	31.3	30.2-32.3	1683	37.2	35.4-39.0	1680	26.7	25.4-28.1
1998	3403	31.4	30.3-32.4	1693	36.9	35.1-38.7	1710	27.2	25.9-28.5
1999	3434	31.3	30.2-32.4	1661	36.0	34.3-37.8	1773	27.8	26.5-29.1
2000	3449	31.2	30.1-32.2	1694	36.2	34.5-38.0	1755	27.4	26.1-28.7
2001	3262	29.3	28.3-30.3	1592	33.8	32.1-35.5	1670	25.9	24.7-27.2
2002	3329	29.5	28.5-30.6	1620	33.8	32.1-35.5	1709	26.3	25.0-27.6
2003	3294	28.9	28.0-30.0	1611	33.1	31.5-34.8	1683	25.7	24.5-27.0
2004	3100	27.1	26.1-28.0	1458	29.5	27.9-31.0	1642	25.1	23.9-26.3
2005	2906	25.1	24.2-26.0	1426	28.7	27.2-30.3	1480	22.4	21.3-23.6
2006	2816	24.2	23.3-25.1	1383	27.6	26.2-29.1	1433	21.5	20.4-22.7
2007	2908	24.8	23.9-25.7	1429	28.0	26.6-29.6	1479	22.3	21.2-23.5
2008	2839	23.9	23.0-24.8	1392	27.0	25.6-28.5	1447	21.5	20.4-22.7
2009	2696	22.5	21.6-23.3	1342	25.6	24.2-27.0	1354	19.9	18.8-21.0
2010	2827	23.4	22.5-24.3	1369	25.7	24.3-27.1	1458	21.5	20.4-22.7
2011	2788	22.9	22.0-23.8	1390	25.7	24.4-27.2	1398	20.6	19.5-21.8
2012	2704	22.1	21.2-22.9	1361	24.8	23.4-26.2	1343	19.8	18.7-20.9
2013	2706	21.9	21.0-22.7	1386	24.8	23.5-26.2	1320	19.4	18.3-20.5
2014	2715	21.7	20.9-22.5	1380	24.5	23.2-25.8	1335	19.3	18.2-20.4
2015	2674	21.4	20.6-22.3	1347	23.8	22.5-25.1	1327	19.3	18.3-20.5
2016	2868	22.7	21.8-23.6	1464	25.5	24.2-26.9	1404	20.2	19.1-21.3
2017	2805	22.3	21.4-23.2	1467	25.4	24.1-26.8	1338	19.6	18.5-20.8
2018	2792	22.0	21.1-22.8	1423	24.5	23.2-25.9	1369	19.7	18.6-20.8
2019	2741	21.5	20.7-22.4	1465	25.1	23.8-26.5	1276	18.4	17.3-19.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Colorectal, Blacks**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	78	25.6	19.6-32.6	27	16.6	10.4-25.1	51	31.2	22.6-41.6
1977	101	31.2	25.0-38.5	37	28.2	19.0-39.9	64	33.9	25.7-43.7
1978	116	34.7	28.2-42.1	51	34.4	24.9-46.1	65	34.4	26.1-44.3
1979	126	37.6	30.8-45.4	63	42.4	31.4-55.7	63	33.8	25.5-43.7
1980	129	38.0	31.2-45.8	59	45.5	33.1-60.6	70	34.2	26.3-43.5
1981	134	40.0	33.1-47.8	67	50.2	37.8-65.0	67	33.5	25.7-42.9
1982	126	34.7	28.5-41.8	55	37.6	27.1-50.4	71	33.5	25.9-42.5
1983	120	32.7	26.8-39.4	49	31.5	22.3-42.7	71	33.4	25.9-42.3
1984	125	32.4	26.6-38.9	56	36.1	26.1-48.2	69	30.7	23.7-38.9
1985	123	32.3	26.5-38.9	52	35.9	25.7-48.3	71	30.7	23.8-39.0
1986	130	33.3	27.5-39.9	61	37.7	27.9-49.6	69	30.1	23.1-38.3
1987	126	29.7	24.5-35.6	61	35.2	25.9-46.5	65	26.3	20.2-33.7
1988	147	33.5	28.0-39.6	65	35.3	26.4-46.0	82	32.2	25.5-40.1
1989	156	34.6	29.1-40.8	78	44.4	33.9-56.8	78	29.4	23.1-36.8
1990	127	27.6	22.8-33.0	60	32.1	23.7-42.3	67	24.7	19.0-31.5
1991	131	27.4	22.7-32.7	58	27.9	20.6-36.8	73	26.5	20.7-33.4
1992	147	29.1	24.4-34.4	71	33.9	25.7-43.8	76	26.2	20.5-32.8
1993	139	26.6	22.1-31.6	64	27.1	20.3-35.3	75	25.4	19.9-31.9
1994	163	31.9	27.0-37.4	84	39.8	30.8-50.5	79	26.7	21.1-33.4
1995	161	29.7	25.1-34.9	74	34.2	26.0-43.9	87	27.3	21.8-33.8
1996	186	33.7	28.8-39.0	94	40.7	32.2-50.7	92	28.8	23.1-35.4
1997	176	29.5	25.1-34.3	93	37.5	29.6-46.7	83	24.2	19.2-30.0
1998	194	32.2	27.7-37.3	88	35.0	27.3-44.1	106	30.7	25.1-37.2
1999	211	34.3	29.7-39.5	98	36.5	28.9-45.3	113	32.6	26.8-39.2
2000	195	31.0	26.7-35.8	93	35.4	27.9-44.1	102	28.2	22.9-34.3
2001	203	31.0	26.7-35.7	107	42.3	34.0-51.8	96	24.6	19.8-30.1
2002	220	33.4	29.0-38.2	89	33.9	26.6-42.4	131	33.6	28.1-40.0
2003	214	30.5	26.4-35.0	110	38.6	31.0-47.3	104	26.1	21.2-31.6
2004	231	32.3	28.1-36.9	130	42.9	35.1-51.7	101	25.1	20.3-30.5
2005	227	30.7	26.7-35.1	117	37.1	30.1-45.1	110	26.2	21.5-31.7
2006	220	28.5	24.8-32.7	112	31.8	25.7-38.8	108	25.3	20.7-30.5
2007	211	26.5	22.9-30.4	104	29.6	23.8-36.3	107	24.2	19.8-29.3
2008	230	28.7	25.0-32.8	114	34.3	27.8-41.8	116	25.1	20.7-30.2
2009	215	25.2	21.8-28.9	109	29.2	23.6-35.6	106	22.0	18.0-26.7
2010	211	24.3	21.1-28.0	112	31.4	25.4-38.2	99	19.7	16.0-24.1
2011	221	25.2	21.9-28.9	117	31.0	25.2-37.5	104	21.0	17.1-25.5
2012	206	21.9	19.0-25.3	106	25.3	20.4-31.0	100	19.2	15.6-23.5
2013	203	21.3	18.4-24.5	104	25.0	20.2-30.6	99	18.3	14.8-22.4
2014	246	25.4	22.2-28.9	112	26.1	21.2-31.6	134	24.4	20.4-29.0
2015	237	22.8	19.9-26.0	122	26.7	21.9-32.2	115	19.8	16.3-23.8
2016	242	23.3	20.4-26.6	137	31.3	26.0-37.2	105	17.8	14.5-21.6
2017	260	24.9	21.9-28.3	150	33.3	27.9-39.5	110	19.1	15.7-23.1
2018	264	24.3	21.4-27.5	131	28.4	23.5-34.0	133	21.5	17.9-25.5
2019	270	23.8	21.0-26.9	139	28.4	23.6-33.7	131	20.5	17.1-24.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Lung and Bronchus, All Races**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	5693	32.4	31.6-33.3	4031	54.0	52.3-55.8	1662	16.9	16.1-17.7
1977	5776	32.8	32.0-33.7	4054	54.7	53.0-56.5	1722	17.5	16.6-18.3
1978	5859	33.3	32.5-34.2	4141	55.9	54.1-57.7	1718	17.5	16.6-18.3
1979	6021	34.2	33.3-35.1	4118	55.5	53.7-57.2	1903	19.4	18.5-20.3
1980	6535	37.0	36.1-37.9	4525	60.7	58.9-62.6	2010	20.2	19.3-21.1
1981	6452	36.5	35.6-37.4	4332	58.4	56.6-60.2	2120	21.2	20.3-22.2
1982	6676	37.6	36.7-38.5	4378	58.6	56.8-60.4	2298	23.0	22.0-23.9
1983	6805	38.1	37.2-39.0	4468	59.6	57.8-61.4	2337	23.1	22.1-24.0
1984	6872	38.5	37.6-39.5	4477	59.5	57.7-61.4	2395	23.8	22.9-24.8
1985	7203	40.2	39.2-41.1	4627	61.3	59.5-63.2	2576	25.4	24.4-26.5
1986	7143	39.8	38.9-40.8	4525	60.1	58.3-61.9	2618	25.8	24.8-26.8
1987	7373	40.9	40.0-41.9	4679	61.8	60.0-63.7	2694	26.4	25.4-27.4
1988	7096	39.3	38.4-40.2	4346	57.4	55.7-59.2	2750	26.9	25.8-27.9
1989	7186	39.8	38.9-40.8	4326	56.8	55.0-58.5	2860	28.0	27.0-29.1
1990	7134	39.4	38.5-40.3	4382	57.9	56.1-59.6	2752	26.6	25.6-27.7
1991	7100	39.3	38.4-40.2	4226	55.3	53.6-57.0	2874	27.9	26.9-29.0
1992	6871	37.8	36.9-38.7	4117	53.7	52.0-55.4	2754	26.7	25.7-27.7
1993	6617	36.2	35.3-37.1	3832	49.7	48.1-51.3	2785	26.9	25.9-27.9
1994	6956	37.9	37.0-38.8	3951	50.9	49.3-52.5	3005	28.9	27.9-30.0
1995	7256	39.3	38.4-40.2	4151	52.9	51.3-54.6	3105	29.6	28.6-30.7
1996	7473	40.4	39.5-41.3	4248	54.2	52.6-55.9	3225	30.7	29.7-31.8
1997	7443	39.9	39.0-40.8	4219	53.4	51.7-55.0	3224	30.5	29.5-31.6
1998	7376	39.2	38.3-40.1	4095	50.8	49.2-52.4	3281	30.7	29.7-31.8
1999	7504	39.3	38.4-40.2	4165	51.2	49.7-52.8	3339	30.9	29.8-31.9
2000	7867	40.6	39.7-41.5	4281	51.9	50.3-53.5	3586	32.6	31.6-33.7
2001	7588	38.8	37.9-39.7	4065	48.9	47.4-50.5	3523	31.7	30.6-32.8
2002	8080	40.8	39.9-41.7	4377	52.1	50.6-53.7	3703	33.0	32.0-34.1
2003	8469	42.3	41.4-43.3	4607	53.9	52.4-55.5	3862	34.1	33.0-35.2
2004	9332	46.2	45.2-47.1	4966	57.6	56.0-59.2	4366	38.0	36.9-39.1
2005	9397	46.2	45.2-47.1	5041	58.0	56.4-59.6	4356	37.8	36.7-39.0
2006	9567	46.8	45.8-47.7	4935	56.2	54.7-57.8	4632	40.2	39.0-41.3
2007	9609	46.3	45.4-47.3	5008	56.3	54.7-57.9	4601	39.3	38.2-40.5
2008	9847	46.8	45.9-47.8	5099	56.4	54.9-58.0	4748	40.1	38.9-41.2
2009	9729	45.6	44.7-46.6	5154	56.1	54.5-57.7	4575	38.1	37.0-39.3
2010	9847	45.4	44.5-46.3	5144	54.7	53.2-56.3	4703	38.8	37.6-39.9
2011	10027	45.4	44.5-46.3	5182	54.2	52.7-55.7	4845	39.1	38.0-40.2
2012	10036	44.6	43.7-45.5	5167	52.8	51.3-54.3	4869	38.7	37.6-39.9
2013	9910	43.2	42.4-44.1	5028	50.2	48.8-51.7	4882	38.2	37.1-39.3
2014	9680	41.3	40.4-42.1	5010	48.9	47.5-50.3	4670	35.7	34.6-36.7
2015	9864	41.4	40.6-42.2	5092	49.0	47.6-50.3	4772	36.0	34.9-37.0
2016	9264	38.1	37.3-38.9	4719	44.3	43.1-45.7	4545	33.6	32.6-34.6
2017	9254	37.5	36.7-38.2	4731	43.4	42.1-44.7	4523	33.0	32.1-34.0
2018	8935	35.5	34.7-36.3	4538	40.8	39.6-42.0	4397	31.6	30.6-32.5
2019	8954	34.8	34.1-35.6	4446	39.0	37.9-40.2	4508	31.7	30.8-32.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Lung and Bronchus, Whites**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	5041	31.8	30.9-32.7	3547	52.6	50.8-54.4	1494	16.9	16.0-17.8
1977	5090	32.1	31.2-33.1	3541	52.9	51.1-54.7	1549	17.7	16.8-18.6
1978	5152	32.6	31.7-33.6	3628	54.4	52.6-56.2	1524	17.4	16.5-18.3
1979	5263	33.4	32.5-34.3	3577	53.5	51.7-55.4	1686	19.4	18.4-20.4
1980	5681	36.0	35.1-37.0	3906	58.4	56.5-60.3	1775	20.1	19.2-21.1
1981	5617	35.8	34.9-36.8	3727	56.2	54.4-58.1	1890	21.6	20.6-22.6
1982	5787	36.8	35.8-37.7	3759	56.4	54.5-58.3	2028	23.1	22.1-24.2
1983	5940	37.7	36.8-38.7	3848	57.9	56.0-59.8	2092	23.6	22.6-24.7
1984	5950	38.1	37.1-39.1	3847	58.0	56.1-59.9	2103	24.1	23.1-25.2
1985	6238	39.7	38.7-40.7	3955	59.4	57.5-61.4	2283	26.1	25.0-27.2
1986	6166	39.5	38.5-40.5	3872	58.6	56.7-60.5	2294	26.3	25.2-27.4
1987	6366	40.8	39.8-41.8	3991	60.3	58.4-62.3	2375	27.2	26.1-28.4
1988	6171	39.6	38.6-40.6	3730	56.5	54.7-58.4	2441	27.9	26.8-29.1
1989	6181	39.8	38.8-40.9	3660	55.4	53.5-57.2	2521	29.0	27.8-30.2
1990	6139	39.5	38.5-40.5	3743	56.9	55.1-58.8	2396	27.4	26.2-28.5
1991	6129	39.7	38.7-40.8	3590	54.4	52.6-56.3	2539	29.2	28.0-30.4
1992	5868	37.9	36.9-38.9	3458	52.4	50.7-54.2	2410	27.8	26.7-29.0
1993	5617	36.3	35.4-37.3	3199	48.3	46.6-50.1	2418	28.0	26.8-29.1
1994	5892	38.0	37.1-39.0	3316	50.0	48.3-51.8	2576	29.8	28.7-31.0
1995	6172	39.8	38.8-40.8	3484	52.2	50.5-54.0	2688	30.8	29.7-32.0
1996	6318	40.8	39.8-41.9	3571	53.8	52.1-55.7	2747	31.6	30.4-32.8
1997	6281	40.4	39.4-41.4	3530	52.8	51.1-54.6	2751	31.6	30.4-32.8
1998	6188	39.5	38.6-40.5	3400	50.2	48.5-51.9	2788	31.7	30.6-33.0
1999	6421	40.7	39.7-41.7	3525	51.8	50.1-53.5	2896	32.8	31.6-34.0
2000	6692	42.0	41.0-43.0	3584	52.1	50.4-53.9	3108	34.8	33.6-36.1
2001	6449	40.2	39.2-41.2	3399	49.3	47.6-50.9	3050	33.8	32.6-35.1
2002	6919	42.8	41.8-43.8	3719	53.4	51.7-55.2	3200	35.4	34.2-36.7
2003	7164	44.1	43.0-45.1	3878	55.1	53.3-56.8	3286	36.1	34.9-37.4
2004	7889	48.1	47.0-49.1	4141	58.4	56.7-60.3	3748	40.7	39.4-42.1
2005	7937	48.3	47.2-49.3	4211	59.1	57.3-61.0	3726	40.5	39.2-41.8
2006	8133	49.5	48.4-50.6	4137	57.8	56.0-59.6	3996	43.7	42.4-45.1
2007	8077	48.6	47.5-49.7	4169	57.6	55.9-59.4	3908	42.2	40.8-43.5
2008	8313	49.5	48.4-50.6	4258	58.1	56.3-59.9	4055	43.4	42.1-44.8
2009	8155	48.0	47.0-49.1	4269	57.5	55.7-59.2	3886	41.2	39.9-42.6
2010	8194	47.8	46.7-48.8	4237	56.1	54.4-57.8	3957	41.8	40.5-43.2
2011	8296	47.6	46.6-48.7	4235	55.3	53.6-57.0	4061	42.2	40.8-43.5
2012	8224	46.5	45.5-47.5	4190	53.6	52.0-55.3	4034	41.5	40.2-42.8
2013	8210	45.9	44.9-46.9	4113	51.7	50.1-53.4	4097	41.6	40.3-42.9
2014	7924	43.4	42.4-44.4	4041	49.9	48.3-51.5	3883	38.6	37.3-39.8
2015	8063	43.8	42.8-44.7	4081	49.9	48.4-51.5	3982	39.4	38.1-40.6
2016	7474	39.8	38.9-40.7	3734	44.7	43.3-46.2	3740	36.3	35.1-37.5
2017	7466	39.4	38.5-40.3	3739	44.0	42.5-45.4	3727	35.9	34.7-37.1
2018	7184	37.3	36.4-38.2	3565	41.2	39.8-42.6	3619	34.4	33.3-35.6
2019	7126	36.3	35.4-37.1	3475	39.3	38.0-40.7	3651	34.1	33.0-35.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Lung and Bronchus, Blacks**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	575	35.4	32.4-38.6	428	64.0	57.7-70.9	147	15.7	13.2-18.6
1977	629	39.0	35.8-42.4	464	72.7	65.5-80.4	165	17.0	14.4-19.9
1978	645	38.6	35.5-41.8	473	69.8	63.1-77.0	172	17.8	15.1-20.7
1979	694	41.4	38.2-44.8	500	74.3	67.3-81.8	194	19.9	17.0-23.0
1980	784	44.4	41.2-47.8	573	81.9	74.7-89.5	211	20.0	17.3-23.0
1981	774	42.2	39.2-45.4	558	77.5	70.6-84.8	216	19.3	16.8-22.1
1982	784	42.5	39.4-45.7	552	75.4	68.8-82.5	232	21.0	18.3-24.0
1983	789	41.4	38.5-44.5	569	76.0	69.4-83.0	220	19.3	16.8-22.1
1984	840	43.5	40.5-46.7	570	74.0	67.7-80.8	270	23.7	20.9-26.8
1985	879	44.2	41.2-47.3	612	77.8	71.3-84.6	267	22.6	19.9-25.6
1986	888	43.8	40.8-46.8	597	76.6	70.1-83.5	291	23.5	20.9-26.4
1987	906	43.5	40.7-46.6	621	76.3	70.0-83.0	285	22.9	20.2-25.7
1988	846	40.3	37.5-43.2	559	69.2	63.2-75.6	287	22.6	20.0-25.4
1989	914	42.3	39.5-45.2	606	72.9	66.8-79.3	308	23.6	21.0-26.4
1990	887	41.0	38.3-43.9	571	69.6	63.5-76.0	316	23.8	21.3-26.7
1991	868	39.8	37.1-42.6	568	68.6	62.6-74.9	300	22.5	20.0-25.3
1992	884	38.7	36.1-41.4	582	66.2	60.5-72.2	302	21.9	19.4-24.5
1993	808	35.7	33.2-38.3	506	59.5	54.0-65.4	302	21.8	19.4-24.5
1994	876	37.5	35.0-40.1	524	58.3	53.1-63.9	352	24.8	22.2-27.6
1995	887	37.6	35.2-40.3	532	57.7	52.6-63.2	355	25.1	22.5-27.8
1996	1015	42.1	39.5-44.9	589	63.6	58.2-69.3	426	29.3	26.5-32.2
1997	990	40.8	38.3-43.5	586	63.2	57.8-68.8	404	27.4	24.8-30.3
1998	985	39.4	37.0-42.0	558	57.0	52.1-62.2	427	28.5	25.8-31.3
1999	918	36.1	33.8-38.6	538	54.7	49.9-59.8	380	24.8	22.3-27.4
2000	990	37.7	35.4-40.2	581	56.6	51.8-61.6	409	25.9	23.5-28.6
2001	954	36.1	33.8-38.5	540	52.7	48.1-57.6	414	25.9	23.4-28.5
2002	968	35.6	33.3-37.9	543	52.1	47.6-56.9	425	25.7	23.3-28.3
2003	1045	37.6	35.3-40.0	554	51.1	46.7-55.8	491	29.3	26.7-32.0
2004	1168	41.1	38.8-43.6	658	57.7	53.2-62.5	510	30.1	27.5-32.9
2005	1193	41.9	39.5-44.3	653	59.6	54.9-64.6	540	31.5	28.9-34.3
2006	1157	40.0	37.7-42.5	629	55.5	51.0-60.3	528	30.4	27.8-33.1
2007	1169	39.4	37.1-41.7	630	53.9	49.5-58.5	539	30.2	27.7-32.9
2008	1176	39.4	37.1-41.8	637	54.5	50.1-59.1	539	30.0	27.5-32.7
2009	1219	39.5	37.2-41.8	663	54.4	50.1-58.9	556	30.1	27.6-32.8
2010	1273	40.3	38.0-42.6	674	54.1	49.9-58.6	599	31.7	29.2-34.4
2011	1312	40.4	38.2-42.8	693	54.4	50.2-58.8	619	31.8	29.3-34.4
2012	1337	40.2	38.0-42.5	702	53.4	49.3-57.7	635	32.0	29.5-34.6
2013	1213	35.3	33.3-37.4	638	46.9	43.2-50.9	575	27.9	25.7-30.4
2014	1266	36.1	34.1-38.2	671	47.9	44.1-51.8	595	28.4	26.1-30.8
2015	1259	34.6	32.7-36.6	683	47.3	43.7-51.2	576	26.5	24.4-28.8
2016	1242	33.7	31.8-35.7	652	43.8	40.4-47.5	590	26.9	24.7-29.2
2017	1208	31.8	30.0-33.7	654	43.0	39.7-46.6	554	24.7	22.6-26.9
2018	1186	30.7	29.0-32.6	623	40.4	37.1-43.8	563	24.4	22.4-26.6
2019	1181	29.9	28.2-31.7	607	37.8	34.7-41.0	574	24.6	22.6-26.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Lung and Bronchus, All Races**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	2475	32.1	30.8-33.4	1725	53.8	51.2-56.5	750	16.9	15.7-18.2
1977	2472	32.5	31.2-33.9	1714	55.2	52.5-57.9	758	17.2	15.9-18.5
1978	2504	33.5	32.1-34.8	1750	56.6	53.9-59.4	754	17.7	16.4-19.0
1979	2504	33.6	32.2-34.9	1694	55.6	52.9-58.4	810	18.7	17.4-20.1
1980	2588	35.1	33.8-36.6	1775	58.5	55.8-61.4	813	19.2	17.8-20.6
1981	2534	34.4	33.0-35.8	1673	56.1	53.3-58.9	861	19.7	18.4-21.1
1982	2698	36.7	35.3-38.2	1717	57.4	54.7-60.3	981	22.8	21.4-24.4
1983	2632	35.7	34.3-37.1	1709	57.4	54.6-60.2	923	21.2	19.8-22.6
1984	2676	36.5	35.1-37.9	1750	58.7	55.9-61.6	926	21.7	20.3-23.2
1985	2750	37.3	35.9-38.7	1780	59.2	56.4-62.1	970	22.5	21.1-24.0
1986	2703	36.7	35.3-38.2	1694	56.5	53.8-59.3	1009	23.4	22.0-25.0
1987	2854	38.7	37.2-40.1	1810	60.6	57.7-63.5	1044	24.0	22.5-25.5
1988	2621	35.9	34.5-37.3	1601	54.3	51.6-57.1	1020	23.7	22.2-25.2
1989	2596	36.0	34.6-37.4	1531	51.9	49.3-54.6	1065	25.3	23.8-26.9
1990	2547	35.4	34.1-36.8	1519	52.4	49.8-55.2	1028	24.3	22.8-25.9
1991	2547	35.8	34.4-37.2	1497	51.4	48.8-54.1	1050	25.1	23.6-26.7
1992	2614	36.6	35.2-38.1	1603	55.3	52.6-58.1	1011	24.1	22.7-25.7
1993	2434	34.2	32.8-35.5	1442	49.8	47.2-52.5	992	23.9	22.4-25.4
1994	2505	35.1	33.8-36.5	1417	48.6	46.1-51.2	1088	26.1	24.6-27.7
1995	2566	35.8	34.4-37.2	1469	50.0	47.4-52.6	1097	25.9	24.4-27.5
1996	2690	37.5	36.1-38.9	1508	51.6	48.9-54.3	1182	28.1	26.5-29.7
1997	2606	36.2	34.8-37.6	1460	49.6	47.1-52.3	1146	27.1	25.6-28.8
1998	2626	36.0	34.6-37.4	1446	47.9	45.4-50.4	1180	27.6	26.0-29.2
1999	2511	33.8	32.4-35.1	1394	45.8	43.4-48.3	1117	25.6	24.2-27.2
2000	2745	36.3	35.0-37.7	1504	48.5	46.1-51.1	1241	28.0	26.4-29.6
2001	2539	33.2	32.0-34.6	1388	44.3	42.0-46.7	1151	25.6	24.1-27.1
2002	2670	34.5	33.2-35.8	1447	45.4	43.1-47.9	1223	27.0	25.5-28.5
2003	2727	35.1	33.8-36.4	1519	47.5	45.1-49.9	1208	26.5	25.0-28.0
2004	3025	38.5	37.2-39.9	1658	51.2	48.7-53.8	1367	29.7	28.2-31.3
2005	3023	38.2	36.8-39.6	1652	50.8	48.3-53.3	1371	29.8	28.2-31.4
2006	3010	38.1	36.7-39.5	1625	50.0	47.5-52.5	1385	29.9	28.4-31.6
2007	3114	39.0	37.6-40.4	1663	50.4	47.9-52.9	1451	31.1	29.6-32.8
2008	3038	37.7	36.3-39.0	1598	47.7	45.3-50.1	1440	30.8	29.2-32.4
2009	3020	36.9	35.6-38.3	1608	47.4	45.1-49.8	1412	29.7	28.2-31.3
2010	3056	36.7	35.4-38.1	1645	47.3	45.0-49.7	1411	29.2	27.7-30.8
2011	3152	37.1	35.8-38.4	1682	47.7	45.4-50.1	1470	29.8	28.3-31.4
2012	3112	35.8	34.5-37.1	1609	44.2	42.0-46.5	1503	30.1	28.6-31.7
2013	3011	33.9	32.7-35.2	1579	42.4	40.3-44.6	1432	28.0	26.6-29.6
2014	2986	32.7	31.5-33.9	1602	41.5	39.4-43.6	1384	26.4	25.0-27.8
2015	2953	31.8	30.6-33.0	1620	41.3	39.2-43.4	1333	25.1	23.8-26.5
2016	2879	30.3	29.2-31.4	1489	37.1	35.2-39.0	1390	25.5	24.1-26.9
2017	2892	30.0	28.9-31.2	1553	37.9	36.0-39.9	1339	24.4	23.1-25.7
2018	2732	27.7	26.7-28.8	1432	33.8	32.0-35.6	1300	23.2	22.0-24.6
2019	2789	27.9	26.8-28.9	1418	33.0	31.2-34.7	1371	24.1	22.8-25.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Lung and Bronchus, Whites**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	1976	30.7	29.3-32.1	1356	50.7	47.9-53.6	620	16.8	15.5-18.3
1977	1983	31.5	30.1-33.0	1355	52.4	49.6-55.4	628	17.4	16.0-18.9
1978	1956	32.0	30.6-33.5	1360	53.8	50.9-56.8	596	17.3	15.9-18.8
1979	1935	31.8	30.3-33.3	1292	51.8	48.9-54.8	643	18.4	17.0-20.0
1980	1971	33.2	31.7-34.8	1326	54.0	51.1-57.1	645	19.0	17.5-20.6
1981	1927	32.8	31.3-34.3	1244	52.0	49.0-55.0	683	19.8	18.3-21.4
1982	2067	35.6	34.0-37.2	1289	54.1	51.1-57.3	778	23.2	21.5-24.9
1983	1997	34.7	33.2-36.3	1259	54.0	50.9-57.1	738	21.9	20.3-23.6
1984	2001	35.5	33.9-37.2	1296	56.2	53.1-59.4	705	21.8	20.2-23.6
1985	2037	36.0	34.4-37.6	1289	55.4	52.3-58.6	748	22.9	21.2-24.7
1986	1969	35.3	33.7-37.0	1216	53.0	50.0-56.1	753	23.3	21.6-25.1
1987	2065	37.4	35.7-39.1	1280	56.5	53.4-59.7	785	24.4	22.7-26.3
1988	1954	36.3	34.6-37.9	1164	52.7	49.6-55.9	790	25.2	23.4-27.1
1989	1883	35.8	34.1-37.5	1065	48.8	45.9-51.9	818	26.9	25.1-28.9
1990	1819	35.0	33.4-36.7	1064	49.9	46.9-53.1	755	25.1	23.3-27.1
1991	1853	36.5	34.8-38.2	1043	49.5	46.5-52.6	810	27.3	25.4-29.4
1992	1891	37.4	35.7-39.2	1130	54.3	51.1-57.6	761	26.0	24.1-28.0
1993	1725	34.7	33.0-36.4	1003	48.4	45.4-51.5	722	25.4	23.6-27.4
1994	1791	36.2	34.6-38.0	982	47.6	44.6-50.7	809	28.7	26.7-30.8
1995	1810	36.7	35.0-38.5	1012	49.3	46.3-52.5	798	27.6	25.7-29.7
1996	1860	38.0	36.3-39.8	1026	50.6	47.5-53.8	834	29.4	27.4-31.5
1997	1776	36.4	34.7-38.1	981	48.3	45.3-51.5	795	28.1	26.2-30.2
1998	1782	36.3	34.6-38.0	959	46.7	43.8-49.8	823	28.6	26.7-30.7
1999	1747	35.3	33.7-37.0	937	45.5	42.6-48.5	810	28.4	26.4-30.4
2000	1918	38.6	36.8-40.3	1029	49.5	46.5-52.6	889	30.8	28.8-33.0
2001	1749	35.0	33.4-36.7	931	44.6	41.8-47.6	818	28.1	26.2-30.1
2002	1842	36.7	35.1-38.4	978	46.5	43.6-49.5	864	29.6	27.7-31.7
2003	1817	36.3	34.6-38.0	1016	48.5	45.5-51.6	801	27.3	25.4-29.3
2004	2012	40.1	38.3-41.9	1090	51.9	48.8-55.1	922	31.8	29.8-34.0
2005	2006	40.1	38.3-41.9	1085	51.7	48.7-54.9	921	32.0	30.0-34.2
2006	1994	40.3	38.5-42.1	1063	51.1	48.1-54.3	931	32.6	30.5-34.8
2007	2027	41.0	39.2-42.8	1077	51.7	48.6-54.9	950	33.3	31.2-35.5
2008	1972	39.9	38.1-41.7	1017	48.4	45.5-51.5	955	33.8	31.7-36.1
2009	1935	38.7	37.0-40.5	1013	47.9	44.9-50.9	922	32.3	30.2-34.5
2010	1919	38.4	36.6-40.1	1016	47.7	44.8-50.8	903	31.5	29.5-33.7
2011	1918	37.8	36.1-39.5	1014	47.0	44.2-50.1	904	31.2	29.1-33.3
2012	1872	36.3	34.7-38.0	963	43.4	40.7-46.3	909	31.3	29.3-33.5
2013	1858	35.5	33.9-37.2	962	42.8	40.1-45.6	896	30.3	28.3-32.4
2014	1764	33.0	31.5-34.6	917	39.8	37.2-42.5	847	28.0	26.1-30.0
2015	1727	32.2	30.7-33.8	938	40.6	38.1-43.4	789	26.2	24.3-28.1
2016	1672	30.6	29.1-32.1	832	35.4	33.0-38.0	840	27.2	25.4-29.2
2017	1634	29.8	28.4-31.3	849	35.8	33.4-38.3	785	25.3	23.5-27.2
2018	1491	26.7	25.3-28.1	726	29.6	27.5-31.9	765	24.6	22.9-26.5
2019	1503	26.6	25.3-28.0	745	30.1	27.9-32.4	758	24.0	22.3-25.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Lung and Bronchus, Blacks**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	441	35.0	31.6-38.6	324	63.8	56.4-71.8	117	16.0	13.1-19.3
1977	448	35.5	32.1-39.2	325	66.7	58.8-75.3	123	15.8	13.1-19.0
1978	497	38.0	34.5-41.6	356	68.1	60.6-76.2	141	18.5	15.5-22.0
1979	516	39.8	36.2-43.6	370	71.3	63.6-79.7	146	19.5	16.3-23.2
1980	564	41.3	37.8-45.0	414	78.1	70.1-86.8	150	18.2	15.3-21.5
1981	562	39.4	36.1-43.0	396	72.3	64.7-80.5	166	18.8	16.0-22.0
1982	564	39.4	36.1-43.0	382	68.5	61.3-76.2	182	21.1	18.1-24.5
1983	575	39.0	35.8-42.5	408	71.6	64.3-79.5	167	18.8	16.0-22.0
1984	610	40.6	37.3-44.0	405	69.6	62.5-77.3	205	22.6	19.5-26.0
1985	649	42.4	39.1-45.9	446	75.5	68.1-83.3	203	22.1	19.1-25.4
1986	660	41.8	38.6-45.3	430	72.3	65.1-80.1	230	23.7	20.6-27.0
1987	702	43.9	40.6-47.4	471	77.8	70.4-85.7	231	23.6	20.6-26.9
1988	616	38.0	35.0-41.2	399	65.8	59.0-73.1	217	21.7	18.9-24.9
1989	643	38.8	35.8-42.0	417	67.3	60.6-74.5	226	22.1	19.3-25.3
1990	649	38.7	35.7-41.8	402	64.3	57.7-71.4	247	23.8	20.9-27.0
1991	613	36.6	33.6-39.7	401	64.6	58.0-71.7	212	20.4	17.7-23.4
1992	634	36.3	33.5-39.3	412	62.3	56.0-69.0	222	20.8	18.1-23.8
1993	592	34.5	31.7-37.5	362	58.0	51.7-64.9	230	21.6	18.8-24.6
1994	618	35.1	32.4-38.1	376	56.9	51.0-63.4	242	22.3	19.5-25.3
1995	633	35.5	32.8-38.5	376	55.6	49.8-61.9	257	23.5	20.7-26.6
1996	721	39.8	36.9-42.9	411	60.4	54.3-66.9	310	27.9	24.8-31.2
1997	691	37.9	35.1-40.9	393	57.8	51.9-64.2	298	26.5	23.6-29.7
1998	680	36.3	33.6-39.2	376	52.4	46.9-58.2	304	26.6	23.7-29.8
1999	637	33.3	30.7-36.0	373	51.8	46.4-57.6	264	22.4	19.8-25.3
2000	679	34.7	32.1-37.4	380	51.2	46.0-56.9	299	24.8	22.1-27.8
2001	635	32.3	29.8-34.9	351	47.2	42.2-52.6	284	23.4	20.8-26.3
2002	670	32.9	30.4-35.5	371	49.0	43.9-54.5	299	23.8	21.2-26.7
2003	702	34.2	31.6-36.8	359	46.2	41.4-51.5	343	27.1	24.3-30.2
2004	790	37.9	35.3-40.7	428	52.4	47.4-57.9	362	28.5	25.6-31.6
2005	804	38.2	35.5-40.9	423	53.4	48.2-59.0	381	29.5	26.6-32.7
2006	798	37.3	34.7-40.0	420	51.2	46.2-56.5	378	28.7	25.9-31.8
2007	791	36.3	33.7-38.9	411	48.8	44.0-54.0	380	28.6	25.8-31.7
2008	791	36.4	33.8-39.1	417	50.1	45.1-55.4	374	28.2	25.4-31.2
2009	803	35.8	33.3-38.5	417	48.3	43.6-53.4	386	28.3	25.5-31.3
2010	833	36.3	33.9-38.9	433	49.2	44.5-54.4	400	28.7	25.9-31.7
2011	898	38.3	35.8-40.9	457	51.3	46.4-56.4	441	30.6	27.8-33.7
2012	872	36.3	33.9-38.9	428	46.4	41.9-51.2	444	30.4	27.6-33.4
2013	777	31.6	29.3-33.9	400	41.8	37.6-46.4	377	25.2	22.7-27.9
2014	833	33.0	30.8-35.4	433	44.3	40.0-48.8	400	26.1	23.6-28.8
2015	783	29.7	27.7-31.9	409	39.2	35.4-43.4	374	23.7	21.3-26.3
2016	787	29.6	27.5-31.8	394	37.2	33.5-41.2	393	24.5	22.1-27.0
2017	797	29.2	27.2-31.4	426	39.5	35.7-43.6	371	22.7	20.4-25.2
2018	779	28.2	26.3-30.3	411	37.8	34.1-41.7	368	22.1	19.8-24.5
2019	771	27.6	25.6-29.7	380	34.1	30.7-37.9	391	23.2	21.0-25.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Lung and Bronchus, All Races**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	3218	32.8	31.6-34.0	2306	54.4	52.1-56.8	912	16.9	15.8-18.0
1977	3304	33.1	32.0-34.3	2340	54.5	52.3-56.9	964	17.7	16.6-18.9
1978	3355	33.3	32.2-34.5	2391	55.4	53.1-57.8	964	17.4	16.3-18.5
1979	3516	34.6	33.5-35.8	2423	55.4	53.1-57.7	1093	19.8	18.6-21.0
1980	3946	38.4	37.2-39.7	2750	62.4	60.0-64.9	1196	21.0	19.8-22.3
1981	3916	38.1	36.9-39.3	2657	60.1	57.8-62.6	1259	22.3	21.1-23.6
1982	3978	38.2	37.0-39.4	2661	59.3	57.0-61.7	1317	23.0	21.7-24.3
1983	4173	39.8	38.6-41.1	2759	61.0	58.6-63.4	1414	24.4	23.1-25.8
1984	4194	40.0	38.7-41.2	2726	60.0	57.7-62.4	1468	25.4	24.1-26.8
1985	4448	42.2	40.9-43.5	2842	62.7	60.3-65.1	1606	27.6	26.3-29.0
1986	4435	42.0	40.7-43.3	2828	62.5	60.1-64.9	1607	27.5	26.2-28.9
1987	4516	42.4	41.2-43.7	2869	62.7	60.3-65.1	1647	28.1	26.7-29.5
1988	4473	41.6	40.4-42.9	2745	59.4	57.2-61.8	1728	29.1	27.7-30.5
1989	4589	42.4	41.2-43.7	2794	59.9	57.6-62.2	1795	30.0	28.6-31.4
1990	4584	42.0	40.7-43.2	2862	61.2	58.9-63.6	1722	28.2	26.9-29.6
1991	4551	41.5	40.3-42.7	2727	57.6	55.4-59.9	1824	29.9	28.5-31.3
1992	4257	38.5	37.4-39.7	2514	52.6	50.5-54.7	1743	28.5	27.1-29.8
1993	4182	37.5	36.4-38.7	2390	49.5	47.5-51.6	1792	29.0	27.6-30.3
1994	4449	39.6	38.4-40.8	2534	52.2	50.2-54.3	1915	30.9	29.5-32.3
1995	4690	41.5	40.3-42.7	2682	54.5	52.5-56.7	2008	32.1	30.7-33.6
1996	4782	42.3	41.1-43.5	2739	55.8	53.7-57.9	2043	32.5	31.1-34.0
1997	4837	42.3	41.2-43.6	2759	55.6	53.5-57.7	2078	32.8	31.4-34.3
1998	4750	41.2	40.0-42.4	2649	52.4	50.4-54.4	2101	33.0	31.6-34.4
1999	4992	42.8	41.6-44.0	2771	54.5	52.4-56.5	2221	34.4	33.0-35.9
2000	5121	43.4	42.2-44.6	2776	53.9	51.9-55.9	2345	35.9	34.4-37.4
2001	5048	42.4	41.2-43.6	2676	51.7	49.7-53.7	2372	35.9	34.5-37.4
2002	5410	44.9	43.7-46.1	2930	56.1	54.1-58.2	2480	37.2	35.8-38.7
2003	5742	47.0	45.8-48.3	3088	57.8	55.8-59.9	2654	39.4	37.9-40.9
2004	6303	51.0	49.7-52.3	3305	61.2	59.1-63.4	2998	43.6	42.1-45.2
2005	6374	51.2	49.9-52.5	3389	62.2	60.1-64.4	2985	43.2	41.7-44.8
2006	6557	52.3	51.0-53.6	3310	60.0	57.9-62.1	3247	47.0	45.4-48.7
2007	6494	50.9	49.7-52.2	3344	59.6	57.6-61.7	3150	44.7	43.2-46.3
2008	6809	52.5	51.3-53.8	3501	61.5	59.4-63.6	3308	46.2	44.7-47.9
2009	6709	51.1	49.8-52.3	3546	61.0	59.0-63.1	3163	43.7	42.2-45.3
2010	6791	50.8	49.6-52.1	3499	58.9	57.0-61.0	3292	45.1	43.6-46.7
2011	6875	50.5	49.3-51.8	3500	57.9	56.0-59.9	3375	45.3	43.7-46.9
2012	6923	50.1	48.9-51.3	3557	57.9	55.9-59.8	3366	44.5	43.0-46.0
2013	6899	49.2	48.0-50.4	3449	54.9	53.0-56.8	3450	45.0	43.4-46.5
2014	6694	46.7	45.6-47.9	3408	53.2	51.4-55.1	3286	41.9	40.4-43.4
2015	6911	47.6	46.4-48.7	3472	53.6	51.8-55.4	3439	43.3	41.8-44.8
2016	6385	43.1	42.0-44.2	3230	48.7	47.0-50.5	3155	39.1	37.7-40.5
2017	6362	42.2	41.2-43.3	3178	46.7	45.1-48.4	3184	38.9	37.6-40.3
2018	6201	40.5	39.5-41.5	3105	45.0	43.4-46.7	3096	37.1	35.8-38.5
2019	6164	39.3	38.3-40.3	3027	42.7	41.1-44.3	3137	36.9	35.6-38.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Lung and Bronchus, Whites**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	3065	32.5	31.3-33.7	2191	53.8	51.5-56.2	874	16.8	15.7-18.0
1977	3107	32.5	31.3-33.7	2186	53.1	50.8-55.5	921	17.7	16.5-18.9
1978	3196	33.1	31.9-34.3	2268	54.7	52.4-57.1	928	17.4	16.3-18.6
1979	3327	34.2	33.0-35.4	2284	54.4	52.1-56.8	1043	19.7	18.5-21.0
1980	3709	37.7	36.5-39.0	2580	61.1	58.7-63.6	1129	20.8	19.6-22.1
1981	3688	37.5	36.3-38.8	2481	58.8	56.4-61.2	1207	22.5	21.2-23.8
1982	3720	37.3	36.1-38.6	2470	57.6	55.2-60.0	1250	22.9	21.6-24.3
1983	3943	39.4	38.2-40.7	2589	60.0	57.6-62.4	1354	24.5	23.2-25.9
1984	3947	39.5	38.2-40.7	2550	58.9	56.5-61.3	1397	25.5	24.1-26.9
1985	4197	41.8	40.5-43.1	2662	61.6	59.2-64.0	1535	27.9	26.4-29.3
1986	4193	41.8	40.5-43.1	2653	61.6	59.2-64.1	1540	27.9	26.5-29.3
1987	4298	42.6	41.3-43.9	2711	62.4	60.0-64.9	1587	28.6	27.2-30.1
1988	4215	41.3	40.1-42.6	2566	58.5	56.2-60.9	1649	29.3	27.9-30.8
1989	4297	41.9	40.7-43.2	2594	58.7	56.4-61.1	1703	30.1	28.6-31.6
1990	4318	41.7	40.5-43.0	2678	60.3	58.0-62.7	1640	28.5	27.1-29.9
1991	4274	41.3	40.1-42.6	2545	56.9	54.6-59.2	1729	30.1	28.7-31.6
1992	3977	38.1	37.0-39.4	2328	51.5	49.4-53.7	1649	28.6	27.3-30.1
1993	3892	37.1	36.0-38.3	2196	48.3	46.2-50.4	1696	29.2	27.8-30.7
1994	4100	38.9	37.7-40.1	2334	51.1	49.0-53.3	1766	30.4	29.0-31.9
1995	4362	41.2	40.0-42.4	2472	53.5	51.4-55.7	1890	32.3	30.9-33.8
1996	4457	42.1	40.9-43.4	2544	55.3	53.1-57.5	1913	32.6	31.2-34.2
1997	4505	42.2	41.0-43.5	2549	54.9	52.7-57.0	1956	33.2	31.7-34.7
1998	4406	41.0	39.8-42.2	2441	51.6	49.6-53.7	1965	33.2	31.7-34.7
1999	4673	43.1	41.9-44.3	2588	54.5	52.4-56.6	2085	34.8	33.3-36.4
2000	4773	43.6	42.3-44.8	2554	53.2	51.2-55.4	2219	36.7	35.2-38.2
2001	4699	42.6	41.4-43.8	2467	51.3	49.3-53.4	2232	36.5	35.0-38.1
2002	5077	45.5	44.3-46.8	2741	56.5	54.4-58.7	2336	38.1	36.5-39.7
2003	5347	47.5	46.2-48.8	2862	57.9	55.8-60.1	2485	40.1	38.5-41.7
2004	5875	51.6	50.3-52.9	3050	61.2	59.0-63.4	2825	44.7	43.1-46.4
2005	5931	51.8	50.5-53.2	3126	62.2	60.1-64.5	2805	44.3	42.6-46.0
2006	6139	53.4	52.1-54.8	3074	60.5	58.4-62.7	3065	48.6	46.9-50.4
2007	6049	51.8	50.5-53.1	3091	60.0	57.9-62.2	2958	46.0	44.4-47.8
2008	6341	53.5	52.2-54.9	3241	62.0	59.8-64.2	3100	47.6	45.9-49.4
2009	6220	51.9	50.6-53.2	3256	61.3	59.1-63.4	2964	45.0	43.4-46.7
2010	6275	51.7	50.4-53.0	3221	59.4	57.3-61.5	3054	46.2	44.6-48.0
2011	6378	51.7	50.4-53.0	3221	58.5	56.5-60.6	3157	46.9	45.2-48.6
2012	6351	50.7	49.4-52.0	3226	57.7	55.7-59.8	3125	45.7	44.1-47.4
2013	6352	50.1	48.9-51.4	3151	55.3	53.3-57.3	3201	46.4	44.7-48.0
2014	6160	47.6	46.4-48.9	3124	53.9	52.0-55.9	3036	43.0	41.5-44.6
2015	6336	48.5	47.3-49.8	3143	53.6	51.7-55.6	3193	45.0	43.4-46.6
2016	5802	43.5	42.4-44.7	2902	48.4	46.6-50.3	2900	40.2	38.7-41.7
2017	5832	43.3	42.1-44.4	2890	47.2	45.5-49.0	2942	40.3	38.8-41.9
2018	5691	41.6	40.5-42.7	2838	45.8	44.1-47.5	2853	38.5	37.0-40.0
2019	5622	40.2	39.1-41.3	2729	43.0	41.3-44.7	2893	38.3	36.9-39.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Lung and Bronchus, Blacks**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	134	37.0	30.7-44.1	104	65.1	52.5-79.7	30	14.7	9.7-21.2
1977	181	51.7	43.9-60.3	139	91.8	75.7-110.2	42	21.3	15.0-29.2
1978	148	40.6	34.0-48.0	117	75.3	60.9-91.8	31	14.9	10.0-21.3
1979	178	47.1	40.0-54.9	130	84.0	68.5-101.7	48	21.1	15.5-28.0
1980	220	55.3	47.9-63.4	159	93.8	78.6-110.9	61	26.9	20.4-34.7
1981	212	52.0	44.9-59.8	162	94.4	79.2-111.5	50	21.1	15.5-27.9
1982	220	53.1	46.0-60.9	170	97.5	82.2-114.7	50	20.9	15.4-27.6
1983	214	49.7	43.0-57.0	161	89.8	75.5-105.8	53	20.8	15.5-27.3
1984	230	53.7	46.7-61.4	165	87.9	74.1-103.4	65	27.9	21.3-35.8
1985	230	50.3	43.8-57.5	166	84.8	71.7-99.6	64	24.7	18.9-31.6
1986	227	50.6	44.0-57.9	167	90.4	76.0-106.5	60	22.9	17.4-29.6
1987	204	42.1	36.4-48.5	150	71.4	59.8-84.4	54	20.0	14.9-26.2
1988	230	47.9	41.7-54.7	160	79.6	66.9-93.9	70	25.5	19.8-32.3
1989	271	54.0	47.5-61.0	189	89.8	76.5-104.7	82	28.7	22.8-35.7
1990	238	48.7	42.5-55.6	169	85.8	72.0-101.1	69	24.1	18.7-30.6
1991	255	50.8	44.5-57.6	167	80.7	67.7-95.1	88	30.3	24.2-37.4
1992	250	46.4	40.7-52.8	170	77.7	65.4-91.6	80	25.4	20.1-31.7
1993	216	39.4	34.2-45.2	144	63.6	52.9-75.8	72	22.8	17.7-28.7
1994	258	45.0	39.5-51.0	148	62.4	52.1-74.1	110	33.2	27.2-40.1
1995	254	44.3	38.8-50.2	156	63.8	53.4-75.4	98	30.2	24.4-36.9
1996	294	49.2	43.6-55.3	178	72.5	61.4-84.9	116	33.8	27.8-40.5
1997	299	49.8	44.1-55.9	193	78.1	66.6-90.8	106	30.5	24.9-37.0
1998	305	49.0	43.5-55.0	182	69.9	59.3-81.7	123	34.6	28.7-41.3
1999	281	45.1	39.9-50.9	165	62.8	52.9-73.9	116	32.8	27.0-39.4
2000	311	46.8	41.6-52.5	201	71.1	60.9-82.4	110	29.6	24.2-35.7
2001	319	47.3	42.1-52.9	189	67.9	57.8-79.0	130	33.5	27.9-39.8
2002	298	43.8	38.8-49.2	172	60.8	51.5-71.3	126	32.2	26.7-38.3
2003	343	47.5	42.4-52.9	195	64.2	54.9-74.5	148	35.9	30.3-42.3
2004	376	50.0	44.9-55.5	228	71.3	61.7-81.8	148	34.6	29.2-40.8
2005	389	52.6	47.3-58.2	230	76.1	65.7-87.6	159	37.5	31.8-43.9
2006	359	48.2	43.2-53.6	209	67.3	57.8-77.7	150	35.5	30.0-41.7
2007	378	48.2	43.3-53.4	219	66.8	57.7-76.9	159	35.1	29.8-41.1
2008	385	47.8	43.0-53.0	220	66.1	57.0-76.0	165	35.5	30.2-41.4
2009	416	49.4	44.6-54.5	246	69.8	60.7-79.7	170	35.5	30.3-41.3
2010	440	50.9	46.1-56.0	241	66.1	57.5-75.6	199	40.4	34.9-46.5
2011	414	46.0	41.5-50.8	236	61.6	53.4-70.5	178	35.1	30.0-40.7
2012	465	50.6	45.9-55.5	274	70.3	61.7-79.7	191	36.5	31.4-42.1
2013	436	44.9	40.7-49.5	238	59.2	51.4-67.6	198	35.1	30.3-40.4
2014	433	44.1	39.9-48.6	238	56.4	49.1-64.6	195	34.9	30.1-40.3
2015	476	47.9	43.6-52.5	274	67.8	59.6-76.8	202	34.4	29.8-39.6
2016	455	44.7	40.6-49.2	258	60.1	52.6-68.3	197	33.8	29.1-38.9
2017	411	38.5	34.8-42.6	228	52.3	45.4-59.9	183	29.7	25.5-34.4
2018	407	37.3	33.7-41.3	212	47.1	40.6-54.2	195	30.9	26.6-35.6
2019	410	36.0	32.5-39.7	227	46.8	40.6-53.6	183	28.4	24.4-32.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Female Breast, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	4333	46.5	45.1-47.9
1977	4326	46.7	45.3-48.1
1978	4554	48.8	47.4-50.3
1979	4366	46.8	45.3-48.2
1980	4545	48.4	47.0-49.9
1981	4630	49.4	48.0-50.9
1982	4681	49.6	48.2-51.1
1983	4814	50.7	49.3-52.2
1984	4769	49.9	48.4-51.3
1985	4852	50.7	49.2-52.2
1986	4745	49.5	48.0-50.9
1987	4692	48.5	47.1-49.9
1988	4466	46.1	44.7-47.5
1989	4054	41.9	40.6-43.2
1990	4164	42.8	41.5-44.1
1991	4183	42.7	41.4-44.0
1992	4029	41.2	39.9-42.5
1993	3874	39.3	38.1-40.6
1994	4157	41.8	40.5-43.1
1995	4265	42.5	41.2-43.8
1996	4432	43.9	42.6-45.2
1997	4423	43.5	42.2-44.8
1998	4385	42.7	41.4-43.9
1999	4367	41.9	40.6-43.1
2000	4555	43.2	41.9-44.5
2001	4369	40.9	39.7-42.2
2002	4543	42.1	40.9-43.3
2003	4771	43.9	42.6-45.1
2004	5003	45.7	44.4-47.0
2005	4970	44.9	43.7-46.2
2006	5093	45.9	44.7-47.2
2007	5091	45.5	44.2-46.7
2008	5088	45.3	44.0-46.5
2009	5202	45.7	44.4-47.0
2010	4908	42.9	41.7-44.2
2011	5166	44.7	43.4-45.9
2012	4850	41.5	40.3-42.7
2013	5075	43.1	41.9-44.4
2014	5284	44.3	43.1-45.6
2015	5200	43.7	42.5-45.0
2016	5033	42.0	40.8-43.2
2017	5019	42.0	40.8-43.3
2018	4950	41.0	39.8-42.2
2019	5013	41.1	40.0-42.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Female Breast, Whites**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	3902	47.0	45.5-48.6
1977	3871	47.2	45.7-48.8
1978	4078	49.5	47.9-51.1
1979	3906	47.5	46.0-49.1
1980	4018	48.9	47.3-50.4
1981	4104	50.0	48.5-51.7
1982	4094	50.0	48.4-51.6
1983	4243	51.9	50.3-53.6
1984	4171	50.7	49.1-52.3
1985	4249	51.9	50.3-53.5
1986	4115	50.3	48.8-52.0
1987	4094	50.0	48.4-51.6
1988	3839	47.0	45.4-48.5
1989	3452	42.5	41.0-44.0
1990	3536	43.5	42.1-45.0
1991	3516	43.2	41.8-44.7
1992	3372	41.9	40.4-43.4
1993	3228	40.0	38.6-41.4
1994	3446	42.3	40.9-43.8
1995	3519	43.1	41.6-44.5
1996	3638	44.7	43.2-46.2
1997	3611	44.1	42.6-45.6
1998	3554	43.2	41.8-44.7
1999	3570	43.0	41.6-44.5
2000	3692	44.3	42.8-45.7
2001	3590	42.8	41.4-44.3
2002	3630	42.9	41.5-44.3
2003	3831	45.2	43.8-46.7
2004	3951	46.3	44.9-47.8
2005	3894	45.4	44.0-46.9
2006	3978	46.5	45.0-48.0
2007	3991	46.6	45.2-48.1
2008	3920	45.7	44.2-47.2
2009	4030	46.5	45.1-48.0
2010	3761	43.6	42.2-45.1
2011	3866	44.3	42.9-45.8
2012	3651	41.7	40.3-43.1
2013	3789	43.2	41.8-44.7
2014	3920	44.3	42.9-45.8
2015	3770	42.9	41.5-44.4
2016	3620	41.2	39.8-42.6
2017	3595	41.2	39.8-42.6
2018	3560	40.2	38.8-41.6
2019	3517	39.5	38.2-41.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Female Breast, Blacks**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	382	40.8	36.6-45.4
1977	394	41.2	37.0-45.6
1978	439	44.9	40.6-49.4
1979	421	42.0	37.9-46.4
1980	479	45.6	41.5-50.0
1981	489	46.2	42.1-50.7
1982	522	48.1	43.9-52.6
1983	511	45.4	41.4-49.6
1984	533	45.7	41.8-49.9
1985	527	44.8	40.9-48.9
1986	549	44.4	40.7-48.4
1987	536	42.4	38.8-46.2
1988	570	44.3	40.7-48.2
1989	547	41.8	38.3-45.5
1990	541	40.1	36.7-43.7
1991	572	42.0	38.6-45.7
1992	558	40.0	36.7-43.5
1993	566	39.6	36.4-43.1
1994	591	40.4	37.2-43.9
1995	633	42.5	39.3-46.0
1996	662	44.1	40.7-47.6
1997	687	44.1	40.8-47.5
1998	689	43.6	40.3-47.0
1999	657	40.4	37.3-43.6
2000	694	41.6	38.5-44.8
2001	607	36.1	33.2-39.1
2002	750	43.3	40.3-46.6
2003	758	43.2	40.2-46.4
2004	840	47.5	44.3-50.9
2005	865	48.2	45.0-51.5
2006	883	48.5	45.3-51.8
2007	864	46.9	43.8-50.1
2008	931	50.1	46.9-53.5
2009	907	48.5	45.4-51.8
2010	873	45.6	42.6-48.8
2011	970	49.7	46.6-53.0
2012	926	46.7	43.7-49.8
2013	978	48.6	45.6-51.8
2014	1029	50.1	47.0-53.3
2015	1034	49.9	46.9-53.1
2016	1028	48.9	45.9-52.0
2017	1025	48.6	45.6-51.7
2018	998	46.9	44.0-50.0
2019	1063	49.6	46.6-52.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Female Breast, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	2012	48.8	46.7-51.1
1977	1938	48.0	45.8-50.2
1978	1945	48.2	46.0-50.5
1979	1911	47.7	45.5-50.0
1980	1920	47.9	45.7-50.2
1981	2021	50.7	48.4-53.0
1982	2010	50.4	48.2-52.7
1983	1986	49.2	47.0-51.5
1984	2056	51.3	49.1-53.7
1985	1989	49.3	47.1-51.6
1986	1917	47.5	45.4-49.8
1987	1899	46.7	44.6-48.9
1988	1781	44.3	42.2-46.5
1989	1716	43.1	41.0-45.2
1990	1702	42.5	40.5-44.6
1991	1849	46.4	44.3-48.6
1992	1699	42.9	40.9-45.1
1993	1594	39.9	38.0-41.9
1994	1759	43.6	41.6-45.7
1995	1796	44.2	42.2-46.3
1996	1863	45.7	43.6-47.9
1997	1743	42.3	40.3-44.4
1998	1809	43.3	41.4-45.4
1999	1776	41.8	39.8-43.8
2000	1897	44.1	42.2-46.2
2001	1747	40.0	38.2-41.9
2002	1809	41.0	39.1-42.9
2003	1865	42.1	40.2-44.0
2004	1987	44.7	42.8-46.7
2005	1919	42.8	40.9-44.8
2006	2010	44.9	43.0-47.0
2007	2012	44.6	42.6-46.6
2008	2004	44.0	42.1-46.0
2009	2049	44.6	42.7-46.6
2010	1947	42.2	40.3-44.2
2011	2129	45.2	43.3-47.2
2012	1968	41.2	39.4-43.1
2013	2065	42.9	41.0-44.8
2014	2172	44.4	42.5-46.4
2015	2160	43.9	42.0-45.8
2016	2107	42.4	40.5-44.3
2017	2111	42.6	40.8-44.5
2018	2088	41.8	40.0-43.7
2019	2116	41.8	40.0-43.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Female Breast, Whites**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	1678	50.9	48.4-53.5
1977	1590	49.8	47.2-52.5
1978	1574	49.2	46.6-51.8
1979	1555	49.9	47.3-52.6
1980	1519	49.1	46.5-51.7
1981	1611	52.3	49.6-55.1
1982	1561	52.0	49.3-54.9
1983	1553	51.6	49.0-54.4
1984	1584	53.5	50.8-56.4
1985	1513	51.1	48.4-53.9
1986	1435	49.2	46.6-52.0
1987	1439	49.6	47.0-52.4
1988	1315	46.6	44.0-49.3
1989	1253	45.1	42.6-47.8
1990	1244	45.2	42.6-47.9
1991	1333	49.5	46.8-52.3
1992	1199	45.7	43.0-48.4
1993	1118	42.7	40.1-45.3
1994	1227	46.1	43.5-48.8
1995	1229	46.7	44.1-49.5
1996	1238	47.6	44.9-50.4
1997	1136	43.4	40.9-46.1
1998	1171	44.8	42.2-47.5
1999	1177	44.6	42.1-47.3
2000	1229	46.8	44.2-49.5
2001	1159	44.0	41.5-46.6
2002	1148	43.1	40.6-45.7
2003	1172	44.6	42.0-47.2
2004	1239	47.4	44.7-50.1
2005	1157	44.1	41.6-46.8
2006	1199	46.4	43.8-49.2
2007	1217	47.6	44.9-50.4
2008	1167	45.1	42.5-47.8
2009	1210	47.0	44.4-49.8
2010	1132	44.3	41.7-47.0
2011	1194	45.7	43.0-48.4
2012	1086	41.8	39.3-44.5
2013	1140	43.4	40.9-46.1
2014	1191	45.0	42.4-47.7
2015	1123	42.3	39.8-45.0
2016	1107	41.9	39.4-44.6
2017	1112	42.3	39.8-45.0
2018	1044	39.1	36.6-41.6
2019	1039	38.5	36.1-41.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Female Breast, Blacks**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	306	41.2	36.5-46.4
1977	306	40.1	35.5-45.0
1978	346	44.9	40.1-50.1
1979	336	42.4	37.8-47.4
1980	369	44.8	40.2-49.8
1981	383	46.1	41.5-51.2
1982	419	49.5	44.7-54.7
1983	396	45.0	40.6-49.9
1984	425	46.9	42.4-51.7
1985	419	45.4	41.0-50.1
1986	421	43.5	39.4-48.0
1987	421	42.3	38.3-46.6
1988	433	43.3	39.2-47.7
1989	426	41.7	37.8-46.0
1990	404	38.4	34.7-42.4
1991	448	42.7	38.8-46.9
1992	429	40.1	36.3-44.1
1993	416	37.7	34.1-41.5
1994	446	40.0	36.3-43.9
1995	492	43.3	39.5-47.4
1996	525	46.1	42.2-50.2
1997	513	43.3	39.6-47.2
1998	529	44.2	40.5-48.2
1999	485	39.3	35.9-43.0
2000	530	42.1	38.6-45.9
2001	450	35.6	32.3-39.0
2002	539	41.7	38.2-45.4
2003	557	42.5	39.0-46.2
2004	592	44.9	41.3-48.7
2005	607	45.2	41.7-49.0
2006	636	47.1	43.5-50.9
2007	620	45.6	42.0-49.3
2008	668	49.0	45.3-52.9
2009	645	47.1	43.5-50.9
2010	621	44.5	41.1-48.2
2011	698	49.1	45.5-53.0
2012	685	47.1	43.6-50.8
2013	709	48.3	44.8-52.1
2014	731	49.4	45.8-53.2
2015	741	49.6	46.1-53.4
2016	725	48.0	44.5-51.8
2017	706	46.8	43.3-50.5
2018	745	49.4	45.8-53.2
2019	767	50.3	46.7-54.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Female Breast, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	2320	44.7	42.8-46.5
1977	2388	45.8	44.0-47.7
1978	2609	49.4	47.5-51.4
1979	2454	46.1	44.3-48.0
1980	2625	48.7	46.9-50.7
1981	2608	48.5	46.6-50.5
1982	2670	49.1	47.2-51.0
1983	2827	51.8	49.9-53.8
1984	2711	48.8	47.0-50.8
1985	2859	51.7	49.7-53.6
1986	2826	50.9	49.0-52.8
1987	2793	49.8	47.9-51.7
1988	2685	47.3	45.5-49.2
1989	2337	41.0	39.3-42.7
1990	2459	42.9	41.2-44.7
1991	2334	40.1	38.5-41.8
1992	2328	40.0	38.4-41.7
1993	2279	38.9	37.3-40.5
1994	2398	40.6	39.0-42.3
1995	2469	41.3	39.7-43.0
1996	2569	42.8	41.1-44.5
1997	2680	44.3	42.6-46.0
1998	2575	42.1	40.5-43.8
1999	2590	41.9	40.3-43.6
2000	2657	42.5	40.9-44.1
2001	2621	41.5	39.9-43.1
2002	2734	42.9	41.3-44.5
2003	2905	45.1	43.5-46.8
2004	3015	46.3	44.6-48.0
2005	3051	46.3	44.7-48.0
2006	3083	46.6	44.9-48.3
2007	3079	46.0	44.4-47.7
2008	3084	46.1	44.4-47.8
2009	3153	46.4	44.7-48.1
2010	2961	43.4	41.8-45.0
2011	3037	44.2	42.6-45.9
2012	2882	41.6	40.0-43.2
2013	3010	43.3	41.7-44.9
2014	3112	44.1	42.5-45.8
2015	3040	43.6	42.0-45.2
2016	2926	41.7	40.2-43.4
2017	2908	41.6	40.0-43.2
2018	2861	40.4	38.8-42.0
2019	2897	40.8	39.2-42.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Female Breast, Whites**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	2224	44.7	42.8-46.7
1977	2281	45.8	43.9-47.8
1978	2504	49.7	47.7-51.7
1979	2350	46.2	44.3-48.2
1980	2499	48.8	46.8-50.8
1981	2493	48.7	46.8-50.7
1982	2532	49.0	47.1-51.0
1983	2689	52.0	50.1-54.1
1984	2585	49.3	47.3-51.3
1985	2733	52.3	50.3-54.3
1986	2679	51.1	49.1-53.1
1987	2655	50.2	48.3-52.2
1988	2524	47.3	45.4-49.2
1989	2198	41.1	39.4-42.9
1990	2289	42.6	40.8-44.4
1991	2183	40.2	38.5-41.9
1992	2172	40.1	38.4-41.8
1993	2109	38.7	37.0-40.4
1994	2219	40.5	38.8-42.2
1995	2290	41.4	39.7-43.1
1996	2400	43.3	41.6-45.1
1997	2475	44.4	42.6-46.2
1998	2382	42.4	40.7-44.2
1999	2393	42.3	40.6-44.0
2000	2463	43.1	41.4-44.8
2001	2430	42.3	40.6-44.0
2002	2482	42.7	41.1-44.5
2003	2659	45.6	43.8-47.4
2004	2711	45.9	44.2-47.7
2005	2737	46.0	44.2-47.8
2006	2779	46.6	44.8-48.4
2007	2774	46.2	44.5-48.0
2008	2753	45.9	44.2-47.7
2009	2820	46.3	44.6-48.1
2010	2629	43.3	41.6-45.0
2011	2672	43.6	41.9-45.4
2012	2565	41.6	40.0-43.3
2013	2649	43.1	41.4-44.8
2014	2729	44.0	42.3-45.8
2015	2647	43.2	41.5-45.0
2016	2513	40.9	39.2-42.6
2017	2483	40.8	39.1-42.6
2018	2516	40.7	39.1-42.5
2019	2478	40.1	38.4-41.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Female Breast, Blacks**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	75	39.0	30.2-49.4
1977	88	45.1	35.6-56.2
1978	93	45.1	36.1-55.7
1979	85	40.4	31.9-50.4
1980	110	48.7	39.8-59.1
1981	106	46.3	37.6-56.4
1982	103	42.9	34.8-52.2
1983	115	46.2	37.9-55.7
1984	108	41.8	34.0-50.7
1985	108	42.5	34.6-51.6
1986	127	47.1	39.1-56.2
1987	115	42.6	35.0-51.3
1988	137	48.3	40.4-57.2
1989	121	41.6	34.3-50.0
1990	137	46.0	38.5-54.6
1991	124	39.8	33.0-47.6
1992	128	39.6	32.9-47.1
1993	150	46.2	39.0-54.3
1994	145	41.7	35.1-49.2
1995	141	39.8	33.4-47.0
1996	137	37.4	31.3-44.3
1997	174	46.7	39.9-54.3
1998	160	41.5	35.2-48.5
1999	172	43.8	37.4-50.9
2000	164	40.1	34.1-46.8
2001	157	37.3	31.6-43.7
2002	211	47.9	41.6-54.9
2003	200	45.1	39.0-51.9
2004	248	55.3	48.6-62.8
2005	258	56.7	49.9-64.2
2006	247	52.4	46.0-59.4
2007	244	50.4	44.2-57.2
2008	263	53.3	47.0-60.2
2009	262	52.5	46.3-59.4
2010	252	48.8	42.9-55.3
2011	272	51.2	45.2-57.8
2012	241	45.3	39.7-51.4
2013	269	49.8	43.9-56.2
2014	298	51.6	45.8-57.9
2015	293	51.0	45.2-57.3
2016	303	51.0	45.3-57.2
2017	319	53.7	47.8-60.0
2018	252	40.9	35.9-46.4
2019	296	48.2	42.8-54.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Cervix, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	483	5.3	4.8-5.8
1977	461	5.0	4.6-5.5
1978	447	5.0	4.5-5.5
1979	445	4.9	4.4-5.4
1980	471	5.1	4.6-5.6
1981	438	4.7	4.3-5.2
1982	454	5.0	4.5-5.5
1983	424	4.6	4.2-5.1
1984	432	4.8	4.3-5.3
1985	395	4.2	3.8-4.6
1986	406	4.4	3.9-4.8
1987	448	4.7	4.3-5.2
1988	396	4.2	3.8-4.7
1989	406	4.2	3.8-4.6
1990	443	4.7	4.3-5.2
1991	414	4.4	4.0-4.8
1992	412	4.2	3.8-4.6
1993	396	4.1	3.7-4.5
1994	439	4.5	4.0-4.9
1995	430	4.3	3.9-4.8
1996	453	4.5	4.1-5.0
1997	421	4.2	3.8-4.6
1998	379	3.7	3.4-4.1
1999	389	3.8	3.4-4.2
2000	390	3.8	3.4-4.2
2001	376	3.6	3.2-4.0
2002	374	3.5	3.2-3.9
2003	379	3.5	3.2-3.9
2004	395	3.7	3.3-4.1
2005	426	4.0	3.6-4.4
2006	437	4.0	3.6-4.4
2007	442	4.0	3.7-4.4
2008	435	4.0	3.6-4.4
2009	477	4.3	3.9-4.7
2010	437	4.0	3.6-4.4
2011	423	3.7	3.4-4.1
2012	438	3.9	3.5-4.2
2013	450	4.0	3.6-4.3
2014	461	4.0	3.6-4.4
2015	442	3.8	3.4-4.2
2016	400	3.5	3.1-3.8
2017	449	3.9	3.5-4.3
2018	397	3.4	3.0-3.7
2019	407	3.6	3.2-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Cervix, Whites**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	353	4.3	3.8-4.8
1977	318	3.9	3.5-4.4
1978	313	4.0	3.5-4.4
1979	297	3.8	3.4-4.3
1980	340	4.2	3.8-4.7
1981	325	4.1	3.6-4.6
1982	312	4.1	3.6-4.6
1983	309	3.9	3.5-4.4
1984	299	3.9	3.5-4.4
1985	285	3.6	3.2-4.0
1986	268	3.4	3.0-3.8
1987	320	4.0	3.6-4.5
1988	288	3.7	3.3-4.2
1989	287	3.6	3.2-4.1
1990	295	3.8	3.4-4.3
1991	290	3.8	3.3-4.2
1992	274	3.4	3.0-3.9
1993	257	3.3	2.9-3.7
1994	302	3.8	3.4-4.3
1995	309	4.0	3.5-4.4
1996	313	4.0	3.5-4.4
1997	292	3.7	3.3-4.2
1998	263	3.3	2.9-3.7
1999	272	3.4	3.0-3.9
2000	265	3.3	2.9-3.7
2001	245	3.0	2.6-3.4
2002	229	2.8	2.5-3.2
2003	256	3.1	2.8-3.5
2004	249	3.0	2.7-3.4
2005	269	3.3	2.9-3.7
2006	282	3.4	3.0-3.9
2007	284	3.5	3.1-3.9
2008	299	3.7	3.3-4.2
2009	322	3.9	3.5-4.4
2010	268	3.3	2.9-3.7
2011	271	3.3	2.9-3.7
2012	283	3.4	3.0-3.9
2013	279	3.4	3.0-3.9
2014	296	3.5	3.1-3.9
2015	282	3.4	3.0-3.9
2016	242	2.9	2.5-3.3
2017	268	3.3	2.9-3.7
2018	254	3.1	2.7-3.5
2019	234	2.9	2.6-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Cervix, Blacks**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	120	12.8	10.4-15.4
1977	128	13.6	11.3-16.3
1978	121	12.6	10.4-15.2
1979	138	13.5	11.3-16.1
1980	122	12.0	9.9-14.5
1981	101	8.9	7.2-10.9
1982	126	11.4	9.4-13.6
1983	103	9.1	7.4-11.1
1984	116	10.1	8.3-12.1
1985	99	8.4	6.8-10.3
1986	123	10.4	8.6-12.4
1987	109	8.6	7.0-10.4
1988	96	7.5	6.0-9.2
1989	107	8.0	6.5-9.7
1990	133	9.5	7.9-11.2
1991	112	8.2	6.7-9.9
1992	117	8.1	6.7-9.7
1993	107	7.5	6.2-9.1
1994	105	7.3	5.9-8.8
1995	101	6.8	5.5-8.2
1996	114	7.4	6.1-8.9
1997	102	6.7	5.4-8.1
1998	91	5.6	4.5-6.9
1999	94	5.8	4.7-7.1
2000	106	6.5	5.3-7.8
2001	110	6.5	5.3-7.8
2002	116	6.8	5.6-8.2
2003	105	6.0	4.9-7.3
2004	116	6.6	5.4-7.9
2005	128	7.2	6.0-8.6
2006	118	6.6	5.4-7.9
2007	128	7.0	5.8-8.4
2008	112	6.0	4.9-7.2
2009	124	6.5	5.4-7.7
2010	131	6.9	5.8-8.2
2011	115	5.9	4.8-7.1
2012	117	6.0	5.0-7.2
2013	121	6.0	5.0-7.2
2014	118	5.8	4.8-7.0
2015	120	5.8	4.8-6.9
2016	122	5.8	4.8-7.0
2017	122	5.8	4.8-6.9
2018	104	4.8	3.9-5.9
2019	107	5.1	4.2-6.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Cervix, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	254	6.4	5.6-7.2
1977	257	6.6	5.8-7.5
1978	233	6.3	5.5-7.2
1979	248	6.5	5.7-7.3
1980	227	6.0	5.2-6.9
1981	245	6.4	5.6-7.3
1982	254	6.8	6.0-7.7
1983	211	5.4	4.7-6.2
1984	210	5.6	4.8-6.4
1985	192	4.8	4.1-5.6
1986	208	5.4	4.7-6.2
1987	231	5.9	5.1-6.7
1988	209	5.4	4.7-6.2
1989	200	5.0	4.3-5.7
1990	234	6.0	5.2-6.8
1991	218	5.7	5.0-6.6
1992	225	5.6	4.9-6.4
1993	218	5.6	4.9-6.4
1994	257	6.4	5.6-7.2
1995	224	5.6	4.9-6.3
1996	235	5.8	5.1-6.6
1997	228	5.5	4.8-6.3
1998	204	4.9	4.2-5.6
1999	207	4.9	4.2-5.6
2000	214	5.0	4.4-5.7
2001	201	4.6	4.0-5.3
2002	205	4.7	4.1-5.4
2003	203	4.6	4.0-5.3
2004	212	4.8	4.1-5.4
2005	225	5.1	4.4-5.8
2006	229	5.1	4.5-5.8
2007	222	4.9	4.3-5.7
2008	215	4.7	4.1-5.4
2009	230	5.1	4.4-5.8
2010	215	4.7	4.1-5.4
2011	221	4.8	4.1-5.4
2012	225	4.7	4.1-5.4
2013	214	4.4	3.8-5.0
2014	231	4.8	4.2-5.5
2015	220	4.5	3.9-5.1
2016	217	4.5	3.9-5.1
2017	246	5.1	4.4-5.8
2018	210	4.2	3.6-4.8
2019	216	4.4	3.9-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Cervix, Whites**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	150	4.6	3.9-5.5
1977	138	4.4	3.7-5.3
1978	123	4.4	3.6-5.3
1979	127	4.4	3.7-5.3
1980	125	4.5	3.7-5.5
1981	144	5.2	4.3-6.2
1982	136	5.2	4.3-6.2
1983	122	4.4	3.6-5.3
1984	104	3.9	3.1-4.8
1985	102	3.7	3.0-4.5
1986	91	3.3	2.6-4.1
1987	128	4.7	3.9-5.7
1988	116	4.4	3.6-5.4
1989	111	4.1	3.3-5.0
1990	119	4.7	3.8-5.6
1991	117	4.8	3.9-5.7
1992	112	4.3	3.5-5.2
1993	107	4.3	3.5-5.2
1994	145	5.8	4.9-6.8
1995	128	5.2	4.4-6.3
1996	128	5.1	4.3-6.1
1997	116	4.6	3.8-5.6
1998	107	4.3	3.5-5.2
1999	116	4.6	3.8-5.6
2000	110	4.3	3.6-5.2
2001	91	3.6	2.9-4.4
2002	89	3.6	2.9-4.4
2003	105	4.1	3.4-5.0
2004	94	3.6	2.9-4.4
2005	100	4.0	3.2-4.9
2006	102	4.0	3.3-4.9
2007	96	3.9	3.1-4.7
2008	107	4.2	3.5-5.1
2009	110	4.4	3.6-5.3
2010	85	3.5	2.8-4.3
2011	97	3.9	3.1-4.8
2012	105	4.1	3.3-5.0
2013	86	3.3	2.6-4.1
2014	102	4.0	3.2-4.9
2015	100	3.9	3.2-4.8
2016	92	3.6	2.9-4.5
2017	102	4.1	3.3-5.0
2018	99	3.7	3.0-4.5
2019	80	3.1	2.4-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Cervix, Blacks**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	97	12.9	10.4-15.9
1977	106	14.4	11.7-17.6
1978	102	13.3	10.7-16.3
1979	114	14.1	11.5-17.1
1980	95	11.8	9.5-14.6
1981	91	10.2	8.1-12.5
1982	104	12.2	9.9-14.8
1983	82	9.3	7.4-11.7
1984	93	10.4	8.3-12.8
1985	81	8.8	6.9-11.0
1986	104	11.3	9.2-13.7
1987	87	8.7	6.9-10.7
1988	83	8.3	6.6-10.3
1989	80	7.6	6.0-9.5
1990	103	9.4	7.7-11.5
1991	91	8.5	6.8-10.5
1992	95	8.7	7.0-10.6
1993	86	8.0	6.4-9.9
1994	90	8.2	6.6-10.1
1995	82	7.2	5.7-9.0
1996	85	7.2	5.7-8.9
1997	92	7.9	6.4-9.7
1998	74	6.0	4.7-7.6
1999	74	6.0	4.7-7.5
2000	89	7.1	5.7-8.8
2001	91	7.1	5.7-8.8
2002	94	7.3	5.9-9.0
2003	82	6.2	4.9-7.7
2004	91	6.9	5.5-8.4
2005	98	7.4	6.0-9.1
2006	96	7.2	5.8-8.8
2007	101	7.5	6.1-9.1
2008	88	6.4	5.2-8.0
2009	97	6.9	5.5-8.4
2010	100	7.3	5.9-8.9
2011	96	6.8	5.5-8.3
2012	90	6.4	5.1-7.9
2013	84	5.8	4.6-7.2
2014	91	6.3	5.1-7.8
2015	92	6.1	4.9-7.5
2016	92	6.2	4.9-7.6
2017	93	6.1	4.9-7.6
2018	79	5.2	4.1-6.5
2019	85	5.8	4.6-7.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Cervix, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	229	4.4	3.9-5.1
1977	204	3.9	3.4-4.5
1978	214	4.1	3.5-4.7
1979	197	3.8	3.3-4.4
1980	244	4.4	3.9-5.0
1981	193	3.5	3.0-4.1
1982	200	3.7	3.2-4.3
1983	213	4.0	3.5-4.6
1984	222	4.2	3.6-4.8
1985	202	3.7	3.2-4.2
1986	196	3.6	3.1-4.2
1987	217	3.8	3.3-4.4
1988	187	3.4	2.9-3.9
1989	206	3.7	3.2-4.2
1990	208	3.8	3.3-4.3
1991	196	3.4	3.0-4.0
1992	187	3.2	2.8-3.7
1993	177	3.1	2.6-3.6
1994	182	3.1	2.7-3.6
1995	206	3.5	3.1-4.1
1996	218	3.7	3.2-4.2
1997	193	3.3	2.8-3.8
1998	175	2.9	2.5-3.4
1999	182	3.0	2.6-3.5
2000	176	2.9	2.5-3.4
2001	175	2.8	2.4-3.3
2002	169	2.7	2.3-3.2
2003	176	2.8	2.4-3.3
2004	182	2.9	2.5-3.4
2005	201	3.2	2.8-3.7
2006	208	3.3	2.8-3.7
2007	220	3.4	3.0-3.9
2008	220	3.5	3.0-4.0
2009	247	3.8	3.3-4.3
2010	222	3.5	3.1-4.0
2011	202	3.1	2.7-3.6
2012	213	3.3	2.8-3.8
2013	236	3.7	3.2-4.2
2014	230	3.4	3.0-3.9
2015	222	3.4	2.9-3.8
2016	183	2.8	2.4-3.2
2017	203	3.0	2.6-3.5
2018	187	2.8	2.4-3.3
2019	191	3.0	2.6-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Cervix, Whites**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	203	4.1	3.5-4.7
1977	180	3.6	3.1-4.2
1978	190	3.8	3.3-4.4
1979	170	3.4	2.9-4.0
1980	215	4.1	3.6-4.7
1981	181	3.5	3.0-4.1
1982	176	3.5	3.0-4.1
1983	187	3.7	3.2-4.3
1984	195	3.9	3.4-4.5
1985	182	3.5	3.0-4.1
1986	176	3.4	2.9-4.0
1987	192	3.6	3.1-4.2
1988	172	3.3	2.8-3.9
1989	176	3.4	2.9-3.9
1990	176	3.4	2.9-4.0
1991	173	3.3	2.8-3.8
1992	162	3.0	2.6-3.6
1993	149	2.8	2.4-3.3
1994	157	2.9	2.5-3.4
1995	181	3.4	2.9-3.9
1996	185	3.4	2.9-4.0
1997	176	3.3	2.8-3.8
1998	156	2.9	2.4-3.4
1999	156	2.9	2.5-3.4
2000	155	2.8	2.4-3.3
2001	154	2.7	2.3-3.2
2002	140	2.5	2.1-3.0
2003	151	2.7	2.3-3.2
2004	155	2.8	2.4-3.3
2005	169	3.0	2.6-3.5
2006	180	3.2	2.7-3.7
2007	188	3.3	2.8-3.8
2008	192	3.5	3.0-4.0
2009	212	3.7	3.2-4.2
2010	183	3.3	2.8-3.8
2011	174	3.0	2.6-3.6
2012	178	3.1	2.7-3.6
2013	193	3.5	3.0-4.1
2014	194	3.3	2.8-3.9
2015	182	3.2	2.7-3.7
2016	150	2.6	2.2-3.1
2017	166	2.9	2.4-3.4
2018	155	2.8	2.3-3.3
2019	154	2.9	2.5-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Cervix, Blacks**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	23	12.1	7.3-18.8
1977	22	11.0	6.6-17.0
1978	19	10.4	6.0-16.4
1979	24	11.3	7.0-17.2
1980	27	13.0	8.4-19.1
1981	10	4.5	2.1-8.4
1982	22	8.6	5.3-13.0
1983	21	8.1	5.0-12.5
1984	23	9.0	5.6-13.5
1985	18	7.0	4.0-11.1
1986	18	6.5	3.8-10.4
1987	22	8.3	5.1-12.6
1988	13	4.7	2.4-8.0
1989	27	9.5	6.2-13.8
1990	29	9.1	6.0-13.2
1991	21	7.2	4.4-11.1
1992	22	6.0	3.7-9.2
1993	21	6.0	3.6-9.2
1994	15	4.3	2.4-7.1
1995	19	5.2	3.1-8.2
1996	29	8.1	5.4-11.6
1997	10	2.7	1.3-5.0
1998	17	4.3	2.5-6.9
1999	20	5.5	3.3-8.4
2000	17	4.3	2.5-6.9
2001	19	4.6	2.7-7.2
2002	22	5.0	3.1-7.6
2003	23	5.6	3.5-8.4
2004	24	5.4	3.5-8.1
2005	30	6.7	4.5-9.6
2006	22	4.8	3.0-7.2
2007	27	5.7	3.8-8.3
2008	24	4.7	3.0-7.0
2009	27	5.5	3.6-8.1
2010	31	5.9	4.0-8.3
2011	19	3.5	2.1-5.5
2012	27	5.0	3.3-7.3
2013	37	6.7	4.7-9.3
2014	27	4.6	3.0-6.7
2015	28	4.9	3.3-7.2
2016	30	5.0	3.3-7.2
2017	29	4.9	3.3-7.1
2018	25	4.1	2.6-6.1
2019	22	3.4	2.1-5.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Prostate, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	1616	26.1	24.8-27.5
1977	1619	25.5	24.2-26.9
1978	1635	25.9	24.6-27.2
1979	1717	27.8	26.4-29.2
1980	1870	29.3	27.9-30.7
1981	1929	30.4	29.0-31.8
1982	1948	30.5	29.1-31.9
1983	1880	28.6	27.3-30.0
1984	1951	29.5	28.1-30.9
1985	2048	30.3	29.0-31.7
1986	2040	30.4	29.0-31.8
1987	2077	30.5	29.1-31.9
1988	2117	31.0	29.6-32.4
1989	2080	30.1	28.8-31.5
1990	2329	33.0	31.6-34.4
1991	2565	35.9	34.5-37.3
1992	2717	36.4	35.0-37.9
1993	2472	33.1	31.8-34.5
1994	2243	29.7	28.5-31.0
1995	2081	27.1	26.0-28.4
1996	1906	24.7	23.6-25.9
1997	1882	24.4	23.3-25.6
1998	1623	20.8	19.7-21.8
1999	1605	20.1	19.1-21.1
2000	1614	19.9	18.9-20.9
2001	1470	17.7	16.8-18.7
2002	1475	17.5	16.7-18.5
2003	1475	17.3	16.4-18.2
2004	1712	19.3	18.4-20.2
2005	1772	19.5	18.6-20.4
2006	1952	21.3	20.4-22.3
2007	2003	21.4	20.5-22.4
2008	2095	21.7	20.7-22.6
2009	2211	22.3	21.3-23.2
2010	2318	22.9	22.0-23.9
2011	2309	22.4	21.5-23.4
2012	2135	20.1	19.3-21.0
2013	2332	21.7	20.8-22.6
2014	2410	22.0	21.1-22.9
2015	2581	23.1	22.2-24.0
2016	2790	24.7	23.7-25.6
2017	2951	25.5	24.6-26.5
2018	3066	26.3	25.4-27.3
2019	3201	26.8	25.9-27.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Prostate, Whites**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	1384	24.5	23.2-25.9
1977	1360	23.6	22.3-24.9
1978	1355	23.6	22.3-24.9
1979	1434	25.6	24.2-27.0
1980	1507	26.2	24.8-27.6
1981	1595	27.8	26.4-29.3
1982	1625	28.2	26.7-29.6
1983	1559	26.4	25.1-27.8
1984	1609	27.2	25.8-28.6
1985	1654	27.5	26.2-28.9
1986	1646	27.6	26.2-29.0
1987	1695	28.1	26.7-29.5
1988	1723	28.4	27.0-29.8
1989	1696	27.8	26.5-29.2
1990	1896	30.6	29.2-32.1
1991	2100	33.5	32.1-35.0
1992	2222	34.1	32.7-35.6
1993	2006	31.0	29.6-32.4
1994	1706	26.1	24.9-27.4
1995	1566	23.9	22.7-25.1
1996	1483	22.6	21.5-23.8
1997	1398	21.4	20.3-22.6
1998	1204	18.2	17.2-19.3
1999	1194	17.8	16.8-18.9
2000	1196	17.8	16.8-18.8
2001	1104	16.0	15.1-17.0
2002	1105	15.9	15.0-16.9
2003	1135	16.1	15.2-17.1
2004	1308	18.0	17.1-19.1
2005	1384	18.7	17.7-19.7
2006	1504	20.3	19.2-21.3
2007	1542	20.4	19.3-21.4
2008	1619	20.9	19.9-21.9
2009	1682	21.2	20.1-22.2
2010	1795	22.2	21.2-23.3
2011	1744	21.3	20.3-22.3
2012	1611	19.2	18.3-20.2
2013	1719	20.3	19.3-21.3
2014	1778	20.6	19.7-21.6
2015	1903	21.7	20.8-22.8
2016	2075	23.4	22.4-24.5
2017	2209	24.7	23.6-25.7
2018	2288	25.3	24.3-26.4
2019	2318	25.2	24.2-26.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State, Prostate, Blacks**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	216	47.8	40.7-55.6
1977	246	51.5	44.4-59.2
1978	273	57.3	49.8-65.6
1979	275	59.0	51.4-67.2
1980	352	70.6	62.6-79.4
1981	320	63.3	55.8-71.5
1982	314	61.4	54.1-69.3
1983	307	56.2	49.4-63.6
1984	326	57.6	50.8-64.9
1985	370	61.2	54.5-68.5
1986	374	62.6	55.8-70.0
1987	366	58.1	51.8-65.0
1988	372	61.6	54.9-68.8
1989	375	58.3	52.0-65.1
1990	416	61.5	55.2-68.3
1991	427	61.6	55.3-68.3
1992	443	60.8	54.8-67.3
1993	422	58.0	52.1-64.3
1994	473	62.5	56.5-68.9
1995	463	57.5	52.0-63.4
1996	380	45.2	40.4-50.4
1997	440	52.7	47.5-58.3
1998	376	43.9	39.3-48.9
1999	384	42.6	38.1-47.4
2000	384	39.9	35.8-44.4
2001	321	33.6	29.8-37.7
2002	337	33.9	30.2-38.0
2003	302	29.0	25.6-32.7
2004	367	33.8	30.2-37.7
2005	331	29.7	26.4-33.3
2006	389	33.5	30.0-37.2
2007	406	34.8	31.3-38.5
2008	434	34.0	30.7-37.5
2009	444	34.7	31.4-38.3
2010	457	34.7	31.4-38.2
2011	465	34.2	31.0-37.6
2012	437	30.7	27.8-33.9
2013	514	35.8	32.6-39.2
2014	524	35.2	32.1-38.5
2015	526	34.6	31.6-37.8
2016	570	37.0	33.9-40.3
2017	575	35.3	32.3-38.4
2018	597	36.5	33.5-39.7
2019	663	39.6	36.5-42.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Prostate, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	730	26.4	24.4-28.5
1977	720	25.8	23.8-27.8
1978	753	27.4	25.4-29.5
1979	801	29.8	27.7-32.1
1980	861	31.6	29.4-33.9
1981	839	31.2	29.0-33.5
1982	870	32.2	30.0-34.5
1983	811	29.7	27.6-31.9
1984	867	31.5	29.4-33.7
1985	893	32.1	29.9-34.3
1986	863	31.5	29.3-33.7
1987	852	30.5	28.4-32.7
1988	814	30.3	28.2-32.5
1989	845	31.1	29.0-33.3
1990	921	34.0	31.7-36.3
1991	984	35.8	33.5-38.1
1992	962	34.5	32.3-36.8
1993	888	31.9	29.8-34.2
1994	917	32.4	30.3-34.6
1995	847	29.6	27.6-31.7
1996	721	25.0	23.2-26.9
1997	748	26.0	24.1-28.0
1998	643	22.0	20.3-23.8
1999	670	22.6	20.9-24.4
2000	641	20.9	19.3-22.6
2001	598	19.2	17.7-20.8
2002	595	18.9	17.4-20.5
2003	534	16.8	15.4-18.3
2004	613	18.4	16.9-19.9
2005	657	19.5	18.1-21.1
2006	716	21.1	19.6-22.8
2007	753	22.0	20.4-23.6
2008	723	20.3	18.8-21.9
2009	820	22.6	21.0-24.2
2010	823	22.2	20.7-23.8
2011	875	23.2	21.6-24.8
2012	802	20.4	19.0-21.9
2013	923	23.3	21.8-24.9
2014	918	22.6	21.1-24.1
2015	991	23.8	22.3-25.3
2016	1041	24.7	23.2-26.2
2017	1111	25.7	24.2-27.3
2018	1103	25.4	23.9-26.9
2019	1175	26.3	24.8-27.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Prostate, Whites**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	548	23.5	21.4-25.6
1977	527	22.4	20.4-24.5
1978	545	23.4	21.4-25.5
1979	573	25.5	23.4-27.8
1980	568	25.2	23.1-27.4
1981	581	26.2	24.0-28.5
1982	614	27.6	25.4-29.9
1983	576	26.0	23.9-28.3
1984	598	26.9	24.8-29.3
1985	581	26.3	24.1-28.6
1986	568	26.2	24.0-28.5
1987	554	25.2	23.2-27.5
1988	521	24.6	22.5-26.9
1989	539	25.7	23.6-28.0
1990	585	28.4	26.1-30.8
1991	639	30.7	28.4-33.3
1992	597	28.7	26.4-31.2
1993	555	27.1	24.9-29.5
1994	532	25.8	23.7-28.2
1995	462	22.7	20.7-24.9
1996	422	20.7	18.8-22.8
1997	410	20.4	18.5-22.5
1998	341	16.9	15.1-18.8
1999	362	17.9	16.1-19.8
2000	339	16.5	14.8-18.3
2001	330	15.8	14.1-17.6
2002	334	16.0	14.4-17.9
2003	310	14.7	13.1-16.5
2004	340	15.9	14.3-17.7
2005	377	17.5	15.8-19.4
2006	411	19.2	17.4-21.2
2007	435	20.1	18.2-22.1
2008	404	18.5	16.7-20.4
2009	463	20.7	18.9-22.7
2010	461	20.5	18.6-22.5
2011	476	20.9	19.0-22.9
2012	428	18.4	16.6-20.2
2013	480	20.4	18.6-22.3
2014	484	20.2	18.4-22.1
2015	500	20.6	18.8-22.5
2016	551	22.5	20.6-24.5
2017	579	23.6	21.7-25.6
2018	581	23.4	21.5-25.4
2019	581	23.0	21.2-25.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York City, Prostate, Blacks**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	171	48.3	40.2-57.2
1977	185	51.2	43.1-60.1
1978	205	57.7	49.0-67.3
1979	224	63.0	54.1-72.8
1980	285	76.7	67.0-87.3
1981	250	65.7	56.9-75.3
1982	252	66.0	57.3-75.5
1983	229	55.0	47.3-63.4
1984	255	59.3	51.5-67.9
1985	291	63.2	55.4-71.7
1986	276	60.9	53.2-69.2
1987	285	59.9	52.5-67.9
1988	277	62.1	54.3-70.5
1989	299	61.3	53.9-69.3
1990	325	63.6	56.2-71.5
1991	315	60.1	53.0-67.7
1992	329	61.3	54.3-68.9
1993	306	56.3	49.7-63.5
1994	348	61.9	55.1-69.3
1995	342	57.1	50.8-63.9
1996	264	43.0	37.6-48.9
1997	307	50.6	44.7-57.1
1998	264	42.0	36.7-47.7
1999	290	43.4	38.2-49.0
2000	275	38.9	34.2-44.1
2001	239	34.8	30.3-39.7
2002	237	32.1	27.9-36.6
2003	200	26.8	23.0-31.0
2004	247	31.3	27.4-35.7
2005	239	30.1	26.2-34.3
2006	264	32.5	28.5-36.9
2007	283	34.6	30.5-39.1
2008	292	32.7	28.9-36.9
2009	300	33.6	29.7-37.8
2010	313	34.4	30.5-38.6
2011	336	35.7	31.8-39.9
2012	311	31.1	27.6-34.9
2013	369	36.9	33.0-41.0
2014	358	34.6	31.0-38.6
2015	380	35.9	32.3-39.9
2016	387	36.4	32.7-40.3
2017	407	35.8	32.2-39.6
2018	401	35.2	31.7-39.0
2019	443	38.2	34.6-42.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Prostate, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	886	25.9	24.2-27.8
1977	899	25.4	23.7-27.2
1978	882	24.8	23.1-26.6
1979	915	26.2	24.4-28.0
1980	1009	27.6	25.9-29.5
1981	1090	29.8	28.0-31.8
1982	1078	29.3	27.4-31.2
1983	1069	27.8	26.1-29.7
1984	1084	28.1	26.3-29.9
1985	1155	29.2	27.4-31.0
1986	1177	29.7	27.9-31.5
1987	1224	30.5	28.8-32.4
1988	1303	31.4	29.6-33.2
1989	1235	29.5	27.8-31.2
1990	1408	32.4	30.6-34.2
1991	1581	35.9	34.1-37.8
1992	1755	37.5	35.7-39.4
1993	1584	33.8	32.1-35.6
1994	1326	28.1	26.6-29.7
1995	1234	25.7	24.2-27.2
1996	1185	24.6	23.2-26.1
1997	1134	23.5	22.1-24.9
1998	980	20.0	18.8-21.4
1999	934	18.6	17.4-19.8
2000	973	19.4	18.1-20.6
2001	872	16.8	15.7-18.0
2002	880	16.7	15.6-17.9
2003	940	17.6	16.4-18.7
2004	1099	19.8	18.7-21.1
2005	1113	19.4	18.3-20.6
2006	1236	21.4	20.2-22.7
2007	1250	21.1	20.0-22.4
2008	1372	22.4	21.3-23.7
2009	1391	22.1	21.0-23.4
2010	1495	23.3	22.1-24.6
2011	1434	22.0	20.9-23.2
2012	1333	20.0	18.9-21.2
2013	1409	20.8	19.7-21.9
2014	1492	21.7	20.6-22.8
2015	1590	22.7	21.6-23.9
2016	1749	24.6	23.5-25.8
2017	1840	25.4	24.2-26.6
2018	1963	26.9	25.7-28.1
2019	2026	27.2	26.0-28.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Prostate, Whites**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	836	25.3	23.5-27.2
1977	833	24.5	22.8-26.3
1978	810	23.7	22.0-25.5
1979	860	25.6	23.8-27.4
1980	939	26.9	25.1-28.8
1981	1014	28.9	27.1-30.8
1982	1011	28.6	26.8-30.6
1983	983	26.7	24.9-28.5
1984	1011	27.3	25.6-29.2
1985	1073	28.3	26.6-30.1
1986	1078	28.4	26.7-30.3
1987	1140	29.8	28.0-31.7
1988	1202	30.4	28.6-32.2
1989	1157	28.9	27.2-30.7
1990	1311	31.6	29.8-33.4
1991	1461	34.8	33.0-36.7
1992	1625	36.6	34.8-38.5
1993	1451	32.7	31.0-34.4
1994	1174	26.3	24.8-27.9
1995	1104	24.4	22.9-25.9
1996	1061	23.5	22.1-25.0
1997	988	21.9	20.5-23.3
1998	863	18.9	17.6-20.2
1999	831	17.7	16.5-18.9
2000	857	18.4	17.1-19.7
2001	774	16.1	15.0-17.3
2002	771	15.8	14.7-17.0
2003	825	16.7	15.6-17.9
2004	968	19.0	17.8-20.2
2005	1005	19.1	17.9-20.3
2006	1093	20.7	19.5-22.0
2007	1107	20.5	19.3-21.7
2008	1215	21.8	20.6-23.1
2009	1219	21.3	20.1-22.6
2010	1334	22.9	21.7-24.2
2011	1268	21.4	20.3-22.7
2012	1183	19.5	18.4-20.7
2013	1239	20.2	19.1-21.4
2014	1294	20.8	19.7-22.0
2015	1403	22.2	21.0-23.4
2016	1524	23.8	22.6-25.1
2017	1630	25.0	23.8-26.3
2018	1707	26.0	24.8-27.3
2019	1737	26.0	24.8-27.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2019
New York State Excl New York City, Prostate, Blacks**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	45	46.5	32.3-63.9
1977	61	52.5	38.6-69.2
1978	68	56.1	42.0-72.9
1979	51	45.8	32.3-62.2
1980	67	51.2	38.3-66.6
1981	70	55.9	42.2-72.1
1982	62	46.6	34.9-60.6
1983	78	60.1	46.0-76.6
1984	71	52.1	39.5-67.0
1985	79	54.7	42.0-69.6
1986	98	68.2	54.0-84.5
1987	81	52.3	40.5-66.0
1988	95	60.0	47.4-74.5
1989	76	49.0	37.6-62.4
1990	91	54.9	43.1-68.7
1991	112	66.2	53.3-80.9
1992	114	59.0	47.6-72.1
1993	116	62.8	50.8-76.6
1994	125	64.1	52.3-77.6
1995	121	58.9	47.8-71.5
1996	116	51.4	41.7-62.4
1997	133	58.4	48.1-70.0
1998	112	49.5	40.0-60.3
1999	94	40.5	32.0-50.4
2000	109	42.9	34.6-52.5
2001	82	29.9	23.3-37.6
2002	100	39.6	31.4-49.2
2003	102	34.9	27.9-43.0
2004	120	40.6	33.0-49.2
2005	92	28.6	22.5-35.7
2006	125	35.4	29.1-42.6
2007	123	34.7	28.4-41.9
2008	142	36.9	30.7-43.9
2009	144	37.1	30.9-44.2
2010	144	35.1	29.3-41.7
2011	129	30.1	24.9-36.1
2012	126	29.5	24.3-35.5
2013	145	33.0	27.6-39.2
2014	166	36.0	30.4-42.4
2015	146	30.8	25.7-36.6
2016	183	37.9	32.4-44.2
2017	168	33.9	28.7-39.8
2018	196	39.5	33.8-45.7
2019	220	41.9	36.3-48.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by County, All Races, 2015-2019
Colorectal**

County/Region	Males and Females			Males			Females		
	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI
New York State	5135.4	21.7	21.4-21.9	2663.0	25.1	24.6-25.5	2472.4	18.9	18.5-19.2
New York City	2011.6	21.3	20.9-21.7	1047.2	25.4	24.7-26.2	964.4	18.1	17.6-18.6
New York State Excl New York City	3123.8	21.9	21.6-22.3	1615.8	24.9	24.3-25.5	1508.0	19.4	18.9-19.9
Bronx	304.2	21.0	20.0-22.1	154.6	25.3	23.5-27.2	149.6	17.8	16.6-19.2
Kings (Brooklyn)	622.4	22.5	21.7-23.3	326.0	27.5	26.2-28.9	296.4	18.8	17.8-19.8
New York (Manhattan)	374.6	19.3	18.4-20.2	200.0	23.5	22.0-25.0	174.6	15.9	14.8-17.0
Queens	574.4	21.2	20.4-22.0	295.0	24.3	23.1-25.6	279.4	18.6	17.7-19.7
Richmond (Staten Island)	136.0	23.7	21.9-25.6	71.6	27.7	24.8-30.8	64.4	20.4	18.1-22.8
Albany	80.6	21.3	19.2-23.6	38.8	23.6	20.3-27.3	41.8	19.4	16.7-22.4
Allegany	19.2	33.3	26.6-41.2	10.6	37.7	27.9-50.1	8.6	29.2	20.6-40.4
Broome	50.0	19.3	16.9-22.0	25.2	21.5	17.8-25.8	24.8	17.6	14.5-21.3
Cattaraugus	25.4	25.9	21.4-31.2	12.8	27.7	20.9-35.9	12.6	24.1	18.1-31.4
Cayuga	21.0	20.1	16.3-24.6	10.4	21.3	15.7-28.3	10.6	18.2	13.4-24.4
Chautauqua	32.4	19.0	16.0-22.3	16.6	21.4	16.8-26.8	15.8	16.8	13.1-21.3
Chemung	20.0	18.7	15.0-23.0	14.0	27.4	21.2-34.9	6.0	11.7	7.6-17.3
Chenango	15.0	22.2	17.2-28.2	8.0	24.5	17.1-34.1	7.0	20.4	14.0-29.1
Clinton	23.0	23.8	19.5-28.7	12.8	27.5	21.0-35.4	10.2	19.9	14.6-26.7
Columbia	20.8	21.3	17.2-26.2	10.0	21.8	16.0-29.4	10.8	20.5	15.1-27.6
Cortland	13.8	24.6	19.0-31.4	7.2	29.2	20.2-40.9	6.6	20.6	14.0-29.6
Delaware	16.8	23.7	18.5-30.0	9.2	28.0	19.9-38.5	7.6	19.9	13.8-28.5
Dutchess	75.8	19.7	17.7-21.9	38.0	20.9	17.9-24.2	37.8	18.6	15.9-21.6
Erie	266.2	22.4	21.2-23.7	133.8	25.6	23.6-27.6	132.4	19.9	18.4-21.6
Essex	12.0	21.1	15.8-27.9	6.0	23.6	15.6-34.7	6.0	18.3	12.2-27.4
Franklin	14.2	22.6	17.5-28.7	5.4	16.8	10.9-24.8	8.8	26.5	19.0-36.4
Fulton	19.8	26.6	21.4-32.8	9.8	29.3	21.4-39.5	10.0	24.1	17.6-32.6
Genesee	18.8	24.9	19.8-30.8	11.4	32.5	24.3-42.6	7.4	17.9	12.2-25.6
Greene	18.6	26.2	20.9-32.6	10.8	30.7	22.7-40.9	7.8	22.7	15.7-32.2
Hamilton	2.2	19.3	9.5-45.0	1.2	20.9	7.3-70.0	1.0	17.2	5.6-64.8
Herkimer	24.0	27.5	22.6-33.3	12.4	30.4	23.0-39.7	11.6	23.9	18.0-31.5
Jefferson	32.2	28.6	24.2-33.5	14.8	28.2	22.0-35.6	17.4	29.5	23.4-36.7
Lewis	10.8	33.6	24.8-44.6	5.4	34.7	22.4-51.4	5.4	32.7	20.8-49.1
Livingston	19.4	23.8	19.2-29.3	11.8	31.0	23.4-40.5	7.6	18.2	12.7-25.5
Madison	21.6	24.3	19.7-29.7	11.8	29.3	22.0-38.4	9.8	19.6	14.3-26.4
Monroe	199.2	21.4	20.1-22.9	101.8	24.2	22.1-26.5	97.4	19.1	17.4-21.0
Montgomery	16.8	27.3	21.6-34.1	8.6	30.5	21.8-41.6	8.2	24.7	17.5-34.1
Nassau	379.2	21.5	20.5-22.5	188.4	23.9	22.4-25.5	190.8	19.4	18.2-20.8
Niagara	72.6	26.3	23.5-29.3	39.2	31.4	27.0-36.3	33.4	21.9	18.6-25.8
Oneida	58.6	19.3	17.1-21.8	30.0	21.7	18.3-25.7	28.6	17.5	14.6-20.9
Onondaga	120.4	21.1	19.4-22.9	59.0	23.1	20.5-26.0	61.4	19.4	17.2-21.8
Ontario	29.8	20.1	16.8-23.8	15.8	22.7	17.8-28.7	14.0	17.8	13.6-22.9
Orange	91.4	22.0	20.0-24.2	47.2	24.3	21.2-27.7	44.2	19.9	17.3-22.8
Orleans	16.2	30.4	23.8-38.3	8.8	34.0	24.3-46.6	7.4	26.6	18.4-37.6
Oswego	33.0	23.1	19.6-27.0	15.8	22.7	17.8-28.6	17.2	22.8	18.1-28.4
Otsego	22.6	28.5	23.1-34.9	14.2	37.9	29.0-48.7	8.4	20.1	14.0-28.3
Putnam	26.6	20.4	17.0-24.4	12.2	19.1	14.5-25.0	14.4	20.5	15.9-26.2
Rensselaer	41.8	22.0	19.0-25.3	21.8	25.9	21.2-31.5	20.0	18.7	15.1-23.0
Rockland	81.6	21.5	19.4-23.8	44.6	25.9	22.5-29.6	37.0	17.9	15.3-20.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by County, All Races, 2015-2019
Colorectal**

County/Region	Males and Females			Males			Females		
	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI
St. Lawrence	33.2	24.7	21.0-28.9	17.8	28.2	22.5-35.0	15.4	21.4	16.7-27.0
Saratoga	65.2	22.0	19.6-24.6	34.0	25.2	21.4-29.4	31.2	19.1	16.1-22.5
Schenectady	38.0	19.9	17.1-23.0	20.2	23.4	18.9-28.6	17.8	16.7	13.3-20.7
Schoharie	9.0	19.1	13.7-26.3	5.0	22.0	13.9-33.8	4.0	16.1	9.5-26.4
Schuyler	6.6	28.7	19.1-41.6	4.0	35.2	20.6-56.6	2.6	22.7	11.3-41.3
Seneca	10.4	23.4	17.2-31.2	5.6	24.6	16.1-36.4	4.8	22.3	13.8-34.3
Steuben	34.4	25.8	21.9-30.2	17.4	27.5	21.8-34.4	17.0	24.0	18.9-30.1
Suffolk	425.6	22.6	21.6-23.6	230.6	26.9	25.3-28.5	195.0	19.1	17.9-20.4
Sullivan	24.2	25.0	20.5-30.1	12.6	26.0	19.7-33.7	11.6	23.5	17.5-31.0
Tioga	15.8	22.1	17.3-28.0	8.6	25.2	18.0-34.7	7.2	19.7	13.5-28.1
Tompkins	15.6	14.6	11.5-18.4	8.4	17.3	12.3-23.7	7.2	12.2	8.4-17.1
Ulster	58.8	24.2	21.4-27.3	28.8	25.2	21.1-29.9	30.0	23.4	19.6-27.8
Warren	19.0	20.5	16.3-25.6	9.2	20.8	14.9-28.5	9.8	19.8	14.2-27.1
Washington	19.8	24.3	19.5-29.9	10.8	27.1	20.1-35.9	9.0	21.4	15.2-29.4
Wayne	30.8	25.6	21.6-30.3	16.6	28.4	22.4-35.6	14.2	23.2	17.8-29.7
Westchester	237.2	19.2	18.1-20.3	121.0	22.1	20.4-24.0	116.2	16.8	15.4-18.3
Wyoming	12.2	24.6	18.6-31.9	7.4	29.9	20.8-41.8	4.8	19.6	12.2-30.2
Yates	4.6	12.6	7.6-19.9	2.2	12.8	6.1-24.4	2.4	12.7	5.9-24.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by County, All Races, 2015-2019
Lung and Bronchus**

County/Region	Males and Females			Males			Females		
	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI
New York State	9256.0	37.4	37.1-37.8	4705.8	43.2	42.6-43.7	4550.2	33.2	32.7-33.6
New York City	2849.8	29.5	29.0-30.0	1502.8	36.5	35.7-37.3	1347.0	24.5	23.9-25.1
New York State Excl New York City	6405.6	42.5	42.0-42.9	3202.6	47.2	46.5-48.0	3203.0	39.0	38.4-39.7
Bronx	465.6	31.5	30.2-32.8	247.4	40.8	38.5-43.2	218.2	25.1	23.6-26.6
Kings (Brooklyn)	814.4	29.2	28.3-30.1	445.2	37.9	36.3-39.5	369.2	23.1	22.0-24.1
New York (Manhattan)	534.0	26.5	25.5-27.5	253.8	29.4	27.8-31.1	280.2	24.3	23.0-25.7
Queens	796.6	28.6	27.7-29.5	435.8	36.0	34.4-37.6	360.8	23.0	22.0-24.1
Richmond (Staten Island)	239.2	40.6	38.3-43.0	120.6	46.6	42.9-50.6	118.6	36.4	33.5-39.5
Albany	187.6	47.7	44.7-51.0	92.0	53.7	48.8-59.1	95.6	43.7	39.8-48.0
Allegany	34.6	51.3	43.8-59.9	19.2	58.7	47.3-72.3	15.4	45.1	35.3-57.1
Broome	119.6	44.9	41.2-48.8	59.6	48.9	43.4-55.0	60.0	42.2	37.3-47.6
Cattaraugus	54.2	47.5	41.9-53.8	27.4	51.2	42.7-61.0	26.8	44.8	37.3-53.5
Cayuga	58.0	51.8	45.8-58.4	29.2	55.1	46.3-65.3	28.8	49.0	41.1-58.2
Chautauqua	93.0	49.7	45.1-54.6	47.8	55.3	48.3-63.1	45.2	45.6	39.6-52.3
Chemung	67.0	55.3	49.3-61.7	35.2	63.1	53.9-73.5	31.8	49.0	41.4-57.7
Chenango	37.8	50.9	43.6-59.2	19.8	53.4	43.1-65.7	18.0	49.3	39.1-61.5
Clinton	54.6	50.5	44.5-57.1	28.2	55.1	46.2-65.4	26.4	46.8	38.9-56.0
Columbia	41.8	42.0	36.3-48.6	20.6	42.5	34.5-52.3	21.2	42.0	34.0-51.8
Cortland	37.4	61.8	53.1-71.6	19.0	67.7	54.5-83.4	18.4	57.4	46.1-71.0
Delaware	36.2	47.7	40.6-55.9	21.2	57.5	46.4-70.9	15.0	38.5	29.9-49.4
Dutchess	160.6	39.8	37.0-42.7	77.6	41.2	37.1-45.7	83.0	39.1	35.3-43.2
Erie	588.0	46.4	44.7-48.2	284.2	51.8	49.0-54.6	303.8	42.8	40.7-45.1
Essex	31.0	48.4	40.8-57.2	15.4	52.2	40.8-66.2	15.6	46.6	36.6-59.4
Franklin	33.8	51.0	43.4-59.6	20.4	62.2	50.4-76.1	13.4	41.9	32.1-54.1
Fulton	43.2	54.4	47.2-62.4	22.0	59.4	48.5-72.2	21.2	51.3	41.7-62.7
Genesee	46.0	55.2	48.1-63.1	22.6	58.1	47.6-70.4	23.4	54.5	44.8-65.8
Greene	40.0	52.3	45.1-60.4	22.8	60.9	50.0-73.8	17.2	44.0	34.9-55.3
Hamilton	5.0	55.4	34.7-90.2	3.0	58.3	32.3-114.5	2.0	51.0	22.4-112.0
Herkimer	45.4	49.4	43.0-56.6	24.8	57.7	47.7-69.5	20.6	41.3	33.6-50.6
Jefferson	69.2	58.8	52.7-65.5	35.8	66.4	56.8-77.1	33.4	53.0	45.1-61.9
Lewis	13.6	37.9	29.3-48.4	7.6	43.0	30.2-59.7	6.0	33.1	22.1-48.3
Livingston	43.6	49.9	43.3-57.3	22.2	54.7	44.7-66.4	21.4	46.1	37.6-56.3
Madison	45.8	44.9	39.1-51.4	26.2	53.0	44.1-63.4	19.6	37.8	30.5-46.6
Monroe	413.6	42.5	40.6-44.4	210.8	49.3	46.3-52.5	202.8	37.6	35.2-40.0
Montgomery	39.6	56.9	49.1-65.8	21.8	67.5	55.1-82.1	17.8	48.8	38.9-60.7
Nassau	592.6	31.9	30.8-33.1	278.8	34.3	32.5-36.2	313.8	30.5	29.0-32.1
Niagara	153.2	50.3	46.7-54.1	76.6	55.1	49.6-61.2	76.6	47.3	42.5-52.5
Oneida	162.6	51.2	47.6-54.9	86.4	59.9	54.3-66.0	76.2	44.5	40.0-49.4
Onondaga	285.4	47.1	44.7-49.7	139.6	52.2	48.3-56.3	145.8	43.6	40.4-47.0
Ontario	74.6	46.5	41.8-51.6	36.8	50.0	42.8-58.1	37.8	44.5	38.2-51.7
Orange	175.0	41.6	38.8-44.5	82.4	43.1	38.9-47.7	92.6	40.9	37.2-44.9
Orleans	34.6	61.3	52.2-71.6	18.2	70.2	56.0-87.1	16.4	54.8	43.3-68.8
Oswego	101.2	66.1	60.3-72.3	55.8	78.2	69.0-88.5	45.4	55.8	48.6-63.8
Otsego	40.0	43.5	37.5-50.3	21.0	48.9	39.8-59.8	19.0	38.6	30.9-48.1
Putnam	57.0	43.0	38.1-48.5	27.4	44.8	37.3-53.4	29.6	42.0	35.4-49.7
Rensselaer	110.2	52.9	48.4-57.6	53.8	57.1	50.2-64.6	56.4	50.0	44.2-56.4
Rockland	125.4	32.1	29.6-34.8	59.0	34.7	30.8-38.9	66.4	30.2	27.0-33.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by County, All Races, 2015-2019
Lung and Bronchus**

County/Region	Males and Females			Males			Females		
	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI
St. Lawrence	73.4	50.7	45.5-56.3	39.4	57.5	49.5-66.4	34.0	44.9	38.3-52.5
Saratoga	148.0	47.2	43.8-50.8	74.8	52.4	47.1-58.2	73.2	43.6	39.1-48.4
Schenectady	101.0	49.2	44.9-53.8	49.2	55.9	48.9-63.6	51.8	45.2	39.7-51.2
Schoharie	22.2	46.2	37.6-56.4	12.6	54.0	40.9-70.6	9.6	38.3	27.7-52.3
Schuyler	15.8	55.3	43.5-69.9	8.4	61.0	43.5-84.5	7.4	49.5	34.5-70.3
Seneca	26.8	55.7	46.4-66.5	12.8	55.5	42.4-71.6	14.0	57.5	44.2-73.9
Steuben	66.6	47.4	42.3-53.0	35.4	53.0	45.3-61.7	31.2	42.4	35.8-49.9
Suffolk	774.8	39.7	38.5-41.0	382.6	44.3	42.3-46.4	392.2	36.5	34.9-38.2
Sullivan	46.6	42.6	37.2-48.7	25.8	49.0	40.7-58.6	20.8	37.2	30.2-45.5
Tioga	31.8	43.5	36.9-51.3	17.6	50.1	40.0-62.2	14.2	37.8	29.1-48.6
Tompkins	42.6	37.6	32.6-43.2	22.0	43.4	35.4-52.8	20.6	33.7	27.4-41.2
Ulster	123.8	45.9	42.3-49.8	66.0	53.3	47.5-59.6	57.8	39.7	35.1-44.8
Warren	54.0	50.8	44.8-57.5	28.6	57.0	47.8-67.7	25.4	45.3	37.6-54.6
Washington	48.0	51.8	45.3-59.1	21.6	50.0	40.7-60.9	26.4	54.1	45.1-64.6
Wayne	67.8	52.3	46.7-58.4	36.6	59.9	51.3-69.8	31.2	46.0	38.8-54.3
Westchester	366.8	28.4	27.1-29.8	171.8	31.0	28.9-33.2	195.0	26.6	24.9-28.3
Wyoming	25.6	45.6	37.9-54.6	15.0	53.0	41.3-67.2	10.6	38.1	28.3-50.6
Yates	18.0	48.8	38.8-60.7	10.2	59.7	43.8-80.0	7.8	40.6	28.5-56.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by County, All Races, 2015-2019
Female Breast**

County/Region	Avg Ann Cases	Rate per 100,000	95% CI
New York State	5043.0	42.0	41.4-42.5
New York City	2116.4	42.5	41.7-43.3
New York State Excl New York City	2926.4	41.6	40.9-42.3
Bronx	336.6	41.4	39.4-43.5
Kings (Brooklyn)	679.4	45.9	44.4-47.5
New York (Manhattan)	399.6	40.8	39.0-42.7
Queens	574.2	40.9	39.4-42.5
Richmond (Staten Island)	126.6	42.4	39.0-46.0
Albany	79.6	43.2	38.8-48.0
Allegany	8.2	32.0	22.2-44.5
Broome	48.8	41.4	35.9-47.4
Cattaraugus	19.6	38.3	30.6-47.3
Cayuga	16.8	32.4	25.3-41.0
Chautauqua	33.6	40.0	33.7-47.2
Chemung	26.4	45.5	37.6-54.7
Chenango	14.6	45.3	34.8-58.2
Clinton	17.6	37.1	29.3-46.4
Columbia	21.4	49.8	39.9-61.6
Cortland	12.4	47.6	35.9-61.9
Delaware	11.6	35.9	26.4-48.0
Dutchess	74.0	39.9	35.7-44.5
Erie	255.6	43.1	40.6-45.7
Essex	11.0	36.5	26.9-49.1
Franklin	10.6	37.5	27.6-49.9
Fulton	12.6	33.2	25.0-43.4
Genesee	15.8	40.7	31.8-51.5
Greene	13.0	38.7	29.2-50.8
Hamilton	1.6	32.5	13.2-84.8
Herkimer	14.6	33.8	26.0-43.3
Jefferson	23.4	40.5	33.3-48.9
Lewis	6.6	37.4	25.0-54.1
Livingston	13.2	30.7	23.4-39.9
Madison	17.8	41.1	32.4-51.3
Monroe	196.6	42.0	39.3-44.8
Montgomery	13.6	41.6	31.7-53.7
Nassau	371.0	42.4	40.4-44.5
Niagara	57.6	41.1	36.2-46.5
Oneida	48.4	33.4	29.1-38.3
Onondaga	107.4	37.4	34.1-40.9
Ontario	26.6	35.0	29.0-42.1
Orange	92.6	44.1	40.1-48.5
Orleans	8.8	34.4	24.6-47.1
Oswego	25.2	38.2	31.5-46.0
Otsego	16.4	45.4	35.1-57.8
Putnam	27.6	42.5	35.3-50.8
Rensselaer	38.2	37.5	32.2-43.6
Rockland	79.0	43.3	38.9-48.0
St. Lawrence	22.8	34.6	28.2-42.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by County, All Races, 2015-2019
Female Breast**

County/Region	Avg Ann Cases	Rate per 100,000	95% CI
Saratoga	67.2	44.3	39.5-49.6
Schenectady	45.4	46.7	40.5-53.6
Schoharie	8.0	40.8	28.1-57.6
Schuyler	5.4	38.3	24.6-58.2
Seneca	7.0	35.6	24.0-51.0
Steuben	23.8	39.5	32.2-48.0
Suffolk	399.6	42.2	40.3-44.2
Sullivan	24.4	53.1	43.4-64.2
Tioga	14.6	43.9	33.7-56.4
Tompkins	24.6	47.0	38.7-56.5
Ulster	52.2	44.2	38.7-50.3
Warren	21.8	46.1	37.3-56.6
Washington	15.0	38.7	29.8-49.4
Wayne	28.4	50.5	42.0-60.3
Westchester	258.0	41.8	39.5-44.3
Wyoming	13.8	56.3	43.1-72.5
Yates	5.0	34.2	21.0-52.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by County, All Races, 2015-2019**
Cervix

County/Region	Avg Ann Cases	Rate per 100,000	95% CI
New York State	419.0	3.6	3.5-3.8
New York City	221.8	4.5	4.3-4.8
New York State Excl New York City	197.2	3.0	2.8-3.2
Bronx	42.0	5.3	4.6-6.1
Kings (Brooklyn)	72.4	4.9	4.4-5.4
New York (Manhattan)	36.6	3.7	3.2-4.4
Queens	61.8	4.6	4.1-5.1
Richmond (Staten Island)	9.0	3.2	2.3-4.3
Albany	5.2	3.3	2.1-4.9
Allegany	1.0	4.8	1.4-11.3
Broome	3.4	2.9	1.6-4.9
Cattaraugus	2.2	5.2	2.4-9.7
Cayuga	1.2	3.1	1.0-7.1
Chautauqua	2.2	2.8	1.3-5.3
Chemung	1.4	2.3	0.9-5.1
Chenango	1.8	5.4	2.3-11.0
Clinton	1.0	2.4	0.7-5.9
Columbia	1.6	3.3	1.4-7.2
Cortland	0.4	2.2	0.2-7.6
Delaware	1.2	5.4	1.8-12.5
Dutchess	5.0	3.0	1.9-4.5
Erie	17.0	3.3	2.6-4.2
Essex	1.0	5.1	1.4-13.2
Franklin	0.8	2.6	0.7-7.3
Fulton	1.2	4.0	1.4-9.0
Genesee	0.4	0.9	0.1-4.1
Greene	0.6	2.1	0.4-6.9
Hamilton	0.2	2.8	0.1-49.3
Herkimer	1.4	4.3	1.6-9.1
Jefferson	1.4	2.9	1.1-6.0
Lewis	0.6	4.6	0.8-14.0
Livingston	0.6	2.1	0.4-6.5
Madison	0.4	0.6	0.1-3.1
Monroe	12.8	2.9	2.2-3.7
Montgomery	0.6	2.3	0.4-7.0
Nassau	24.0	2.8	2.3-3.4
Niagara	4.8	3.5	2.2-5.4
Oneida	3.6	2.5	1.4-4.0
Onondaga	7.2	2.4	1.7-3.4
Ontario	2.2	3.2	1.5-6.0
Orange	10.4	5.3	3.9-6.9
Orleans	1.2	4.7	1.6-11.0
Oswego	1.4	2.4	0.9-5.1
Otsego	1.2	3.2	0.9-7.9
Putnam	2.0	3.6	1.6-7.0
Rensselaer	2.6	3.2	1.7-5.7
Rockland	5.0	2.5	1.6-3.8
St. Lawrence	2.0	3.2	1.5-6.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by County, All Races, 2015-2019**
Cervix

County/Region	Avg Ann Cases	Rate per 100,000	95% CI
Saratoga	3.2	2.3	1.3-3.9
Schenectady	3.0	3.5	1.9-5.9
Schoharie	0.4	1.6	0.2-7.6
Schuyler	0.4	4.3	0.4-17.1
Seneca	1.2	6.9	2.4-15.7
Steuben	2.0	3.1	1.4-6.1
Suffolk	26.0	2.9	2.4-3.5
Sullivan	2.4	4.6	2.2-8.6
Tioga	1.2	4.0	1.3-9.4
Tompkins	0.6	1.0	0.2-3.4
Ulster	4.4	4.7	2.8-7.3
Warren	0.8	2.0	0.4-5.8
Washington	1.6	5.0	2.0-10.4
Wayne	1.8	3.6	1.5-7.1
Westchester	12.8	2.1	1.6-2.7
Wyoming	0.8	2.7	0.7-8.0
Yates	0.4	3.1	0.3-12.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by County, All Races, 2015-2019
Prostate**

County/Region	Avg Ann Cases	Rate per 100,000	95% CI
New York State	2917.8	25.3	24.9-25.8
New York City	1084.2	25.2	24.5-25.9
New York State Excl New York City	1833.6	25.4	24.9-26.0
Bronx	188.6	30.1	28.1-32.1
Kings (Brooklyn)	313.6	25.5	24.2-26.8
New York (Manhattan)	264.6	29.9	28.3-31.6
Queens	254.2	19.7	18.6-20.8
Richmond (Staten Island)	63.2	22.8	20.3-25.5
Albany	52.4	28.5	25.0-32.3
Allegany	9.8	28.6	21.0-38.5
Broome	42.8	32.3	28.0-37.1
Cattaraugus	12.8	22.5	17.2-29.1
Cayuga	12.6	22.8	17.4-29.5
Chautauqua	15.6	16.4	12.9-20.7
Chemung	13.8	22.8	17.7-29.2
Chenango	11.4	30.3	22.8-39.8
Clinton	12.2	22.5	17.1-29.3
Columbia	14.6	27.9	21.8-35.7
Cortland	6.4	24.0	16.3-34.4
Delaware	10.6	26.7	19.8-35.9
Dutchess	47.0	23.7	20.7-27.1
Erie	121.4	20.4	18.8-22.2
Essex	6.6	19.4	13.3-28.3
Franklin	10.8	31.8	23.7-42.0
Fulton	11.2	28.8	21.6-38.0
Genesee	10.6	24.9	18.5-33.0
Greene	8.2	20.4	14.5-28.3
Hamilton	2.0	40.0	18.4-94.0
Herkimer	12.6	25.8	19.7-33.6
Jefferson	16.0	27.6	21.7-34.5
Lewis	4.0	22.5	13.6-35.5
Livingston	19.4	42.7	34.5-52.6
Madison	9.0	17.1	12.4-23.3
Monroe	177.4	38.8	36.2-41.5
Montgomery	9.4	26.8	19.5-36.2
Nassau	182.6	20.9	19.5-22.3
Niagara	31.6	21.7	18.3-25.6
Oneida	36.2	23.7	20.3-27.5
Onondaga	58.6	20.1	17.8-22.6
Ontario	37.2	46.2	39.6-53.7
Orange	48.4	23.4	20.4-26.6
Orleans	6.6	23.8	16.2-34.1
Oswego	13.2	17.8	13.6-22.9
Otsego	11.8	25.8	19.5-33.8
Putnam	18.2	25.6	20.4-31.9
Rensselaer	24.6	24.3	20.1-29.3
Rockland	34.2	18.9	16.2-22.1
St. Lawrence	15.0	22.1	17.3-28.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry
Regional/Distant Stage Cancer Incidence by County, All Races, 2015-2019
Prostate**

County/Region	Avg Ann Cases	Rate per 100,000	95% CI
Saratoga	41.6	27.3	23.6-31.5
Schenectady	23.6	24.8	20.4-29.9
Schoharie	10.8	43.5	32.4-57.8
Schuyler	4.2	32.3	19.3-51.8
Seneca	10.0	39.8	29.3-53.3
Steuben	23.2	32.0	26.3-38.8
Suffolk	235.6	25.9	24.4-27.4
Sullivan	9.8	18.2	13.3-24.5
Tioga	9.6	25.0	18.3-33.7
Tompkins	13.6	25.6	19.7-32.9
Ulster	26.0	19.8	16.5-23.7
Warren	13.0	24.8	19.0-32.1
Washington	12.6	28.3	21.6-36.7
Wayne	25.6	39.9	33.1-47.8
Westchester	161.0	27.7	25.7-29.7
Wyoming	7.2	24.2	16.6-34.3
Yates	7.4	39.2	27.3-55.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2021. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.